








SITE 852 HOLE B CORE 6H


Meter	Graphic Lith.	Section	Age	Structure	Disturb	Sample	Color	Description
0.5		1		///		S	10YR 6/3	NANNOFOSSIL OOZE WITH FORAMINIFERS and NANNOFOSSIL OOZE WITH FORAMINIFERS AND RADIOLARIANS
1.0		2		~			10YR 8/3	Major Lithologies:
		3		~			10YR 6/3	This upper two sections of this core contain light brown (10YR 6/3 - 10YR 8/3) NANNOFOSSIL OOZE WITH FORAMINIFERS AND RADIOLARIANS. The lower three sections contain very pale brown (10YR 8/1) to pale gray (N8 and 5Y 8/1) NANNOFOSSIL OOZE WITH FORAMINIFERS.
	Pliocene	4		~			10YR 8/2	Minor Lithology:
		5		~			10YR 8/3	The minor lithology in the core is very pale brown (10YR 8/2 to 10YR 8/1) FORAMINIFER NANNOFOSSIL OOZE.
		6		~			10YR 7/2	General Description:
		7		~			10YR 8/2	The first sediments in the core that are "green" in color occur in Section 5 at 55 cm.
	CC						10YR 8/1	TRACE FOSSILS: The core is slightly to moderately bioturbated. The most common trace fossils are filled burrows.
							N8	BANDING CONTRAST: The banding contrast in the upper half of the core is moderate; the contrast in the lower half of the core is very slight.
							10YR 8/1	
							N8	
							5Y 7/1	











1363

