

13. DATA REPORT: NEOGENE DIATOMS RECOVERED ON LEG 150X¹

Lloyd H. Burckle²

INTRODUCTION AND METHODS

This data report lists Neogene diatoms recovered from three drill sites along the New Jersey coastal plain (Atlantic City, Cape May, and Island Beach; Tables 1–3; see papers by Miller et al., 1994a, 1994b; Miller, et al., 1996). Our purpose was to characterize the diatoms at these sites and compare their occurrence with sediments of similar age from Leg 150 (Mountain, Miller, Blum, et al., 1994) and from the East Coast of the United States (Andrews, 1978, 1988). Ninety-seven samples were examined from Atlantic City, 103 from Cape May, and 79 from Island Beach. Sampling intervals ranged from 3 ft (0.93 m) to as much as 100 ft (31.2 m), but were generally within the range of 10 ft (3.1 m); sample size varied, but was on the order of 2–3 cm³. Diatom slides were prepared using the method described by Schrader (1974) and taxonomy followed Abbott, (1987a, 1987b), Andrews, (1978, 1988), Baldauf, (1984, 1986), and Burckle, (1972, 1978). Where a zonal designation was warranted, the schemes of Andrews (1988) and Burckle (1996) were used.

RESULTS

Atlantic City: The interval from 14 to 291 ft (4.3–90.9 m) is either barren of diatoms or contains few diatoms, none of which are age diagnostic. At and below a depth of 399 ft (124 m), diatoms are generally abundant. The joint occurrence of *Delphineis novaecaesarea*, *Delphineis penelliptica*, and *Denticulopsis hustedtii* indicate an interval equivalent to the *Delphineis novaecaesarea* and *Coscinodiscus* (now *Thalassiosira*) *yabei* Zones of Burckle (1996), even though the latter named taxon is not present. Further, this interval is equivalent to much of East Coast Diatom Zone (ECDZ) 6 and all of ECDZ 7 of Andrews (1988).

The top of the *Coscinodiscus lewisiatus* Zone of Burckle (1996) occurs at a depth of 454 ft (141 m). This zone extends to at least 637 ft (199 m), but may very well occur below that. Below 199 m, diatom occurrence becomes sporadic, with most samples being barren. Andrews (1988), using second-order correlation, tied this last occurrence to the lower part of foraminiferal Zone N11 of Blow (1969), approximately equivalent to somewhere in Magnetochron 14 (C5AA; Berggren et al., 1985). The last occurrence of *Raphidodiscus marylandicus* at a depth of 473 ft (147 m) and the co-occurrence of *Delphineis novaecaesarea* and *Delphineis penelliptica* suggest that the interval between 454 and 473 ft (141 and 147 m) ranges from the upper part of ECDZ 2 to the lowermost part of ECDZ 6 of Andrews (1988).

Island Beach: Diatoms are present in the Island Beach hole from the surface to a depth below 347 ft (108 m). Below this level, diatoms either are not present, are present in low numbers, or are in fragments. The topmost sample (79 ft [24 m]) contains abundant diatoms, but no age-diagnostic forms other than *Actinocyclus ingens* were identified.

This sample is tentatively placed in the upper Miocene. The samples below that level (88 ft [27 m]) to a depth of 347 ft (108 m) generally contain abundant diatoms, including sporadic occurrences of *C. lewisiatus*, *D. novaecaesarea*, *D. penelliptica*, and *Raphidodiscus marylandicus*. Burckle (1996) reported this latter species in his *C. lewisiatus* Zone. The interval between 88 and 347 ft (27 and 108 m) is placed in the *C. lewisiatus* Zone of Burckle (1996) and in the lower part of ECDZ 2 and possible ECDZ 1 of Andrews (1988).

Cape May: Diatoms are generally present in the upper part of the Cape May hole, although some samples were barren. Below a level of 750 ft (234 m), diatoms are generally not present, occurring in only a few samples. Those present in samples between the top of the hole and a depth of 412 ft (128 m) are generally near-shore, brackish water forms. Below that level, a number of zonal markers appear. The interval between 412 and 658 ft (128 and 205 m) contains *D. novaecaesarea* and *D. penelliptica* as well as sporadic occurrences of *D. hustedtii*. This interval is probably equivalent to the *D. novaecaesarea* and *C. yabei* Zones of Burckle (1996). The absence of *C. lewisiatus* and the presence of the aforementioned taxa suggest that it is equivalent to most of the ECDZ 6 and all of ECDZ 7 of Andrews (1988). The last occurrence of *C. lewisiatus* and *R. marylandicus* at 658 ft (205 m) indicates that this site penetrated into the *C. lewisiatus* Zone of Burckle (1996) and ECDZ 1 and the lower half of ECDZ 2 of Andrews (1988).

REFERENCES

- Abbott, W.H., 1987a. Diatom occurrences, Deep Sea Drilling Project Site 604. In van Hinte, J.E., Wise, S.W., Jr., et al. *Init. Repts. DSDP*, 93: Washington (U.S. Govt. Printing Office), 789–792.
- _____, 1987b. Diatom occurrences, Deep Sea Drilling Project Sites 612 and 613. In Poag, C.W., Watts, A.B., et al., *Init. Repts. DSDP*, 95: Washington (U.S. Govt. Printing Office), 417–418.
- Andrews, G.W., 1978. Marine diatom sequence in Miocene strata of the Chesapeake Bay region, Maryland. *Micropaleontology*, 24:371–406.
- _____, 1988. A revised marine diatom zonation for Miocene strata of the southeastern United States. *Geol. Surv. Prof. Pap. U.S.*, 1481:1–29.
- Baldauf, J.G., 1984. Cenozoic diatom biostratigraphy and paleoceanography of the Rockall Plateau region, North Atlantic, Deep Sea Drilling Project Leg 81. In Roberts, D.G., Schnitker, D., et al., *Init. Repts. DSDP*, 81: Washington (U.S. Govt. Printing Office), 439–478.
- _____, 1986. Diatom biostratigraphic and paleoceanographic interpretations for the middle to high latitude North Atlantic Ocean. In Summerhayes, C.P., and Shackleton, N.J. (Eds.), *North Atlantic Paleoceanography*. Geol. Soc. Spec. Publ. London, 21:243–254.
- Berggren, W.A., Kent, D.V., and Van Couvering, J.A., 1985. The Neogene, Part 2. Neogene geochronology and chronostratigraphy. In Snelling N.J. (Ed.), *The Chronology of the Geological Record*. Geol. Soc. London Mem., 10:211–260.
- Burckle, L.H., 1972. Late Cenozoic planktonic diatom zones from the eastern equatorial Pacific. In Simonsen, R. (Ed.), *First Symposium on Recent and Fossil Marine Diatoms*. Nova Hedwigia Beih., 39:217–250.
- _____, 1978. Early Miocene to Pliocene diatom datum levels for the equatorial Pacific. Proc. Second Working Group Mtg. Biostratigraphic Datum Planes, Pacific Neogene, IGCP Proj. 114, Spec. Publ., Geol. Res. Dev. Ctr., 1:25–44.
- _____, 1996. Diatom biostratigraphy, Leg 150. In Mountain, G.S., Miller, K.G., Blum, P., Poag, C.W., and Twichell, D.C. (Eds.), *Proc. ODP, Sci. Results*, 150: College Station, TX (Ocean Drilling Program), 17–35.

¹Miller, K.G., and Snyder, S.W. (Eds.), 1997. *Proc. ODP, Sci. Results*, 150X: College Station, TX (Ocean Drilling Program).

²Lamont-Doherty Earth Observatory, Palisades, NY 10964, U.S.A.
burckle@LDEO.Columbia.edu

- Miller, K.G., et al., 1996. *Proc. ODP, Init. Repts.* 150X (Suppl.): College Station, TX (Ocean Drilling Program).
- Miller, K.G., Browning, J.V., Liu, C., Sugarman, P., Kent, D.V., Van Fossen, M., Queen, D., Goss, M., Gwynn, D., Mullikin, L., Feigenson, M.D., Aubry, M.-P., Burckle, L.H., 1994a. Atlantic City site report. In Miller, K.G., et al., *Proc. ODP, Init. Repts.* 150X: College Station (Ocean Drilling Program), 35–55.
- Miller, K.G., Sugarman, P., Van Fossen, M., Liu, C., Browning, J.V., Queen, D., Aubry, M.-P., Burckle, L.H., Goss, M., and Bukry, D., 1994b. Island Beach site report. In Miller, K.G., et al., *Proc. ODP, Init. Repts.*, 150X: College Station (Ocean Drilling Program), 5–33.
- Mountain, G.S., Miller, K.G., Blum, P., et al., 1994. *Proc. ODP, Init. Repts.*, 150: College Station, TX, (Ocean Drilling Program).
- Schrader, H.-J., 1973. Proposal for a standardized method of cleaning diatom-bearing deep-sea and land-exposed marine sediments. *Nova Hedwigia Beih.*, 45:403–409.

Date of initial receipt: 17 September 1996

Date of acceptance: 20 December 1996

Ms 150XSR-309

Table 1. Atlantic City diatoms.

Depth (ft)	Taxonomic comments	Depth (ft)	Taxonomic comments	Depth (ft)	Taxonomic comments
14.0	<i>Cocconeis</i> spp. <i>Diploneis bombus</i> <i>Nitzschia granulata</i> <i>Thalassiosira oestrupii</i>	469.5	<i>Actinocyclus ingens</i> <i>Actinocyclus tenellus</i> <i>Actinoptychus senarius</i> <i>Cymatosira belgica</i> <i>Delphineis novaecaesarea</i> <i>Delphineis penelliptica</i> <i>Melosira westii</i> <i>Paralia sulcata</i>	569.7	<i>Actinoptychus senarius</i> <i>Melosira westii</i> <i>Paralia coronata</i> <i>Paralia sulcata</i>
19.5	<i>Nitzschia granulata</i>	473.5	<i>Actinoptychus senarius</i> <i>Delphineis novaecaesarea</i> <i>Delphineis penelliptica</i> <i>Melosira westii</i> <i>Paralia coronata</i> <i>Paralia sulcata</i>	575.8	<i>Melosira westii</i> <i>Paralia sulcata</i>
28.6	<i>Cocconeis</i> spp.	481.7	<i>Actinoptychus senarius</i> <i>Delphineis novaecaesarea</i> <i>Delphineis penelliptica</i> <i>Melosira westii</i> <i>Paralia coronata</i> <i>Paralia sulcata</i>	583.7	<i>Actinocyclus ingens</i> <i>Actinoptychus senarius</i> <i>Delphineis penelliptica</i> <i>Paralia coronata</i> <i>Paralia sulcata</i>
42.9	<i>Nitzschia granulata</i>	489.8	<i>Actinoptychus senarius</i> <i>Delphineis penelliptica</i>	591.4	<i>Diploneis bombus</i> <i>Paralia coronata</i> <i>Paralia sulcata</i>
48.9	<i>Cocconeis</i> sp.	495.9	<i>Actinoptychus senarius</i> <i>Coscinodiscus marginatus</i> <i>Melosira westii</i> <i>Cavitatus jouseana</i>	607.8	<i>Melosira westii</i> <i>Paralia coronata</i> <i>Paralia sulcata</i>
67 to 291.1	No or few diatoms	503.9	<i>Actinocyclus ingens</i> <i>Actinoptychus senarius</i> <i>Denticulopsis norvegica</i> <i>Diploneis bombus</i> <i>Paralia sulcata</i>	619.8	<i>Actinoptychus senarius</i> <i>Melosira westii</i> <i>Paralia sulcata</i> <i>Xanthiopxis oblonga</i>
399	<i>Actinocyclus ingens</i> ? <i>Actinocyclus tenellus</i> <i>Actinoptychus senarius</i> <i>Craspedodiscus coscinodiscus</i> <i>Delphineis novaecaesarea</i> <i>Melosira westii</i> <i>Paralia sulcata</i> <i>Xanthiopxis oblonga</i>	521.1	<i>Paralia coronata</i> <i>Paralia sulcata</i>	629.9	<i>Actinocyclus ehrenbergii</i> <i>Melosira westii</i> <i>Paralia coronata</i> <i>Paralia sulcata</i>
406.1	<i>Actinoptychus senarius</i> <i>Craspedodiscus coscinodiscus</i> <i>Delphineis novaecaesarea</i> <i>Delphineis penelliptica</i> <i>Melosira westii</i> <i>Paralia coronata</i> <i>Thalassiosira lineata</i> <i>Xanthiopxis oblonga</i>	526	<i>Actinocyclus tenella</i> <i>Actinoptychus senarius</i> <i>Delphineis penelliptica</i> <i>Melosira westii</i> <i>Paralia sulcata</i> <i>Synedra aducens</i>	637.4	<i>Actinocyclus tenella</i> <i>Melosira westii</i> <i>Paralia coronata</i> <i>Paralia sulcata</i>
413.6	<i>Delphineis novaecaesarea</i> <i>Delphineis penelliptica</i> <i>Melosira westii</i> <i>Paralia coronata</i>	509.6	<i>Actinocyclus tenella</i> <i>Coscinodiscus lewisi</i> <i>Melosira westii</i> <i>Paralia sulcata</i> <i>Rhizosolenia barbata</i>	644.1	Few diatom fragments
421.9	<i>Cymatosira belgica</i> (?) <i>Delphineis novaecaesarea</i> <i>Delphineis penelliptica</i> <i>Denticulopsis hustedtii</i> <i>Melosira westii</i> <i>Paralia coronata</i> <i>Triceratium subtrotundum</i>	539	<i>Actinocyclus tenella</i> <i>Melosira westii</i> <i>Xanthiopxis oblonga</i>	650.9	<i>Melosira westii</i>
427.7	<i>Delphineis penelliptica</i> Few diatoms	551.1	<i>Actinocyclus ingens</i> <i>Actinoptychus senarius</i> <i>Delphineis penelliptica</i> <i>Melosira westii</i> <i>Paralia coronata</i> <i>Paralia sulcata</i>	659.6	<i>Actinocyclus ingens</i> <i>Actinocyclus tenellus</i> <i>Delphineis penelliptica</i> <i>Melosira westii</i> <i>Paralia coronata</i>
433	<i>Delphineis novaecaesarea</i> <i>Delphineis penelliptica</i> <i>Melosira westii</i> <i>Paralia coronata</i> <i>Paralia sulcata</i> <i>Thalassiosira lineata</i>	564.1	<i>Delphineis penelliptica</i> <i>Melosira westii</i> <i>Paralia coronata</i> <i>Synedra aducens</i> (?)	663.8 to 750	No diatoms
441.6	Few diatom fragments Sponge spicules	576	<i>Actinocyclus tenella</i> <i>Melosira westii</i> <i>Xanthiopxis oblonga</i>	755.9	<i>Actinoptychus senarius</i> <i>Delphineis novaecaesarea</i> <i>Denticulopsis hustedtii</i> <i>Diploneis bombus</i>
448.4	<i>Delphineis novaecaesarea</i> Few diatoms	591.1	<i>Actinocyclus ingens</i> <i>Actinoptychus senarius</i> <i>Delphineis penelliptica</i> <i>Diploneis bombus</i> <i>Melosira westii</i> <i>Paralia coronata</i>	768.7	<i>Actinoptychus senarius</i> <i>Melosira westii</i> <i>Paralia coronata</i> <i>Paralia sulcata</i>
454.1	<i>Coscinodiscus lewisi</i> few broken diatoms	591.1	<i>Actinocyclus ingens</i> <i>Actinoptychus senarius</i> <i>Delphineis penelliptica</i> <i>Diploneis bombus</i> <i>Melosira westii</i> <i>Paralia coronata</i>	776 to 977	No diatoms or few diatoms
457.8	<i>Actinocyclus ingens</i> <i>Actinoptychus senarius</i> <i>Coscinodiscus lewisi</i> <i>Delphineis novaecaesarea</i> <i>Delphineis penelliptica</i> <i>Melosira westii</i> <i>Paralia coronata</i> <i>Xanthiopxis oblonga</i>	591.1	<i>Delphineis penelliptica</i> <i>Melosira westii</i> <i>Paralia coronata</i> <i>Synedra aducens</i> (?)		

Table 2. Island Beach diatoms, Hole 1B.

Depth (ft)	Taxonomic comments	Depth (ft)	Taxonomic comments	Depth (ft)	Taxonomic comments		
79.8 to 79.9	<i>Actinocyclus tenellus</i> <i>Actinocyclus ingens</i> <i>Actinptychus senarius</i> <i>Coscinodiscus radiatus</i> <i>Coscinodiscus oculus-iridus</i> <i>Craspedodiscus</i> sp. <i>Delphineis biseriata</i> <i>Delphineis ovata</i> <i>Delphineis surirella</i> <i>Diploneis bombus</i> <i>Melosira westii</i> <i>Paralia sulcata</i> <i>Thalassionema obtusa</i>	111.2 to 111.3	<i>Actinptychus senarius</i> <i>Cymatogonium amblyoceras</i> <i>Diploneis bombus</i> <i>Paralia sulcata</i> <i>Stephanopyxis turris</i> <i>Thalassionema nitzschiooides</i> <i>Thalassionema obtusa</i>	143.5 to 143.6	<i>Actinocyclus octonarius</i> <i>Asteromphalus imbricatus</i> <i>Aulacodiscus</i> sp. <i>Aulocoseira italicica</i> <i>Coscinodiscus asteromphalus</i> <i>Coscinodiscus cf. C. lewisianus</i> <i>Coscinodiscus lewisianus</i> <i>Coscinodiscus perforatus</i> <i>Cymatosira belgica</i> <i>Diploneis bombus</i> <i>Melosira westii</i> <i>Paralia complexa</i> <i>Paralia coronata</i> <i>Pleurosigma manni</i> <i>Pyxilla johnsoniana</i> <i>Raphoneis gemmifera</i> <i>Raphoneis scalaris</i> <i>Rhizosolenia barboi</i> <i>Stephanopyxis turris</i> <i>Thalassiosira eccentrica</i>		
88	<i>Actinocyclus tenellus</i> <i>Actinocyclus ingens</i> <i>Actinocyclus octonarius</i> <i>Actinptychus senarius</i> <i>Azpeitia nodulifera</i> <i>Coscinodiscus radiatus</i> <i>Coscinodiscus oculus-iridus</i> <i>Craspedodiscus</i> sp. <i>Delphineis biseriata</i> <i>Delphineis ovata</i> <i>Delphineis penelliptica</i> <i>Delphineis surirella</i> <i>Diploneis bombus</i> <i>Melosira westii</i> <i>Paralia sulcata</i> <i>Pleurosigma affine</i> var. <i>marylandica</i> <i>Raphidodiscus marylandicus</i> <i>Rhizosolenia styliformis</i> <i>Stephanopyxis turris</i> <i>Thalassionema obtusa</i> <i>Triceratium condecorum</i>	115.2 to 115.3	<i>Actinocyclus ingens</i> <i>Actinptychus senarius</i> <i>Paralia sulcata</i> <i>Raphoneis scalaris</i> <i>Sceptroneis grandis</i> <i>Stephanopyxis</i> sp. <i>Thalassionema nitzschiooides</i> <i>Thalassionema obtusa</i>	119.5 to 119.6	<i>Actinptychus senarius</i> <i>Coscinodiscus perforatus</i> <i>Diploneis bombus</i> <i>Epithemia</i> sp. <i>Paralia sulcata</i> <i>Podosira stelligera</i> <i>Raphoneis scalaris</i> <i>Thalassionema obtusa</i>	144.0 to 144.1	<i>Actinptychus marylandicus</i> <i>Actinptychus senarius</i> <i>Cavatatus jouseana</i> <i>Coscinodiscus lewisianus</i> <i>Coscinodiscus perforatus</i> <i>Cymatosira immunitis</i> <i>Delphineis penelliptica</i> <i>Denticulopsis norwegica</i> <i>Diploneis bombus</i> <i>Paralia sulcata</i> <i>Raphoneis scalaris</i> <i>Sceptroneis grandis</i> <i>Stephanopyxis turris</i> <i>Thalassionem nitzschiooides</i> <i>Thalassionema obtusa</i>
91.8 to 91.9	<i>Actinptychus senarius</i> <i>Coscinodiscus oculus-iridus</i> <i>Diploneis bombus</i> <i>Paralia sulcata</i> <i>Pleurosigma</i> sp. <i>Thalassionema obtusa</i>	127.9 to 128	<i>Actinocyclus ingens</i> <i>Actinocyclus tenellus</i> <i>Actinocyclus senarius</i> <i>Asterolampra marylandica</i> <i>Coscinodiscus perforatus</i> <i>Delphineis penelliptica</i> <i>Diploneis bombus</i> <i>Epithemia</i> sp. <i>Lithodesmium</i> cf. <i>L. undulatum</i> <i>Raphidodiscus marylandicus</i> <i>Thalassionema nitzschiooides</i> <i>Thalassionema obtusa</i>	153.4	<i>Actinocyclus ingens</i> <i>Actinocyclus octonarius</i> <i>Actinptychus senarius</i> <i>Aulocoseira granulata</i> <i>Aulocoseira italicica</i> <i>Cavatatus jouseana</i> <i>Coscinodiscus endoi</i> <i>Coscinodiscus marginatus</i> <i>Craspedodiscus coscinodiscus</i> <i>Cymatosira belgica</i> <i>Delphineis ovata</i> <i>Diploneis bombus</i> <i>Melosira westii</i> <i>Paralia coronata</i> <i>Pyxilla johnsoniana</i> <i>Raphidodiscus</i> cf. <i>R. marylandicus</i> <i>Raphoneis diamontella</i> <i>Raphoneis rhombica</i> <i>Rhizosolenia barboi</i> <i>Stephanopyxis turris</i>		
95.0 to 95.1	No diatoms	132 to 132.1	<i>Asterolampra marylandica</i> <i>Coscinodiscus asteromphalus</i> <i>Coscinodiscus perforatus</i> <i>Delphineis penelliptica</i> <i>Denticula norwegica</i> <i>Diploneis bombus</i> <i>Paralia sulcata</i> <i>Raphoneis scalaris</i> <i>Raphoneis amphiceras</i> <i>Sceptroneis grandis</i> <i>Thalassionema nitzschiooides</i> <i>Thalassionema obtusa</i>	161.5 to 161.6	<i>Melosira westii</i> <i>Podosira stelligera</i> <i>Xanthiopyxis oblonga</i> <i>Coscinodiscus perforatus</i> <i>Triceratium condecorum</i>		
99.1 to 99.2	<i>Actinocyclus ingens</i> <i>Actinptychus senarius</i> <i>Amphiceras</i> sp. <i>Anaulus</i> sp. <i>Azpeitia nodulifera</i> <i>Coscinodiscus marginatus</i> <i>Delphineis penelliptica</i> <i>Diploneis bombus</i> <i>Paralia sulcata</i> <i>Raphidodiscus marylandicus</i> <i>Raphoneis gemmifera</i> <i>Raphoneis</i> cf. <i>R. amphiceras</i> <i>Raphoneis amphiceras</i> <i>Sceptroneis grandis</i> <i>Thalassionema nitzschiooides</i> <i>Thalassionema obtusa</i>	136.3 to 136.4	<i>Actinptychus senarius</i> <i>Biddulphia</i> cf. <i>B. aurita</i> <i>Coscinodiscus lewisianus</i> <i>Coscinodiscus perforatus</i> <i>Delphineis penelliptica</i> <i>Denticulopsis norwegica</i> <i>Diploneis bombus</i> <i>Rhizosolenia miocenica</i> <i>Rhizosolenia styliformis</i> <i>Thalassionema nitzschiooides</i> <i>Thalassionema obtusa</i>	168.6 to 168.7	<i>Actinocyclus ehrenbergii</i> <i>Aulacodiscus</i> sp. <i>Coscinodiscus lewisianus</i> <i>Coscinodiscus</i> cf. <i>C. perforatus</i> <i>Delphineis</i> cf. <i>D. ovata</i> <i>Melosira westii</i> <i>Paralia sulcata</i> <i>Xanthiopyxis oblonga</i> <i>Xanthiopyxis ovalis</i> <i>Bolboforma</i> sp. (not a diatom)		
103.2 to 103.3	<i>Actinocyclus ingens</i> <i>Actinptychus senarius</i> <i>Coscinodiscus argus</i> <i>Diploneis bombus</i> <i>Delphineis</i> cf. <i>D. surirella</i> <i>Paralia sulcata</i> <i>Raphoneis amphiceras</i> <i>Raphoneis scalaris</i> <i>Stephanopyxis turris</i> <i>Thalassionema nitzschiooides</i> <i>Thalassionema obtusa</i>	140.2 to 140.3	<i>Actinocyclus tenellus</i> <i>Actinptychus senarius</i> <i>Asterolampra marylandica</i> <i>Delphineis penelliptica</i> <i>Denticulopsis norwegica</i> <i>Epithemia</i> sp. <i>Raphoneis gemmifera</i> <i>Raphoneis amphiceras</i> <i>Rhizosolenia</i> cf. <i>R. miocenica</i> <i>Thalassionema nitzschiooides</i> <i>Thalassionema obtusa</i>	175.6 to 175.7 and 181.5 to 181.6	Diatom fragments or no diatoms		
107.5 to 107.6	<i>Actinptychus australis</i> <i>Actinptychus senarius</i> <i>Coscinodiscus marginatus</i> <i>Coscinodiscus radiatus</i> <i>Diploneis bombus</i> <i>Paralia sulcata</i> <i>Raphoneis gemmifera</i> <i>Sceptroneis grandis</i> <i>Stephanopyxis</i> sp. <i>Thalassionema nitzschiooides</i> <i>Thalassionema obtusa</i>						

Table 2 (continued).

Depth (ft)	Taxonomic comments	Depth (ft)	Taxonomic comments
191.0 to 191.1	<i>Actinopychus australis</i> <i>Cavatius jouseana</i> <i>Coscinodiscus marginatus</i> <i>Delphineis cf. D. ovata</i> <i>Delphineis penelliptica</i> <i>Denticulopsis norwegica</i> <i>Paralia complexa</i> <i>Paralia coronata</i> <i>Pleurosigma marylandica</i> <i>Stephanopyxis turris</i> <i>Xanthiopyxis oblonga</i> <i>Xanthiopyxic ovalis</i>	340.9 to 341.0	<i>Cymatogonia amblyoceros</i> <i>Thalasionema nitzschiooides</i> <i>Thalassionema obtusa</i> <i>Diploneis cf. D. bombus</i> <i>Paralia sulcata</i>
200.3 to 200.4 and 265.0 to 265.1	Diatom fragments or no diatoms	347.8 to 348.0	<i>Actinocyclus ingens</i> <i>Diploneis sp.</i> <i>Rhyzosolenia styliformis</i> <i>Thalassionema nitzschiooides</i> <i>Thalassionema obtusa</i>
274.5 to 274.6	<i>Actinocyclus octonarius</i> <i>Actinocyclus tenellus</i> <i>Diplomenora cocconeiformis</i> <i>Coscinodiscus cf. C. monicae</i> <i>Cymatosira belgica</i> <i>Delphineis novaecaesarea</i> <i>Dimmerogramma elliptica</i> <i>Diplomenora cocconeiformis</i> <i>Paralia complexa</i> <i>Paralia coronata</i> <i>Pyxilla johnsoniana</i> <i>Rhizosolenia praebarboi</i> <i>Stephanopyxis turris</i> <i>Triceratium condecorum</i> <i>Triceratium tesselatum</i> <i>Xanthiopyxis oblonga</i> <i>Bolboforma sp. (not a diatom)</i>	355.0 to 355.1	No diatoms
284.2 to 284.3 and 317.0 to 317.1	Diatom fragments or no diatoms	365.0 to 365.1	<i>Paralia sp.</i> <i>Thalassionema nitzschiooides</i> Few diatoms
320.0 to 320.2	<i>Rhyzosolenia styliformis</i> Diatom fragments	375.0 to 375.1 and 488.0 to 488.1	Diatom fragments or no diatoms
326.0 to 326.1	<i>Actinopychus senarius</i> <i>Paralia sulcata</i> <i>Raphoneis gemmifera</i> <i>Raphoneis rhombicus</i> <i>Raphidodiscus cf. R. marylandicus</i> <i>Thalassionema nitzschiooides</i> <i>Thalassionema obtusa</i>	494.8 to 495.0 and 509.0 to 509.1	No diatoms
333.0 to 333.2	<i>Actinocyclus tenellus</i> <i>Actinopychus senarius</i> <i>Coccineis sp.</i> <i>Coscinodiscus lewisanus</i> <i>Coscinodiscus perforatus</i> <i>Delphineis penelliptica</i> <i>Diploneis bombus</i> <i>Epithemia sp.</i> <i>Paralia sulcata</i> <i>Raphoneis scalaris</i> <i>Sceptroneis grandis</i> <i>Thalassionema nitzschiooides</i> <i>Thalassionema obtusa</i> <i>Thalassiosira sp.</i>	517.0 to 517.1 and 529.4 to 529.5	<i>Thalassionema nitzschiooides</i> <i>Thalassionema obtusa</i> Diatom fragments Pollen and spores common
		539.0 to 539.1	<i>Coscinodiscus perforatus</i> <i>Diploneis cf. D. bombus</i> <i>Raphoneis scalaris</i> <i>Raphoneis sp.</i> <i>Thalassionema nitzschiooides</i>
		542.8 to 542.9	<i>Diploneis cf. D. bombus</i> <i>Paralia sulcata</i> <i>Raphoneis sp.</i> <i>Thalassionema nitzschiooides</i> <i>Thalassionema obtusa</i>
		548.2 to 548.4 and 667.7 to 667.8	No diatoms

Table 3. Cape May diatoms.

Depth (ft)	Taxonomic comments	Depth (ft)	Taxonomic comments
19.0	<i>Actinptychus senarius</i> <i>Coscinodiscus marginatus</i> <i>Paralia sulcata</i> <i>Raphoneis amphiceras</i> <i>Thalassiosira eccentrica</i>	451.0	<i>Delphineis novacaesarea</i> <i>Denticulopsis hustedtii</i> <i>Melosira westii</i>
34.2	Rare diatom fragments Few sponge spicules Opal phytoliths	463.0	<i>Delphineis novacaesarea</i> <i>Paralia sulcata</i>
44.8	<i>Actinptychus senarius</i> <i>Coscinodiscus marginatus</i> <i>Paralia coronata</i> <i>Paralia sulcata</i> <i>Raphoneis amphiceras</i>	469.5	<i>Actinptychus senarius</i> <i>Delphineis novacaesarea</i>
54.2 to 84	Few diatom fragments	485.0	<i>Delphineis novacaesarea</i> <i>Denticulopsis hustedtii</i> <i>Diplogeis bombus</i> <i>Melosira westii</i> <i>Raphoneis amphiceras</i>
91.9	<i>Actinptychus senarius</i> <i>Coscinodiscus obscurus</i> <i>Diplogeis bombus</i> <i>Paralia sulcata</i> <i>Raphoneis amphiceras</i>	492.0	<i>Actinptychus senarius</i> <i>Coscinodiscus marginatus</i> <i>Delphineis novacaesarea</i> <i>Denticulopsis hustedtii</i> <i>Diplogeis bombus</i> <i>Melosira westii</i> <i>Paralia sulcata</i> <i>Thalassiosira lineata</i>
101.9	<i>Diplogeis bombus</i> <i>Paralia sulcata</i>	532.0	<i>Actinptychus senarius</i> Diatoms rare
111.9	<i>Actinptychus senarius</i> <i>Raphoneis lancetulla</i> <i>Pleurosigma sp.</i> <i>Nitzschia granulata</i>	541.9	Rare diatom fragments
121.9	<i>Actinptychus senarius</i> <i>Nitzschia granulata</i> <i>Raphoneis amphiceras</i> <i>Raphoneis lancetulla</i> <i>Thalassiosira lineata</i>	561.0	<i>Diplogeis bombus</i> Diatom fragments
131.9	<i>Diplogeis bombus</i> <i>Nitzschia granulata</i> <i>Raphoneis lancetulla</i>	593.9	<i>Actinptychus senarius</i> <i>Diplogeis bombus</i> <i>Paralia sulcata</i>
161.9 to 171.9	No diatoms or rare diatom fragments	610.0	<i>Paralia coronata</i>
181.9	<i>Nitzschia granulata</i> Rare diatom fragments	624.5 to 647	No diatoms
185.9	Rare diatom fragments Opal phytoliths	658.2	<i>Actinptychus senarius</i> <i>Coscinodiscus lewisanus</i> <i>Delphineis penelliptica</i> <i>Melosira westii</i> <i>Paralia coronata</i> <i>Raphidodiscus marylandicus</i>
196.1	<i>Raphoneis lancetulla</i> Rare diatom fragments	675.0	No diatoms
211.9 to 262	No diatoms	690.9	<i>Actinptychus senarius</i> <i>Delphineis penelliptica</i> <i>Diplogeis bombus</i> <i>Melosira westii</i> <i>Paralia coronata</i>
277	<i>Actinptychus senarius</i> <i>Raphoneis lancetulla</i>	712.0	No diatoms
288	<i>Raphoneis amphiceras</i> Rare diatoms	741.1	<i>Coscinodiscus marginatus</i> <i>Delphineis novacaesarea</i> <i>Delphineis penelliptica</i> <i>Diplogeis bombus</i> <i>Melosira westii</i>
293.1	<i>Nitzschia granulata</i> Rare diatoms	752.5 to 922.01	No diatoms
302.1	<i>Nitzschia granulata</i> <i>Raphoneis lancetulla</i>	1045	<i>Cavatibus jouseana</i> <i>Diplogeis bombus</i> <i>Melosira westii</i> <i>Paralia sulcata</i> <i>Xanthiopixis bombus</i>
313.3	<i>Diplogeis bombus</i> <i>Raphoneis lancetulla</i>	1061 to 1247.9	No diatoms
325.0 to 391.6	No diatoms or rare diatoms	1253.9	<i>Paralia sulcata</i> <i>Sceptroneis caduceus</i>
402.5	<i>Actinptychus senarius</i> <i>Melosira westii</i> <i>Raphoneis rhombica</i>	1261.9	<i>Melosira westii</i> <i>Paralia sulcata</i> <i>Sceptroneis caduceus</i>
412.6	<i>Actinocyclus tenella</i> <i>Paralia sulcata</i> <i>Raphoneis lancetulla</i> <i>Delphineis novacaesarea</i>	1300 to 1480	No diatoms
428.0	<i>Actinptychus senarius</i>		
440.0	<i>Actinptychus senarius</i> <i>Coscinodiscus marginatus</i> <i>Craspedodiscus coscinodiscus</i> <i>Delphineis novacaesarea</i> <i>Melosira westii</i>		