



						Texture										Minerals										Biogenic										TOT														
Site	H	Core	T	Sec	Int.	Depth (mbsf)	Observer	Lithology	Sand	Silt	Clay	Quartz	Feldspar	Plagioclase	Mica	Biotite	Muscovite	Amphibole	Rock Fragments	Glass volcanic glass	Opques	Fe-Oxide	Pyrite	Zeolite	Glauconite	Phosphate	Carbonate	Dolomite	Micrite	Carbonate grains	Clay	Foraminifers	Pellets	Bioclasts	Echinoids	Diatoms	Radiolarians	Silicoflagelates	Fine Siliceous Fragments	Sponge Spicules	Fish Remains	Nannos	Discoaster	Coccolith	Organic Debris or Matter	Plant Debris	Amorphous Organic Aggregates	Others	Total Percentage	
1013	A	16	X	4	10	141.10	TAD	D	3	17	80	2	T							2		2							7	12	1																			100
1013	B	5	H	4	78	42.38	TAD	M	95	4	1	40	15					10	5									5		15		10																		100
1013	B	9	H	1	5	71.65	PIK	M		90	10	1									T							98																						100
1013	B	11	H	5	27	100.37	PIK	M																				100																						100
1013	B	11	H	3	137-138	98.47-98.48	PIK	M		20	80	2	3						1	1	1							74		5			T								2		10	1					100	
1013	C	1	H	3	91	3.91	PIK	M	5	35	65	2							T	1	1							65		5		5																100		

Previous Chapter

Table of Contents

Next Chapter