

		Nannofossil zones/ subzones	
Martini (1971)	Okada and Bukry (1980)		
		Core, section, interval (cm)	
		Depth (mbsf)	Total abundance
			Sample preservation
			Reworked Cretaceous
			<i>Biantholithus sparsus</i>
			<i>Bomolithus elegans</i>
			<i>Braarudosphaera bigelowii</i>
			<i>Chiasmolithus bidens</i>
			<i>Chiasmolithus consuetus</i>
			<i>Chiasmolithus danicus</i>
			<i>Chiasmolithus spp.</i>
			<i>Coccolithus pelagicus</i>
			<i>Coccolithus robustus</i>
			<i>Coccolithus spp.</i>
			<i>Coccolithus subpertusus</i>
			<i>Cruciplacolithus edwardsii</i>
			<i>Cruciplacolithus intermedius</i>
			<i>Cruciplacolithus primus</i>
			<i>Cruciplacolithus spp.</i>
			<i>Cruciplacolithus tenuis</i>
			<i>Cyclagelosphaera margerelii</i>
			<i>Discoaster falcatus</i>
			<i>Discoaster mohleri</i>
			<i>Discoaster multiradiatus</i>
			<i>Discoaster nobilis</i>
			<i>Discoaster spp.</i>
			<i>Ellipsolithus macellus</i>
			<i>Fasciculithus alanii</i>
			<i>Fasciculithus aubertae</i>
			<i>Fasciculithus biliii</i>
			<i>Fasciculithus bobii</i>
			<i>Fasciculithus clinatus</i>
			<i>Fasciculithus involutus</i>
			<i>Fasciculithus janii</i>
			<i>Fasciculithus lilianae</i>
			<i>Fasciculithus magnus</i>
			<i>Fasciculithus pileatus</i>
			<i>Fasciculithus shaulbii</i>
			<i>Fasciculithus spp.</i>
			<i>Fasciculithus thomasii</i>
			<i>Fasciculithus tympaniformis</i>
			<i>Heliolithus kleinpelli</i>
			<i>Heliolithus nieldii</i>
			<i>Markalius inversus</i>

Notes: Abundance: A = abundant, C = common, F = few, R = rare, B = barren. Preservation: M = moderate, P = poor.

