

Table T7. Eocene benthic foraminiferal occurrences. (Continued on next page.)

[illegible]

Sample depth (ft)	<i>Lenticulina</i> <i>Loxostomoides</i> <i>Marginulina</i> <i>Marginulinopsis</i> <i>Massalina</i>	<i>Melonis</i> <i>Nodosarella attenuata</i> <i>Nodosaria</i> <i>Nodosaria latejugata</i> <i>Nonionellina</i>	<i>Oridarsalis</i> <i>Osangularia</i> <i>Pararotalia</i> <i>Percultazonaria</i> <i>Planulina</i>	<i>Pleurostomella</i> <i>Polymorphina</i> <i>Pullenia quinqueloba</i> <i>Pulsiphorina prima</i> <i>Quinqueloculina</i>	<i>Ramulina globulifera</i> <i>Sigmoidella</i> <i>Sigmomorphina</i> <i>Siphonina claibornensis</i> <i>Spiroplectamina</i>	<i>Spiroplectamina spectabilis</i> <i>Stilosomella</i> <i>Tappanina selmensis</i> <i>Textularis</i> <i>Trifarina</i>	<i>Turrillina robertsi</i> <i>Uvigerina</i> <i>Valvulineria</i> <i>Virginullina</i>
171.0							
176.0	x						
181.0	x						x
186.0	x						x
191.0	x	x					x
196.0	x	x					x
201.0	x						x
206.0	x	x					x
211.0	x	x					x
216.0	x	x					x
221.0	x	x					x
226.0	x						x
231.0	x	x					x
235.0	x	x					x
241.0	x						
246.0	x	x					
251.0	x	x					
256.0	x	x					
261.0	x						
266.0	x						
271.0	x						
276.0	x						
281.0	x						
286.0	x						
290.4	x						
301.0	x	x					
306.0	x						
311.0	x	x					
316.0	x	x					
321.0	x						
326.0	x	x					
331.0	x						
336.0	x						
341.0							
346.0							
351.0							
356.0							
361.0							
366.0		x					
371.0							
376.0							
386.0							
446.0							
456.0		x					
466.0	x						
476.0	x						
486.0							
494.7	x						
501.0							