

Table 4. Relative abundance of benthic foraminiferal species and overall abundance of benthic foraminifers at Hole 1082A.

Notes: The relative abundance of benthic foraminiferal species is given as a percentage, where + = <1% and P = present (the relative abundance was not calculated because of small sample size). Absolute abundance (per ~20 cm³ of sediment) of benthic foraminifers is given as A = abundant (>500 specimens); C = common (250–500 specimens); Lith = lithified (not possible to count or classify benthic foraminifers).

ODP Proceedings, Initial Reports, Volume 175: Chapter 10, Table 4.

Table 4. Relative abundance

Core, section, interval	Depth (mbsf)	Abundance	<i>Anomalinoides globulosus</i>	<i>Astronion novozelandicum</i>	<i>Astronion stelligerum</i>	<i>Bolivina pseudoduplicata</i>	<i>Bolivina seminuda</i>	<i>Bolivina subtenaensis</i>	<i>Bolivina sp. 1</i>	<i>Bolivinopsis cubensis</i>	<i>Bulimina aculeata</i>	<i>Bulimina exilis</i>	<i>Bulimina marginata</i>	<i>Bulimina mexicana</i>	<i>Bulimina truncana</i>	<i>Cassidulina laevigata</i>	<i>Cassidulina minuta</i>	<i>Cassidulinoides cf. bradyi</i>	<i>Chilostomella oolina</i>	<i>Chilostomella ovoida</i>	<i>Chrysaglomium spp.</i>	<i>Cibicidoides bradyi</i>	<i>Cibicidoides pachyderma</i>	<i>Cibicidoides wellerstorfi</i>	<i>Eggerella bradyi</i>	? <i>Eggerella sp. 1</i>
175-1082A-																										
1H-CC	7.84	A																								2
2H-CC	17.37	A																								+
3H-CC	27.16	A																								+
4H-CC	36.71	A																								
5H-CC	46.52	A																								
6H-CC	55.82	A																								
7H-CC	65.47	A																								
8H-CC	75.27	A																								
9H-CC	84.50	A																								
10H-CC	94.34	A																								
11H-CC	103.45	A																								1
12H-CC	112.84	A																								
13H-CC	120.24	A																								
14H-CC	128.58	A																								
15X-CC	135.30	A																								
16X-CC	145.34	A																								
17X-CC	152.80	A																								
18X-CC	163.06	A																								
19X-CC	172.74	A																								
20X-CC	183.51	A	1																							
21X-CC	193.47	A																								
22X-CC	201.31	A																								
23X-CC	213.19	C																								
24X-CC	223.03	C																								
25X-CC	231.46	C																								
26X-CC	244.38	A																								
27X-CC	252.12	A																								
28X-CC	262.14	A																								
29X-CC	273.16	A																								
30X-CC	280.53	A																								
32X-CC	299.62	A																								
33X-CC	309.63	A																								
34X-CC	317.67	A																								
35X-CC	328.60	A																								
36X-CC	328.55	A																								
37X-CC	337.01	A																								
38X-CC	349.23	A																								
39X-CC	356.49	A																								
40X-CC	366.46	A																								
41X-CC	378.29	A	+ +																							
42X-CC	386.97	Lith																								
43X-CC	396.06	A																								
44X-CC	405.07	A																								
45X-CC	414.98	A																								
46X-CC	424.11	A																								
47X-CC	432.96	A																								
48X-CC	442.70	Lith																								
49X-CC	455.85	A	+ +																							
50X-CC	464.01	A																								
51X-CC	472.75	A	2																							
52X-CC	482.40	A																								
53X-CC	491.85	A																								
54X-CC	503.56	A	+ +																							
55X-CC	513.60	A	2																							
56X-CC	523.15	A	+ +																							
57X-CC	532.68	A	+ +																							
58X-CC	541.98	A																								
59X-CC	550.05	A	3																							
60X-CC	561.13	A	2																							
61X-CC	570.48	A	3																							
62X-CC	579.77	A	2	3																						
63X-CC	590.05	A	2																							
64X-CC	599.98	A	+ +																							

Notes: The relative abundance of benthic foraminiferal species is given as a percentage, where + = <1% and P = present (the relative abundance was not calculated foraminifers).

of benthic foraminiferal species and overall abundance of benthic foraminifers at Hole 1082A.

ed because of small sample size). Absolute abundance (per ~20 cm³ of sediment) of benthic foraminifers is given as A = abundant (>500 specimens); C = common (250–500 specimens); Lith = lithi-

				Number of specimens counted
	<i>Uvigerina auberti</i>			
	<i>Uvigerina gallowayi</i>			
	<i>Uvigerina hispida</i>			
	<i>Uvigerina hispidocostata</i>			
	<i>Uvigerina peregrina</i>			
	Unidentified			
8		+	3	348
+		+		311
12		+	3	352
9		+	1	418
12		2	7	407
4		61	2	512
+		20	17	413
		+	+	361
1		14	8	242
+		+	+	455
+		74	2	288
+		+	1	297
3		1	7	245
11	2	2	2	362
+		+	2	325
2		5	1	458
8	4	11	6	275
1		37	1	293
1		8	9	319
1		+	2	268
32			3	248
13		+	10	367
7			3	329
1			7	260
17		+	6	361
4		+	6	273
13			12	392
18		6	3	340
24		+	5	202
			8	278
20	2	+	7	420
15	4		6	361
14	3		7	317
17	8		3	405
3	2	+	8	350
13			9	451
10		6	4	398
18		4	2	326
10		+	13	313
15	+	3	4	323
			0	
17		7	3	419
30	8	+	4	386
5	2		6	365
12			6	352
12	2	+	7	423
			0	
7		+	9	377
17			7	369
17			3	376
14			6	353
15			3	384
16		4	7	305
7	4	3	6	324
7	+		14	353
13			11	282
12	2	7	2	380
12	1	13	8	309
2	+	2	17	243
4		6	2	211
			17	185
9		+	4	338
15	2	1	8	250

fied (not possible to count or classify benthic