

Table T3 (continued).

Core, section, interval (cm)	Depth (mbsf)	Preservation	Group abundance																
				<i>Reticulofenestra reticulata</i>	<i>Reticulofenestra samodurovii</i>	<i>Reticulofenestra</i> spp.	<i>Reticulofenestra umbilicus</i>	<i>Sphenolithus moriformis</i>	<i>Sphenolithus primus</i>	<i>Sphenolithus radians</i>	<i>Sphenolithus spiniger</i>	<i>Sphenolithus</i> spp.	<i>Toweius callosus</i>	<i>Toweius pertusus</i>	<i>Transversopontis pulcher</i>	<i>Transversopontis pulcheroideus</i>	<i>Transversopontis rectipons</i>	<i>Triquetrorhabdulus rugosus</i>	<i>Zygrhablithus bijugatus</i>
189-1171A-																			
1H-CC, 8-13	7.02	G	A																
2H-CC, 18-25	16.10	G	A																
3H-CC, 17-22	24.76	G	A																
4H-CC, 20-25	35.78	G	A																
5H-CC, 18-23	45.36	G	A																
6H-CC, 17-22	55.17	G	A																
7H-CC, 25-30	64.20	G	A																
8H-CC, 14-19	73.05	G	A															F	
9H-CC, 19-24	82.40	G	A																
10H-CC, 14-19	92.66	G	A																
11H-CC, 13-18	102.49	M	A						F										
12H-CC, 5-10	110.62	M	A																
13X-CC, 20-25	113.66	M	A																
14X-CC, 15-20	120.53	M	A																
189-1171B-																			
1H-CC, 9-14	4.21	G	A																
2H-CC, 15-20	13.79	G	A		F														
3H-CC, 14-19	23.17	G	A																
4H-CC, 13-18	32.19	G	A																
5H-CC, 4-9	42.03	M	A																
6H-CC, 13-18	52.04	M	A																
7H-CC, 15-20	60.74	M	A																
8H-CC, 14-19	70.75	G	A															R	
9H-CC, 8-13	80.23	G	A																
10H-CC, 14-19	89.77	G	A																
11H-CC, 13-18	97.93	M	A															R	
12H-CC, 11-18	109.06	M	A																
189-1171C-																			
1H-CC, 6-11	9.65	G	A																
2H-CC, 8-13	19.24	G	A																
3H-CC, 17-22	28.56	G	A																
4H-CC, 14-19	37.80	G	A																
5H-CC, 17-22	47.86	G	A																
6H-CC, 18-22	57.45	G	A																
7H-CC, 23-28	67.13	G	A															R	
8H-CC, 0-6	75.33	G	A																
9H-CC, 12-17	85.64	M	A																
10H-CC, 14-22	94.40	M	A																F

Table T3 (continued).

Core, section, interval (cm)	Depth (mbsf)	Preservation		Group abundance	Reticulofenestra reticulata	Reticulofenestra samodurovii	Reticulofenestra spp.	Reticulofenestra umbilicus	Sphenolithus moriformis	Sphenolithus primus	Sphenolithus radians	Sphenolithus spiniger	Sphenolithus spp.	Toweius callosus	Toweius pertusus	Transversopontis pulcher	Transversopontis pulcheroideus	Transversopontis rectipons	Triquetrorhabdulus rugosus	Zygrhablithus bijugatus
11H-CC, 0-8	104.12	M	A																	
12X-CC, 11-16	109.92	M	A																	
13X-CC, 13-18	115.72	M	A																	
14X-CC, 13-18	123.49	M	A																	
15X-CC, 13-18	134.45	M	A																	
16X-CC, 17-22	139.66	M	A																	
17X-CC, 11-16	150.18	M	A																	
18X-CC, 16-21	162.93	M	A																	
19X-CC, 12-17	172.76	M	A																	
20X-CC, 16-23	181.10	M	A				C													
21X-CC, 18-23	190.11	M	A																	
22X-CC, 16-21	198.93	M	A																	
23X-CC, 18-23	210.12	M	A																	
24X-CC, 23-28	219.74	M	A																	
25X-CC, 19-24	228.48	M	A																	
26X-CC, 14-19	237.56	M	A																	
27X-CC, 36-41	247.33	M	A																	
28X-CC, 29-34	253.79	M	A																	
29X-CC, 35-40	264.83	M	A																	
30X-CC, 13-18	272.87	M	A																	
31X-CC, 28-33	274.43	P	R																	
189-1171D-																				
1R-CC, 15-20	250.27	M	A																	
2R-CC, 17-22	260.68	M	A																	
3R-CC, 14-19	271.34	G	C																	R
4R-CC, 15-20	280.59	G	A																	C
5R-CC, 15-20	287.53	P	F	F																F
6R-CC, 7-12	295.26	P	R	R																C
7R-CC, 15-20	302.30	M	C																	F
8R-CC, 0-5	307.84	M	C			C														C
9R-CC, 12-17	318.77	M	F			F														C
10R-CC, 12-17	330.78	P	R																	C
11R-CC, 12-13	334.02	P	F																	C
12R-CC, 11-13	343.61	M	C			C														C
13R-CC, 8-11	353.28	M	C			C														F
14R-CC, 16-21	363.28	P	C			C														F
15R-CC, 20-25	374.35	M	F			F														F
16R-CC, 18-23	385.18	P	C	F																R
17R-CC, 24-29	398.40	M	C																	F
18R-CC, 0-3	401.20	M	C																	F

Table T3 (continued).

Core, section, interval (cm)	Depth (mbsf)	Preservation	Group abundance	Reticulofenestra				Sphenolithus				Toweius						
				<i>reticulata</i>	<i>samodurovii</i>	spp.	<i>umbilicus</i>	<i>moriformis</i>	<i>primus</i>	<i>radians</i>	<i>spiniger</i>	spp.	<i>callosus</i>	<i>pertusus</i>	<i>pulcher</i>	<i>pulchroides</i>	<i>rectipons</i>	<i>triquetrohabdulus rugosus</i>
19R-CC, 0-5	417.63	P	C															
20R-CC, 6-12	428.65	M	R															
21R-CC, 32-38	430.32		B															
22R-CC, 12-17	448.84	M	A				R											
23R-CC, 10-16	453.57	P	F	F	C	R							R	R				
24R-CC, 29-34	464.32		B															
25R-CC, 0-5	476.81		B															
26R-CC, 15-20	487.98	M	F		F	R												
27R-CC, 7-12	497.45	M	A	F														
28R-CC, 5-10	505.73	P	C	F	C													
29R-CC, 13-18	516.79	M	A	F														
30R-CC, 12-17	524.58	M	A															
31R-CC, 28-33	533.73	M	A															
32R-CC, 0-5	543.96	M	A															
33R-CC, 0-5	555.02	M	A															
34R-CC, 0-5	564.79	M	A															
35R-CC, 7-12	573.86	M	A															
36R-CC, 13-18	584.10	M	A															
37R-CC, 17-23	593.56	M	A															
38R-CC, 0-5	603.22	M	A		C													
39R-CC, 0-5	611.85	M	A	F	C												R	
40R-CC, 19-24	622.70	M	A									C	C					
41R-CC, 12-17	631.14	M	C		F													
42R-CC, 13-18	641.81	M	C		F													
43R-CC, 9-14	651.01	M	A		C							F	F					
44R-CC, 17-22	660.71	M	A									F		F			C	
45R-CC, 16-21	665.86	M	A					F	R			C					A	
46R-CC, 0-5	677.79	M	A					F		R				C	F		A	
47R-CC, 17-22	690.24	M	A					F	R								F	
48R-CC, 0-5	699.50	P	R		R													
49R-CC, 0-7	709.35	P	R		R													
50R-CC, 23-28	718.48	M	R		R													
51R-CC, 0-6	726.94	M	F		R													
52R-CC, 8-13	738.11	M	R		R													
53R-CC, 16-21	745.80	M	R		R													
54R-CC, 9-14	757.26	M	C		R												R	
55R-CC, 11-16	766.95	M	C		R												R	
56R-CC, 17-22	771.40	P	R		R					R				R				
57R-CC, 16-21	781.73	P	R					R										
58R-CC, 11-16	795.49		B															

Table T3 (continued).

Core, section, interval (cm)	Depth (mbsf)	Preservation		Group abundance	<i>Reticulofenestra reticulata</i> <i>Reticulofenestra samodurovii</i> <i>Reticulofenestra</i> spp. <i>Reticulofenestra umbilicus</i> <i>Sphenolithus moriformis</i> <i>Sphenolithus primus</i> <i>Sphenolithus radians</i> <i>Sphenolithus spiniger</i> <i>Sphenolithus</i> spp. <i>Toweius callosus</i> <i>Toweius pertusus</i> <i>Transversopontis pulcher</i> <i>Transversopontis pulcheroideus</i> <i>Transversopontis rectipons</i> <i>Triquetrorhabdulus rugosus</i> <i>Zygrhablithus bijugatus</i>
59R-CC, 10-15	805.32		B		
60R-CC, 10-15	814.60		B		
61R-CC, 6-11	822.60		B		
62R-CC, 9-14	834.15		B		
63R-CC, 15-20	843.81		B		
64R-CC, 9-14	852.01	P	R		
65R-CC, 20-25	863.00	P	R		R
66R-CC, 0-8	870.71	P	R		
67R-CC, 21-26	882.22	P	R		R
68R-CC, 0-5	889.42		B		
69R-CC, 27-33	901.08		B		
70R-CC, 17-22	908.07		B		
71R-CC, 34-39	919.12		B		
72R-CC, 0-5	929.29		B		
73R-CC, 0-5	939.51		B		
74R-CC, 0-8	939.50		B		
75R-CC, 23-28	954.94		B		