

Core Photo

Site 1202 Hole A Core 1H Cored 0.0-7.1 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
0.0						SS		<p>CLAYEY SILT</p> <p>This core consists of homogeneous, dark greenish gray CLAYEY SILT with minor bioturbation. The core is calcareous and bioturbation consists of faint, black mottled intervals.</p>
2.0						SS IW		
4.0						IW	dk gn GY	
6.0						SS SS IW		
						IW		
						IW		
						IW PAL		

Core Photo

Site 1202 Hole A Core 2H Cored 7.1-16.6 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
8					ooo	IW	dk gn GY	<p>CLAYEY SILT</p> <p>This core consists of homogeneous, dark greenish gray CLAYEY SILT with minor bioturbation. This core is calcareous and bioturbation consists of faint, black mottled intervals. A large (3 mm) white macrofossil occurs in Section 6, 35 cm. Also in Section 6, bioturbated sand-sized intervals with a vigorous reaction to 10% HCl occur at 92-93 cm and 119-124 cm.</p>
10						IW		
12						IW		
						SS		
14						IW		
						IW		
16						SS		
						IW		
						PAL		

Core Photo

Site 1202 Hole A Core 3H Cored 16.6-26.1 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
18						IW		<p>CLAYEY SILT</p> <p>This core consists of homogeneous, dark greenish gray CLAYEY SILT with moderate bioturbation. This core is calcareous and bioturbation consists of faint, black mottled intervals. White shell fragments are rare but visible in most sections. Short (< 5 cm) fining-upward intervals occur in Sections 4 and 5. The latter half of the core is slightly disturbed by horizontal gas voids.</p>
20						IW		
22						IW	dk gn GY	
24						IW		
						SS		
						SS		
						IW		
26						IW		

Core Photo

Site 1202 Hole A Core 4H Cored 26.1-35.6 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
28						IW	dk gn GY	<p>CLAYEY SILT</p> <p>This core consists of homogeneous, dark greenish gray CLAYEY SILT with moderate to rare bioturbation. This core is calcareous and bioturbation consists of faint, black mottled intervals. White shell fragments are rare, but visible in most sections. Short (< 5 cm) fining-upward intervals occur in Section 1. The bases of these fining-upward intervals, and rare sand-sized intervals, react strongly with 10% HCl. The core is slightly disturbed by horizontal gas voids.</p>
30					IW			
					SS			
					IW	dk gn GY		
32					SS			
					IW			
34								
							dk gn GY	
36						PAL		




Core Photo

Site 1202 Hole A Core 5H Cored 35.6-45.1 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
36								<p>CLAYEY SILT</p> <p>This core consists of homogeneous, dark greenish gray CLAYEY SILT with moderate to rare bioturbation. A black colored interval begins in Section 4 and extends through to Section 6. This core is calcareous and bioturbation consists of faint, black mottled intervals. White shell fragments are rare, but visible in most sections. Rare, thin (< 5 mm-thick) sand-sized intervals, react strongly with 10% HCl. The core is slightly disturbed by horizontal gas voids.</p>
38							dk gn GY	
40								
42						SS	BK	
44							dk gn GY	
						PAL		

Core Photo

Site 1202 Hole A Core 6H Cored 45.1-54.6 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
46						— IW — SS — PAL	GY	<p>CLAYEY SILT</p> <p>This core consists of homogeneous, gray and dark greenish gray CLAYEY SILT with moderate to rare bioturbation. This core is calcareous and bioturbation consists of faint, black mottled intervals. White shell fragments are rare, but visible in most sections. Rare, thin (< 5 mm-thick) sand-sized intervals, react strongly with 10% HCl. Two short (< 2 cm) fining-upward intervals occur in Sections 3 and 4. Basal contacts are sharp and erosive. The core is slightly disturbed by horizontal gas voids.</p>
48								
50								
52								
54							dk gn GY	

Core Photo

Site 1202 Hole A Core 7H Cored 54.6-64.1 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
56						IW	GY	<p>CLAYEY SILT</p> <p>This core consists of homogeneous, gray CLAYEY SILT with moderate to rare bioturbation. This core is calcareous and bioturbation consists of faint, black mottled intervals. White shell fragments are rare, but visible in most sections. Rare, thin (< 5 mm-thick), sand-sized intervals react strongly with 10% HCl. The core is slightly disturbed by horizontal gas voids.</p>
58								
60								
62								
64						PAL		

Core Photo

Site 1202 Hole A Core 8H Cored 64.1-73.6 mbsf									
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION	
66								<p>CLAYEY SILT</p> <p>This core consists of homogeneous, gray CLAYEY SILT with moderate bioturbation. This core is calcareous and bioturbation consists of faint, black mottled intervals. Faint black horizontal horizons (<5 mm thick) occur in Sections 2, 4, 6, and 7, and are bioturbated. White shell fragments are rare, but visible in most sections. Rare, thin (< 5 mm-thick) sand-sized intervals, react strongly with 10% HCl. The core is slightly disturbed by horizontal gas voids.</p>	
68									
70									
72									
74									
								<p>SS</p> <p>IW</p> <p>PAL</p> <p>GY</p>	




Core Photo

Site 1202 Hole A Core 9H Cored 73.6-83.1 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
74							GY	<p>CLAYEY SILT</p> <p>This core consists of homogeneous, gray CLAYEY SILT with moderate bioturbation. This core is calcareous and bioturbation consists of faint, black mottled intervals. Faint black horizontal horizons (<1 cm thick) occur in Sections 1-5, and are bioturbated. White shell fragments are rare, but visible in most sections. Rare, thin (< 5 mm-thick), sand-sized intervals react strongly with 10% HCl. The core is slightly disturbed by horizontal gas voids.</p>
76						SS		
78						IW		
80						SS		
82						PAL		

Core Photo

Site 1202 Hole A Core 11H Cored 92.6-102.1 mbsf									
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION	
94							GY	CLAYEY SILT This core consists of homogeneous, gray to black colored CLAYEY SILT with common to moderate bioturbation. This core is calcareous and bioturbation consists of black mottled intervals. White shell fragments are rare, but visible in some sections. The core is slightly disturbed by horizontal gas voids.	
96									
98									IW
100									SS
102									PAL


Core Photo

Site 1202 Hole A Core 12H Cored 102.1-110.0 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
104 106 108						IW SS PAL	GY	<p>CLAYEY SILT</p> <p>This core consists of homogeneous, gray CLAYEY SILT with moderate to rare bioturbation. This core is calcareous and bioturbation consists of black mottled intervals. Discrete burrows are less well-defined downcore. Sections 1-3 contain sand-sized, black fragments with a clayey silt texture. White shell fragments are rare, but visible in some sections. The core is slightly disturbed by horizontal gas voids.</p>


Core Photo

Site 1202 Hole A Core 13H Cored 110.0-119.5 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
112						SS	GY	<p>CLAYEY SILT</p> <p>This core consists of homogeneous, gray CLAYEY SILT with common bioturbation. This core is calcareous and bioturbation consists of black mottled intervals. The core is finer grained and reacts less with 10% HCL starting in Section 2, 29 cm. White shell fragments are rare, but visible in some sections. Rare, thin (< 5 cm-thick), sand-sized intervals react strongly with 10% HCl. Faint black horizontal horizons (< 1 cm-thick) occur in Sections 1-3. The core is slightly disturbed by horizontal gas voids.</p>
114			SS					
116			SS					
118			IW					
120			PAL	GY				


Core Photo

Site 1202 Hole B Core 1H Cored 0.0-2.9 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
2							dk GY	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of faint black mottled intervals. Hydrogen sulfide smell.</p>


Core Photo

Site 1202 Hole B Core 2H Cored 2.9-12.4 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
4							dk GY	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of faint black mottled intervals. Isolated calcareous sand layers appear throughout the core. Some of them are discontinuous. Shell fragments are rare, but occur throughout the core. Hydrogen sulfide smell.</p>
6								
8								
10								
12								



Core Photo

Site 1202 Hole B Core 4H Cored 21.9-31.4 mbsf									
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION	
24								<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of faint black mottled intervals. Horizons of dark gray and isolated calcareous sand layers appear throughout the core. Some of the sand layers are discontinuous. Shell fragments are rare, but occur throughout the core. Hydrogen sulfide smell.</p>	
26									
28									
30									
32									




Core Photo

Site 1202 Hole B Core 5H Cored 31.4-40.9 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
32 34 36 38 40							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of dark gray mottles and faint black mottled intervals which diminish downcore. Horizons of dark gray and isolated calcareous sand layers appear throughout the core. Some of them are discontinuous. Shell fragments are rare, but occur throughout the core. Hydrogen sulfide smell.</p>




Core Photo

Site 1202 Hole B Core 6H Cored 40.9-50.4 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
42							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of dark gray mottles. Horizons of dark gray and isolated calcareous sand layers appear throughout the core. Some sand layers are discontinuous. Shell fragments are rare, but occur throughout the core. Hydrogen sulfide smell.</p>
44								
46								
48								
50								



Core Photo

Site 1202 Hole B Core 7H Cored 50.4-59.9 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
52							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of dark gray mottles. Horizons of dark gray and isolated calcareous sand layers appear throughout the core. Some sand layers are discontinuous and sand-filled burrows are common. Shell fragments are rare, but occur throughout the core. The core is moderately disrupted by horizontal gas voids.</p>
54								
56								
58								
60								



Core Photo

Site 1202 Hole B Core 8H Cored 59.9-69.4 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
62							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of dark gray mottles. Horizons of dark gray and isolated calcareous sand layers appear throughout the core. Some sand layers are discontinuous and sand-filled burrows are common. Shell fragments are rare, but occur throughout the core. The core is moderately disrupted by horizontal gas voids.</p>
64								
66								
68								
70								




Core Photo

Site 1202 Hole B Core 9H Cored 69.4-78.9 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
70							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of dark gray mottles. Horizons of dark gray and isolated calcareous sand layers appear throughout the core. Some sand layers are discontinuous and sand-filled burrows are common. Shell fragments are rare, but occur throughout the core. The core is moderately disrupted by horizontal gas voids.</p>
72								
74								
76								
78								



Core Photo

Site 1202 Hole B Core 11H Cored 88.4-97.9 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
90							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of dark gray mottles. Horizons of dark gray and isolated calcareous sand layers appear throughout the core. Some sand layers are discontinuous and sand-filled burrows are common. Shell fragments are rare, but occur throughout the core. The core is moderately disrupted by horizontal gas voids.</p>
92								
94								
96								
98								



Core Photo

Site 1202 Hole B Core 13H Cored 104.2-111.6 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
106 108 110							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of dark gray mottles. Horizons of dark gray and isolated calcareous sand layers appear throughout the core. Some sand layers are discontinuous and sand-filled burrows are common. Shell fragments are rare, but occur throughout the core. The core is moderately disrupted by horizontal gas voids.</p>




Core Photo

Site 1202 Hole B Core 16X Cored 130.9-140.5 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
132 134 136 138							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of dark gray mottles. Horizons of dark gray and isolated calcareous sand layers appear throughout the core. Some sand layers are discontinuous and sand-filled burrows are common. Shell fragments are rare, but occur throughout the core. The core is moderately disrupted by horizontal gas voids.</p>



Core Photo

Site 1202 Hole C Core 1H Cored 0.0-5.7 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
2 4							dk GY	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of faint black mottled intervals. Hydrogen sulfide smell.</p>



Core Photo

Site 1202 Hole C Core 4H Cored 24.7-34.2 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
26							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of dark gray mottles and faint black mottled intervals which diminish downcore. Horizons of dark gray and isolated calcareous sand layers appear throughout the core. Some of them are normally-graded and/or discontinuous. Shell fragments are rare, but occur throughout the core. Hydrogen sulfide smell. Core is disrupted by horizontal gas voids.</p>
28								
30								
32								
34								



Core Photo

Site 1202 Hole C Core 5H Cored 34.2-43.7 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
36 38 40 42 44							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of dark gray mottles. Horizons of dark gray and isolated calcareous sand layers appear throughout the core. Some of the sand layers are normally-graded and/or discontinuous. Shell fragments are rare, but occur throughout the core. The core is disrupted by horizontal gas voids.</p>



Core Photo

Site 1202 Hole C Core 7H Cored 53.2-62.7 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
54 56 58 60 62							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of dark gray mottles. Horizons of dark gray and isolated calcareous sand layers appear throughout the core. Some of the sand layers are normally-graded and/or discontinuous. Shell fragments are rare, but occur throughout the core. The core is disrupted by horizontal gas voids.</p>




Core Photo

Site 1202 Hole C Core 8H Cored 62.7-72.2 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
64 66 68 70 72							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of dark gray mottles. Horizons of dark gray and isolated calcareous sand layers appear throughout the core. Some of the sand layers are normally-graded and/or discontinuous. Shell fragments are rare, but occur throughout the core. The core is disrupted by horizontal gas voids.</p>



Core Photo

Site 1202 Hole C Core 9H Cored 72.2-81.0 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
74.							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of dark gray mottles. Horizons of dark gray and isolated calcareous sand layers appear throughout the core. Some of the sand layers are normally-graded and/or discontinuous. Shell fragments are rare, but occur throughout the core. The core is disrupted by horizontal gas voids.</p>
76								
78								
80								

Core Photo

Site 1202 Hole C Core 10H Cored 81.0-90.5 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
82							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of dark gray mottles. Horizons of dark gray and isolated calcareous sand layers appear throughout the core. Some of the sand layers are normally-graded and/or discontinuous. Shell fragments are rare, but occur throughout the core. The core is disrupted by horizontal gas voids.</p>
84								
86								
88								
90								

Core Photo

Site 1202 Hole C Core 11H Cored 90.5-97.5 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
92							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of dark gray mottles. Horizons of dark gray and isolated calcareous sand layers appear throughout the core. Some of the sand layers are normally-graded and/or discontinuous. Shell fragments are rare, but occur throughout the core. The core is disrupted by horizontal gas voids.</p>
94								
96								

Core Photo

Site 1202 Hole D Core 1H Cored 0.0-0.2 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
								<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark greenish gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of faint, black mottled intervals. Hydrogen sulfide smell.</p>

Core Photo

Site 1202 Hole D Core 2H Cored 0.2-9.7 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
2						— SS	dk gn GY	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark greenish gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of faint, black mottled intervals, particularly frequent in Section 4. Layers with white shell fragments occur in Section 4 at 110 cm and in Section 5 at 139 cm. A fine sand interval appears in Section 6 at 120 cm. Hydrogen sulfide smell.</p>
4						— SS		
6								
8								
10						— SS		




Core Photo

Site 1202 Hole D Core 4H Cored 19.2-28.7 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
20						SS	ol GY	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, olive gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of faint, black mottled intervals. A shell fragment occurs in Section 5 at 84 cm. Sand layers - some of them are discontinuous - appear throughout the core and become abundant in Sections 6 and 7. Hydrogen sulfide smell.</p>
22								
24								
26								
28								

Core Photo

Site 1202 Hole D Core 5H Cored 28.7-38.2 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
30							ol GY	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, olive gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of faint, black and olive mottled intervals. Shell fragments occur in Section 3 at 46 cm and in Section 6 at 20 cm. Sections 4 and 5 include brown calcareous nodules, <2 mm in diameter. Sand layers appear throughout the core. The core is disrupted by up to 2 cm thick horizontal gas voids. Hydrogen sulfide smell.</p>
32								
34								
36								
38								



Core Photo

Site 1202 Hole D Core 6H Cored 38.2-47.7 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
40							ol GY	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, olive gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of faint, black and olive mottled intervals. Sand layers appear throughout the core. The core is disrupted by up to 2 cm thick horizontal gas voids. Hydrogen sulfide smell.</p>
42								
44								
46								
48								



Core Photo

Site 1202 Hole D Core 7H Cored 47.7-57.2 mbsf									
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION	
48							ol GY	CLAYEY SILT This sediment core consists of homogeneous, olive gray and olive CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of faint, black mottled intervals. Black mottles are absent in Sections 5 and 6. Sand layers appear throughout the core. One calcareous sand layer is present in Section 1 at 141 cm. The core is disrupted by up to 2 cm thick horizontal gas voids. Hydrogen sulfide smell.	
50									
52									
54									
56									OL
58									SS


Core Photo

Site 1202 Hole D Core 8H Cored 57.2-66.7 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
58 60 62 64 66							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, olive CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation consists of faint, black mottled intervals, particularly in Section 6 and in the core catcher. Sand layers appear throughout the core with a 1 cm thick layer in Section 6 at 26 cm. The core is disrupted by up to 2 cm thick horizontal gas voids. Hydrogen sulfide smell.</p>
						SS		




Core Photo

Site 1202 Hole D Core 9H Cored 66.7-76.2 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
68						SS	OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, olive CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation is partly displayed by black mottled intervals. Shell fragments occur in Section 2 at 116 cm, in Section 3 at 114 cm, in Section 4 at 95 cm, and in Section 6 at 35 cm and 52 cm. Sand layers, some calcareous, appear throughout the core. The core is disrupted by up to 2 cm thick horizontal gas voids. Hydrogen sulfide smell.</p>
70								
72								
74								
76								




Core Photo

Site 1202 Hole D Core 10X Cored 76.2-85.7 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
78						SS	BK OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, black and olive CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation is displayed by dark sandy and calcareous mottles. Isolated sand layers appear throughout the core. Hydrogen sulfide smell.</p>




Core Photo

Site 1202 Hole D Core 11X Cored 85.7-95.3 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
86. 88. 90.						SS	OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, olive CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation is displayed by dark sandy and calcareous mottles. Isolated sand layers appear throughout the core. The core is disrupted by up to 2 cm thick horizontal gas voids. Hydrogen sulfide smell.</p>

Core Photo

Site 1202 Hole D Core 12X Cored 95.3-104.9 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
96							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, olive and black CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation is displayed by dark mottles. Section 3 is more calcareous. Some isolated sand layers appear throughout the core. The core is disrupted by <2 cm thick horizontal gas voids. Hydrogen sulfide smell.</p>
98							BK	
100							OL	



Core Photo

Site 1202 Hole D Core 13X Cored 104.9-114.5 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
106 108 110 112							ol BR	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, olive brown CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation is displayed by dark sandy and calcareous mottles. Isolated sand laminae appear throughout the core, which exceed 2 cm in thickness in Section 2. The core is disrupted by up to 2 cm thick horizontal gas voids. Hydrogen sulfide smell.</p>




Core Photo

Site 1202 Hole D Core 15X Cored 124.1-133.8 mbsf									
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION	
126							ol GY	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, olive gray and dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation is displayed by black mottles. Isolated sand layers and faint sandy laminae appear throughout the core. Shell fragments occur in Section 3 at 22 cm, in Section 4 at 89 cm, and in Section 6 in a sand interval between 48 and 51 cm. The core is disrupted by horizontal gas voids.</p>	
128							dk GY		
130									ol GY
132									dk GY



Core Photo

Site 1202 Hole D Core 16X Cored 133.8-143.4 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
134							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, olive and dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation is displayed by dark gray mottles. Isolated sand layers appear throughout the core. Calcareous sand intervals occur in Section 1 between 82 and 86 cm, in Section 2 between 40 and 43 cm, and in Section 6 between 9 and 11 cm. The core is disrupted by horizontal gas voids with a thick void in Section 2 between 85 and 105 cm.</p>
136								
138								
140								
142								




Core Photo

Site 1202 Hole D Core 21X Cored 182.0-191.7 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
184							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray and olive CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation is displayed by dark gray mottles. Isolated sand layers, which are partly graded, appear throughout the core and reach several cm thickness in Section 1. Shell fragments are present in Section 1 between 24 and 26 cm. A black nodule occurs in Section 3 at 39 cm. The core is disrupted by horizontal gas voids.</p>
186							dk GY	
188							OL	



Core Photo

Site 1202 Hole D Core 22X Cored 191.7-201.3 mbsf										
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION		
192						SS	OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray and olive CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation is displayed by dark gray mottles. Isolated calcareous sand layers, some are graded, appear throughout the core and are abundant in Section 3. The core is disrupted by horizontal gas voids.</p>		
194										
196										
198										
							<table border="1"> <tr> <td>dk GY</td> </tr> <tr> <td>OL</td> </tr> </table>	dk GY	OL	
dk GY										
OL										




Core Photo

Site 1202 Hole D Core 23X Cored 201.3-210.9 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
202							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray and olive CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation is displayed by dark gray mottles. Isolated sand layers, some are graded, appear throughout the core and are up to 3 cm thick in Section 3. Section 5 includes shell fragments at 107 cm. The core is disrupted by horizontal gas voids.</p>
204								
206								
208						SS		



Core Photo

Site 1202 Hole D Core 24X Cored 210.9-220.5 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
212 214 216 218							OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray and olive CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation is displayed by dark gray mottles. Abundant calcareous sand layers appear throughout the core. Some of them are graded and/or show sharp basal contacts. The core is disrupted by horizontal gas voids.</p>



Core Photo

Site 1202 Hole D Core 25X Cored 220.5-230.1 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
222 224						SS	dk GY OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, dark gray and olive CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation is displayed by dark gray mottles. Calcareous sand layers appear throughout the core and become increasingly abundant and thicker (several cm) downcore. Some of them are graded and/or show sharp basal contacts. The core is disrupted by horizontal gas voids.</p>

Core Photo



Site 1202 Hole D Core 26X Cored 230.1-239.7 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
								CLAYEY SILT The core catcher of this sediment core recovered slightly calcareous, dark gray CLAYEY SILT with moderate bioturbation. It includes a sand layer between 5 and 8 cm. The core is moderately disturbed.

Core Photo

Site 1202 Hole D Core 28X Cored 249.3-258.9 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
250							dk GY	<p>INTERBEDDED SAND AND CLAYEY SILT</p> <p>This sediment core consists of dark gray and olive INTERBEDDED SAND AND CLAYEY SILT. Clayey silt is slightly calcareous, while sand is calcareous. The sand beds are several cm thick and often show grading and erosional basal contacts. The core occasionally is disrupted by horizontal gas voids and shows extreme disturbance in the core catcher.</p>
252							OL	

1202D-29X NO RECOVERY

Core Photo

Site 1202 Hole D Core 30X Cored 268.5-278.1 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
270							OL	<p>INTERBEDDED SAND AND CLAYEY SILT</p> <p>This sediment core consists of olive INTERBEDDED SAND AND CLAYEY SILT. Clayey silt is slightly calcareous, while sand is calcareous. The sand beds are several cm thick and often show grading and erosional basal contacts. The core occasionally is disrupted by horizontal gas voids and shows extreme disturbance in the core catcher.</p>


Core Photo

Site 1202 Hole D Core 35X Cored 316.6-326.2 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
318 320 322 324						SS SS	OL	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, olive CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation is displayed by dark gray mottles. Isolated calcareous sand layers appear throughout the core and are frequent in Section 1. Some of them are graded and/or show erosional basal contacts. A microfault is visible in Section 5 at 58 cm. The core occasionally is disrupted by horizontal gas voids. Strongly disturbed intervals appear in Section 4 between 86 cm and 98 cm and in Section 6 between 90 cm and 110 cm.</p>

Core Photo

Site 1202 Hole D Core 39X Cored 355.3-364.9 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
356						SS	dk GY	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, olive and dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation is displayed by dark gray mottles and sandy burrows. Isolated calcareous sand layers appear throughout the core. Many of them are discontinuous. The core occasionally is disrupted by horizontal gas voids. Section 6 exhibits extreme core disturbance.</p>
358							OL	
360							dk GY	
362							OL	

Core Photo

Site 1202 Hole D Core 44X Cored 403.5-410.0 mbsf								
METERS	GRAPHIC LITH.	BIOTURB.	STRUCTURE	ACCESSORIES	DISTURB.	SAMPLE	COLOR	DESCRIPTION
404						— SS	dk GY	<p>CLAYEY SILT</p> <p>This sediment core consists of homogeneous, olive and dark gray CLAYEY SILT with moderate bioturbation. The core is slightly calcareous and bioturbation is displayed by dark gray mottles. Isolated calcareous sand layers and intervals appear throughout the core. Some of them are graded and/or show erosional basal contacts.</p>
406							OL	
							dk GY	
408							OL	

Sample	Depth (mbsf)	Lithology	Texture (vol %)			Component										
			Sand	Silt	Clay	Nannofossils	Foraminifers	Biosiliceous Remains	Quartz & Feldspar	Detrital Carbonate	Clay Minerals	Lithoclasts	Mica	Heavy Minerals	Opagues	Volcanic Glass Shards
Hole D																
195-1202D-02-H-01, 105 cm	1.25	Black Mottle	10	80	10	co		pr	do	co	pr		pr	pr	ab	
195-1202D-02-H-02, 040 cm	2.10	Clayey Silt	5	80	15	co		co	ab	co	co		co	pr	co	
195-1202D-02-H-06, 120 cm	8.90	Sand	70	25	5	co	pr	pr	do	ab	pr	co	pr	co	pr	
195-1202D-04-H-02, 100 cm	21.70	Silt	5	90	5	pr		pr	do	ab	pr		co	pr	co	pr
195-1202D-05-H-07, 040 cm	38.10	Clayey Silt	2	88	10	pr		co	do	co	co		pr	pr	pr	
195-1202D-06-H-06, 040 cm	46.10	Clayey Silt	1	79	20	ra		pr	ab	co	co		pr	pr	pr	
195-1202D-07-H-06, 040 cm	55.60	Clayey Silt	5	85	10	pr		pr	do	co	co		pr	pr	pr	
195-1202D-08-H-06, 090 cm	65.60	Clayey Silt	1	79	20	pr		co	do	co	co		pr	ra	ra	
195-1202D-09-H-04, 132 cm	72.52	Clayey Silt	1	79	20	pr		pr	do	co	co		ra	ra	ra	
195-1202D-10-X-02, 055 cm	78.25	Clayey Silt	2	83	15	co		pr	do	ab	pr		ra	ra	pr	
195-1202D-11-X-02, 090 cm	88.10	Silt	2	93	5	pr		pr	ab	ab	pr		pr	pr	pr	
195-1202D-14-X-06, 100 cm	123.00	Clayey Silt	5	70	25	co		pr	ab	ab	co		pr	pr	pr	
195-1202D-14-X-06, 120 cm	123.20	Sandy Silt	20	75	5	ra		ra	ab	ab	pr	co	co	co	co	ra
195-1202D-17-X-03, 113 cm	147.53	Clayey Silt	2	88	10	co		pr	ab	ab	pr		pr	pr	pr	
195-1202D-17-X-03, 119 cm	147.59	Sandy Silt	35	55	10	co		pr	ab	co	pr	co	co	co	pr	
195-1202D-19-X-06, 020 cm	170.40	Clayey Silt		70	30	co		pr	ab	ab	ab		ra	ra	ra	
195-1202D-22-X-02, 090 cm	194.10	Clayey Silt	2	73	25	co		pr	ab	ab	ab		ra	ra	ra	
195-1202D-23-X-05, 060 cm	207.42	Clayey Silt		70	30	pr		pr	ab	ab	ab		ra	ra	ra	
195-1202D-25-X-01, 060 cm	221.10	Clayey Silt		70	30	pr		ra	ab	ab	ab		ra	ra	ra	
195-1202D-27-X-01, 070 cm	240.40	Sandy Silt	40	55	5	ra		ra	ab	ab	ra		pr	co	pr	
195-1202D-34-X-02, 090 cm	309.40	Silty Clay		85	15	pr		ra	do	ab	co		ra	ra	ra	
195-1202D-35-X-03, 055 cm	320.15	Sandy Silt	25	65	10	pr	ra	ra	ab	ab	pr		co	co	pr	
195-1202D-35-X-03, 080 cm	320.40	Clayey Silt	2	68	30	pr		ra	ab	ab	ab		ra	ra	pr	
195-1202D-37-X-03, 070 cm	339.60	Clayey Silt	1	64	35	ra		ra	ab	ab	ab		ra	ra	pr	
195-1202D-42-X-02, 100 cm	386.70	Clayey Silt	2	68	30	pr		ra	ab	ab	ab		ra	ra	ra	
195-1202D-44-X-02, 120 cm	406.20	Clayey Silt	1	74	25	ra		ra	ab	ab	ab		pr	ra	ra	