

Table T3. Location of calcareous nannofossil events.

Site	Event depth (mcd)											
	LCO <i>Gephyrocapsa caribbeanica</i>	FO <i>Emiliania huxleyi</i>	LO <i>Pseudoemiliania lacunosa</i>	FCO <i>Gephyrocapsa caribbeanica</i>	LO <i>Reticulofenestra asanoi</i>	RE Medium <i>Gephyrocapsa</i>	FO <i>Reticulofenestra asanoi</i>	LO <i>Gephyrocapsa</i> >5.5 µm	FO <i>Gephyrocapsa</i> >5.5 µm	LO <i>Calcidiscus macintyreii</i>	LO <i>Discoaster broweri</i>	
202-												
1237	5.03	5.35	13.38	14.88	21.98	27.82	27.82	30.30	31.81	36.61	37.90	
1238	10.00	11.00	27.44	30.15	49.42	49.42	68.52	68.52	77.54	82.43	98.92	
1240	16.80	20.52	42.87	74.15*	85.93	88.01	99.27	108.74	122.35	136.65	145.77	
1241	3.85	5.37	8.12	14.68*	21.16	22.66	24.00	26.17	28.34	37.19	49.57	
1242	29.64	34.48	52.37	62.01*	126.44	128.79	131.16	180.79	192.31	209.78	258.23	

Notes: * = depth is approximate. LCO = last common occurrence, FO = first occurrence, LO = last occurrence, FCO = first common occurrence, RE = reentrance.