

Table T8 (continued).

Hole, core, section, interval (cm)	Depth (mbsf)	Depth (mcd)	Preparation	Abundance	Preservation	Comment	<i>Globorotalia truncatulinoides</i>	<i>Globorotalia tosaensis</i>	<i>Globorotalia crassiformis</i>	<i>Globocerella inflata</i>	<i>Globigerina bullata</i>	<i>Globigerina quinqueloba</i>	<i>Globigerinella glutinata</i>	<i>Globigerinella scitula</i>	<i>Hirsutella universa</i>	<i>Globigerinoides sacculifer</i>	<i>Globigerinoides trilobus</i>	<i>Globigerinoides ruber</i>	<i>Globigerinella siphonifera</i>	<i>Globigerinoides conglobatus</i>
1266A-22H-CC	208.67	241.60	S	A	M	Reworking														
1266A-23H-CC	215.02	249.37	S	R	P	Downhole contamination														
1266A-24H-CC	215.88	251.27	S	R	P															
1266A-25H-CC	217.13	252.66	S	A	P															
1266A-26X-CC	218.66	254.37	S	R	P															
1266B-2X-CC	229.56	262.56	S	A	M-G															
1266A-27X-CC	230.94	267.32	S	A	M															
1266B-3X-CC	239.17	273.59	S	A	G															
1266A-28X-CC	237.33	275.14	S	A	G															
1266B-4X-CC	248.63	284.48	S	A	G															
1266A-29X-CC	250.06	289.30	S	A	G															
1266B-5H-CC	258.44	295.72	S	A	G															
1266A-30X-CC	259.93	300.62	S	A	G															
1266B-6H-7, 100–101	267.40	306.10	38	A	M	Some fragmentation														
1266B-6H-7, 113–114	267.53	306.23	38	A	M	Minor dissolution														
1266B-6H-7, 128–129	267.68	306.38	38	A	P-M	Heavy fragmentation														
1266B-6H-7, 139–140	267.79	306.49	38	C	P-M	Heavy fragmentation														
1266B-6H-7, 148–149	267.88	306.58	38	R	P-M	Pyrite?, abundant benthics														
1266B-6H-CC, 6–7	267.96	306.66	38	R	P-M															
1266B-6H-CC, 22–23	268.12	306.82	38	R	P	Chalky, heavy fragmentation														
1266B-6H-CC, 26–27	268.16	306.86	38	C	M-G															
1266B-6H-CC	268.17	306.87	S	A	G															
1266A-31X-CC	265.26	308.39	S	A	G															
1266B-7X-CC	275.84	315.96	S	A	M-G															
1266A-32X-CC	277.47	321.07	S	A	G															
1266B-8X-CC	281.46	325.32	S	A	M-G															
1266A-33X-CC	287.44	332.47	S	A	G															
1266B-9X-CC	292.32	336.49	S	A	M-G															
1266A-34X-CC	295.14	343.92	S	A																
1266B-10X-CC	302.08	347.69	S	A	M-G															
1266B-11X-CC	311.23	358.28	S	A	M															
1266C-20X-CC	320.93	367.21	S	A	P-M	Chalky														
1266C-21X-CC	333.08	379.36	S	A	M															

Notes: Preparation: S = >63-µm size fraction, 38 = >38-µm size fraction. Abundance: A = abundant , C = common, F = few, R = rare,

SHIPBOARD SCIENTIFIC PARTY
CHAPTER 7, SITE 1266

10

Table T8 (continued).

Hole, core, section, interval (cm)	Depth (mbsf)	Depth (mcd)	Preparation	Abundance	Preservation	Comment	<i>Hirsutella hirsuta</i>	<i>Menardella menardi</i>	<i>Globorotalia tumida</i>	<i>Neogloboquadrina pachyderma</i> (dextral)	<i>Neogloboquadrina dutertrei</i>	<i>Globigerinoides obliquus</i>	<i>Sphaeroidinella defhsicens</i>	<i>Globigerina apertura</i>	<i>Globigerinoides extremus</i>	<i>Pulvinatina praecursor</i>	<i>Globconella punctulata</i>	<i>Globigerina decolorata</i>	<i>Globconella conomiozea</i>	<i>Globigerina woodi</i>	<i>Hirsutella marginata</i>
1266A-22H-CC	208.67	241.60	S	A	M	Reworking															
1266A-23H-CC	215.02	249.37	S	R	P	Downhole contamination															
1266A-24H-CC	215.88	251.27	S	R	P																
1266A-25H-CC	217.13	252.66	S	A	P																
1266A-26X-CC	218.66	254.37	S	R	P																
1266B-2X-CC	229.56	262.56	S	A	M-G																
1266A-27X-CC	230.94	267.32	S	A	M																
1266B-3X-CC	239.17	273.59	S	A	G																
1266A-28X-CC	237.33	275.14	S	A	G																
1266B-4X-CC	248.63	284.48	S	A	G																
1266A-29X-CC	250.06	289.30	S	A	G																
1266B-5H-CC	258.44	295.72	S	A	G																
1266A-30X-CC	259.93	300.62	S	A	G																
1266B-6H-7, 100-101	267.40	306.10	38	A	M	Some fragmentation															
1266B-6H-7, 113-114	267.53	306.23	38	A	M	Minor dissolution															
1266B-6H-7, 128-129	267.68	306.38	38	A	P-M	Heavy fragmentation															
1266B-6H-7, 139-140	267.79	306.49	38	C	P-M	Heavy fragmentation															
1266B-6H-7, 148-149	267.88	306.58	38	R	P-M	Pyrite?, abundant benthics															
1266B-6H-CC, 6-7	267.96	306.66	38	R	P-M																
1266B-6H-CC, 22-23	268.12	306.82	38	R	P	Chalky, heavy fragmentation															
1266B-6H-CC, 26-27	268.16	306.86	38	C	M-G																
1266B-6H-CC	268.17	306.87	S	A	G																
1266A-31X-CC	265.26	308.39	S	A	G																
1266B-7X-CC	275.84	315.96	S	A	M-G																
1266A-32X-CC	277.47	321.07	S	A	G																
1266B-8X-CC	281.46	325.32	S	A	M-G																
1266A-33X-CC	287.44	332.47	S	A	G																
1266B-9X-CC	292.32	336.49	S	A	M-G																
1266A-34X-CC	295.14	343.92	S	A																	
1266B-10X-CC	302.08	347.69	S	A	M-G																
1266B-11X-CC	311.23	358.28	S	A	M																
1266C-20X-CC	320.93	367.21	S	A	P-M	Chalky															
1266C-21X-CC	333.08	379.36	S	A	M																

B = barren. Preservation: G = good, M = moderate, P = poor. Occurrence: a = abundant, c = common, f = frequent, r = rare, b = barren.

Table T8 (continued).

Table T8 (continued).

Table T8 (continued).

Table T8 (continued).

Table T8 (continued).

Hole, core, section, interval (cm)	Depth (mbsf)	Depth (mcd)	Preparation	Abundance	Preservation	Comment	<i>Morozovella gracilis</i>	<i>Morozovella subbotinae</i>	<i>Morozovella aequa</i>	<i>Morozovella acuta</i>	Biserials	<i>Morozovella acutispira</i>	<i>Subbotina patagonica</i>	<i>Subbotina triangularis</i>	<i>Subbotina relascoensis</i>	<i>Parasubbotina variospirae</i>	<i>Subbotina triloculinoidea</i>	<i>Morozovella velascensis</i>	<i>Morozovella conicotruncata</i>	<i>Morozovella angulata</i>	<i>Morozovella occulta</i>
1266A-22H-CC	208.67	241.60	S	A	M	Reworking															
1266A-23H-CC	215.02	249.37	S	R	P	Downhole contamination															
1266A-24H-CC	215.88	251.27	S	R	P																
1266A-25H-CC	217.13	252.66	S	A	P																
1266A-26X-CC	218.66	254.37	S	R	P																
1266B-2X-CC	229.56	262.56	S	A	M-G																
1266A-27X-CC	230.94	267.32	S	A	M																
1266B-3X-CC	239.17	273.59	S	A	G																
1266A-28X-CC	237.33	275.14	S	A	G																
1266B-4X-CC	248.63	284.48	S	A	G																
1266A-29X-CC	250.06	289.30	S	A	G																
1266B-5H-CC	258.44	295.72	S	A	G																
1266A-30X-CC	259.93	300.62	S	A	G																
1266B-6H-7, 100–101	267.40	306.10	38	A	M	Some fragmentation	f	f	f	f	f	f	f	f	f				r	r	
1266B-6H-7, 113–114	267.53	306.23	38	A	M	Minor dissolution	r	c	f	r	f	f	r	r	f				r	r	
1266B-6H-7, 128–129	267.68	306.38	38	A	P-M	Heavy fragmentation	c	f	r	r	r	r	r	r	r						
1266B-6H-7, 139–140	267.79	306.49	38	C	P-M	Heavy fragmentation	c	r	r	r	r	r	r	r	r						
1266B-6H-7, 148–149	267.88	306.58	38	R	P-M	Pyrite?, abundant benthics	r	r	r	r	r	r	r	r	r						
1266B-6H-CC, 6–7	267.96	306.66	38	R	P-M																
1266B-6H-CC, 22–23	268.12	306.82	38	R	P	Chalky, heavy fragmentation	r	r	r	r	r	r	r	r	r						
1266B-6H-CC, 26–27	268.16	306.86	38	C	M-G		f	r	r	r	f	r	r	f	f		r	r	f	f	
1266B-6H-CC	268.17	306.87	S	A	G		c	f	f	r	f	f	f	f	f		f	f	f	f	
1266A-31X-CC	265.26	308.39	S	A	G		f	f	f	f	f	f	f	f	f		r	r	f	f	
1266B-7X-CC	275.84	315.96	S	A	M-G		f	f	f	f	f	f	f	f	f		f	f	f	f	
1266A-32X-CC	277.47	321.07	S	A	G		f	f	r	f	f	f	f	f	f		r	r	f	f	
1266B-8X-CC	281.46	325.32	S	A	M-G		f	f	r	f	f	f	f	f	f		r	r	f	f	
1266A-33X-CC	287.44	332.47	S	A	G		f	f	r	f	f	f	f	f	f		r	r	f	f	
1266B-9X-CC	292.32	336.49	S	A	M-G		f	f	r	f	f	f	f	f	f		r	r	f	f	
1266A-34X-CC	295.14	343.92	S	A			f	f	r	f	f	f	f	f	f		r	r	r	r	
1266B-10X-CC	302.08	347.69	S	A	M-G		f	f	r	f	f	f	f	f	f		r	r	f	f	
1266B-11X-CC	311.23	358.28	S	A	M												r	r	r	r	
1266C-20X-CC	320.93	367.21	S	A	P-M	Chalky											r	r	r	r	
1266C-21X-CC	333.08	379.36	S	A	M												r	r	f	c	r

SHIPBOARD SCIENTIFIC PARTY
CHAPTER 7, SITE 1266

16

Table T8 (continued).

Hole, core, section, interval (cm)	Depth (mbsf)	Depth (mcd)	Preparation	Abundance	Preservation	Comment	Acarinina coalingensis	Acarinina nitida	Acarinina mckannai	Morozovella edgari	Globanomalina planocanica	Globanomalina imitata	Globanomalina chapmani	Globanomalina ehrenbergi	Globanomalina pseudomenardi	Igorina albeani	Igorina tadjikistanensis	Acarinina subsphaerica		
1266A-22H-CC	208.67	241.60	S	A	M	Reworking														
1266A-23H-CC	215.02	249.37	S	R	P	Downhole contamination														
1266A-24H-CC	215.88	251.27	S	R	P															
1266A-25H-CC	217.13	252.66	S	A	P															
1266A-26X-CC	218.66	254.37	S	R	P															
1266B-2X-CC	229.56	262.56	S	A	M-G															
1266A-27X-CC	230.94	267.32	S	A	M															
1266B-3X-CC	239.17	273.59	S	A	G															
1266A-28X-CC	237.33	275.14	S	A	G															
1266B-4X-CC	248.63	284.48	S	A	G															
1266A-29X-CC	250.06	289.30	S	A	G															
1266B-5H-CC	258.44	295.72	S	A	G															
1266A-30X-CC	259.93	300.62	S	A	G															
1266B-6H-7, 100–101	267.40	306.10	38	A	M	Some fragmentation	f				f	r								
1266B-6H-7, 113–114	267.53	306.23	38	A	M	Minor dissolution	f		r											
1266B-6H-7, 128–129	267.68	306.38	38	A	P-M	Heavy fragmentation	c		r											
1266B-6H-7, 139–140	267.79	306.49	38	C	P-M	Heavy fragmentation	c													
1266B-6H-7, 148–149	267.88	306.58	38	R	P-M	Pyrite?, abundant benthics	c													
1266B-6H-CC, 6–7	267.96	306.66	38	R	P-M		r													
1266B-6H-CC, 22–23	268.12	306.82	38	R	P	Chalky, heavy fragmentation	r			r		r								
1266B-6H-CC, 26–27	268.16	306.86	38	C	M-G		f				r	r								
1266B-6H-CC	268.17	306.87	S	A	G		r		r	r	r	r								
1266A-31X-CC	265.26	308.39	S	A	G		f	f	f	f	r	r								
1266B-7X-CC	275.84	315.96	S	A	M-G		r		f									r?		
1266A-32X-CC	277.47	321.07	S	A	G															
1266B-8X-CC	281.46	325.32	S	A	M-G		f		f									f		
1266A-33X-CC	287.44	332.47	S	A	G											r		f		
1266B-9X-CC	292.32	336.49	S	A	M-G															
1266A-34X-CC	295.14	343.92	S	A																
1266B-10X-CC	302.08	347.69	S	A	M-G		f		f							f		f		
1266B-11X-CC	311.23	358.28	S	A	M					f						f		f		
1266C-20X-CC	320.93	367.21	S	A	P-M	Chalky				r		r	r	r	c		f	r		
1266C-21X-CC	333.08	379.36	S	A	M					r		f	f							