

INDEX TO VOLUME 113

This index provides coverage for both the *Initial Reports* and *Scientific Results* portions of Volume 113 of the *Proceedings of the Ocean Drilling Program*. Index entries with the suffix *a* refer to pages in the *Initial Reports*, and those with *b*, to pages in the *Scientific Results* (this book).

The index is presented in three parts: (1) a Subject Index, (2) a Site Index, and (3) a Paleontological Index. In addition to this printed version, the index is also available in the form of a machine-readable, ASCII-encoded, 9-track magnetic tape, 1600 bpi.

The index was prepared by Wm. J. Richardson Associates, Inc., under subcontract to the Ocean Drilling Program. It follows the concept developed by the Deep Sea Drilling Project at Scripps Institution of Oceanography for a comprehensive, cumulative index of DSDP volumes. Both of these indexes are based on a hierarchy of entries: (1) a main entry, defined as a key word or concept followed by a reference to the page on which that word or concept appears; (2) a subentry, defined as a further elaboration on the main entry followed by a page reference; and (3) a sub-subentry, defined as an even further elaboration on the main entry or subentry followed by a page reference.

The Subject Index follows a standard format. Geographic and individual names are referenced in the index only if they are subjects of discussion. This index also includes broad fossil groups, such as foraminifers and radiolarians, which also appear in the Paleontological Index.

The Site Index is structured to contain entries for the sites discussed in the volume. Site entries are modified by subject subentries.

The Paleontological Index is an index relating to significant findings and/or substantive discussions, not of species names *per se*. This index covers three varieties of information: (1) broad fossil groups, including individual genera and species that have been erected or emended formally; (2) biostratigraphic zones; and (3) fossils depicted in illustrations.

The indexes cover text figures and tables but not core description forms ("barrel sheets") or core photographs. Also excluded are bibliographic references, names of individuals, and routine front and back matter.

For further information, contact the Chief Production Editor, Ocean Drilling Program, 1000 Discovery Drive, College Station, Texas 77845-9547.

SUBJECT INDEX

- Acoustic basement, South Orkney microcontinent: Site 696, location, 28b
 Agulhas Plateau, Maestrichtian water temperatures, 821b
 Alaska, glacial diamictites, 953b
 Alkalinity
 Dronning Maud Land margin: Site 693, 374a–375a
 depth and, 138b, 141b, 380a
 Jane Basin, 730a, 735a
 depth and, 140b, 143b, 736a–737a
 Maud Rise: Site 689, depth and, 129a–130a, 138b, 140b
 Maud Rise: Site 690, 230a
 depth and, 138b, 141b, 237a
 South Orkney microcontinent: Site 695, 560a
 depth and, 139b, 142b, 561a–562a
 South Orkney microcontinent: Site 696, 647a
 depth and, 139b, 143b, 650a–651a
 Weddell Basin: Site 694, depth and, 139b, 142b
 Alkane, Weddell Sea, 201b–202b, 206b
 Alteration
 benthic foraminiferal isotope record and, 836b
 volcanic ash, 144b
 See also Calcite
 Alteration, volcanic
 clay, 53b
 interstitial-water chemistry and, 140b–145b
 Ammonia concentrations
 Dronning Maud Land margin: Site 693, 375a
 depth and, 380a
 Jane Basin, depth and, 735a–737a
 Maud Rise: Site 690, 230a
 depth and, 237a
 South Orkney microcontinent: Site 695, depth and, 561a–562a
 South Orkney microcontinent: Site 696, 647a–648a
 depth and, 650a–651a
 Weddell Basin: Site 694, 484a
 depth and, 485a–486a
 Ammonium, as evidence of production rates, 889b
 Amphibole
 Dronning Maud Land margin: Site 693, dating, 55b
 Jane Basin, dating, 62b, 63b
 Maud Rise: Site 690, dating, 58b
 South Orkney microcontinent: Site 696
 dating, 61b
 depth and, 30b
 peak areas and heights, 29b
 Weddell Basin: Site 694, dating, 59b
 Anoxia, cyclic, Cretaceous, 195b
 Antarctic Bottom Water (AABW), 122b, 938b
 in Jane Basin, 803b
 Jane Basin area, 111b
 Maud Rise area, 865b–866b
 See also Proto-AABW
 Antarctic Circumpolar Current
 development of, 65b, 938b, 952b
 initiation of, 803b
 Antarctic margin
 glaciation, 385a
 location, 122b
 morphology, 43b
 three-dimensional view, 43b
 Antarctic margin E
 geologic provinces, 90b
 glaciation, 658a
 ice sheets, development, 462a, 489a, 955b
 as source of quartz sand, 98b–99b
 Antarctic margin W
 geologic provinces, 90b
 glaciation, 658a–659a
 ice sheets, 892b
 development, 462a, 489a–490a
 geochemical signals of formation, 881b
 Antarctic Ocean
 climatic change before Cretaceous/Tertiary boundary, 842b
 as CO₂ sink, 127b
 drill sites, 902b
 Miocene cooling, isotope changes and, 890b
 planktonic foraminiferal seasonal succession, Paleogene, 850b
 sea-ice formation, in Oligocene, 875b
 seasonal thermocline
 Cretaceous changes, 941b
 Eocene, 948b
 Paleocene-Eocene changes, 861b–862b
 stable isotopes
 benthic foraminiferal data, 865b
 Paleogene anomalies, 875b–876b
 surface-water temperatures, 850b
 Cretaceous, 843b, 844b
 Cretaceous-Eocene, 937b
 Eocene, 947b
 Paleocene-Eocene changes, 859b–861b, 863b
 Warm Saline Deep Waters in Paleogene, 874b, 875b
 See also Weddell Sea
 Antarctic Peninsula
 biostratigraphic correlations, 902b
 cooling, Miocene, 955b
 ice-rafterd dropstones, 463a
 See also Seymour Island
 Antarctica
 Aptian land masses, 345b
 bottom water production and, Maestrichtian, 821b
 carbon isotopes at Cretaceous/Tertiary boundary, 844b
 climate, Maestrichtian, 941 b
 conifer forests, Cretaceous, 454b–455b
 continental shelves, as source of cold deep waters, 842b
 cryosphere
 development, 937b
 Miocene, 954b–955b
 glaciation
 differential rates of, 952b
 evolution of, 881b
 initiation of, 950b
 Pliocene, 803b, 807b–808b
 ice sheets
 Antarctic water column and, 866b
 development, 462a, 489a–490a, 939b, 953b–954b
 Eocene growth, 948b
 formation of, 889b–890b
 isotopic composition and, 872b–873b
 ice-shelf formation, reflector correlations, 125b
 Nothofagus forests
 Eocene, 604b, 937b, 949b
 Oligocene, 952b
 Paleocene, 945b
 “Strangeglove Ocean” theory and, 844b
 Antarctica E
- cryosphere, Oligocene development of, 952b–953b
 ice-sheet development, 937b, 938b
 ice-sheet formation, 876b
 Antarctica W, ice-sheet development, 938b, 955b–956b
 Apatite
 Dronning Maud Land margin: Site 692, formation method, 75b
 Maud Rise: Site 689, LREE content, 10b
 Maud Rise: Site 690
 cleavage traces, 14b
 LREE content, 10b
 micropore analysis, 7b
 texture, 13b
 Argentine Basin, oxygen isotope values, 821b
 Ash, volcanic
 Dronning Maud Land margin, 299a–302a
 Jane Basin, 708a–710a
 Maud Rise, 194a
 South Orkney microcontinent, 533a, 535a, 620a
 Atlantic Ocean N
 carbon isotope values, 873b
 ice rafting, 803b
 Atlantic Ocean NE, “black shales,” 194b–195b
 Atlantic Ocean NW, “black shales” 195b
 Atlantic Ocean S, bottom water production, Maestrichtian, 821b
 Baffin Bay, carbon isotope changes, sources of, 881b–882b
 Basalt, Maud Rise: Site 690, dating, 189a
 Beaufort Sea, terrigenous matter transport, 881b
 Bellingshausen Basin, *Bolboforma*, 954b
 Biotite, Maud Rise: Site 690, micropore analysis, 7b
 “Black shales,” locations, 194b–195b
 Blake Event, Jane Basin, 727a
 Blake-Bahama Formation, “black shales,” 194b
 Bouguer Basin
 reflectors, two-way travelttime, 34b
 seismic stratigraphy, P-wave velocity, 35b
 See also South Orkney microcontinent
 Bransfield Strait
 carbon isotopes, 890b
 perylene, 206b
 Butane concentrations
 Dronning Maud Land margin: Site 692, depth and, 313a
 Dronning Maud Land margin: Site 693, 377a
 depth and, 382a
 retention times, 383a
 Weddell Basin: Site 694, depth and, 487a
 C/N ratio
 Dronning Maud Land margin: Site 693, 883b
 Maud Rise: Site 689, 883b, 890b, 891b
 Maud Rise: Site 690, 883b, 891b
 Weddell Basin: Site 694, 883b–884b
 Calcite
 alteration, isotopic composition and, 814b
 Dronning Maud Land margin: Site 692, depth and, 75b, 76b
 foraminiferal
 chemical preservation, 816b
 isotopic preservation, 816b
 textural alteration, 816b
 precipitation, Cretaceous vs. Recent
 foraminiferal species, 815b
 preservation, foraminiferal data, 827b

SUBJECT INDEX

- recrystallization, 815b-816b
secondary, foraminiferal data, 824b-826b
- Calcium carbonate
Maud Rise
 accumulation rates, 841b
 Oligocene abundances, 954b
Maud Rise: Site 689, depth and, 103a-104a
Maud Rise: Site 690, depth and, 207a-208a
- Calcite compensation depth
Weddell Sea, 951b, 952b
 Oligocene-Neogene, 954b
- Calcium concentrations
Dronning Maud Land margin: Site 693, 375a
 depth and, 138b, 141b, 380a
Jane Basin, depth and, 140b, 143b, 736a-737a
Maud Rise: Site 689, depth and, 129a-130a, 138b, 140b
Maud Rise: Site 690, 230a
 depth and, 138b, 141b, 237a
South Orkney microcontinent: Site 695, depth and, 139b, 142b, 561a-562a
- South Orkney microcontinent: Site 696, 648a-649a
 depth and, 139b, 143b, 650a-651a
Weddell Basin: Site 694, 484a
 depth and, 139b, 142b, 485a-486a
- Carbon. *See also* C/N ratio
Carbon burial, Cretaceous climate change and, 843b-844b
Carbon, inorganic, global reservoir, Cretaceous changes in, 941b
Carbon isotope ratio
 disequilibrium in foraminiferal test secretion, 849b
Maud Rise, 836b, 839b-841b
 foraminiferal data, 814b, 815b, 818b, 819b, 820b, 831b-832b, 837b
 Sites 689/690 compared, 947b
 water temperatures and, 813b
Maud Rise: Hole 690, 943b
Paleocene/Eocene boundary, negative shift, 945b
- Carbon isotopes
Antarctic Ocean, as climatic indicators, 866b
calcite recrystallization and, 815b-816b
Cenozoic excursions, 843b
in clastic sediments, 889b
Cretaceous/Tertiary boundary, global changes, 944b
Maestrichtian foraminiferal records compared, 821b
- Maud Rise, 861b-862b
 Cretaceous climate changes and, 941b
 Sites 689/690 comparisons, 868b, 869b-870b
 surface-to-deep-water gradients, 871b-872b
Maud Rise: Site 689, 858b-859b
Maud Rise: Site 690, 853b-854b, 856b-858b
Weddell Sea, values, 884b-889b, 890b, 891b, 894b-897b
- Carbon, organic
 accumulation in modern deep-sea sediments, 844b
DOC in interstitial waters, 173b-174b
 Weddell Sea, 173b, 176b
- Dronning Maud Land margin: Site 692
 black shale, 316a
 depth and, 314a
 Rock-Eval and coulometric comparison, 315a
- terrigenous versus marine sources of, 88 lb
- TOC levels, Weddell Sea, 189b-190b
- Carbon, total dissolved (TDC), global change at Cretaceous/Tertiary boundary, 829b, 844b-845b
- Carbonates
- Maud Rise
 accumulation rates, 834b
 alteration debated, 874b
- chalk
Dronning Maud Land margin, dating, 299a
Maud Rise: Site 689
 chert with, 98a
 dating, 95a
Maud Rise: Site 690, dating, 189a
- Chalk, nannofossil, Dronning Maud Land margin: Site 692, depth and, 74b
- Chert
Dronning Maud Land margin: Site 692, depth and, 74b
Dronning Maud Land margin: Site 693, color, 77b-78b
Maud Rise: Site 689, chalk with, 98a
- Chloride concentrations
Dronning Maud Land margin: Site 693, depth and, 380a
Jane Basin, 730a
 depth and, 736a-737a
Maud Rise: Site 690, depth and, 237a
- South Orkney microcontinent: Site 695, depth and, 561a-562a
- South Orkney microcontinent: Site 696, depth and, 650a-651a
- Chlorinity
Dronning Maud Land margin: Site 693, 374a
 depth and, 379a-380a
Maud Rise: Site 689, depth and, 128a-130a
Maud Rise: Site 690, 230a
South Orkney microcontinent: Site 695, 560a
South Orkney microcontinent: Site 696, 643a, 646a
- Weddell Basin: Site 694, depth and, 481a, 485a-486a
- Chlorite
Dronning Maud Land margin: Site 693, dating, 344a
Jane Basin
 dating, 62b, 63b
 TEM photomicrograph, 231b
Maud Rise: Site 689, dating, 57b, 99a
Maud Rise: Site 690, dating, 197a
- South Orkney microcontinent: Site 695
 chemical weathering and, 538a
 dating, 539a
- South Orkney microcontinent: Site 696
 dating, 61b, 623a
 peak areas and heights, 29b
Weddell Basin: Site 694, dating, 59b
- Chondrite
Maud Rise: Site 690, composition, 8b
South Orkney microcontinent: Site 696, size, 621a
- Weddell Basin: Site 694, mudstone and, 462a
- Chromite, Maud Rise: Site 690, microprobe analysis, 7b
- Circulation, oceanic
 Antarctic role in, 938b-939b
 climate and, in Eocene, 874b
 evolution of, 875b
 in Paleogene, 876b
 halothermal, 865b
 Paleogene, 950b-951b
 salinity and, 865b
 in Paleogene, 877b
 thermohaline, 865b
 Antarctic sea-ice formation and, 875b
- Circumpolar Deep Water, Maud Rise, 866b
- Clay
 Antarctica E, major events, 64b
 Antarctica W, major events, 64b
- Dronning Maud Land margin
 diagenesis, 53b
 major events, 64b
 mineralogy, 57b
- Dronning Maud Land margin: Site 691, dating, 299a
- Dronning Maud Land margin: Site 692
 carbonate and, 77b, 79b-80b
 color, 73b-75b
 dating, 299a, 300a
 depth and, 74b, 75b, 76b
 mineralogy, 301a, 303a
 texture, 73b-75b
- Dronning Maud Land margin: Site 693
 bioturbation, 83b
 dating, 335a, 336a
 depth and, 84b, 335a, 336a
 mineralogy, 68b
- Jane Basin
 clay, 63b
 dating, 63b, 711a
 depth and, 711a
 microfabric analysis, 227b-230b
 microfabric techniques, 226b-227b
 mineralogy, 62b, 710a-712a, 715a
 physico-chemical forces, 225b-226b
 size, 712a
- Lord Howe Rise, volcanic alteration, 53b
- Maud Rise
 major events, 64b
 volcanic alteration, 53b
- Maud Rise: Site 689, mineralogy, 57b
- Maud Rise: Site 690
 depth and, 190a-191a
 mineralogy, 58b
- mineralogy, Maud Rise: Site 689, 67b
- South Orkney microcontinent
 diagenesis, 53b
 weathering, 53b-54b
- South Orkney microcontinent: Site 695
 dating, 532a
 mineralogy, 537a-539a
- South Orkney microcontinent: Site 696
 dating, 614a-615a
 depth and, 30b, 614a, 616a
 mineralogy, 61b, 620a-623a
- Tasman Sea, volcanic alteration, 53b
- Weddell Basin: Site 694
 dating, 59b
 depth and, 460a
- Weddell Sea
 diagenesis, 53b
 volcanic alteration, 53b
 weathering, 53b-54b
- Clay minerals
Antarctic Ocean, Miocene climate indicators, 955b
porosity and, 244b
- South Orkney microcontinent, Oligocene climate and, 953b
- Weddell Basin: Site 694, 59b, 461a-462a, 465a
- Weddell Sea, 51b-52b
 Cretaceous/Tertiary boundary section, 944b
 Eocene climate indicators, 949b
 Maestrichtian climate and, 941b
 Paleogene changes, 951b
- See also* particular minerals
- Consolidation
clay, 230b
inference from stress tests, 212b
processes involved, 211b
shear strength and, 211b
- Cooling, global

- Cretaceous, 820b-821b
isotope data and, 819b
- Corundum
South Orkney microcontinent: Site 696
depth and, 30b
X-ray diffractogram, 31b
- Cretaceous/Tertiary boundary
Antarctica
carbon isotopic temperature gradient, 839b-841b
climate changes preceding, 829b, 844b-845b
oxygen isotopic temperature gradient, 836b-839b
benthic foraminiferal assemblage, 575b, 585b
calcareous dinoflagellate assemblages, 533b-548b
carbonate accumulation rates, 844b
mass extinction event, climatic change preceding, 842b, 844b-845b
- Maud Rise
basis for age model, 833b
magnetostratigraphy, 255b, 258b-259b
- Maud Rise: Site 689, unconformity, 903b
nannofossil assemblages, 515b-526b
“survivor” microfossil species defined, 519b
- Weddell Sea
climatic changes, 941b
microfossil population changes, 944b
surface-to-deep-water gradients, 944b
- See also* Iridium
- Cryosphere, global, elements in, 952b
- Density
Dronning Maud Land margin: Site 692, depth and, 304a
- Dronning Maud Land margin: Site 693, depth and, 351a-352a
- Jane Basin, depth and, 716a
- Maud Rise: Site 689
depth and, 102a-103a, 105a
large variations in, 101a
- Maud Rise: Site 690, depth and, 205a-206a
- South Orkney microcontinent: Site 695, depth and, 180b, 541a-543a
- South Orkney microcontinent: Site 696
depth and, 30b, 625a
range, 624a, 626a
- Weddell Basin: Site 694, depth and, 466a-467a
- Diagenesis
dissolved organic carbon, 169b
opal, physical properties associated with, 32b
- Diamictites, glacial, Prdzy Bay, 953b
- Dissolution, diatoms, as marker for sea-ice limits, 804b
- Dolomitization, Weddell Sea, 140b
- Drake Passage
deepening and widening, 65b
opening, 658a
- Dronning Maud Land margin
mineralogy, clay, 57b
paleoenvironment, thermal subsidence, 54b
- Dronning Maud Land margin: Site 691
bathymetry, 40b, 296a
lithostratigraphy, 298a
summary, 299a
- See also* Dronning Maud Land margin: Site 692
- location, 296a
seismic reflection profile, 295a
seismic stratigraphy, 305a
volcanic ash, dating, 299a
- Dronning Maud Land margin: Site 692
bathymetry, 40b, 296a
butane concentrations, depth and, 313a
- calcium carbonate content, 74b
depth and, 75b
- density, depth and, 304a
- depositional environment, 75b-77b
- ethane concentrations, depth and, 313a
- kerogen concentrations, 189b-190b, 314a-315a
carbon isotope ratios, 195b-196b
data summary, 191b
sulfur isotope ratios, 195b-196b
vitrinite reflectance measurements, 192b-193b
- lithology, Site 693 compared with, 86b
- lithostratigraphy
ice-rafterd dropstones, 301a-303a, 304a
Miocene-Quaternary, 298a
sedimentary structures and, 73b, 74b, 78b
summary, 299a
Valanginian-Hauterivian, 298a-301a
- location, 72b, 296a
- magnetic properties, 311a-312a
demagnetization, 312a
natural remanent magnetization, 312a-313a
- methane concentrations, depth and, 313a
- mineralogy, 75b
clay, 301a, 303a
depth and, 76b
- organic carbon concentrations
black shale, 316a
depth and, 314a
Rock-Eval and coulometric comparison, 315a
- organic carbon content, 75b
- paleoenvironment, clay, 301a
- paleolatitude, 85b
- porosity, depth and, 304a-305a
- propane concentrations, depth and, 313a
- reflectors, dating, 316a-317a
- Rock-Eval procedure, 191b-192b, 313a-315a
- seismic reflection profile, 295a
- seismic stratigraphy, 305a
lithology correlated with, 46b
P-wave velocity, 304a
sequences, 317a
- velocity, depth and, 304a-305a
- volcanic ash
composition, 300a
dating, 299a-300a
size, 301a-302a
- water content, depth and, 304a
- Dronning Maud Land margin: Site 693
age-depth plots, 927b
alkalinity, 374a-375a
depth and, 138b, 141b, 380a
- ammonia concentrations, 375a
depth and, 380a
- bathymetry, 40b, 331a
- butane concentrations, 377a
depth and, 382a
retention times, 383a
- calcium carbonate content, 83b
depth and, 974b-975b
- calcium concentrations, 375a
depth and, 138b, 141b, 380a
- chloride concentrations, depth and, 380a
- chlorinity, 374a
depth and, 379a-380a
- consolidation
data summary, 214b, 222b
field curve, 217b
- density, depth and, 351a-352a
- depositional environment, 79b-80b
- dissolved organic carbon
depth and, 172b
- pore water concentrations, 174b
- downhole measurements, 378a, 381a
gamma-ray log, 242b-243b, 383a
- resistivity log, 242b-243b, 383a
- sonic log, 242b-243b, 383a
- synthetic seismogram, 384a
- ethane concentrations, 377a
depth and, 382a
retention times, 383a
- hydrography, 122b
- interstitial-water chemistry
data summary, 138b, 379a
procedure, 136b-137b
- kerogen concentrations, 190b-193b, 377a
vitrinite reflectance measurements, 192b-193b
- lithology
dating, 92b
depth and, 92b
Site 692 compared with, 86b
- lithostratigraphy
Albian-Santonian, 339a-340a
Aptian-Albian, 340a
ice-rafterd dropstones, 345a-349a, 350a
Miocene-Oligocene, 240b
Oligocene, 338a-339a, 341a
Oligocene-Pliocene, 336a-338a, 339a
Pleistocene, 334a
Pliocene-Miocene, 239b-240b
Pliocene-Pleistocene, 334a, 336a
sedimentary structures and, 82b
summary, 333a
- location, 72b, 331a
- magnesium concentrations, 375a
depth and, 138b, 141b, 380a
- magnetic properties
demagnetization, 375a
natural remanent magnetization, 368a-369a, 370a-371a, 373a
- magnetostratigraphy
geomagnetic polarity time scale, 372a, 376a, 378a
glacial-interglacial cycles correlated with, 127b
natural remanent magnetization, 384a
- methane concentrations, 377a
depth and, 382a
retention times, 383a
- mineralogy
clay, 68b, 342a, 344a
depth and, 84b
- oxygen isotope distribution, depth and, 138b, 141b
- paleoclimate
Milankovitch cycles and logs, 245b-250b
Miocene-Pliocene, 248b-250b
Oligocene-Miocene; 248b
Pliocene, 250b
spectral analysis, 244b-248b
- paleoenvironment, 343a, 345a
- paleolatitude, 85b
- pH, 374a
depth and, 380a
- phosphate concentrations, 375a
depth and, 380a
- porosity, 240b, 242b-244b
depth and, 215b-217b, 243b, 351a-353a
- diatom relationship with, 215b-217b
- potassium concentrations, 375a-376a
depth and, 138b, 141b, 380a
- propane concentrations
depth and, 382a
retention times, 383a
- reflectors
dating, 357a-358a, 385a
depth and, 358a
- ice shelf formation correlated with, 125b

SUBJECT INDEX

- two-way travertime, 45b
unconformity boundary, 44b-45b
- Rock-Eval procedure, 377a, 381a
salinity, 374a
- sedimentation processes, 122b-123b
distribution, 124b, 126b
Oligocene-Miocene, 123b-125b
Pleistocene, 126b-128b
Pliocene-Pleistocene, 125b-126b, 129b
sedimentation rates
 biostratigraphic data, 369a, 377a
 hiatus, 55b, 369a, 372a, 383a
 magnetostratigraphic data, 369a, 377a
seismic reflection profile, 332a
seismic stratigraphy, 353a-356a
 Cretaceous-Oligocene hiatus, 357a-358a
 lithology correlated with, 46b
 Miocene hiatus, 356a-357a
 P-wave velocity, 44b, 243b, 353a
shear strength, 351a-352a
 depth and, 217b, 355a
 vertical effective stress relationship with, 217b
- strontium isotope distribution
 calculations, 153b, 155b-156b
 depth and, 150b
 sediment age versus, 152b
- sulfate concentrations, 374a-375a
 depth and, 138b, 141b, 380a
 pore water concentrations, 175b
- thermal conductivity, 352a
 depth and, 356a
- total organic carbon, depth and, 190b
- velocity, 240b, 242b-244b, 351a
 depth and, 353a, 354a
- volcanic glass, color, 78b
- water content, depth and, 351a-352a
- See also* Weddell Sea
- Dropstones
 glacial marine, Maud Rise, 96a
ice rafted
 Antarctic Peninsula, 463a
 Dronning Maud Land margin: Site 692, 301a-303a, 304a
 Dronning Maud Land margin: Site 693, 345a-349a, 350a
 Maud Rise: Site 689, 99a-101a
 source of, 952b, 953b
South Orkney microcontinent: Site 695, 539a-540a
South Orkney microcontinent: Site 696, 622a-624a
Weddell Basin: Site 694, 463a, 466a
- East Antarctic ice sheet. *See* Antarctic margin E; Antarctica E
- Eocene/Oligocene boundary
isotopic shift at, 842b
Maud Rise: Site 689, 903b
Weddell Sea, climate cooling, 950b
- Eotvos Basin
reflectors
 strength, 35b
 two-way travertime, 34b
- Ethane concentrations
Dronning Maud Land margin: Site 692, depth and, 313a
Dronning Maud Land margin: Site 693, 377a
 depth and, 382a
 retention times, 383a
Jane Basin, depth and, 738a
Maud Rise: Site 689, depth and, 131a
Maud Rise: Site 690, depth and, 238a
South Orkney microcontinent: Site 695, depth and, 563a-564a
- South Orkney microcontinent: Site 696, 650a
 depth and, 652a-653a
- Weddell Basin: Site 694, 485a
 depth and, 487a
- Explora Escarpment
morphology, 39b-40b
structure, 39b, 40b, 41b-42b, 47b
- Explora Wedge, reflectors, unconformity boundary, 44b
- Falkland Plateau
calcareous nannofossil assemblages, 948b, 952b
calcareous sequences, 939b
cooling event, Pliocene-Pleistocene, 538a
Maud Rise biogeography compared with, 938b
organic-rich facies compared with Weddell Sea facies, 206b-207b
- Feldspar
Dronning Maud Land margin: Site 692
 dating, 300a
 depth and, 75b
 X-ray diffractogram, 79b
- Dronning Maud Land margin: Site 693
 dating, 335a, 336a
 depth and, 335a, 336a
- Maud Rise: Site 689, zeolite replaced by, 10b
Maud Rise: Site 690
 microprobe analysis, 7b
 zeolite replaced by, 10b
- South Orkney microcontinent: Site 695, dating, 532a
- South Orkney microcontinent: Site 696
 depth and, 616a
 X-ray diffractogram, 31b
- Weddell Basin: Site 694, depth and, 460a
- Filchner/Ronne Ice Shelf, Pliocene expansion, 957b
- Fluorapatite, Dronning Maud Land margin: Site 692, X-ray diffractogram, 79b
- Gas hydrates, South Orkney microcontinent: Site 695, 33b-35b
- Gauss/Matuyama chron boundary, diatom floras, 806b
- Geochemistry
inorganic
Dronning Maud Land margin: Site 693, 372a-377a, 380a
Jane Basin, 729a-730a, 735a-737a
Maud Rise: Site 689, 128a-130a
Maud Rise: Site 690, 7b-8b, 199a-200a, 227a, 230a-231a
 alkali basalt, 201a-202a
 major element data, 201a, 203a
 trace element data, 201a, 203a
- South Orkney microcontinent: Site 695, 559a-562a
- South Orkney microcontinent: Site 696, 643a, 646a-651a
- Weddell Basin: Site 694, 478a, 481a, 484a-485a, 486a
- Weddell Sea, 52b
- See also* Alkalinity; Calcium concentrations; Chlorinity; Magnesium concentrations; pH, Phosphate concentrations; Potassium concentrations; Salinity, Silica; Sulfate concentrations
- organic
Dronning Maud Land margin: Site 692, 199b-207b, 312a-315a, 316a
- Dronning Maud Land margin: Site 693, 199b-207b, 377a-378a, 381a-383a
- Jane Basin, 736a, 738a-740a
- Maud Rise: Site 689, 9b, 129a, 131a
- Maud Rise: Site 690, 9b, 231a, 234a-236a, 238a
- South Orkney microcontinent: Site 695, 563a-565a
- South Orkney microcontinent: Site 696, 650a, 652a-655a
- Weddell Basin: Site 694, 199b-207b, 485a, 487a-488a
- See also* Ethane concentrations; Methane concentrations; Propane concentrations; Rock-Eval procedure, specific site
- Geochemistry, organic, Weddell Sea, Neogene climate indicators, 881b
- Glacial-interglacial cycles, Dronning Maud Land margin: Site 693, 127b
- Glacial-interglacial cycles, sedimentation patterns, 128b
- Glaciation
Eucampia antarctica abundances and, 806b
organic matter contents as signals of, 881b
sea-ice build-up as proxy for, 808b
- Glass, volcanic, Dronning Maud Land margin: Site 693, 78b
- Glauconite
Dronning Maud Land margin: Site 692, depth and, 76b
- Dronning Maud Land margin: Site 693
 dating, 335a
 depth and, 84b, 335a
- South Orkney microcontinent: Site 696
 composition, 622a
 dating, 61b
 deposition, 658a
 depth and, 616a
 thickness, 615a
- Gondwanaland, reconstructions, 342b-344b
- Gough Island, incompatible elements, mantle values, 9b
- Halite
South Orkney microcontinent: Site 696
 peak areas and heights, 29b
 X-ray diffractogram, 31b
- Hatteras Formation, "black shales," 194b
- Heat flow
Maud Rise, 17b, 24b
- South Orkney microcontinent: Site 695, 17b-20b, 25b
- South Orkney microcontinent: Site 696, 20b-21b
- Hydrocarbons
aliphatic, Weddell Sea, 201b-202b, 203b
Antarctica, organic matter sources, 199b
aromatic, Dronning Maud Land margin: Site 692, 202b
- isoprenoid, Weddell Sea, 201b, 206b
- steroid, Weddell Sea, 201b, 205b
- terpenoid, Weddell Sea, 201b-202b, 204b
- Ice rafting
Antarctic Ocean, Oligocene evidence for, 953b
Antarctica E, Miocene, 955b
- Dronning Maud Land margin: Site 692, 301a-303a
- Dronning Maud Land margin: Site 693, 345a-349a
 glaciation and, 954b
- Maud Rise, 99a-101a
Oligocene, 875b
- Ross Sea, Oligocene, 875b
- South Orkney microcontinent: Site 695, 539a-540a

- South Orkney microcontinent: Site 696, 622a–624a
 Weddell Basin: Site 694, 466a
- Iddingsite
 Maud Rise: Site 689, olivine replaced by, 10b
 Maud Rise: Site 690
 microprobe analysis, 7b
 olivine replaced by, 10b, 12b–14b
 orthopyroxene replaced by, 12b
- Illite
 Dronning Maud Land margin, abundance, 54b
 Dronning Maud Land margin: Site 693
 dating, 55b, 344a
 weathering conditions suggested by, 124b
- Jane Basin
 dating, 62b, 63b
 TEM photomicrograph, 231b, 235b
 Maud Rise: Site 689, dating, 57b, 99a
 Maud Rise: Site 690, dating, 58b, 197a
 South Orkney microcontinent: Site 695
 chemical weathering and, 538a
 dating, 539a
 South Orkney microcontinent: Site 696
 dating, 61b, 623a
 peak areas and heights, 29b
 Weddell Basin: Site 694, dating, 59b
- Indian Ocean, sources of low-latitude warm saline intermediate waters, Miocene, 866b
- Indian Ocean S, sea-ice cover, diatom abundances and, 806b
- Interstitial waters, dissolved organic carbon, sources of, 173b–174b
- Interstitial-water chemistry
 Dronning Maud Land margin: Site 693
 data summary, 138b, 379a
 oxygen isotope distribution, 138b, 141b
 procedure, 136b–137b
 strontium isotope distribution, 150b
 Jane Basin
 data summary, 140b
 oxygen isotope distribution, 140b, 143b
 procedure, 137b
 strontium isotope distribution, 151b
 Maud Rise: Site 689
 data summary, 129a, 138b
 oxygen isotope distribution, 138b, 140b
 procedure, 135b–136b
 strontium isotope distribution, 149b
 Maud Rise: Site 690
 data summary, 138b, 236a
 oxygen isotope distribution, 138b, 141b
 procedure, 136b
 strontium isotope distribution, 150b
 strontium isotope distribution, 150b
 South Orkney microcontinent, oxygen isotope distribution, 139b
 South Orkney microcontinent: Site 695
 amino acids, 179b–186b
 data summary, 139b
 oxygen isotope distribution, 139b, 142b
 procedure, 137b
 strontium isotope distribution, 150b
 South Orkney microcontinent: Site 696, 650a
 data summary, 139b
 oxygen isotope distribution, 143b
 procedure, 137b
 strontium isotope distribution, 151b
 Weddell Basin: Site 694
 data summary, 139b, 485a
 oxygen isotope distribution, 139b, 142b
 procedure, 137b
 strontium isotope distribution, 150b
 Weddell Sea
 calculations, 139b–144b
 oxygen isotope distribution, 145b
- Iridium
 Maud Rise, 903b
 abundance, 161b–163b, 164b–168b
 at Cretaceous/Tertiary boundary, 829b, 833b
 bioturbation and, 161b–162b
 volcanism and, 161b–162b
 Maud Rise: Site 689, calculations, 162b
- Jane Basin
 age-depth plots, versus magnetostratigraphy, 935b
 alkalinity, 730a, 735a
 depth and, 140b, 143b, 736a–737a
 ammonia concentrations, depth and, 735a–737a
 Antarctic Bottom Water and, 803b
 bathymetry, 706a–707a
 calcium concentrations, depth and, 140b, 143b, 736a–737a
 chloride concentrations, 730a
 depth and, 736a–737a
 climatic evolution, 119b–120b
 consolidation
 data summary, 214b, 222b
 field curve, 221b
 deep-water circulation, 112b
 density, depth and, 716a
 dissolved organic carbon
 depth and, 173b
 pore water concentrations, 174b
 ethane concentrations, depth and, 738a
 grain-size analyses
 diatom content and, 117b–118b
 Sedigraph cumulative frequency curve, 116b
 interstitial-water chemistry
 data summary, 140b
 procedure, 137b
 kerogen concentrations, 736a, 738a
 vitrinite reflectance measurements, 192b–194b
 lithology, 710a
 dating, 95b
 depth and, 95b
 thickness and, 114b
 lithostratigraphy
 Pleistocene–Pliocene, 708a–709a, 738a
 Pliocene, 709a, 738a–740a
 location, 706a–707a
 magnesium concentrations, depth and, 140b, 143b, 736a–737a
 magnetic properties
 magnetic susceptibility, 727a, 732a–734a
 natural remanent magnetization, 726a, 728a–729a, 731a
 magnetostratigraphy, 726a–727a
 age identification and, 114b
 age-depth curve, 115b
 geomagnetic polarity time scale, 730a, 735a
 methane concentrations, depth and, 738a
 mineralogy, clay, 62b, 63b, 710a–712a, 715a
 oxygen isotope distribution, depth and, 140b, 143b
 paleoenvironment, 711a–712a
 pH, 730a
 depth and, 737a
 phosphate concentrations, depth and, 735a–737a
 Pliocene sea-ice cover, 803b, 808b
 porosity
 depth and, 226b, 716a–717a
 diatom relationship with, 220b
 TEM photomicrograph, 229b–232b, 234b–237b
 potassium concentrations, depth and, 140b, 143b, 736a–737a
 propane concentrations, depth and, 738a
 reflectors
- sedimentology suggested by, 113b
 strength, 716a–717a
 two-way traveltime, 113b
 Rock-Eval procedure, 736a, 738a–740a
 salinity, 730a
 depth and, 736a–737a
 sedimentation rates
 biostratigraphic data, 727a–729a, 735a, 740a–741a
 magnetostratigraphic data, 727a–729a, 740a
 seismic reflection profile, 113b, 708a, 721a
 seismic stratigraphy, 716a–717a
 P-wave velocity, 719a
 shear strength
 calculations, 713a–714a
 depth and, 716a, 718a
 strontium isotope distribution
 calculations, 153b, 156b
 depth and, 151b
 sediment age versus, 152b
 sulfate concentrations, 730a, 735a
 depth and, 140b, 143b, 736a–737a
 pore water concentrations, 175b
 thermal conductivity, 715a
 depth and, 719a–720a
 velocity
 calculations, 714a
 depth and, 719a
 volcanic ash
 color, 708a–710a
 depth and, 710a
 water content, depth and, 716a
See also Weddell Sea
- Kaolinite
 Dronning Maud Land margin: Site 693, dating, 55b, 344a
 East Antarctic climate indicator, Paleocene, 945b
 Jane Basin
 dating, 62b, 63b
 TEM photomicrograph, 231b
 Maud Rise: Site 689, dating, 57b, 99a
 Maud Rise: Site 690, dating, 197a
 South Orkney microcontinent: Site 695, dating, 539a
 South Orkney microcontinent: Site 696, dating, 61b, 623a
 Weddell Basin: Site 694, dating, 59b
 Kerguelen Plateau, ice-raftered debris, Oligocene, 953b
- Kerogen concentrations
 Dronning Maud Land margin, sulfur isotope ratios, 195b–196b
 Dronning Maud Land margin: Site 692, 189b–190b, 314a–315a
 carbon isotope ratios, 195b–196b
 data summary, 191b
 vitrinite reflectance measurements, 192b–193b
 Dronning Maud Land margin: Site 693, 190b–193b, 377a
 vitrinite reflectance measurements, 192b–193b
 Jane Basin, 736a, 738a
 vitrinite reflectance measurements, 192b–194b
 South Orkney microcontinent: Site 695
 depth and, 563a–564a
 temperature measurements, 563a–564a, 565a
 vitrinite reflectance measurements, 192b–194b
 South Orkney microcontinent: Site 696
 temperature measurements, 655a
 vitrinite reflectance measurements, 192b–194b
 Weddell Basin: Site 694, Site 692 compared with, 488a
 Weddell Sea, site comparisons, 194b–197b

SUBJECT INDEX

Labrador Sea, carbon isotope changes, 881b-882b
 Laterite, Maud Rise, 945b
 Layer II, duration of seawater interactions, 139b
 Limestone
 Dronning Maud Land margin: Site 692, depth and, 74b
 Dronning Maud Land margin: Site 693
 color, 78b
 pyrite with, 83b
 Lord Howe Rise
 clay, volcanic alteration, 53b
 strontium, 154b
 Magnesium concentrations
 Dronning Maud Land margin: Site 693, 375a
 depth and, 138b, 141b, 380a
 Jane Basin, depth and, 140b, 143b, 736a-737a
 Maud Rise: Site 689, depth and, 129a-130a, 138b, 140b
 Maud Rise: Site 690, 230a
 depth and, 138b, 141b, 237a
 South Orkney microcontinent: Site 695, depth and, 139b, 142b, 561a-562a
 South Orkney microcontinent: Site 696, 648a-649a
 depth and, 139b, 143b, 650a-651a
 Weddell Basin: Site 694, 484a
 depth and, 139b, 142b, 485a-486a
 Magnetic properties
 Dronning Maud Land margin: Site 692, 311a-312a
 demagnetization, 312a
 natural remanent magnetization, 312a-313a
 Dronning Maud Land margin: Site 693
 demagnetization, 375a
 natural remanent magnetization, 368a-369a, 370a-371a, 373a
 Jane Basin
 magnetic susceptibility, 727a, 732a-734a
 natural remanent magnetization, 726a, 728a-729a, 731a
 Maud Rise: Site 689, natural remanent magnetization, 121a-123a
 Maud Rise: Site 690, natural remanent magnetization, 225a-227a, 230a
 South Orkney microcontinent: Site 695, natural remanent magnetization, 554a-557a
 South Orkney microcontinent: Site 696, natural remanent magnetization, 641a-642a, 644a-645a
 Weddell Basin: Site 694
 magnetic susceptibility, 477a, 482a-483a
 natural remanent magnetization, 477a-480a
 Magnetite, Maud Rise: Site 690, shape, 13b
 Magnetostратigraphy
 Dronning Maud Land margin: Site 693
 geomagnetic polarity time scale, 372a, 376a, 378a
 glacial-interglacial cycles correlated with, 127b
 natural remanent magnetization, 384a
 Jane Basin, 726a-727a, 740a
 age identification and, 114b
 age-depth curve, 115b
 geomagnetic polarity time scale, 730a, 735a
 Maud Rise, biostratigraphic correlations, 915b-936b
 Maud Rise: Site 689
 age-depth curve, 122a-127a
 geomagnetic polarity time scale, 262b
 hiatus, 123a, 128a
 magnetostratigraphy, 268b-273b
 natural remanent magnetization, 256b-259b, 264b-267b, 281b-284b, 295b-304b

Maud Rise: Site 690
 age-depth curve, 225a-227a
 geomagnetic polarity time scale, 225a, 228a-229a, 232a-234a, 262b, 290b-292b
 magnetostratigraphy, 274b-280b
 natural remanent magnetization, 256b-259b, 267b, 286b-289b, 305b-315b
 South Orkney microcontinent: Site 695, 555a
 geomagnetic polarity time scale, 558a-559a
 South Orkney microcontinent: Site 696,
 geomagnetic polarity time scale, 646a-649a
 Weddell Basin: Site 694, 477a
 geomagnetic polarity time scale, 481a, 484a
 Weddell Sea
 biostratigraphic correlations, 915b-936b
 rounded end members based on, 101b
 Manganese
 Dronning Maud Land margin: Site 693, dissolved, 172b
 Jane Basin, dissolved, 173b
 Maud Rise, dissolved, 171b
 South Orkney microcontinent, dissolved, 173b
 Weddell Basin: Site 694, dissolved, 172b
 Maud Rise
 age-depth plots, 907b, 908b
 versus magnetostratigraphy, 922b, 924b
 Antarctic Bottom Water (AABW), 865b-866b
 benthic foraminiferal extinctions, 876b
 carbon isotopes, depletion, Maestrichtian, 821b
 Cretaceous temperature changes, environmental implications, 842b
 Cretaceous/Tertiary boundary, stable isotope values, 829b, 831b-833b
 Falkland Plateau biogeography compared with, 938b
 hiatuses, 874b
 location, 6a
 Maestrichtian water temperatures, isotope ratios and, 813b
 paleoenvironment
 cooling, 56b
 deep-water circulation, 56b
 paleotemperatures, 821b
 Maestrichtian, 822b
 sea-ice absence in Paleocene-Eocene, 863b
 stable isotopes, reversals, 877b
 stratigraphy,
 biostratigraphic-magnetostratigraphic correlations, 901b-909b
 Maud Rise: Site 689
 alkalinity, depth and, 129a-130a, 138b, 140b
 basement, dating, 90a
 bathymetry, 91a
 calcium carbonate content, depth and, 103a-104a, 975b-976b, 980b-981b, 984b
 calcium concentrations, depth and, 129a-130a, 138b, 140b
 carbon isotope stratigraphy
 depth and, 986b-987b
 time and, 989b
 chlorinity, depth and, 128a-130a
 consolidation
 data summary, 214b, 222b
 field curve, 215b
 density
 depth and, 102a-103a, 105a
 large variations in, 101a
 dissolved organic carbon
 depth and, 171b
 pore water concentrations, 174b
 downhole temperature measurements, 131a-132a
 ethane concentrations, depth and, 131a
 glacial history, 90a
 heat flow measurements, 17b
 depth and, 24b
 time and, 19b
 incompatible elements, mantle values, 9b
 interstitial-water chemistry
 data summary, 129a, 138b
 procedure, 135b-136b
 lithostratigraphy
 boundaries, 106a
 Campanian-Eocene, 97a-98a
 Eocene-Miocene, 95a-97a
 Miocene-Quaternary, 94a-95a, 96a
 Site 690 compared with, 193a-194a
 location, 6b, 91a
 magnesium concentrations, depth and, 129a-130a, 138b, 140b
 magnetic properties, natural remanent magnetization, 121a-123a
 magnetostratigraphy
 age-depth curve, 122a-127a
 geomagnetic polarity time scale, 262b
 hiatus, 123a, 128a
 magnetostratigraphy, 268b-273b
 natural remanent magnetization, 256b-259b, 264b-267b, 281b-284b, 295b-304b
 methane concentrations, depth and, 131a
 mineralogy
 clay, 57b, 67b
 dating, 99a
 oxygen isotope distribution, depth and, 138b, 140b
 paleoenvironment, 133a-134a
 ice-rafterd dropstones, 99a-101a
 pH, depth and, 128a-130a
 phosphate concentrations, depth and, 129a-130a
 porosity
 depth and, 102a-103a, 213b
 diatom relationship with, 213b
 large variations in, 101a
 potassium concentrations, depth and, 129a-130a, 138b, 140b
 propane concentrations, depth and, 131a
 reflectors
 dating, 108a
 depth and, 107a-108a
 Rock-Eval procedure, 131a-132a
 salinity, depth and, 128a-129a
 sedimentation rates, 121a-122a
 biostratigraphic data, 128a
 sedimentology
 Campanian-Eocene, 97a-98a
 Eocene-Miocene, 95a-97a
 Miocene-Quaternary, 94a-95a, 96a
 seismic reflection profile, 92a-93a
 seismic stratigraphy, *P*-wave velocity, 107a
 shear strength, 103a-104a
 depth and, 106a, 214b
 vertical effective stress relationship with, 214b
 strontium isotope distribution
 calculations, 151b, 153b-155b
 depth and, 149b
 sediment age versus, 151b
 sulfate concentrations
 depth and, 129a-130a, 138b, 140b
 pore water concentrations, 175b
 thermal conductivity, 104a-105a
 calculations, 21b
 depth and, 23b, 25b, 106a
 water content versus, 22b
 total organic carbon, depth and, 190b
 velocity, 101a-102a
 depth and, 104a-105a
 water content

- depth and, 102a-103a
large variations, 101a
- Maud Rise: Site 690
alkalinity, 230a
depth and, 138b, 141b, 237a
- ammonia concentrations, 230a
depth and, 237a
- basement
alteration, 198a-199a
basalt stratigraphy, 198a-199a
geochemistry, 199a-200a, 201a-203a
petrography, 198a-199a, 200a
- bathymetry, 91a
- calcium carbonate content, depth and, 207a-208a, 977b-979b, 982b-983b
- calcium concentrations, 230a
depth and, 138b, 141b, 237a
- carbon isotope stratigraphy
depth and, 987b-989b
time and, 989b
- chloride concentrations, depth and, 237a
- chlorinity, 230a
- consolidation
data summary, 214b, 222b
field curve, 216b
- density, depth and, 205a-206a
- dissolved organic carbon
depth and, 172b
pore water concentrations, 174b
- downhole temperature measurements, 231a, 239a
- ethane concentrations, depth and, 238a
- heat flow measurements, 17b
depth and, 24b
time and, 19b
- incompatible elements, mantle values, 9b
- interstitial-water chemistry
data summary, 138b, 236a
procedure, 136b
- lithostratigraphy, 186a-187a
boundaries, 188a, 196a
Campanian-Maestrichtian, 192a-193a
Eocene-Paleocene, 189a
glacial marine dropstones, 192a
Maestrichtian-Paleocene, 189a-192a
Miocene-Quaternary, 187a-188a
Oligocene-Pliocene, 188a-189a
Site 689 compared with, 193a-194a
summary, 189a
- location, 6b, 91a, 185a
- magnesium concentrations, 230a
depth and, 138b, 141b, 237a
- magnetic properties, natural remanent magnetization, 225a-227a, 230a
- magnetotriatigraphy
age-depth curve, 225a-227a
geomagnetic polarity time scale, 225a, 228a-229a, 232a-234a, 262b, 290b-292b
magnetochronology, 274b-280b
natural remanent magnetization, 256b-259b, 267b, 286b-289b, 305b-315b
- major element data, 8b
- methane concentrations, depth and, 238a
- mineralogy
clay, 58b
dating, 197a
- oxygen isotope distribution, depth and, 138b, 141b
- paleoenvironment, 195a-196a, 198a, 236a, 238-240a
- pH, 230a
depth and, 237a
- phosphate concentrations, 230a
- porosity
- depth and, 205a, 207a, 215b
diatom relationship with, 215b
- potassium concentrations, 231a
depth and, 138b, 141b, 237a
- propane concentrations, depth and, 238a
- reflectors
dating, 207a-208a, 216a
depth and, 207a-208a
- Rock-Eval procedure, 231a, 238a
- salinity, 230a
- sedimentation rates, biostratigraphic data, 225a, 235a
- seismic reflection profile, 186a
- seismic stratigraphy, 205a-208a
P-wave velocity, 209a, 215a
- shear strength, 204a
depth and, 212a, 216b
vertical effective stress relationship with, 216b
- strontium isotope distribution
calculations, 151b, 153b-155b
depth and, 150b
sediment age versus, 151b
- sulfate concentrations, 230a
depth and, 138b, 141b, 237a
pore water concentrations, 175b
- thermal conductivity, 204a
calculations, 21b
depth and, 23b, 25b, 212a-213a
water content versus, 22b
- total organic carbon, depth and, 190b
- trace element data, 8b
- velocity, 202a, 204a
depth and, 209a-211a
- volcanic glass, depth and, 190a-191a
- water content, depth and, 205a-206a
- Methane concentrations
Dronning Maud Land margin: Site 692, depth and, 313a
- Dronning Maud Land margin: Site 693, 377a
depth and, 382a
retention times, 383a
- Jane Basin, depth and, 738a
- Maud Rise: Site 689, depth and, 131a
- Maud Rise: Site 690, depth and, 238a
- South Orkney microcontinent: Site 695
calculations, 35b
depth and, 563a-564a
- South Orkney microcontinent: Site 696, 650a
depth and, 652a-653a
- Weddell Basin: Site 694, 485a
depth and, 487a
- Mexico, Gulf of, carbon sources, 881b
- Mica
Dronning Maud Land margin: Site 692, depth and, 75b
- Jane Basin
dating, 711a
depth and, 711a
- Maud Rise: Site 690, depth and, 190a-191a
- South Orkney microcontinent: Site 695, dating, 532a
- South Orkney microcontinent: Site 696, depth and, 616a
- Weddell Basin: Site 694, depth and, 460a
- Milankovitch cycles
Dronning Maud Land margin: Site 693, 246b-248b
logging detection of, 245b-246b
- Mineralogy
clay, Maud Rise: Site 689, 57b, 67b
- Dronning Maud Land margin, clay, 57b
- Dronning Maud Land margin: Site 692, 75b
clay, 301a, 303a
depth and, 76b
- Dronning Maud Land margin: Site 693
clay, 68b, 342a, 344a
depth and, 84b
- Jane Basin, clay, 62b, 63b, 710a-712a, 715a
- Maud Rise: Site 689
basalts, 6b-7b
See also specific types
dating, 99a
- Maud Rise: Site 690
basalts, 6b-7b
See also specific types
clay, 58b
dating, 197a
- South Orkney microcontinent: Site 695, clay, 537a-539a
- South Orkney microcontinent: Site 696, clay, 61b, 620a-623a
- Weddell Basin: Site 694, clay, 59b, 461a-462a, 465a
- Weddell Sea, clay, 51b-52b
- Miocene/Pliocene boundary
Maud Rise, hiatus, 917b
Weddell Sea, climate change, 956b
- Mn/Ca ratio, Maud Rise, 833b
- Mozambique Basin, opening, 343b, 344b
- Mud
Dronning Maud Land margin: Site 692, depth and, 74b
- Jane Basin, dating, 62b, 63b
- Weddell Basin: Site 694, dating, 59b
- Mud, clayey
Jane Basin
dating, 710a
depth and, 710a
- South Orkney microcontinent: Site 695, color, 534a
- Mud, silty
Dronning Maud Land margin: Site 691, dating, 299a
- Dronning Maud Land margin: Site 692, dating, 299a
- Jane Basin
color, 714a
dating, 710a
depth and, 710a
- South Orkney microcontinent: Site 695, color, 534a
- South Orkney microcontinent: Site 696
color, 617a
dating, 614a-615a
depth and, 614a
lithified, 619a
thickness, 615a
- Mudstone, Maud Rise: Site 690, dating, 189a
- Niger delta, organic carbon sources, 881b
- Nitrogen
recycling, Pleistocene-present, 889b
terrestrial versus marine sources, 882b
See also C/N ratio
- Nitrogen fixation, 882b
- Nitrogen isotopes
as indicators of organic matter sources, 882b
Weddell Sea, values, 884b-889b, 890b, 891b, 894b-897b
- North Atlantic Deep Water
production of, 952b
silica source, 127b-128b
- Northern Hemisphere
glaciation
Pliocene, 803b, 957b
versus Southern Hemisphere, 808b-809b

SUBJECT INDEX

- Oceanus
Cenozoic circulation patterns, 875b, 876b
See also Circulation, oceanic; Proto-oceanus
- Olivine
Maud Rise: Site 689, iddingsite replacing, 10b
Maud Rise: Site 690
calcite replacing, 14b
iddingsite replacing, 10b, 12b-14b
microprobe analysis, 7b
shape, 200a
- South Orkney microcontinent: Site 696
depth and, 30b
peak areas and heights, 29b
- Ontong-Java Plateau, strontium, 154b
- Ooze. *See also* Paleontological Index
- Ooze, calcareous, Maud Rise: Site 689, carbonate compensation depth, 90a
- Ooze, carbonate, South Orkney microcontinent: Site 695, color, 536a
- Ooze, diatom
Dronning Maud Land margin: Site 693, composition, 342a
Jane Basin, dating, 62b, 713a
Maud Rise: Site 689
burrow structures, 96a
color, 97a
dating, 57b, 95a
density influenced by, 101a
glacial marine dropstone, 96a
porosity influenced by, 101a
- Maud Rise: Site 690
dating, 58b, 189a
foraminiferal transition with, 192a
- South Orkney microcontinent: Site 695
color, 536a
depth and, 535a
porosity, 546a
- South Orkney microcontinent: Site 696
dating, 61b, 614a-615a
depth and, 614a
thickness, 615a
- Ooze, diatom-nannofossil
Dronning Maud Land margin: Site 693
dating, 334a
depth and, 334a
- Ooze, diatom-radiolarian, Dronning Maud Land margin: Site 692, depth and, 74b
- Ooze, diatomaceous
Dronning Maud Land margin, consolidation characteristics, 215b-217b
Maud Rise, consolidation characteristics, 213b-215b, 216b
- South Orkney microcontinent, consolidation characteristics, 217b-221b
- Ooze, foraminiferal
Maud Rise: Site 690
dating, 58b
diatom transition with, 192a
- Ooze, nannofossil
Dronning Maud Land margin: Site 693, composition, 340a
Maud Rise: Site 689, dating, 57b, 95a
Maud Rise: Site 690
color, 193a, 194a
dating, 58b, 189a
- Ooze, nannofossil chalk
Maud Rise: Site 690
bioturbated ash in, 194a-195a
dating, 189a
- Ooze, nannofossil foraminifers, Maud Rise: Site 690, dating, 189a
- Ooze, radiolarian
Maud Rise: Site 689
dating, 95a
- density influenced by, 101a
porosity influenced by, 101a
- Ooze, siliceous
Dronning Maud Land margin: Site 692, depth and, 74b
Maud Rise: Site 689, dating, 95a
Maud Rise: Site 690, dating, 189a
- Ooze, siliceous nannofossil, Maud Rise: Site 690, dating, 58b
- Opal
abundances, sea-ice production and, 803b
biogenic, diatom productivity and, 804b
Dronning Maud Land margin: Site 693, dating, 55b
Maud Rise: Site 690, dating, 58b
- South Orkney microcontinent: Site 695, siliceous transformation, 34b
- South Orkney microcontinent: Site 696
bladed, 37b
dating, 61b
depth and, 30b
peak areas and heights, 29b
siliceous transformation, 34b
X-ray diffractogram, 31b
- Southern Ocean, abundances, 807b
- Orthopyroxene
Maud Rise: Site 690
iddingsite replacing, 12b
microprobe analysis, 7b
- Oxygen isotope ratio
equilibrium in foraminiferal test secretion, 849b
Maud Rise, 836b-839b
foraminiferal data, 814b, 815b, 818b, 819b, 820b, 831b-832b, 835b
Sites 689/690 compared, 946b, 948b
water temperatures and, 813b
- Maud Rise: Hole 690, 942b
- Oxygen isotopes
calcite recrystallization and, 815b-816b
influence of salinity on values, 876b
- Maud Rise, 860b-861b
reversed Paleogene gradient, 949b-950b
Sites 689/690 comparisons, 867b-868b, 868b-869b, 874b
surface-to-deep-water gradients, 870b-871b, 874b-875b
surface-water temperatures and, 945b
- Maud Rise: Site 689, 852b-854b, 856b
Maud Rise: Site 690, 850b, 853b-854b, 855b
offsets in foraminiferal tests, 834b-836b
Paleogene ocean temperatures reflected in, 873b
relation to polar ice accumulations, 939b-940b
- Pacific Ocean N
carbon isotope values, 873b
foraminifers production in subarctic, 842b, 843b
- Pacific Ocean, tropical, Maestrichtian cooling, 821b
- Paleocene/Eocene boundary
Antarctic Ocean
surface-water temperatures, 859b
temperature excursions and foraminifers extinctions, 849b
warming interval, 877b
isotopic shift at, 842b
- Maud Rise: Site 690, oxygen isotope temperature excursion, 851b-852b
microfossil extinctions, 946b-947b
Weddell Sea, warm interval, 937b, 945b
- Paleoenvironment. *See* Paleontological Index; particular locations
- Pedernales delta, organic carbon sources, 881b
- Perlite
- Maud Rise: Site 690, 12b
glass with, 199a
- Permeability
determination of, 213b
Weddell Sea, calculations, 213b
- Peru continental margin, calcium carbonate contents, 76b
- Perylene, Weddell Sea, 206b
- Petroleum-generating potential, Dronning Maud Land margin: Site 692, 196b
- pH
Dronning Maud Land margin: Site 693, 374a
depth and, 380a
- Jane Basin, 730a
depth and, 737a
- Maud Rise: Site 689, depth and, 128a-130a
- Maud Rise: Site 690, 230a
depth and, 237a
- South Orkney microcontinent: Site 695, 560a
South Orkney microcontinent: Site 696, 646a-647a
depth and, 650a-651a
- Phosphate concentrations
Dronning Maud Land margin: Site 693, 375a
depth and, 380a
- Jane Basin, depth and, 735a-737a
- Maud Rise: Site 689, depth and, 129a-130a
- Maud Rise: Site 690, 230a
- South Orkney microcontinent: Site 695, depth and, 561a-562a
- South Orkney microcontinent: Site 696, 647a-648a
depth and, 650a-651a
- Weddell Basin: Site 694, 481a, 485a-486a
- Phosphorite, depositional environment, 76b, 77b
- Plagioclase
Maud Rise: Site 689, absence, 9b-10b
Maud Rise: Site 690, absence, 9b-10b
- South Orkney microcontinent: Site 696
depth and, 30b
peak areas and heights, 29b
X-ray diffractogram, 31b
- Planolite, South Orkney microcontinent: Site 696, size, 621a
- Polar oceans, water temperatures, global climate and, 813b
- Porosity
Dronning Maud Land margin: Site 692, depth and, 304a-305a
- Dronning Maud Land margin: Site 693, 240b, 242b-244b
depth and, 215b-217b, 243b, 351a-353a
diatom relationship with, 215b-217b
mineralogical and paleoceanographical factors, 248b-250b
- Jane Basin
depth and, 226b, 716a-717a
diatom relationship with, 220b
- TEM photomicrographs, 229b-232b, 234b-237b
- Maud Rise: Site 689
depth and, 102a-103a, 213b
diatom relationship with, 213b
large variations in, 101a
- Maud Rise: Site 690
depth and, 205a, 207a, 215b
diatom relationship with, 215b
sediment components affecting, 244b
- South Orkney microcontinent: Site 695
depth and, 180b, 541a-543a
diatom relationship with, 218b
- South Orkney microcontinent: Site 696
depth and, 30b, 625a-626a

- diatom relationship with, 219b
range, 624a, 626a
Weddell Basin: Site 694
depth and, 218b, 466a-468a
diatom relationship with, 218b
- Potassium concentrations
Dronning Maud Land margin: Site 693, 375a-376a
depth and, 138b, 141b, 380a
Jane Basin, depth and, 140b, 143b, 736a-737a
Maud Rise: Site 689, depth and, 129a-130a, 138b, 140b
Maud Rise: Site 690, 231a
depth and, 138b, 141b, 237a
South Orkney microcontinent: Site 695, depth and, 139b, 142b, 561a-562a
South Orkney microcontinent: Site 696, depth and, 139b, 143b, 650a-651a
Weddell Basin: Site 694, 484a-485a
depth and, 139b, 142b, 485a-486a
- Powell Basin, formation, 658a
propane concentrations
Dronning Maud Land margin: Site 692, depth and, 313a
Dronning Maud Land margin: Site 693
depth and, 382a
retention times, 383a
Jane Basin, depth and, 738a
Maud Rise: Site 689, depth and, 131a
Maud Rise: Site 690, depth and, 238a
South Orkney microcontinent: Site 695, depth and, 563a-564a
South Orkney microcontinent: Site 696, 650a
depth and, 652a-653a
Weddell Basin: Site 694, depth and, 487a
Proteus, 950b
Cenozoic circulation patterns, 875b, 876b
Proto-AABW, formation of, 876b, 877b
Proto-oceanus, 950b
Cenozoic circulation patterns, 875b, 876b
Prydz Bay, glacial diamictites, 953b, 954b
Pyrite
Dronning Maud Land margin: Site 692
dating, 300a
depth and, 75b, 76b
X-ray diffractogram, 79b
Dronning Maud Land margin: Site 693, depth and, 84b
South Orkney microcontinent: Site 696
depth and, 30b
peak areas and heights, 29b
X-ray diffractogram, 31b
- Quartz
Dronning Maud Land margin: Site 692
dating, 300a
depth and, 75b, 76b
X-ray diffractogram, 79b
Dronning Maud Land margin: Site 693
dating, 55b, 335a, 336a
depth and, 84b, 335a, 336a
Jane Basin
dating, 62b, 63b, 711a
depth and, 711a
Maud Rise: Site 690
dating, 58b
depth and, 190a-191a
South Orkney microcontinent: Site 695
dating, 532a
size, 538a
South Orkney microcontinent: Site 696
dating, 61b
depth and, 30b, 616a
- peak areas and heights, 29b
X-ray diffractogram, 31b
Weddell Basin: Site 694
dating, 59b
depth and, 460a
Weddell Sea, end-member identification, 107b-109b
- Recrystallization, carbonate, silica amounts and, 154b
Reflectors
Bouguer Basin, two-way travelttime, 34b
Dronning Maud Land margin: Site 692, dating, 316a-317a
Dronning Maud Land margin: Site 693
dating, 357a-358a, 385a
depth and, 358a
ice shelf formation correlated with, 125b
two-way travelttime, 45b
unconformity boundary, 44b-45b
- Eotvos Basin
strength, 35b
two-way travelttime, 34b
Explora Wedge, unconformity boundary, 44b
Jane Basin
sedimentology suggested by, 113b
strength, 716a-717a
two-way travelttime, 113b
Maud Rise: Site 689
dating, 108a
depth and, 107a-108a
Maud Rise: Site 690
dating, 207a-208a, 216a
depth and, 207a-208a
South Orkney microcontinent: Site 695
bottom simulating, 17b, 22b-23b
boundaries, 33b
depth and, 542a-543a, 550a
diagenesis and, 32b-33b
gas hydrate corresponding with, 33b-35b
South Orkney microcontinent: Site 696
dating, 631a
depth and, 631a
diagenesis and, 32b
strength, 631a
unconformity boundary, 27b-28b
Weddell Basin: Site 694
dating, 470a-471a
strength, 470a-471a
Rio Grande Rise, Paleocene nannofossil assemblages, 945b
Ross Ice Shelf, Pliocene expansion, 957b
Ross Sea
ice rafting, 875b
ice-rafterd detritus, Miocene, 125b
- Salinity
deep-water circulation and, in Paleogene, 877b
Dronning Maud Land margin: Site 693, 374a
effect on isotopic composition, 873b
Jane Basin, 730a
depth and, 736a-737a
Maud Rise: Site 689, depth and, 128a-129a
Maud Rise: Site 690, 230a
role in deep ocean circulation, 865b
South Orkney microcontinent: Site 695, depth and, 561a
South Orkney microcontinent: Site 696, 643a, 646a
depth and, 650a
temperature and, influence on oxygen isotopic composition, 876b
Weddell Basin: Site 694, depth and, 481a, 485a
- Sand
glauconite, Antarctic ice-sheet development and, 953b
Sand, quartz
Fourier grain shape analyses, 89b-90b, 96b
Weddell Sea, types of, 103b
Scotia Sea
deep-water circulation, 112b
deepening and widening, 65b
Sedimentation rates
organic matter preservation and, 889b
Weddell Sea, 925b
See also particular locations
Seismic stratigraphy
Bouguer Basin, *P*- wave velocity, 35b
Dronning Maud Land margin: Site 691, 305a
Dronning Maud Land margin: Site 692, 305a
lithology correlated with, 46b
P- wave velocity, 304a
sequences, 317a
Dronning Maud Land margin: Site 693, 353a-356a
Cretaceous-Oligocene hiatus, 357a-358a
lithology correlated with, 46b
Miocene hiatus, 356a-357a
P- wave velocity, 44b, 243b, 353a
Jane Basin, 716a-717a
P- wave velocity, 719a
Maud Rise: Site 689, *P*- wave velocity, 107a
Maud Rise: Site 690, 205a-208a
P- wave velocity, 209a, 215a
South Orkney microcontinent: Site 695, 542a-546a
P- wave velocity, 544a-545a, 547a, 550a
South Orkney microcontinent: Site 696, 628a-631a
P- wave velocity, 30b, 32b, 628a-629a, 631a
Weddell Basin: Site 694, 469a-471a
P- wave velocity, 470a-471a, 472a
Wegener Canyon, lithology correlated with, 44b, 46b
- Seymour Island
calcareous sequences, 939b
carbon isotope values, 821b-822b
Cretaceous/Tertiary boundary, microfossil extinctions, 944b
isotope ratios, 813b
oxygen isotope values, 821b, 941b
paleotemperatures, Maestrichtian, 821b
Shatsky Rise, oxygen isotope values, 821b
Shear strength
Dronning Maud Land margin: Site 693, 351a-352a
depth and, 217b, 355a
vertical effective stress relationship with, 217b
Jane Basin
calculations, 713a-714a
depth and, 716a, 718a
Maud Rise: Site 689, 103a-104a

SUBJECT INDEX

- depth and, 106a, 214b
vertical effective stress relationship with, 214b
- Maud Rise: Site 690, 204a
depth and, 212a, 216b
vertical effective stress relationship with, 216b
- South Orkney microcontinent: Site 695, 541a
depth and, 543a–544a
vertical effective stress relationship with, 218b
- South Orkney microcontinent: Site 696, 626a
depth and, 627a
vertical effective stress relationship with, 220b
- Weddell Basin: Site 694
depth and, 218b, 467a–468a
vertical effective stress relationship with, 218b
- Silica**
Dronning Maud Land margin: Site 693, 376a–377a
depth and, 380a
- Jane Basin, depth and, 736a–737a
- Maud Rise: Site 689, depth and, 129a–130a
- Maud Rise: Site 690, 231a
depth and, 237a
- North Atlantic Deep Water, 127b–128b
- South Orkney microcontinent: Site 695, depth and, 561a–562a
- South Orkney microcontinent: Site 696, 649a–650a
depth and, 650a–652a
- Weddell Basin: Site 694
depth and, 486a
reversed weathering, 485a
- Silt**
Dronning Maud Land margin: Site 693
dating, 337a
depth and, 337a
- Jane Basin
dating, 713a, 714a
size, 712a
- South Orkney microcontinent: Site 695, color, 537a
- Weddell Basin: Site 694
composition, 457a
lamination, 463a
location, 456a, 459a, 462a
- Smectite**
Dronning Maud Land margin, abundance, 54b
Dronning Maud Land margin: Site 693, dating, 55b, 344a
- Jane Basin
dating, 62b, 63b
TEM photomicrographs, 231b–232b, 235b
- Maud Rise: Site 689
dating, 57b, 99a
temperature associated with, 98a
water depth associated with, 99a
- Maud Rise: Site 690
abundance, 194a–195a
dating, 58b, 197a
- South Orkney microcontinent: Site 695, dating, 539a
- South Orkney microcontinent: Site 696, dating, 61b, 623a
- Weddell Basin: Site 694, dating, 59b
- Weddell Sea, Paleogene climate indicator, 951b
- South Orkney microcontinent, paleoenvironment, cooling, 60b–61b, 64b
- South Orkney microcontinent: Site 695
age-depth plots, versus magnetostratigraphy, 931b
alkalinity, 560a
depth and, 139b, 142b, 561a–562a
ammonia concentrations, depth and, 561a–562a
bathymetry, 530a
- calcium concentrations, depth and, 139b, 142b, 561a–562a
chloride concentrations, depth and, 561a–562a
chlorinity, 560a
consolidation
data summary, 214b, 222b
field curve, 219b
density, depth and, 180b, 541a–543a
dissolved organic carbon
depth and, 173b
pore water concentrations, 174b
downhole measurements, temperature measurements, 564a–568a
ethane concentrations, depth and, 563a–564a
heat flow measurements, 17b–20b
depth and, 25b
time and, 19b–20b
- ice-raftered detritus (IRD). *See* South Orkney microcontinent: Site 695, lithostratigraphy
- interstitial-water chemistry
amino acids, 179b–186b
data summary, 139b
procedure, 137b
- kerogen concentrations
depth and, 563a–564a
temperature measurements, 563a–564a, 565a
vitrinite reflectance measurements, 192b–194b
- lithology
dating, 93b, 531a
depth and, 93b
- lithostratigraphy
ice-raftered dropstones, 539a–540a
Miocene–Pliocene, 537a, 538a, 568a
Pliocene, 535a–537a, 568a
Pliocene–Quaternary, 533a–536a, 568a
- location, 528a, 530a
- magnesium concentrations, depth and, 139b, 142b, 561a–562a
- magnetic properties, natural remanent magnetization, 554a–557a
- methane concentrations
calculations, 35b
depth and, 563a–564a
- mineralogy, clay, 537a–539a
- oxygen isotope distribution, depth and, 139b, 142b
- paleoenvironment, 538a–539a
- pH, 560a
- phosphate concentrations, depth and, 561a–562a
- porosity
depth and, 180b, 541a–543a
diatom relationship with, 218b
- potassium concentrations, depth and, 139b, 142b, 561a–562a
- propane concentrations, depth and, 563a–564a
- reflectors
bottom simulating, 17b, 22b–23b
boundaries, 33b
depth and, 542a–543a, 550a
diagenesis and, 32b–33b
gas hydrate corresponding with, 33b–35b
- Rock-Eval procedure, 565a
- salinity, South Orkney microcontinent: Site 695, depth and, 561a
- sedimentation rates
biostratigraphic data, 555a, 557a, 560a, 569a
hiatus, 557a–558a
magnetostratigraphic data, 555a, 557a, 569a
- seismic reflection profile, 529a, 549a–550a
- seismic stratigraphy, 542a–546a
P-wave velocity, 544a–545a, 547a, 550a
- shear strength, 541a
depth and, 543a–544a
- vertical effective stress relationship with, 218b
- strontium isotope distribution
calculations, 153b, 155b–156b
depth and, 150b
sediment age versus, 152b
- sulfate concentrations, 560a
depth and, 139b, 142b, 561a–562a
pore water concentrations, 175b
- thermal conductivity, 541a
calculations, 21b–23b, 35b, 567a–568a
depth and, 23b, 26b, 543a, 545a
water content versus, 22b
- velocity, 541a–542a
depth and, 543a, 546a–548a
- volcanic ash
color, 537a
composition, 535a
depth and, 533a
- water content, depth and, 541a–543a
See also Weddell Sea
- South Orkney microcontinent: Site 696
age-depth plots, 933b
alkalinity, 647a
depth and, 139b, 143b, 650a–651a
ammonia concentrations, 647a–648a
depth and, 650a–651a
bathymetry, 28b, 611a
calcium concentrations
depth and, 139b, 143b, 650a–651a
South Orkney microcontinent: Site 696, 648a–649a
- chloride concentrations, depth and, 650a–651a
chlorinity, 643a, 646a
- consolidation
data summary, 214b, 222b
field curve, 220b
- density
depth and, 30b, 625a
range, 624a, 626a
- depth and, 94b
- dissolved organic carbon
depth and, 173b
pore water concentrations, 174b
- downhole measurements
gamma-ray log, 657a
resistivity log, 657a
sonic log, 657a
temperature measurements, 650a, 652a–653a
- ethane concentrations, 650a
depth and, 652a–653a
heat flow measurements, 20b–21b
depth and, 25b
time and, 21b
- interstitial-water chemistry, 650a
data summary, 139b
procedure, 137b
- kerogen concentrations
temperature measurements, 655a
vitrinite reflectance measurements, 192b–194b
- lithology, 614a–615a
dating, 94b
- lithostratigraphy
ice-raftered dropstones, 622a–624a
Miocene, 617a–618a
Miocene–Eocene, 620a
Pliocene, 615a–617a
Pliocene–Miocene, 617a
Quaternary–Pliocene, 613a–615a
- location, 609a
- magnesium concentrations, 648a–649a
depth and, 139b, 143b, 650a–651a
- magnetic properties, natural remanent magnetization, 641a–642a, 644a–645a
- magnetostratigraphy, geomagnetic polarity time scale, 646a–649a

- methane concentrations, 650a
depth and, 652a-653a
mineralogy, clay, 61b, 620a-623a
oxygen isotope distribution, depth and, 139b, 143b
paleoenvironment, 622a, 655a-656a, 658a-659a
pH, 646a-647a
depth and, 650a-651a
phosphate concentrations, 647a-648a
depth and, 650a-651a
porosity
depth and, 30b, 625a-626a
diatom relationship with, 219b
range, 624a, 626a
potassium concentrations, depth and, 139b, 143b, 650a-651a
propane concentrations, 650a
depth and, 652a-653a
reflectors
dating, 631a
depth and, 631a
diagenesis and, 32b
strength, 631a
unconformity boundary, 27b-28b
Rock-Eval procedure, 650a, 654a-655a
salinity, 643a, 646a
depth and, 650a
sedimentation rates
biostratigraphic data, 642a-643a, 648a
hiatus, 659a
magnetostratigraphic data, 642a-643a
seismic reflection profile, 610a, 630a
seismic stratigraphy, 628a-631a
P-wave velocity, 30b, 628a-629a, 631a
South Orkney microcontinent: Site 696, *P*-wave velocity, 32b
shear strength, 626a
depth and, 627a
vertical effective stress relationship with, 220b
strontium isotope distribution
calculations, 153b, 155b-156b
depth and, 151b
sediment age versus, 152b
sulfate concentrations, 647a
depth and, 139b, 143b, 650a-651a
pore water concentrations, 175b
thermal conductivity
calculations, 21b-23b
depth and, 23b, 26b, 630a-631a
water content versus, 22b
velocity, depth and, 30b, 626a-629a, 631a
volcanic ash, composition, 620a
water content
depth and, 625a
range, 624a, 626a
See also Weddell Sea
Southern Hemisphere
plate reconstruction, 85b
Pliocene climatic cooling, 803b
Southern Ocean
Antarctic glaciation and, 938b
climatic evolution, 849b
diatom abundances, 806b
ice rafting, 954b
magnetostratigraphy, diatom abundances
correlated with, 806b-807b
map, 805b
Pliocene sea-ice cover
compared with modern, 804b
diatom preservation and, 803b, 804b
sedimentation patterns, 803b-804b
See also Antarctic Ocean
Spathite, Dronning Maud Land margin: Site 692, depth and, 75b
Sr/Ca ratio, Maud Rise, 817b, 833b
"Strangelove Ocean" theory, 844b, 845b
Antarctic evidence, 944b
Strontium, reservoirs, Weddell Sea, 154b-155b, 156b
Sulfate concentrations
Dronning Maud Land margin: Site 693, 374a-375a
depth and, 138b, 141b, 380a
pore water concentrations, 175b
Jane Basin, 730a, 735a
depth and, 140b, 143b, 736a-737a
pore water concentrations, 175b
Maud Rise: Site 689
depth and, 129a-130a, 138b, 140b
pore water concentrations, 175b
Maud Rise: Site 690, 230a
depth and, 138b, 141b, 237a
pore water concentrations, 175b
South Orkney microcontinent: Site 695, 560a
depth and, 139b, 142b, 561a-562a
pore water concentrations, 175b
South Orkney microcontinent: Site 696, 647a
depth and, 139b, 143b, 650a-651a
pore water concentrations, 175b
Weddell Basin: Site 694
depth and, 139b, 142b
pore water concentrations, 175b
Tasman Sea, clay, volcanic alteration, 53b
Tethys, warm saline deep waters in Paleogene ocean and, 876b, 950b
Thermal conductivity
Dronning Maud Land margin: Site 693, 352a
depth and, 356a
Jane Basin, 715a
depth and, 719a-720a
Maud Rise: Site 689, 104a-105a
calculations, 21b
depth and, 23b, 25b, 106a
water content versus, 22b
Maud Rise: Site 690, 204a
calculations, 21b
depth and, 23b, 25b, 212a-213a
water content versus, 22b
South Orkney microcontinent, depth and, 23b
South Orkney microcontinent: Site 695, 541a
calculations, 21b-23b, 35b, 567a-568a
depth and, 23b, 26b, 543a, 545a
water content versus, 22b
South Orkney microcontinent: Site 696
calculations, 21b-23b
depth and, 26b, 630a-631a
water content versus, 22b
Weddell Basin: Site 694, depth and, 469a-470a
Thermocline, seasonal, environmental significance in open ocean, 842b
Turbidites
Weddell Abyssal Plain
source of, 955b-956b
W Antarctic ice sheet and, 938b
Weddell Basin: Site 694
depth and, 455a
thickness, 489a
Upwelling
Eocene/Oligocene boundary, Weddell Sea, 875b
Maud Rise, relation to Cretaceous temperature changes, 842b
mineral characteristics of sediments and, 76b-77b
Southern Ocean, katabatic winds and, 128b
Weddell Sea
Eocene increase, 949b
Oligocene, 951b
Velocity
Dronning Maud Land margin: Site 692, depth and, 304a-305a
Dronning Maud Land margin: Site 693, 240b, 242b-244b, 351a
depth and, 353a, 354a
Jane Basin
calculations, 714a
depth and, 719a
Maud Rise: Site 689, 101a-102a
depth and, 104a-105a
Maud Rise: Site 690, 202a, 204a
depth and, 209a
South Orkney microcontinent: Site 695, 541a-542a
depth and, 543a, 546a-548a
South Orkney microcontinent: Site 696, depth and, 30b, 626a-629a, 631a
Weddell Basin: Site 694, depth and, 467a-469a
Vermiculite, Weddell Basin: Site 694, dating, 59b
Victoria Land, glacial diamictites, 953b-954b
Vitrinite, South Orkney microcontinent, 193b-194b
Volcanism, iridium and, Maud Rise, 161b-163b
Warm Saline Deep Water
Antarctic Ocean
formation of, in Paleogene, 873b-874b
upward displacement of, 875b
Paleogene evidence for, Maud Rise, 865b
Warming, global, Paleocene-Eocene, Antarctic evidence for, 945b
Water content
Dronning Maud Land margin: Site 692, depth and, 304a
Dronning Maud Land margin: Site 693, depth and, 351a-352a
Jane Basin, depth and, 716a
Maud Rise: Site 689
depth and, 102a-103a
large variations, 101a
Maud Rise: Site 690, depth and, 205a-206a
South Orkney microcontinent: Site 695, depth and, 541a-543a
South Orkney microcontinent: Site 696
depth and, 625a
range, 624a, 626a
Weddell Basin: Site 694, depth and, 466a-467a
Weddell Abyssal Plain, Turbidites, W Antarctic ice sheet and, 938b
Weddell Basin
age-depth plots, 929b
paleoenvironment, 56b, 58b-60b
Weddell Basin: Site 694
alkalinity, depth and, 139b, 142b
ammonia concentrations, 484a
depth and, 485a-486a
butane concentrations, depth and, 487a
calcium concentrations, 484a
depth and, 139b, 142b, 485a-486a
chlorinity, depth and, 481a
density, depth and, 466a-467a
dissolved organic carbon
depth and, 172b
pore water concentrations, 174b
ethane concentrations, 485a
depth and, 487a
interstitial-water chemistry
data summary, 139b, 485a
procedure, 137b

SUBJECT INDEX

kerogen concentrations, Site 692 compared with, 488a
 lithologic units, 454a-455a
 comparison, 473a
 lithology
 dating, 93b
 depth and, 93b
 lithostratigraphy, 489a
 ice-rafted dropstones, 463a, 466a
 Miocene, 460a-461a
 Miocene-Pliocene, 455a-460a, 461a
 Pliocene, 455a, 458a
 Pliocene-Pleistocene, 453a-455a, 456a-457a
 location, 451a
 magnesium concentrations, 484a
 depth and, 139b, 142b, 485a-486a
 magnetic properties
 magnetic susceptibility, 477a, 482a-483a
 natural remanent magnetization, 477a-480a
 magnetostratigraphy, 477a
 geomagnetic polarity time scale, 481a, 484a
 methane concentrations, 485a
 depth and, 487a
 mineralogy, clay, 59b, 461a-462a, 465a
 oxygen isotope distribution, depth and, 139b,
 142b
 paleoenvironment, 462a-463a
 phosphate concentrations, 481a, 485a-486a
 porosity
 depth and, 218b, 466a-468a
 diatom relationship with, 218b
 potassium concentrations, 484a-485a
 depth and, 139b, 142b, 485a-486a
 propane concentrations, depth and, 487a
 reflectors
 dating, 470a-471a
 strength, 470a-471a
 Rock-Eval procedure, 487a-488a
 salinity, depth and, 481a, 485a
 sedimentation rates, 478a
 biostratigraphic data, 485a
 depth and, 484a

seismic reflection profile, 452a, 471a
 seismic stratigraphy, 469a-471a
P-wave velocity, 470a-471a, 472a
 shear strength
 depth and, 218b, 467a-468a
 vertical effective stress relationship with, 218b
 strontium isotope distribution
 calculations, 153b, 156b
 depth and, 150b
 sediment age versus, 152b
 sulfate concentrations
 depth and, 139b, 142b
 pore water concentrations, 175b
 thermal conductivity, depth and, 469a-470a
 velocity, depth and, 467a-469a
 water content, depth and, 466a-467a
See also Weddell Sea
 Weddell Gyre, 803b
 Weddell Sea
 carbon and nitrogen recycling,
 Miocene/Pliocene boundary changes, 956b
 depth distribution of sites, 940b
 dissolved organic carbon
 average concentration, 174b
 rate of accumulation logarithm versus, 175b
 fractionation factors, minerals and water, 144b
 grain-shape analyses, 96b
 end-member identification, 96b-103b, 107b-
 109b
 entropy values, 97b
 Fourier procedure, 103b
 Krumbein roundness values, 98b-99b
 interstitial-water chemistry, calculations,
 139b-144b
 kerogen concentrations, site comparisons,
 194b-197b
 location, 882b, 939b
 magnetostratigraphy, rounded end members
 based on, 101b
 mineralogy, clay, 51b-52b
 nepheloid layer transport, 116b-119b
 nitrogen data, 884b-889b
 organic carbon data, 884b-889b
 organic matter preservation, Neogene, 881b
 oxygen isotope distribution, magnesium
 concentrations versus, 145b
 permeability, calculations, 213b
 preconsolidation stress, calculations, 212b-213b
 regional geology, 90b-91b
 sea ice
 expansion, 956b-957b
 formation, Oligocene, 951b
 production in, 803b
 sedimentation rates, 925b
 as source of AABW, 938b
 stratigraphy,
 biostratigraphic-magnetostratigraphic
 correlations, 915b-936b
 surface-water temperatures, Eocene, 949b
 upwelling, Eocene/Oligocene boundary, 875b
 volcanic ash, alteration, 144b
 warm saline deep water, 949b-950b
 Eocene, 937
 water masses dominating, 938b-939b
See also Maud Rise
 Weddell Sea Bottom Water, 122b, 571b, 938b
 Wegener Canyon
 bathymetry, 40b, 42b
 SEABEAM measurements, 41b
 evolution, 45b-47b, 315a-316a, 317a
 rock dredging, 42b-44b
 seismic stratigraphy, lithology correlated with,
 44b, 46b
 West Antarctic ice sheet. *See* Antarctic margin
 W; Antarctica W
 Yukataga Formation, 953b
 Zeolites
 Jane Basin
 dating, 711a
 depth and, 711a
 Maud Rise, feldspar replacing, 10b

SITE INDEX

- Site 4, "black shales," 195b
 Site 5, "black shales," 195b
 Site 99, "black shales," 194b
 Site 100, "black shales," 194b
 Site 101, "black shales," 195b
 Site 105, "black shales," 194b, 195b
 Site 135, "black shales," 195b
 Site 137, "black shales," 195b
 Site 167, "black shales," 195b
 Site 249
 lithologies, 80b-82b
 nanofossil assemblage, 335b
 oxic/anoxic transition, 337b
 Site 264, sporomorphs, 595b
 Site 266
 diatom zones, 761b
 radiolarian zone definitions, 710b
 sporomorphs, 595b
 Site 268, sporomorphs, 595b
 Site 270, sporomorphs, 595b
 Site 278, diatom zones, 761b
 Site 327
 Maestrichtian hiatus, 489b
 nanofossil assemblage, 333b, 335b
 nanofossil assemblages, 469b
 palynoflora, 451b
 Site 328, palynoflora, 451b
 Site 330
 hydrocarbons, 206b
 nanofossil assemblage, 335b
 palynoflora, 451b
 Site 356, nanofossil assemblages, 519b, 520b, 521b
 Site 358, oxygen isotope ratios, Maestrichtian, 821b
 Site 361, 84b
 lithologies, 82b-83b
 nanofossil assemblage, 335b
 oxic/anoxic transition, 336b
 Site 364, petroleum-generating potential, 196b
 Site 367
 "black shales," 194b-195b
 petroleum-generating potential, 196b
 Site 368, "black shales," 195b
 Site 384
 nanofossil assemblages, 536b
 Site 386, "black shales," 195b
 Site 387
 "black shales," 194b
 Nephrolithus frequens, 472b
 Reinhardtites levis, 473b
 Site 391
 "black shales," 194b, 195b
 Site 398, "black shales," 195b
 Site 400, "black shales," 195b
 Site 402, "black shales," 195b
 Site 465, oxygen isotope ratios, Maestrichtian, 821b
 Site 474
 dissolved organic carbon, 169b, 176b
 sulfate reduction, 174b
 Site 479
 dissolved organic carbon, 169b, 176b
 sulfate reduction, 174b
 Site 511, 83b-84b
 benthic foraminifers, 575b
 "black shales," nanofossil assemblage, 333b, 335b
 hydrocarbons, 206b
 Maestrichtian hiatus, 489b
 nanofossil occurrence comparisons, 655b, 656b
 oxic/anoxic transition, 336b
 sporomorphs, 595b
 Site 512
 naonofossil occurrence comparisons, 656b
 sporomorphs, 595b
 Site 513
 nanofossil occurrence comparisons, 655b, 656b
 sporomorphs, 595b
 Site 514
 Cosmiodiscus insignis abundances, 806b
 sporomorphs, 595b
 Site 516
 Chiasmolithus solitus, 654b
 nanofossil occurrence comparisons, 652b, 655b, 656b
 Nephrolithus frequens, 472b
 Site 522, nanofossil occurrence comparisons, 652b
 Site 523
 Chiasmolithus solitus, 654b
 nanofossil occurrence comparisons, 652b, 655b
 Site 524
 nanofossil assemblages, 519b, 520b
 Site 525
 Abathomphalus mayaroensis, 499b
 Reinhardtites levis, 471b
 Site 527
 carbon isotope ratios, 843b
 surface-water temperatures, early Paleogene, 859b
 Site 528, nannofossil occurrence comparisons, 652b
 Site 530
 Nephrolithus frequens absence, 472b
 Reinhardtites levis, 471b
 Site 535, "black shales," 194b
 Site 558, nannofossil occurrence comparisons, 652b
 Site 563, nannofossil occurrence comparisons, 652b
 Site 577, carbon isotope ratios, 843b
 Site 592, smectite laths, 53b
 Site 605, nannofossil biostratigraphy, 472b
 Site 618, dissolved organic carbon, 169b, 176b
 Site 619, dissolved organic carbon, 169b, 176b
 Site 623
 dissolved organic carbon, 169b, 176b
 sulfate reduction, 174b
 Site 689
 age-depth curve, 123a
 benthic foraminifers, 112a-117a
 carbon isotope data, 878b-879b, 904b
 distribution, 491b, 574b, 580b
 oxygen isotope data, 878b-879b, 904b
 See also Paleontological Index
 biostratigraphic chart, 110a-111a, 466b
 calcareous nanofossils, 114a-116a
 distribution, 468b-469b, 616b-620b, 644b-645b
 See also Paleontological Index
 calcium carbonate
 accumulation rates, 841b
 percentages, 974b-976b, 980b-981b, 984b
 carbon isotope data, 986b-987b
 clay mineralogy, 57b, 99a
 major events, 64b
 consolidation test data, 214b
 diatoms, 116a-117a
 distribution, 770b, 921b
 drilling
 objectives, 90a-91a
 operations, 91a-94a
 summary, 9a-11a, 132a-134a
 foraminifers
 carbon isotope data, 814b
 insoluble accumulation rate, 834b
 oxygen isotope data, 814b

SITE INDEX

geochemistry, inorganic, 128a–129a
 geochemistry, organic, 129a–131a
 data, 884b–885b
 Rock-Eval pyrolysis, 131a, 132a
 geomagnetic polarity time scale, 124a–127a,
 262b, 290b–292b
 hiatuses, 123a–128a
 incompatible elements, 9b
 interstitial-water data, 129a, 138b, 149b
 lithology, site comparison, 614b–615b, 641b,
 973b
 lithostratigraphy, 95a–98a
 location, 6b, 91a, 465b, 516b, 534b, 573b,
 902b
 paleogeographic reconstruction, 489b
 magnetic properties, 121a–123a, 295b–304b
 magnetostratigraphy, 258b, 268b–273b
 navigation data, 44a–45a
 organic carbon levels, 190b
 oxygen isotope data, 986b–987b
 petrographic description, 100a
 physical properties, 101a–107a
 planktonic foraminifers, 109a–112a
 carbon isotope data, 851b–852b
 distribution, 491b–492b
 oxygen isotope data, 851b–852b
 See also Paleontological Index
 pollen and spores, 119a–121a
 radiolarians, distribution, 117a–119a,
 682b–687b, 714b–715b, 921b
 sediment composition, 96a
 sedimentation rates, 121a–123a
 seismic stratigraphy, 107a–108a
 silicoflagellates, distribution, 119a,
 734b–735b, 921b
 sponge spicules
 distribution, 964b
 See also Paleontological Index
 stratigraphy, 9a
 temperature measurements, 19b, 131a–132a

Site 690
 age-depth curve, 225a–227a
 basalt, 198a–202a
 geochemistry, 199a–202a
 petrography, 198a–199a
 benthic foraminifers, 213a–216a
 carbon isotope data, 879b–880b, 904b
 distribution, 494b, 574b, 580b
 oxygen isotope data, 879b–880b, 904b
 See also Paleontological Index
 biostratigraphic chart, 217a–218a, 466b
 calcareous nannofossils, 216a–220a
 distribution, 470b–471b, 520b–521b, 622b–
 628b, 648b–649b
 See also Paleontological Index
 calcium carbonate
 accumulation rates, 841b
 percentages, 977b–979b, 982b–983b
 carbon isotope data, 987b–989b
 clay mineralogy, 57b, 194a–198a
 major events, 64b
 consolidation test data, 214b
 diatoms, 221a–222a
 distribution, 770b, 923b
 dinoflagellates, distribution, 535b–536b,
 539b, 541b
 drilling
 objectives, 184a
 operations, 185a–186a
 summary, 9a–11a, 231a, 234a–240a
 foraminifers
 carbon isotope data, 815b

insoluble accumulation rate, 834b
 oxygen isotope data, 815b
 geochemistry, inorganic, 227a–231a
 geochemistry, organic
 carbon levels, 190b
 data, 885b–886b
 Rock-Eval pyrolysis, 231a, 238a
 geomagnetic polarity time scale, 228a–229a,
 232a–234a, 262b, 290b–292b
 ice-raftered detritus, 953b
 incompatible elements, 9b
 interstitial-water data, 138b, 150b
 lithology, site comparison, 614b–615b, 641b,
 973b
 lithostratigraphy, 189a–194a
 Site 689 comparison, 193a–194a
 location, 6b, 185a, 465b, 516b, 534b, 573b,
 902b
 paleogeographic reconstruction, 489b
 magnetic properties, 225a, 305b–315b
 magnetochronology, 258b, 274b–280b
 major element data, 8b, 167b–168b, 201a
 microprobe analyses, 7b
 navigation data, 44a–45a
 oxygen isotope data, 987b–989b
 physical properties, 202a–205a
 planktonic foraminifers, 209a–213a
 carbon isotope data, 853b–854b
 distribution, 494b–495b, 554b
 oxygen isotope data, 853b–854b
 tropical zonations correlated with, 557b
 See also Paleontological Index
 pollen and spores, 223a
 radiolarians, 222a–223a
 distribution, 689b, 716b–717b, 923b
 sediment composition, 190a–191a
 sedimentation rates, 225a
 seismic stratigraphy, 205a–208a
 silicoflagellates, 223a
 distribution, 736b, 923b
 “Strangloove Ocean,” 944b
 stratigraphy, 9a
 temperature measurements, 19b, 231a
 trace element data, 8b, 201a

Site 691
 drilling
 objectives, 294a, 296a
 operations, 296a–297a
 summary, 9a–11a, 315a–317a
 ice conditions, 39a
 lithostratigraphy, 298a–301a
 location, 40b, 72b, 121b
 navigation data, 45a
 sediment composition, 300a
 stratigraphy, 9a
 three-dimensional view, 43b

Site 692
 benthic foraminifers, 307a–308a
 biostratigraphic chart, 311a
 calcareous nannofossils, 309a
 distribution, 329b–331b
 See also Paleontological Index
 clay mineralogy, 301a, 303a
 diatoms, 309a–310a
 dinocysts
 dating, 452b
 distribution, 454b
 See also Paleontological Index
 drilling
 objectives, 294a, 296a
 operations, 297a–298a
 summary, 9a–11a, 315a–317a

geochemistry, organic, 312a–315a
 organic carbon levels, 191b
 parameters, 202b
 Rock-Eval pyrolysis, 313a–314a
 ice conditions, 39a
 ice-raftered dropstones, 301a–303a
 lithology, site comparison, 86b–87b
 lithostratigraphy, 298a–301a
 summary, 74b, 299a, 327b, 445b
 location, 40b, 72b, 121b, 444b, 449b
 magnetic properties, 311a–312a
 molluscs, 310a
 navigation data, 45a
 petrographic description, 304a
 physical properties, 303a–305a
 planktonic foraminifers, 307a
 pollen and spores, 310a
 distribution, 453b
 See also Paleontological Index
 radiolarians, 310a
 sediment composition, 300a
 seismic stratigraphy, 305a–306a
 stratigraphy, 9a
 thin section descriptions, 76b
 three-dimensional view, 43b
 vitrinite reflectance measurements,
 192b–193b
 X-ray diffraction analysis, 79b–80b

Site 693
 benthic foraminifers, 359a–360a
 biostratigraphic chart, 360a–367a
 calcareous nannofossils, 360a–361a
 distribution, 332b–334b, 650b
 See also Paleontological Index
 clay mineralogy, 55b, 342a–343a, 344a
 major events, 64b
 consolidation test data, 214b
 diatoms, 361a–363a
 distribution, 367b, 776b–777b, 926b
 See also Paleontological Index
 drilling
 operations, 330a–332a
 summary, 9a–11a, 383a–386a
 geochemistry, inorganic, 372a–377a
 geochemistry, organic, 377a–378a
 data, 886b–888b
 parameters, 202b
 Rock-Eval pyrolysis, 377a, 381a
 geomagnetic polarity time scale, 372a, 374a,
 376a
 grain shape analysis, 100b–102b
 ice conditions, 39a
 ice-raftered dropstones, 345a–349a
 interstitial-water data, 138b, 150b, 379a
 lithology
 sand fraction, 341a–342a
 site comparison, 86b–87b, 92b–95b, 641b
 lithostratigraphy, 332a–340a
 summary, 82b, 241b, 328b, 333a, 597b
 location, 40b, 72b, 121b, 240b, 427b, 449b
 logging, 378a–381a
 magnetic properties, 368a–369a, 370a–371a,
 373a
 Milankovitch cycles, 245b–246b, 249b–250b
 navigation data, 45a
 organic carbon levels, 190b
 petrographic description, 346a–349a
 physical properties, 349a–353a
 planktonic foraminifers, 359a
 dating, 319b–320b
 See also Paleontological Index
 pollen and spores, 365a
 radiolarians, 363a–364a

distribution, 354b, 718b-719b, 926b
See also Paleontological Index
 sand fraction analysis, 343a
 sediment classification, 17a-22a
 sediment composition, 335a-336a
 sedimentation rates, 369a-372a
 seismic reflectors, 357a-358a
 seismic stratigraphy, 353a-358a
 silicoflagellates, 364a-365a
 distribution, 737b, 926b
 phylogeny, 431b
See also Paleontological Index
 sponge spicules
 distribution, 965b
See also Paleontological Index
 stratigraphy, 9a
 thin section descriptions, 84b
 three-dimensional view, 43b
 vitrinite reflectance measurements, 192b-193b
 warming episodes, 954b

Site 694

benthic foraminifers, 471a
 biostratigraphic chart, 474a-475a
 clay mineralogy, 59b, 461a-463a, 465a
 diatoms, 472a-476a
 distribution, 778b, 928b
 drilling
 objectives, 450a
 operations, 450a-451a, 453a
 summary, 9a-11a, 488a-490a
 geochemistry, inorganic, 478a-485a
 geochemistry, organic, 485a-488a
 data, 888b-889b
 parameters, 202b
 Rock-Eval pyrolysis, 487a, 488a
 geomagnetic polarity time scale, 481a, 484a
 grain shape analysis, 100b-102b
 ice-rafterd dropstones, 463a
 interstitial-water data, 139b, 150b, 485a
 lithology, site comparison, 92b-95b
 lithostratigraphy, 453a-461a
 magnetic properties, 477a, 479a-481a
 navigation data, 45a
 petrographic description, 466a
 physical properties, 464a-469a
 pollen and spores, 476a-477a
 radiolarians, 476a
 distribution, 928b
 sediment composition, 460a
 sedimentation rates, 478a
 seismic stratigraphy, 469a-471a
 silicoflagellates, 476a
 distribution, 928b
 sponge spicules
 distribution, 965b
See also Paleontological Index
 stratigraphy, 9a

Site 695

biostratigraphic chart, 552a-553a
 clay mineralogy, 62b, 537a-539a
 consolidation test data, 214b
 diatoms, 548a-550a
 distribution, 778b-779b, 930b
See also Paleontological Index
 drilling
 objectives, 527a-528a

operations, 528a, 530a-531a
 summary, 9a-11a, 568a-569a
 geochemistry, inorganic, 558a-562a
 geochemistry, organic, 563a-564a
 Rock-Eval pyrolysis, 565a
 grain shape analysis, 100b-102b
 ice conditions, 39a
 ice-rafterd dropstones, 539a-540a
 interstitial-water data, 139b, 150b, 561a
 amino acids, 181b, 183b-184b
 lithology, site comparison, 92b-95b
 lithostratigraphy, 531a-537a
 location, 28b, 180b, 528a
 magnetic properties, 554a-555a, 556a-559a
 navigation data, 45a
 petrographic description, 540a
 physical properties, 540a-542a
 planktonic foraminifers, 547a
 radiolarians, 551a-554a
 distribution, 930b
 sediment composition, 532a
 sedimentation rates, 555a-558a
 seismic stratigraphy, 542a-546a
 silicoflagellates, 548a
 distribution, 738b, 930b
 sponge spicules
 distribution, 966b
See also Paleontological Index
 stratigraphy, 9a
 temperature measurements, 19b-20b, 25b,
 564a-568a
 vitrinite reflectance measurements, 192b,
 194b

Site 696

benthic foraminifers, 632a-637a
 biostratigraphic chart, 633a-635a
 calcareous nannofossils, 637a-638a
 distribution, 651b
 clay mineralogy, 61b, 620a-622a, 623a
 major events, 64b
 consolidation test data, 214b
 core recovery, 30b
 diatoms, 638a-639a
 distribution, 781b, 932b
See also Paleontological Index
 dinoflagellate cysts
 distribution, 600b
 stratigraphic ranges, 602b
See also Paleontological Index
 drilling
 objectives, 608a-609a
 operations, 609a, 611a-613a
 summary, 9a-11a, 653a-656a, 658a-659a
 geochemistry, inorganic, 643a-650a
 geochemistry, organic, 650a
 Rock-Eval pyrolysis, 650a, 654a
 geomagnetic polarity time scale, 646a-649a
 grain shape analysis, 100b-102b
 ice conditions, 39a
 ice-rafterd dropstones, 622a-624a
 interstitial-water data, 139b, 151b, 650a
 lithology, site comparison, 92b-95b, 641b
 lithostratigraphy, 613a-620a
 summary, 597b, 614a
 location, 28b
 logging, 653a
 magnetic properties, 641a-642a, 644a-647a
 navigation data, 45a
 petrographic description, 624a

physical properties, 624a-628a
 planktonic foraminifers, 632a
 pollen and spores, 640a
 radiolarians, 639a-640a
 distribution, 932b
 sediment composition, 616a
 seismic stratigraphy, 628a-631a
 silicoflagellates, 638a
 distribution, 739b, 932b
 sponge spicules
 distribution, 967b
See also Paleontological Index
 stratigraphy, 9a
 temperature measurements, 21b, 25b,
 650a-652a
 vitrinite reflectance measurements, 192b,
 194b

Site 697

benthic foraminifers, 717a
 biostratigraphic chart, 722a-723a
 clay mineralogy, 63b, 710a-712a, 715a
 consolidation test data, 214b
 deep-water circulation, 112b
 diatoms, 718a-725a
 distribution, 932b
 drilling
 objectives, 705a-708a
 summary, 9a-11a, 738a-741a
 geochemistry, inorganic, 729a-736a
 geochemistry, organic, 736a-738a
 Rock-Eval pyrolysis, 736a-737a, 739a
 geomagnetic polarity time scale, 730a, 735a
 grain shape analysis, 100b-102b
 grain size analysis, 117b-118b
 ice conditions, 39a
 interstitial-water data, 140b, 151b, 736a
 lithology, site comparison, 92b-95b
 lithostratigraphy, 708a-710a
 magnetic properties, 726a-727a, 728-734a
 magnetic reversals, 808b
 microfabric analyses, 227b-228b
 navigation data, 45a
 physical properties, 713a-716a
 radiolarians, 725a
 distribution, 932b
 sediment composition, 711a
 sedimentation rates, 727a-729a
 seismic stratigraphy, 716a-717a
 silicoflagellates, 725a
 distribution, 740b, 932b
 sponge spicules
 distribution, 967b
See also Paleontological Index
 stratigraphy, 9a
 vitrinite reflectance measurements, 192b,
 194b

Site 698, Cretaceous foraminifers, 489b

Site 700

Abathomphalus mayaroensis, 498b
 Cretaceous foraminifers, 489b

Site 738, Cretaceous/Tertiary boundary section, 536b

Site 748, ice-rafterd detritus, 953b

Site 751, radiolarians, 688b

PALEONTOLOGICAL INDEX

- Abathomphalus intermedius*, Maud Rise: Site 690, 505b, 512b
- Abathomphalus mayaroensis*
Maud Rise: Site 689, first occurrence, 499b
Maud Rise: Site 690, 505b, 512b
first occurrence, 499b
- Abathomphalus mayaroensis* Total Range Zone
Maud Rise: Site 689, distribution, 497b
Maud Rise: Site 690, distribution, 497b
- Acanthocircus* sp., Dronning Maud Land margin: Site 693, 355b, 359b, 362b
- Acarinina bullbrooki*, Maud Rise: Site 689, 568b
- Acarinina bullbrooki* Interval Zone
Maud Rise: Site 689, 553b
Maud Rise: Site 690, 553b
- Acarinina collectae*, Maud Rise: Site 690, 568b
- Acarinina collectae* Partial Range Zone
Maud Rise: Site 689, 553b
Maud Rise: Site 690, 553b
- Acarinina esnaensis*, Maud Rise: Site 689, 566b
- Acarinina interposita*, Maud Rise: Site 689, 568b
- Acarinina matthewsae*, Maud Rise: Site 689, 568b
- Acarinina nitida*, Maud Rise: Site 690, 566b
- Acarinina pentacamerata* Interval Subzone
Maud Rise: Site 689, 553b
Maud Rise: Site 690, 553b
- Acarinina praepentacamerata* Interval Zone
Maud Rise: Site 689, 553b
Maud Rise: Site 690, 553b
- Acarinina primitiva*, Maud Rise: Site 689, 568b
- Acarinina primitiva* Interval Zone
Maud Rise: Site 689, 553b
Maud Rise: Site 690, 553b
- Acarinina pseudotopolensis*, Maud Rise: Site 689, 566b
- Acarinina velascoensis*, Maud Rise: Site 690, 566b
- Acarinina wilcoxensis berggreni*, Maud Rise: Site 689, 566b
- Acarinina wilcoxensis berggreni* Interval Zone
Maud Rise: Site 689, 553b
Maud Rise: Site 690, 553b
- Acarinina wilcoxensis wilcoxensis*, Maud Rise: Site 690, 566b
- Acrosphaera australis*, Maud Rise: Site 689, 712b-713b, 721b-722b
- Acrosphaera murrayana*, Maud Rise: Site 689, 713b, 722b
- Acrosphaera* sp., Maud Rise: Site 689, 690b, 701b, 713b, 722b
- Actinocyclus ingens*, Maud Rise: Site 689, 791b-792b, 797b
- Actinocyclus ingens* Partial-Range Zone, 766b
- Actinomma golownini*
Maud Rise: Site 689, 690b, 701b
Maud Rise: Site 690, 690b
- Actinomma golownini* Zone
Maud Rise: Site 689, 679b
Maud Rise: Site 690, 679b
- Actinomma medusa*
Maud Rise: Site 689, 690b-691b, 701b
Maud Rise: Site 690, 690b-691b
- Actinomma* sp.
Dronning Maud Land margin: Site 693, 355b, 362b
Maud Rise: Site 689, 359b, 691b, 701b
Maud Rise: Site 690, 691b
- Acuturris scotus*, Maud Rise: Site 690, 532b
- Adonnanonna primadonna*, Maud Rise: Site 689, 751b, 760b
- Ahmuellerella octoradiata*, Maud Rise: Site 690, 479b
- Alisporites grandis*, Dronning Maud Land margin: Site 692, 460b
- Alisporites similis*, Dronning Maud Land margin: Site 692, 460b
- Amblypyrgus campanellus* n. sp., Dronning Maud Land margin: Site 693, 369b-370b, 395b, 398b, 402b
- Amithalithina sigmundii* n. sp., Maud Rise: Site 690, 626b-627b, 637b
- Ancylopyrgus reticulatus* n. sp., Dronning Maud Land margin: Site 693, 370b, 387b, 389b, 400b
- Angiosperms, South Orkney microcontinent:
Site 696, taxonomic list, 601b
- Angulogerina szajnochae*, Maud Rise: Site 690, 593b
- Antarctissa deflandrei* n. comb.
Maud Rise: Site 689, 694b, 704b, 713b, 723b
Maud Rise: Site 690, 694b
- Antarctissa deflandrei* Zone
Maud Rise: Site 689, 678b
Maud Rise: Site 690, 678b
- Archaeodictyonitra* sp.
Dronning Maud Land margin: Site 693, 363b
Maud Rise: Site 689, 360b
- Archaeoglobigerina australis* n. sp.
Maud Rise: Site 689, 827b
Maud Rise: Site 690, 504b-505b, 509b-510b, 513b, 826b
- Archaeoglobigerina mateola* n. sp.
Maud Rise: Site 689, new species, 490b
Maud Rise: Site 690, 505b, 510b-511b, 513b
- Archeopyrgus melosioides* n. sp., Dronning Maud Land margin: Site 693, 370b-371b, 393b, 396b
- Arkhangelskiella cymbiformis*, Maud Rise: Site 690, 532b
- Arkhangelskiella* sp., Maud Rise: Site 690, abundance, 523b
- Asteromphalus inaequabilis* n. sp., Maud Rise: Site 689, 792b, 798b, 802b
- Asteromphalus kennettii* n. sp., Maud Rise: Site 689, 793b, 798b, 802b
- Asteromphalus kennettii* Partial-Range Zone, 768b
- Bachmannocena apiculata*, Maud Rise: Site 689, 747b
- Bachmannocena occidentalis* n. comb., Maud Rise: Site 689, 747b
- Baculatisporites comaumensis*, Dronning Maud Land margin: Site 692, 459b
- Baculatisporites* sp., South Orkney microcontinent: Site 696, 607b
- Basilicostephanus mirabilis* n. sp., Dronning Maud Land margin: Site 693, 371b, 385b-387b, 402b
- Basilicostephanus ornatus* n. sp., Dronning Maud Land margin: Site 693, 371b, 386b-388b
- Batiacasphaera asperata*, Dronning Maud Land margin: Site 692, 462b
- Batiacasphaera compta*, South Orkney microcontinent: Site 696, 609b
- Batioladinium* sp., Dronning Maud Land margin: Site 692, 461b
Benthic foraminifers
Dronning Maud Land margin: Site 692, abundance, 307a-308a
Dronning Maud Land margin: Site 693, abundance, 321b, 359a-360a
- Jane Basin
abundance, 717a
biostratigraphy, 722a-723a
- Maud Rise: Site 689
abundance, 112a-114a, 574b-575b
biostratigraphy, 110a-111a, 572b-573b, 904b
carbon burial, 843b-844b
carbon isotope data, 818b-819b, 878b-879b, 904b
carbon isotopic temperature gradients, 839b-841b
dating, 571b-572b, 575b-578b
diversity, 579b-586b
extinction, 844b
faunal list, 589b-591b
first appearance, 578b-579b, 911b
last appearance, 578b-579b, 911b
magnetostratigraphy, 904b
oxygen isotope correction factors, 834b-836b
oxygen isotope data, 878b-879b, 904b
oxygen isotopic temperature gradients, 836b, 838b-839b
Site 690 depths compared with, 906b-908b
- Maud Rise: Site 690
abundance, 213a-215a, 574b-575b
biostratigraphy, 217a-218a, 572b, 905b
carbon burial, 843b-844b
carbon isotope data, 879b-880b, 904b, 943b
carbon isotopic temperature gradients, 839b-841b
dating, 571b-572b, 575b-578b
diversity, 579b-586b
extinction, 844b
faunal list, 589b-591b
first appearance, 578b-579b, 911b
last appearance, 578b-579b, 911b
magnetostratigraphy, 905b
oxygen isotope correction factors, 834b-836b
oxygen isotope data, 818b-819b, 879b-880b, 904b, 942b
oxygen isotopic temperature gradients, 836b, 838b-839b
Site 689 correlated with, 219a, 906b
- South Orkney microcontinent: Site 695
abundance, 547a
biostratigraphy, 552a-553a
- South Orkney microcontinent: Site 696
abundance, 632a, 635a-637a
biostratigraphy, 633a-635a
- Weddell Basin: Site 694, abundance, 471a
- Biantholithus sparsus*
Maud Rise: Site 690, 528b
abundance, 523b
- Bicolumnus ovatus* n. sp., Maud Rise: Site 689, 658b, 661b
- Bidiscus rotatorius*, Maud Rise: Site 690, 479b
- Bilingua rossii* n. sp., Dronning Maud Land margin: Site 693, 371b-372b, 400b-401b
- Biscutum castrorum*

- Maud Rise: Site 690, 530b
abundance, 523b
- Biscutum constans*, Maud Rise: Site 690, 480b
- Biscutum coronum*, Maud Rise: Site 690, 480b
- Biscutum coronum* Zone
Maud Rise: Site 689, 467b
Maud Rise Site 690, 467b
- Biscutum dissimilis*, Maud Rise: Site 690, 479b
- Biscutum magnum*, Maud Rise: Site 690, 480b
- Biscutum magnum* Zone
Maud Rise: Site 689, 467b
Maud Rise: Site 690, 467b
- Biscutum neocoronum* n. sp., Maud Rise: Site 690, 627b-628b, 633b
- Biscutum notaculum*, Maud Rise: Site 690, 480b
- Bivalves, Dronning Maud Land margin: Site 692, 445b-446b
- Blackites spinosus*, Maud Rise: Site 689, 665b
- Blackites spinosus* Zone, Maud Rise: Site 689, 643b
- Bolboforma antarctica* n. sp.
Maud Rise: Site 689, 668b, 670b, 673b
abundance, 669b
first occurrence, 670b
last occurrence, 670b
- Maud Rise: Site 690, 668b, 670b, 673b
abundance, 670b
first occurrence, 670b
last occurrence, 670b
- Bolboforma irregularis*
Maud Rise: Site 689, 670b-671b, 673b
abundance, 669b
first occurrence, 670b
last occurrence, 670b
- Bolboforma reticulata*
Maud Rise: Site 689, 670b, 673b
abundance, 669b
first occurrence, 670b
last occurrence, 670b
- Bolboforma* sp.
Maud Rise: Site 689, 671b, 673b
abundance, 668b-669b
hiatus, 670b
- Maud Rise: Site 690
abundance, 668b
hiatus, 670b
- Bolboforma spinosa*
Maud Rise: Site 690, 670b, 673b
abundance, 670b
first occurrence, 670b
last occurrence, 670b
- Brigantedinium* sp., South Orkney microcontinent: Site 696, 611b
- Brownsonia enormis*, Maud Rise: Site 690, 481b
- Broomea* sp., Dronning Maud Land margin: Site 692, 461b
- Bulimina callahani*, Maud Rise: Site 689, 592b
- Bulimina elongata*, Maud Rise: Site 690, 592b
- Bulimina midwayensis*, Maud Rise: Site 690, 593b
- Bulimina semicostata*, Maud Rise: Site 690, 592b
- Bulimina simplex*, Maud Rise: Site 690, 592b
- Bulimina thanetensis*, Maud Rise: Site 690, 592b
- Calcareous dinoflagellates
Maud Rise: Site 689
abundance, 534b-537b
Cretaceous/Tertiary boundary, 533b-543b
distribution, 534b-537b
- Maud Rise: Site 690
abundance, 534b-537b
distribution, 534b-537b
- Calcareous nannofossils
Dronning Maud Land margin: Site 692
abundance, 309a, 329b-331b
biostratigraphy, 311a
dating, 326b-327b, 328b-331b
distribution, 329b-331b
ecology, 332b-334b
paleoenvironment, 332b-334b
regional correlations, 334b-335b
- Dronning Maud Land margin: Site 693
abundance, 332b-334b, 336b, 360a-361a, 647b, 650b
black shale, 336b-339b
dating, 330b-334b, 641b
distribution, 332b-334b, 650b
ice patterns related to, 657b
regional correlations, 334b-335b
zonation, 642b
- Maud Rise: Site 689
abundance, 114a-116a, 467b, 468b-469b, 616b-617b, 643b-646b
biogeography, 470b-473b
biostratigraphy, 110a-111a, 468b-470b, 632b
dating, 467b, 468b-469b, 614b, 641b
distribution, 618b-620b, 644b-645b
Falkland Plateau taxa compared with, 472b
first occurrence, 629b
hiatus, 616b, 632b, 657b
holococcoliths, 473b-474b
See also specific types
ice patterns related to, 657b
last occurrence, 629b
magnetostratigraphy, 620b-621b, 632b, 650b, 652b-654b
oldest, 473b
oxygen isotope data, 836b
paleobiogeography, 654b-656b
Site 690 depths compared with, 906b-908b
zonation, 468b-469b, 486b, 642b
- Maud Rise: Site 690
abundance, 216a, 218a-221a, 467b-468b, 470b-471b, 518b, 520b-521b, 646b-647b, 648b-649b
biogeography, 470b-473b
biostratigraphy, 217a-218a, 468b-470b, 516b-518b, 632b
bioturbation, 522b, 618b
boundary comparisons, 519b-522b
boundary selection criteria, 517b-518b
burrows, 522b, 525b
Cretaceous/Tertiary boundary section, 515b-524b
dating, 467b-468b, 470b-471b, 641b
distribution, 622b-628b, 648b-649b
Falkland Plateau taxa compared with, 472b
first occurrence, 629b
hiatus, 522b, 524b, 657b
holococcoliths, 473b-474b
See also specific types
ice patterns related to, 657b
last occurrence, 629b
magnetostratigraphy, 620b-621b, 632b, 650b, 652b-654b
oldest, 473b
oxygen isotope data, 836b
paleobiogeography, 654b-656b
Site 689 depths compared with, 906b-908b
zonation, 470b-471b, 486b, 520b, 615b, 616b, 642b
- South Orkney microcontinent: Site 696
abundance, 637a-638a, 647b, 650b, 651b
biostratigraphy, 633a-635a
dating, 641b
- distribution, 651b
ice patterns related to, 657b
ice-sheet development, 955b-956b
zonation, 642b
See also specific types
- Calcarous nannofossils
Maud Rise: Site 689, zonation, 615b-616b
Maud Rise: Site 690, dating, 615b
- Calcidiscus leptoporus*, Maud Rise: Site 689, 666b
- Calyptosporium carinatum* n. sp., Dronning Maud Land margin: Site 693, 407b, 420b, 423b
- Calyptosporium cyclacaenum* n. sp., Dronning Maud Land margin: Site 693, 407b, 420b
- Calyptosporium effalum* n. sp., Dronning Maud Land margin: Site 693, 407b, 417b, 421b
- Calyptosporium exasperatum* n. sp., Dronning Maud Land margin: Site 693, 407b-408b, 419b
- Calyptosporium galeolum* n. sp., Dronning Maud Land margin: Site 693, 408b, 416b
- Calyptosporium lentiginosum* n. sp., Dronning Maud Land margin: Site 693, 408b, 418b, 425b
- Calyptosporium maximum* n. sp., Dronning Maud Land margin: Site 693, 408b, 416b, 422b
- Calyptosporium peregrinum* n. sp., Dronning Maud Land margin: Site 693, 408b, 415b
- Calyptosporium pustulatum* n. sp., Dronning Maud Land margin: Site 693, 408b, 423b
- Calyptosporium* sp.
Dronning Maud Land margin: Site 693
calyptra, 406b
confossa zone, 406b
fala, 406b
infundibular operculum, 406b
infundibulum, 406b
lacinia, 406b-407b
- Cardiifolia toxon* n. sp., Dronning Maud Land margin: Site 693, 411b, 424b
- Carpocanistrum* sp.
Maud Rise: Site 689, 695b, 705b
Maud Rise: Site 690, 695b
- Centrosphaera barbata*
Maud Rise: Site 689, 541b, 548b
Maud Rise: Site 690, 481b, 541b, 548b
- Cephalopods, Dronning Maud Land margin: Site 692, 445b
- Ceratocyrtis mashae*
Maud Rise: Site 689, 694b, 704b
Maud Rise: Site 690, 694b
- Ceratocyrtis* sp.
Maud Rise: Site 689, 694b, 704b
Maud Rise: Site 690, 694b
- Chiasmolithus altus*
Maud Rise: Site 689, 662b
abundance, 656b
Maud Rise: Site 690, abundance, 656b
- Chiasmolithus altus* Zone
Maud Rise: Site 689, 642b
Maud Rise: Site 690, 642b
- Chiasmolithus bidens*, Maud Rise: Site 690, 633b
- Chiasmolithus consuetus*, Maud Rise: Site 690, 633b
- Chiasmolithus danicus*, Maud Rise: Site 690, 528b, 633b
- Chiasmolithus expansus*
Maud Rise: Site 689, 665b
Maud Rise: Site 690, 633b
- Chiasmolithus oamaruensis*

PALEONTOLOGICAL INDEX

- Maud Rise: Site 689, 665b
magnetostratigraphy, 654b
- Maud Rise: Site 690, magnetostratigraphy, 654b
- Chiasmolithus solitus*
Maud Rise: Site 689, magnetostratigraphy, 655b
- Maud Rise: Site 690, 634b
magnetostratigraphy, 655b
- Chiasmolithus* sp., Maud Rise: Site 690, 634b
- Chiastozygus garrisonii*, Maud Rise: Site 690, 481b
- Chiastozygus* sp., Maud Rise: Site 690, 532b
- Chiloguembelina cubensis*, Maud Rise: Site 689, 567b
- Chiloguembelina cubensis* Interval Subzone
Maud Rise: Site 689, 554b
Maud Rise: Site 690, 554b
- Chiloguembelina taurica*, Maud Rise: Site 689, 564b
- Chiloguembelina wilcoxensis*, Maud Rise: Site 689, 564b
- Chrysophycean cysts, Dronning Maud Land margin: Site 693, 423b-424b
- Cibicidoides*
Maud Rise: Site 689
carbon isotope data, 868b-870b
oxygen isotope data, 867b-869b
- Maud Rise: Site 690
carbon isotope data, 868b-870b
oxygen isotope data, 867b-869b
- Cribrosphaerella danaie*
Maud Rise: Site 690, 532b
abundance, 523b
- Clausicoccus fenestratus*
Maud Rise: Site 689, abundance, 656b
Maud Rise: Site 690, abundance, 656b
- Coccolithus formosus*
Maud Rise: Site 689, abundance, 656b-657b
Maud Rise: Site 690, abundance, 656b-657b
- Coccolithus pelagicus*
Maud Rise: Site 689, 662b
Maud Rise: Site 690, 662b
- Codoniella campanulata*, Dronning Maud Land margin: Site 692, 464b
- Collospaenid* sp.
Maud Rise: Site 689, 690b, 701b
Maud Rise: Site 690, 690b
- Cometodinium comatum*, Dronning Maud Land margin: Site 692, 463b
- Contingisporites* sp., Dronning Maud Land margin: Site 692, 458b
- Corbisema apiculata*, Maud Rise: Site 689, 748b
- Corbisema hastata*, Maud Rise: Site 689, 748b
- Corbisema inermis* *inermis*, Maud Rise: Site 689, 748b
- Corbisema triacantha*, Maud Rise: Site 689, 748b, 754b
- Cornua*
Dronning Maud Land margin: Site 693, 432b
historical background, 428b-429b
- Corollina* sp., Dronning Maud Land margin: Site 692, 460b
- Corollithion covingtonii* n. sp., Dronning Maud Land margin: Site 692, 338b, 349b
- Corollithion rhombicum*, Maud Rise: Site 690, 481b
- Coryphostoma incrassata*
Maud Rise: Site 689, 826b
Maud Rise: Site 690, 825b
- Corythospyris fiscella*
Maud Rise: Site 689, 695b, 705b
Maud Rise: Site 690, 695b
- Corythospyris* sp.
Maud Rise: Site 689, 695b, 705b
Maud Rise: Site 690, 695b
- Cosmiodiscus insignis*, Jane Basin, abundance, 806b-808b
- Cosmiodiscus insignis* Zone, Jane Basin, 807b
- Cosmiodiscus intersectus* Partial-Range Zone, 768b
- Cretarhabdus conicus*, Maud Rise: Site 690, 481b, 486b
- Cretarhabdus crenulatus*, Dronning Maud Land margin: Site 692, 348b
- Cretarhabdus surirellus*, Maud Rise: Site 690, 481b
- Cribrosphaerella danae*, Maud Rise: Site 690, 482b
- Cribrosphaerella danae* Subzone, Maud Rise Site 690, 466b-467b
- Cribrosphaerella ehrenbergii*, Maud Rise Site 690, 482b
- Cromyodruppa concentrica*, Maud Rise: Site 689, 355b, 359b
- Crossophialus glabrus* n. sp., Dronning Maud Land margin: Site 693, 409b, 417b
- Crossophialus gyroscolus* n. sp., Dronning Maud Land margin: Site 693, 409b, 417b
- Crucidenticula nicobarica*, Maud Rise: Site 689, 787b
- Crucidenticula punctata*, Maud Rise Site 689, 788b
- Cruciplacolithus primus/tenuis*, Maud Rise: Site 690, abundance, 523b
- Cruciplacolithus* sp.
Maud Rise: Site 690, 483b
abundance, 524b
- Cyathidites australis*, Dronning Maud Land margin: Site 692, 459b
- Cyathidites* sp., Dronning Maud Land margin: Site 692, 460b
- Cycadopites nitidus*, Dronning Maud Land margin: Site 692, 459b
- Cycadopites* sp., Dronning Maud Land margin: Site 692, 458b
- Cycladophora antiqua* n. sp.
Maud Rise: Site 689, 698b, 707b
Maud Rise: Site 690, 698b
- Cycladophora antiqua* Zone
Maud Rise: Site 689, 680b
Maud Rise: Site 690, 680b
- Cycladophora golli regipileus* Zone
Maud Rise: Site 689, 679b
Maud Rise: Site 690, 679b
- Cycladophora humerus* Zone
Maud Rise: Site 689, 679b
Maud Rise: Site 690, 679b
- Cycladophora spongotorax* Zone, Maud Rise: Site 690, 710b-711b
- Cyclagelosphaera margerelii*, Dronning Maud Land margin: Site 692, 347b
- Cyclagelosphaera reinhardtii*, Maud Rise: Site 690, 531b
- Cyclargolithus abisectus*, Maud Rise: Site 689, 664b
- Cyclargolithus floridanus*
Maud Rise: Site 689, 663b, 664b
magnetostratigraphy, 655b
Maud Rise: Site 690, magnetostratigraphy, 655b
- Cymatosphaera* sp., Dronning Maud Land margin: Site 692, 464b
- Cyphellachetes intricatus* n. sp., Dronning Maud Land margin: Site 693, 409b, 415b
- Cyrtocapsella longithorax* Zone
Maud Rise: Site 689, 679b
Maud Rise: Site 690, 679b
- Cyrtocapsella robusta*
Maud Rise: Site 689, 696b, 705b
Maud Rise: Site 690, 696b
- Cyrtocapsella robusta* Zone, Maud Rise: Site 689, 680b
- Dacrydiumites florinii*, South Orkney microcontinent: Site 696, 607b
- Dasyangea dactylethra* n. sp., Dronning Maud Land margin: Site 693, 409b-410b, 416b
- Deflandrea antarctica*, South Orkney microcontinent: Site 696, 612b
- Dendrospyris megalcephalis* Zone
Maud Rise: Site 689, 679b
Maud Rise: Site 690, 679b
- Denticulopsis hustedtii/Nitzschia gross punctata* Concurrent-Range Zone, 769b
- Denticulopsis macCollumii* Partial-Range Zone, 769b
- Denticulopsis praedimorpha* Partial-Range Zone, 768b
- Diacyanthocapsa* sp., Dronning Maud Land margin: Site 693, 355b-356b, 361b
- Diatoms
Antarctica
zonation, 762b-766b
See also specific zones
- Dronning Maud Land margin: Site 692
abundance, 309a
biostratigraphy, 311a
- Dronning Maud Land margin: Site 693
abundance, 361a-363a, 773b-774b, 776b-777b
age-depth range, 927b
Australian diatoms similar to, 378b
biostratigraphy, 366a-367a, 775b, 926b
dating, 365b-368b, 404b-405b
location, 368b
opaline, 366b, 378b
origin, 411b-412b
perforate process, 369b, 380b
pseudoloculate structure, 379b, 380b
punctate-haplo process, 368b-369b
pyritized, 366b
rhyncho-shaped process, 369b
scissure-haplo process, 369b, 379b, 380b-381b
siphon-shaped process, 369b
- Jane Basin
abundance, 718a-721a, 724a-725a, 778b-779b, 784b
age-depth range, 935b
biostratigraphy, 722a-723a, 782b, 934b
sea-ice expansion, 956b-957b
- Maud Rise: Site 689
abundance, 116a-117a, 769b-772b
age range, 767b
biostratigraphy, 110a-111a
geomagnetic polarity pattern correlated with, 922b
stratigraphic datum points, 771b
- Maud Rise: Site 690
abundance, 221a-222a, 772b-773b
age range, 767b
biostratigraphy, 217a-218a
geomagnetic polarity pattern correlated with, 923b-924b
stratigraphic datum points, 774b
- South Orkney microcontinent: Site 695
abundance, 548a-551a, 775b, 779b

- age-depth range, 931b
 biostratigraphy, 552a-553a, 778b, 930b
 South Orkney microcontinent: Site 696
 abundance, 638a-639a, 775b, 778b, 781b
 age-depth range, 933b
 biostratigraphy, 633a-635a, 780b, 932b
 Weddell Basin: Site 694
 abundance, 472a-473a, 476a, 774b-775b
 age-depth range, 929b
 biostratigraphy, 474a-475a, 778b, 928b
 zonation
 first abundant appearance, 918b
 first appearance, 918b
 geomagnetic polarity pattern correlated with, 921b
 geomagnetic time scale correlated with, 917b
 last abundant appearance, 918b
 last appearance, 918b
See also specific types
Dichotriaenes
 Maud Rise: Site 689, 965b-966b, 969b
 South Orkney microcontinent: Site 696, 965b-966b, 970b
Diconodinium davidii, Dronning Maud Land margin: Site 692, 464b
Dictyocha aspera, Maud Rise: Site 690, 748b, 755b
Dictyocha calida, Maud Rise: Site 689, 748b, 755b
Dictyocha deflandrei, Maud Rise: Site 689, 748b, 755b
Dictyocha frenguelli, Maud Rise: Site 689, 748b, 755b
Dictyocha pentagona, Maud Rise: Site 689, 748b, 754b
Dictyocha spinosa, Maud Rise: Site 689, 748b, 755b
Dictyomitra sp.
 Dronning Maud Land margin: Site 693, 356b, 360b
 Maud Rise: Site 689, 356b, 360b
Didymocystis sp.
 Maud Rise: Site 689, 692b, 703b
 Maud Rise: Site 690, 692b
Dingodinium cerviculum, Dronning Maud Land margin: Site 692, 464b
Dinoflagellate cysts
 Dronning Maud Land margin: Site 692
 dating, 452b
 depth and, 454b
 Dronning Maud Land margin: Site 693, taxonomic list, 601b
 South Orkney microcontinent: Site 696
 abundance, 600b
 dating, 596b-597b, 602b
 kerogen analyses, 603b-604b
 taxonomic list, 601b
Dinoflagellates, Dronning Maud Land margin: Site 693, siliceous microfossil taxa, 405b
Discoaster bifax, Maud Rise: Site 690, 635b
Discoaster deflandrei, Maud Rise: Site 690, 634b
Discoaster kuepperi, Maud Rise: Site 690, 634b
Discoaster lenticularis, Maud Rise: Site 690, 634b
Discoaster lodoensis, Maud Rise: Site 690, 634b
Discoaster multiradiatus, Maud Rise: Site 690, 634b
Discoaster nobilis, Maud Rise: Site 690, 635b
Discoaster sublodoensis, Maud Rise: Site 690, 635b
Discoasters, Maud Rise: Site 689, abundance, 621b-623b
Disolenia sp.
 Maud Rise: Site 689, 690, 701b
 Maud Rise: Site 690, 690b
Distephanus boliviensis
 Maud Rise: Site 689, 748b-749b, 756b
 Maud Rise: Site 690, 748b-749b
Distephanus speculum hemisphaericus, Maud Rise: Site 689, 749b, 757b
Distephanus speculum pentagonus, Maud Rise: Site 689, 749b-750b, 756b
Distephanus speculum speculum
 Maud Rise: Site 689, 756b-759b
 Maud Rise: Site 690, 749b-750b, 757b
Druppatractus hastatus
 Maud Rise: Site 689, 691b, 702b
 Maud Rise: Site 690, 691b
Ebridians
 Dronning Maud Land margin: Site 693, abundance, 737b
 Maud Rise: Site 689, abundance, 735b
 Maud Rise: Site 690, abundance, 736b
 South Orkney microcontinent: Site 695, abundance, 738b
Eiffeliithus turriseiffeli, Maud Rise: Site 690, 483b, 532b
Eoglobigerina edita, Maud Rise: Site 690, 563b
Eoglobigerina eobulloides, Maud Rise: Site 690, 563b
Eoglobigerina fringa Partial Range Zone
 Maud Rise: Site 689, 551b
 Maud Rise: Site 690, 551b
Eucampia antarctica
 Jane Basin
 abundance, 804b, 806b-808b
 opal correlated with, 808b
 sea-ice expansion, 957b
Eucyrtidium calvertense
 Maud Rise: Site 689, 696b, 706b
 Maud Rise: Site 690, 696b
Eucyrtidium punctatum Zone
 Maud Rise: Site 689, 679b
 Maud Rise: Site 690, 679b
Fasciculinthus
 Maud Rise: Site 689, 621b
 Maud Rise: Site 690, 621b
Fasciculinthus involutus, Maud Rise: Site 690, 635b
Foraminifers
 Dronning Maud Land margin: Site 693, organic geochemistry data, 886b-888b, 894b-895b
 Maud Rise: Site 689
 calcium carbonate accumulation rates, 841b
 carbon isotope data, 814b, 831b-832b
 carbon isotopic differences, 818b
 carbon isotopic temperature gradients, 871b, 873b
 chemical preservation, 816b
 isotopic preservation, 816b, 818b
 organic geochemistry data, 884b-885b
 oxygen isotope data, 814b, 831b-832b
 oxygen isotopic differences, 818b
 oxygen isotopic temperature gradients, 870b-872b
 paleotemperature, 821b
 Proteus Ocean, 876b-877b
 saline deep waters, 874b
 sea-surface temperatures, 843b
 Sr/Ca ratios, 817b
 surface water temperature, 822b
 textural preservation, 816b
 thermocline, 842b-843b
Maud Rise: Site 690
 calcium carbonate accumulation rates, 841b
 carbon isotope data, 815b, 831b
 carbon isotopic differences, 818b
 carbon isotopic temperature gradients, 871b-873b
 chemical preservation, 816b
 isotopic preservation, 816b, 818b
 organic geochemistry data, 885b-886b
 oxygen isotope data, 815b, 831b, 833b
 oxygen isotopic differences, 818b
 oxygen isotopic temperature gradients, 871b-872b
 paleotemperature, 821b
 Proteus Ocean, 876b-877b
 saline deep waters, 874b
 sea-surface temperatures
 Sr/Ca ratios, 817b
 surface water temperature, 822b
 textural preservation, 816b
 thermocline, 842b-843b
Weddell Basin: Site 694, organic geochemistry data, 888b-889b, 896b-897b
Foveoporites subtriangularis, Dronning Maud Land margin: Site 692, 459b
Gartnerago diversum, Maud Rise: Site 690, 483b
Gavelinella beccariformis, Maud Rise: Site 690, 824b-825b
Gavelinella sp.
 Maud Rise: Site 689, 847b
 Maud Rise: Site 690, 847b
Gladiopsis ellipsoidea n. sp., Dronning Maud Land margin: Site 693, 372b
Gladiopsis jouseana n. comb., Dronning Maud Land margin: Site 693, 372b
Gladiopsis lagenoides n. sp., Dronning Maud Land margin: Site 693, 372b, 393b-394b, 400b
Gladiopsis maxima n. comb., Dronning Maud Land margin: Site 693, 372b
Gladiopsis modica n. sp., Dronning Maud Land margin: Site 693, 372b, 393b-394b
Gladiopsis pacifica n. comb., Dronning Maud Land margin: Site 693, 372b
Gladiopsis speciosa n. comb., Dronning Maud Land margin: Site 693, 372b
Gladius antiquus, Dronning Maud Land margin: Site 693, 373b, 390b-391b, 398b, 400b
Gladius perfectus, Dronning Maud Land margin: Site 693, 373b-374b, 387b, 398b, 402b
Gladius rhizophorus n. sp., Dronning Maud Land margin: Site 693, 374b, 390b, 392b, 400b, 402b
Globigerina eocaena, Maud Rise: Site 690, 569b
Globigerina euapertura, Maud Rise: Site 690, 569b
Globigerina euapertura Partial Range Subzone
 Maud Rise: Site 689, 554b
 Maud Rise: Site 690, 554b
Globigerina index, Maud Rise: Site 690, 569b
Globigerina labiacrassata, Maud Rise: Site 690, 569b
Globigerina labiacrassata Interval Zone
 Maud Rise: Site 689, 554b
 Maud Rise: Site 690, 554b
Globigerina lozanoi, Maud Rise: Site 689, 569b

PALEONTOLOGICAL INDEX

Globigerina sp., Maud Rise: Site 689, 567b
Globigerinatetha index Partial Range Zone
 Maud Rise: Site 689, 553b
 Maud Rise: Site 690, 553b
Globigerinelloides blowi, Dronning Maud Land margin: Site 693, 321b, 323b
Globigerinelloides impensus
 Maud Rise: Site 689, last occurrence, 499b
 Maud Rise: Site 690, 503b, 508b
 last occurrence, 499b
Globigerinelloides impensus Partial Range Zone
 Maud Rise: Site 689, distribution, 496b
 Maud Rise: Site 690, distribution, 496b
Globigerinelloides multispinatus, Maud Rise: Site 690, 503b, 508b, 848b
Globigerinelloides sp.
 Maud Rise: Site 689, depth habitat, 820b
 Maud Rise: Site 690, depth habitat, 820b
Globigerinelloides subcarinatus, Maud Rise: Site 690, 503b, 508b
Globoconusa daubjergensis Partial Range Sub-zone
 Maud Rise: Site 689, 551b
 Maud Rise: Site 690, 551b
Globorotalia polycamera, Maud Rise: Site 689, 563b
Globorotalites conicus, Maud Rise: Site 690, 593b
Globorotaloides suteri, Maud Rise: Site 689, 569b
Globorotaloides suteri Partial Range Zone
 Maud Rise: Site 689, 553b
 Maud Rise: Site 690, 553b
Globotruncana bulloides, Maud Rise: Site 690, 506b, 512b
Globotruncana subcircumnodifer, Maud Rise: Site 690, 506b, 512b
Globotruncanella havanensis
 Maud Rise: Site 689, 505b, 512b
 first occurrence, 499b
 Maud Rise: Site 690, first occurrence, 499b
Globotruncanella havanensis Interval Zone
 Maud Rise: Site 689, distribution, 497b
 Maud Rise: Site 690, distribution, 497b
Globotruncanella petaloidea, Maud Rise: Site 690, 505b, 511b
Gondwanaria sp.
 Maud Rise: Site 689, 697b, 707b
 Maud Rise: Site 690, 697b
Gorgansium sp., Dronning Maud Land margin: Site 693, 356b, 362b
Gubkinella graysonensis, Dronning Maud Land margin: Site 693, 321b, 323b
Gublerina robusta, Maud Rise: Site 690, 503b, 508b
Guembelitria cretacea
 Maud Rise: Site 689, 502b-503b
 Maud Rise: Site 690, 502b-503b, 508b
Haloragacidites harrisii, South Orkney microcontinent: Site 696, 608b
Hapsocysta peridictya, Dronning Maud Land margin: Site 692, 464b
Hedbergella holmdelensis, Maud Rise: Site 690, 503b, 509b, 513b
Hedbergella infracretacea, Dronning Maud Land margin: Site 693, 321b, 324b
Hedbergella monmouthensis, Maud Rise: Site 690, 503b-504b, 509b, 513b
Hedbergella sliteri n. sp., Maud Rise: Site 690, 504b, 509b, 513b
Hedbergella trocoidea, Dronning Maud Land margin: Site 693, 321b-322b, 323b

Heliolithus cantabriae, Maud Rise: Site 690, 635b
Heliolithus riedelii, Maud Rise: Site 690, 635b
Heterohelix dentata, Maud Rise: Site 690, 503b, 508b
Heterohelix globulosa, Maud Rise: Site 690, 503b, 508b
Heterohelix planata, Maud Rise: Site 690, 503b, 508b
Heterosphaeridium heteracanthum, South Orkney microcontinent: Site 696, 610b
Hexastylus sp.
 Maud Rise: Site 689, 691b, 701b
 Maud Rise: Site 690, 691b
Hibolithes sp., Dronning Maud Land margin: Site 692, 443b-444b, 448b
Hornbrookina
 Maud Rise: Site 689, 621b
 Maud Rise: Site 690, 621b
Hornbrookina australis, Maud Rise: Site 690, 636b
Hornbrookina edwardsii
 Maud Rise: Site 690, 525b, 529b
 abundance, 523b
Hornbrookina teuriensis, Maud Rise: Site 690, 636b
Hyalotrochus incompositus n. sp., Dronning Maud Land margin: Site 693, 410b, 418b
Hyalotrochus radiatus n. sp., Dronning Maud Land margin: Site 693, 410b, 418b, 425b
Hystrichosphaeridium astertes, South Orkney microcontinent: Site 696, 609b
Igorina spiralis, Maud Rise: Site 689, 565b
Inoceramus sp., Dronning Maud Land margin: Site 692, 444b, 448b
Ischyosporites gremius, South Orkney microcontinent: Site 696, 607b
Isthmolithus recurvus
 Maud Rise: Site 689, magnetostratigraphy, 653b
 Maud Rise: Site 690, magnetostratigraphy, 653b
Isthmolithus recurvus Zone, Maud Rise: Site 689, 643b
Kamptnerius magnificus, Maud Rise: Site 690, 483b, 532b
Kerkis bispinosa n. sp., Dronning Maud Land margin: Site 693, 374b-375b, 386b, 393b
Kleithriaspheeridium sp., Dronning Maud Land margin: Site 692, 461b
Kreagra bifalcata n. sp., Dronning Maud Land margin: Site 693, 375b, 399b
Kreagra forfex n. sp., Dronning Maud Land margin: Site 693, 375b, 399b
Kreagra sp. A, Dronning Maud Land margin: Site 693, 375b
Kreagra tripartita n. sp., Dronning Maud Land margin: Site 693, 375b, 399b
Lapideacassis mariae, Maud Rise: Site 690, 484b
Lapideacassis sp., Maud Rise: Site 690, 530b
Lapideacassis tricornus, Maud Rise: Site 690, 484b
Leiosphaera sp., Dronning Maud Land margin: Site 692, 458b
Leptolepidites, Dronning Maud Land margin: Site 692, 460b
Lithomelissa robusta
 Maud Rise: Site 689, 695b, 705b
 Maud Rise: Site 690, 695b
Lucianorhabdus arborius, Maud Rise: Site 690, 487b
Lyamula, Dronning Maud Land margin: Site 693, abundance, 430b-431b
Lychnocanoma conica
 Maud Rise: Site 689, 697b, 706b-707b
Lychnocanoma sp.
 Maud Rise: Site 689, 697b, 706b-707b
 Maud Rise: Site 690, 697b
Markalius apertus, Maud Rise: Site 690, 636b
Markalius inversus
 Maud Rise: Site 689, 665b, 666b
 Maud Rise: Site 690, 530b
 abundance, 523b
Meristosolen cylindratus n. sp., Dronning Maud Land margin: Site 693, 410b, 416b
Microorbis convexus n. sp., Dronning Maud Land margin: Site 693, 376b, 385b, 397b, 402b
Microorbis undulatus n. sp., Dronning Maud Land margin: Site 693, 376b, 385b, 397b
Micula decussata, Maud Rise: Site 690, 484b
Micula murus, Maud Rise: Site 690, first occurrence, 473b
Misceomarginatus pleniporus, Maud Rise: Site 690, 484b
 Molluscs, Dronning Maud Land margin: Site 692, abundance, 310a
Monomarginatus quaternarius, Maud Rise: Site 690, 484b
Muricoglobigerina aquiensis, Maud Rise: Site 689, 565b
Muricoglobigerina mckannai Interval Zone
 Maud Rise: Site 689, 553b
 Maud Rise: Site 690, 553b
Nannotetraena cristata, Maud Rise: Site 689, 636b
Nannotetraena fulgens, Maud Rise: Site 689, 636b
Navicula ponticulus, Maud Rise: Site 689, 754b
Naviculopsis constricta, Maud Rise: Site 689, 750b, 754b
Naviculopsis eobiapiculata, Maud Rise: Site 689, 750b, 754b
Neococcolithes dubius
 Maud Rise: Site 689, 665b
 Maud Rise: Site 690, 637b
Neococcolithes protenus, Maud Rise: Site 690, 636b-637b
Neocrepidolithus frequens
 Maud Rise: Site 690, 474b
 abundance, 475b
Neocrepidolithus sp.
 Maud Rise: Site 690, 530b-531b
 abundance, 523b
Neocrepidolithus watkinsii, Maud Rise: Site 690, 474b, 485b
Neoepionides hillebrandti, Maud Rise: Site 690, 593b
Neoepionides lunata, Maud Rise: Site 690, 593b
Neogloboquadrina pachyderma, Jane Basin, climatic cycles, 957b
Neoraistrickia levidensis, Dronning Maud Land margin: Site 692, 460b
Nephrolithus corystus, Maud Rise: Site 690, 485b
Nephrolithus frequens
 Maud Rise: Site 690
 abundance, 523b
 first occurrence, 473b

- Nephrolithus frequens*, Maud Rise:
Site 690, 532b
- Nephrolithus frequens miniporus*, Maud Rise:
Site 690, 474b, 475b
- Nephrolithus frequens* Zone
Maud Rise: Site 689, 466b
Maud Rise: Site 690, 466b
- Nitzschia aurica*, Maud Rise: Site 689, 786b
- Nitzschia barronii* Partial-Range Zone, 767b-768b
- Nitzschia cylindrica*, Maud Rise: Site 689, 785b
- Nitzschia denticuloides* Partial-Range Zone, 768b
- Nitzschia fossilis*, Maud Rise: Site 689, 785b
- Nitzschia grossepunctata* Partial-Range Zone, 769b
- Nitzschia interfrigidaria*, Maud Rise: Site 689, 785b
- Nitzschia interfrigidaria/Cosmodiscus insignis* Concurrent Range Zone, 767b
- Nitzschia maleinterpretaria* Partial-Range Zone, 769b
- Nitzschia praecurta* Partial-Range Zone, 768b
- Nitzschia pusilla*, Maud Rise: Site 689, 786b
- Nitzschia reinholdii*, Maud Rise: Site 689, 786b
- Nothofagidites* sp., South Orkney microcontinent: Site 696, 610b
- Nuttallides*
Maud Rise: Site 689
carbon isotope data, 869b-871b
oxygen isotope data, 868b-869b, 871b
- Maud Rise: Site 690
carbon isotope data, 868b-871b
oxygen isotope data, 868b-869b, 871b
- Nuttallides* sp.
Maud Rise: Site 689, 847b
Maud Rise: Site 690, 847b
- Nuttallides truempyi*, Maud Rise: Site 690, 594b, 824b
- Obliquipithonella krasheninnikovii* n. comb.,
Maud Rise: Site 689, 540b, 545b
- Obliquipithonella operculata* n. comb., Maud Rise: Site 689, 540b, 545b
- Octocyclus magnus*, Maud Rise: Site 690, 486b
- Odontochitina porifera*, South Orkney microcontinent: Site 696, 610b
- Oligosphaeridium complex*, Dronning Maud Land margin: Site 692, 458b
- Operculodinium bergmannii*, South Orkney microcontinent: Site 696, 612b
- Oridorsalis nitidus*, Maud Rise: Site 690, 594b
- Orthipithonella aspera*, Maud Rise: Site 690, 548b
- Orthopithonella aspera* n. sp.
Maud Rise: Site 689, 538b, 548b
Maud Rise: Site 690, 538b
- Orthopithonella congruens* n. sp.
Maud Rise: Site 689, 539b, 547b
distribution, 539b
Maud Rise: Site 690, 539b, 547b
distribution, 539b
- Orthopithonella flora* n. sp.
Maud Rise: Site 689, 538b, 546b
Maud Rise: Site 690, 538b, 546b
- Orthopithonella gustafsonii*
Maud Rise: Site 689, 538b-539b, 546b
distribution, 539b
Maud Rise: Site 690, 538b-539b, 546b
distribution, 539b
- Orthopithonella minutia* n. sp., Maud Rise: Site 689, 537b-538b, 544b
- Orthopithonella* sp., Maud Rise: Site 689, 539b-540b, 544b
- Oxea*, Maud Rise: Site 689, 965b, 968b
- Oxyasters*
Maud Rise: Site 689, 965b
South Orkney microcontinent: Site 696, 965b, 970b
- Oxytomidae*, Dronning Maud Land margin: Site 692, 444b, 448b
- Palynomorphs*
Dronning Maud Land margin: Site 692
biostratigraphy, 311a, 452b-453b
dating, 449b-450b
paleoclimate, 454b-455b
palynoflora, 451b-452b
sedimentation, 455b-456b
vegetational cover, 454b-455b
- Dronning Maud Land margin: Site 693
absence, 595b, 598b-599b
biostratigraphy, 453b-454b
dating, 449b-450b
sedimentation, 455b-456b
taxonomic list, 601b
- South Orkney microcontinent: Site 696,
taxonomic list, 601b
- See also specific types*
- Planktonic foraminifers*
Austral Province, paleobiogeography, 500b-501b
- Dronning Maud Land margin: Site 692
abundance, 307a
biostratigraphy, 311a
- Dronning Maud Land margin: Site 693
abundance, 320b-321b, 359a
dating, 319b-320b
distribution, 320b
first appearance, 320b
last appearance, 320b
paleoenvironment, 320b-321b
preservation, 320b
- Maud Rise: Site 689
abundance, 109a-112a, 490b, 492b
age, 498b
benthic ratios, 491b
biochronology, 497b-500b
biostratigraphy, 110a-111a, 490b, 551b-552b
- carbon burial, 843b-844b
carbon isotope changes, 861b-862b
carbon isotope data, 858b-859b, 947b
carbon isotopic temperature gradients, 839b-841b
- chronostratigraphy, 554b-558b
climatic evolution, 859b-861b
depth and, 498b
depth habitat, 820b
distribution, 495b
extinction, 844b
first appearance, 911b
last appearance, 911b
oxygen isotope correction factors, 835b-836b
- oxygen isotope data, 852b, 854b, 856b, 860b, 946b, 948b
oxygen isotopic temperature gradients, 836b, 838b-839b
- paleobiogeography, 500b-501b
paleoenvironment, 501b-502b
sedimentation rates, 501b
- Site 690 depths compared with, 906b-908b
size, 496b-497b
- zonation, 496b-497b, 498b, 549b-550b
- South Orkney microcontinent: Site 695,
abundance, 547a
- South Orkney microcontinent: Site 696
abundance, 632a
biostratigraphy, 633a-635a
- See also specific types*
- Planorotalites australiformis*, Maud Rise: Site 690, 567b
- Planorotalites australiformis* Interval Zone
Maud Rise: Site 689, 553b
Maud Rise: Site 690, 553b
- Planorotalites imitatus* Interval Zone
Maud Rise: Site 689, 551b, 553b
Maud Rise: Site 690, 551b, 553b
- Podocarpidites* sp., South Orkney microcontinent: Site 696, 607b
- Pollen*
Dronning Maud Land margin: Site 692, depth and, 453b
- South Orkney microcontinent: Site 696, taxonomic list, 601b
- See also Palynomorphs*
- Polycingulatisporites striatus*, Dronning Maud Land margin: Site 692, 459b
- Praethalassiosiropora hasleae* n. sp., Dronning Maud Land margin: Site 693, 376b-377b, 384b-385b
- Prediscosphaera*, Maud Rise: Site 690, abundance, 518b-519b
- Prediscosphaera cretacea*, Maud Rise: Site 690, 486b, 532b
- Prediscosphaera spinosa*, Maud Rise: Site 690, 486b
- Prediscosphaera stoveri*
Maud Rise: Site 690, 486b, 525b-526b, 532b
abundance, 523b
- Prinsius dimorphosus*, Maud Rise: Site 690, 529b, 637b
- Prinsius martinii*, Maud Rise: Site 690, 637b
- Propylipollis* sp., South Orkney microcontinent: Site 696, 608b
- Proteacidites rynthius*, South Orkney microcontinent: Site 696, 608b
- Prunopyle* sp.
Maud Rise: Site 689, 693b-694b, 703b

PALEONTOLOGICAL INDEX

Maud Rise: Site 690, 693b-694b
Pseudodictyonitria pentacolaensis, Dronning Maud Land margin: Site 693, 356b, 360b, 363b
Pseudohastigerina micra, Maud Rise: Site 689, 567b
Pseudohastigerina micra Interval Zone
 Maud Rise: Site 689, 553b
 Maud Rise: Site 690, 553b
Pseudohastigerina wilcoxensis, Maud Rise: Site 690, 567b
Pseudopyxilla furcata n. sp., Dronning Maud Land margin: Site 693, 410b, 415b
Pseudopyxilla systema n. sp., Dronning Maud Land margin: Site 693, 410b, 415b
Pseudotextularia elegans, Maud Rise: Site 689, 503b, 508b
Psyktsphaera, Maud Rise: Site 690, 474b, 482b
Psyktsphaera firthii, Maud Rise: Site 690, 474b-476b, 482b
Pterospermella aureolata, Dronning Maud Land margin: Site 692, 463b

Radiolarians
 Dronning Maud Land margin: Site 692 abundance, 310a
 biostratigraphy, 311a
 Dronning Maud Land margin: Site 693 abundance, 354b, 363a-364a, 718b-719b age-depth range, 927b
 biostratigraphy, 366a-367a, 926b
 Jane Basin
 abundance, 725a
 age-depth range, 935b
 biostratigraphy, 722a-723a, 934b
 Maud Rise: Site 689 abundance, 117a-119a, 354b, 680b-688b, 714b-715b
 biostratigraphy, 110a-111a
 geomagnetic polarity pattern correlated with, 922b
 zonation, 676b-678b
 Maud Rise: Site 690 abundance, 222a-223a, 688b-690b
 biostratigraphy, 217a-218a
 geomagnetic polarity pattern correlated with, 681b, 923b-924b
 zonation, 676b-678b
 South Orkney microcontinent: Site 695 abundance, 551a, 554a
 age-depth range, 931b
 biostratigraphy, 552a-553a, 930b
 South Orkney microcontinent: Site 696 abundance, 639a-640a
 age-depth range, 933b
 biostratigraphy, 633a-635a, 932b
 Weddell Basin: Site 694
 abundance, 476a
 age-depth range, 929b
 biostratigraphy, 474a-475a, 928b
 zonation
 first abundant appearance, 919b
 first appearance, 919b
 geomagnetic polarity pattern correlated with, 921b
 geomagnetic time scale correlated with, 917b
 last abundant appearance, 919b
 last appearance, 919b
Radulopecten sp., Dronning Maud Land margin: Site 692, 444b, 448b
Reinhardtites anthophorus, Maud Rise: Site 690, 486b

Reinhardtites levis, Maud Rise: Site 690, last occurrence, 473b
Reticulofenestra bisecta
 Maud Rise: Site 689, 664b, 665b
 abundance, 656b
 Maud Rise: Site 690, abundance, 656b
Reticulofenestra daviesii, Maud Rise: Site 689, 663b, 666b
Reticulofenestra daviesii Zone, Maud Rise: Site 689, 642b-643b
Reticulofenestra gelida, Maud Rise: Site 689, 664b
Reticulofenestra hesslandii, Maud Rise: Site 689, 663b, 664b
Reticulofenestra oamaruensis
 Maud Rise: Site 689, 664b
 magnetostratigraphy, 653b
Reticulofenestra oamaruensis Zone, Maud Rise: Site 689, 643b
Reticulofenestra perplexa, Maud Rise: Site 689, 663b, 664b
Reticulofenestra pseudoumbilica, Maud Rise: Site 689, 664b
Reticulofenestra reticulata
 Maud Rise: Site 689, 666b
 magnetostratigraphy, 654b
 Maud Rise: Site 690, magnetostratigraphy, 654b
Reticulofenestra reticulata Zone, Maud Rise: Site 689, 643b
Reticulofenestra samodurovii, Maud Rise: Site 689, 665b, 666b
Reticulofenestra umbilica
 Maud Rise: Site 689, 665b
 magnetostratigraphy, 652b
 Maud Rise: Site 690, magnetostratigraphy, 652b
Retiriteles, South Orkney microcontinent: Site 696, 607b
Rhagodiscus angustus Zone, Dronning Maud Land margin: Site 693, black shale, 336b
Rhagodiscus asper, Dronning Maud Land margin: Site 692, 349b, 350b
Rhoiphites sphaerica, South Orkney microcontinent: Site 696, 608b
Rhynchopyxis siphonoides n. sp., Dronning Maud Land margin: Site 693, 377b, 385b-386b
Rorularia sp., Dronning Maud Land margin: Site 692, 445b, 448b
Rouxia sp., Maud Rise: Site 689, 789b
Rugotruncana circumnodifer, Maud Rise: Site 690, 505b, 511b, 513b
Satyrodiinium sp., South Orkney microcontinent: Site 696, 610b
Schackoina multispinata, Maud Rise: Site 690, 503b, 509b
Senegaliniun asymmetricum, South Orkney microcontinent: Site 696, 611b
Senoniaphaera ptomatis, Dronning Maud Land margin: Site 692, 462b
Sentusidinium, Dronning Maud Land margin: Site 692, 461b
Sentusidinium sp., Dronning Maud Land margin: Site 692, 463b
Seribiscutum primitivum, Dronning Maud Land margin: Site 693, abundance, 335b
Sigmas
 Maud Rise: Site 689, 966b, 970b
 South Orkney microcontinent: Site 696, 966b, 970b
Siliceous sponge spicules

Dronning Maud Land margin Site 693, abundance, 965b
 Maud Rise: Site 689, abundance, 964b
 Weddell Basin: Site 694, abundance, 965b
Silicoflagellates
 Dronning Maud Land margin: Site 693 abundance, 364a-365a, 732b-733b, 737b age-depth range, 927b
 biostratigraphy, 366a, 926b
 dating, 427b-428b, 730b
 phylogeny, 430b-432b
 possible occurrences, 429b-430b
 Jane Basin
 abundance, 725a, 734b
 biostratigraphy, 723a, 934b
 dating, 730b
 Maud Rise: Site 689 abundance, 731b-732b, 734b-735b
 biostratigraphy, 110a-111a
 dating, 730b
 hiatus, 731b-732b
 Maud Rise: Site 690 abundance, 732b, 736b
 biostratigraphy, 217a-218a
 dating, 730b
 geomagnetic polarity pattern correlated with, 923b-924b
pseudofibula plexus, 743b
 paleoenvironmental relationships, 743b, 746b-747b
 stratigraphy, 743b-747b
 skeletal morphology, 735b, 738b-740b, 742b
 South Orkney microcontinent: Site 695 abundance, 547a, 733b, 738b
 biostratigraphy, 552a-553a, 930b
 dating, 730b
 South Orkney microcontinent: Site 696 abundance, 638a, 733b-734b, 739b age-depth range, 933b
 biostratigraphy, 633a-635a, 932b
 dating, 730b
 variability, 740b-741b, 743b
 Weddell Basin: Site 694
 abundance, 733b
 biostratigraphy, 928b
 dating, 730b
 zonation
 first abundant appearance, 920b
 first appearance, 920b
 geomagnetic polarity pattern correlated with, 921b
 geomagnetic time scale correlated with, 917b
 last abundant appearance, 920b
 last appearance, 920b
Siphogenerinoides eleganta, Maud Rise: Site 690, 592b
Sparganiaceaepollenites irregularis, South Orkney microcontinent: Site 696, 609b
 Sphenoliths, Maud Rise: Site 690, abundance, 621b-623b
Sphenolithus editus, Maud Rise: Site 690, 637b
Spinidinium essoi, South Orkney microcontinent: Site 696, 612b
Spinidinium luciae, South Orkney microcontinent Site 696, 611b
Spiticera sp., Dronning Maud Land margin: Site 692, 443b, 448b
Spores
 Dronning Maud Land margin: Site 692
 dating, 452b
 depth and, 453b
 South Orkney microcontinent: Site 696, taxonomic list, 601b

- Sporomorphs**
- South Orkney microcontinent: Site 696
 - abundance, 604b
 - dating, 596b-597b
 - See also specific types*
- Staplinisporites telatus**, Dronning Maud Land margin: Site 692, 459b
- Stichomitra** sp., Dronning Maud Land margin: Site 693, 356b-357b, 360b-361b, 363b
- Stilostomella annulifera**, Maud Rise: Site 690, 592b
- Strongyles**, Maud Rise: Site 689, 965b, 968b
- Stylatractus neptunus**
- Maud Rise: Site 689, 691b, 702b
 - Maud Rise: Site 690, 691b
- Stylatractus santaeannae**
- Maud Rise: Site 689, 691b-692b, 702b
 - Maud Rise: Site 690, 691b-692b
- Stylosphaera radiosa** Zone
- Maud Rise: Site 689, 680b
 - Maud Rise: Site 690, 680b
- Stylosphaera** sp.
- Maud Rise: Site 689, 692b, 702b
 - Maud Rise: Site 690, 692b
- Subbotina angiporoides**, Maud Rise: Site 689, 569b
- Subbotina angiporoides** Interval Zone
- Maud Rise: Site 689, 553b-554b
 - Maud Rise: Site 690, 553b-554b
- Subbotina eocaenica**, Maud Rise: Site 689, 565b
- Subbotina inaequispira** Partial Range Subzone
- Maud Rise: Site 689, 553b
 - Maud Rise: Site 690, 553b
- Subbotina inconstans**, Maud Rise: Site 690, 564b
- Subbotina inconstans** Interval Subzone
- Maud Rise: Site 689, 551b
 - Maud Rise: Site 690, 551b
- Subbotina linaperta**, Maud Rise: Site 689, 569b
- Subbotina moskovini**, Maud Rise: Site 690, 563b
- Subbotina pseudobulloides**, Maud Rise: Site 690, 564b
- Subbotina pseudobulloides** Interval Zone
- Maud Rise: Site 689, 551b
 - Maud Rise: Site 690, 551b
- Subbotina triangularis**, Maud Rise: Site 690, 565b
- Teichorhabdus ethmos**, Maud Rise: Site 690, 487b
- Testicarinata inconspicua**, Maud Rise: Site 689, 569b
- Thalassiosira complicata**, Maud Rise: Site 689, 788b
- Thalassiosira fraga**, Maud Rise: Site 689, 787b, 793b, 798b, 800b-801b
- Thalassiosira inura**, Maud Rise: Site 689, 789b
- Thalassiosira inura** Partial-Range Zone, 768b
- Thalassiosira kolbei** n. comb.
- Maud Rise: Site 689, 793b, 801b
 - Maud Rise: Site 690, 787b, 793b, 797b
- Thalassiosira kolbei** Partial-Range Zone, 766b-767b
- Thalassiosira lentiginosa** Partial-Range Zone, 766b
- Thalassiosira spinosa**, Maud Rise: Site 689, 788b
- Thalassiosira spumellaroides**, Maud Rise: Site 690, 793b-794b, 801b
- Thalassiosira spumellaroides** Partial-Range Zone, 769b
- Theocorys** sp., Dronning Maud Land margin: Site 693, 357b, 361b
- Tholoniid** sp.
- Maud Rise: Site 689, 694b, 704b
 - Maud Rise: Site 690, 694b
- Thoracosphaera operculata**, Maud Rise: Site 689, 540b-541b, 545b
- Thoracosphaera** sp.
- Maud Rise: Site 690, 531b
 - abundance, 523b
- Toweius crassus**, Maud Rise: Site 690, 638b
- Toweius craticulus**, Maud Rise: Site 690, 637b
- Toweius eminens**, Maud Rise: Site 690, 637b
- Toweius magnicrassus**, Maud Rise: Site 690, 638b
- Toweius tovae**, Maud Rise: Site 690, 638b
- Tribrachiatus**
- Maud Rise: Site 689, 621b
 - Maud Rise: Site 690, 621b
- Tribrachiatus contortus**, Maud Rise: Site 690, 638b
- Tribrachiatus orthostylus**, Maud Rise: Site 690, 638b
- Tricolocapsa** sp., Dronning Maud Land margin: Site 693, 357b, 362b
- Tricolporopollenites** sp., South Orkney microcontinent: Site 696, 609b-610b
- Triods**, Maud Rise: Site 689, 965b, 969b
- Triplopollenites** sp., South Orkney microcontinent: Site 696, 608b
- Trissocyklid** sp.
- Maud Rise: Site 689, 695b, 705b
 - Maud Rise: Site 690, 695b
- Trochus elegans** n. sp., Dronning Maud Land margin: Site 693, 377b, 397b-398b
- Tubodiscus verenae**, Dronning Maud Land margin: Site 692, 348b
- Tubotuberella flammingii**, Dronning Maud Land margin: Site 692, 462b
- Tubulifloridites antipodica**
- South Orkney microcontinent: Site 696, 610b
 - rare occurrence, 603b
- Turrilina alsatica**, Maud Rise: Site 690, 592b
- Tylostyles**, South Orkney microcontinent: Site 695, 965b, 969b
- Vallacerta hannahi**, Dronning Maud Land margin: Site 693, 433b, 441b-442b
- Variramus**
- Dronning Maud Land margin: Site 693, 432b
 - skeletal morphology, 429b
- Variramus aculeifera**, Dronning Maud Land margin: Site 693, 432b, 435b-440b
- Variramus loperi**, Dronning Maud Land margin: Site 693, 432b-433b, 438b, 442b
- Vekshinella guadriarculla**, Dronning Maud Land margin: Site 692, 346b
- Vekshinella** sp., Dronning Maud Land margin: Site 693, 351b
- Vekshinella stradneri**, Dronning Maud Land margin: Site 692, 346b
- Velicucullus altus** n. sp.
- Maud Rise: Site 689, 698b, 708b
 - Maud Rise: Site 690, 698b
- Wallodinium krutzschii**, Dronning Maud Land margin: Site 692, 463b
- Watznaueria barnesae**, Dronning Maud Land margin: Site 692, 347b
- Zeugrhabdotus** sp., Dronning Maud Land margin: Site 692, 350b
- Zygodiscus diplogrammus**, Maud Rise: Site 690, 487b
- Zygodiscus sigmoides**
- Maud Rise: Site 690, 486b, 487b, 531b
 - abundance, 523b
- Zygodiscus spiralis**, Maud Rise: Site 690, 487b