

## INDEX TO VOLUME 118

This index provides coverage for both the *Initial Reports* and *Scientific Results* portions of Volume 118 of the *Proceedings of the Ocean Drilling Program*. Index entries with the suffix *a* refer to pages in the *Initial Reports*, and those with *b*, to pages in the *Scientific Results* (this book).

The index usually is presented in three parts: (1) a Subject Index, (2) a Site Index, and (3) a Paleontological Index. For this volume, however, indexable paleontological material consists only of 6 main entries, 5 subentries, and 4 sub-subentries, so we decided to incorporate these listings within the body of the Subject Index.

The index was prepared by Wm. J. Richardson Associates, Inc., under subcontract to the Ocean Drilling Program. It expands the concept developed by the Deep Sea Drilling Project at Scripps Institution of Oceanography for a comprehensive, cumulative index of DSDP volumes. Both of these indexes are based on a hierarchy of entries: (1) a main entry, defined as a key word or concept followed by a reference to the page on which that word or concept appears; (2) a subentry, defined as a further elaboration on the main entry followed by a page reference; and (3) a sub-subentry, defined as an even further elaboration on the main entry or subentry followed by a page reference.

The Subject Index follows a standard format. Geographic and individual names are referenced in the index only if they are subjects of discussion. This index also includes a few paleontological listings, as noted in the second paragraph of this description.

The Site Index is structured to contain entries for the sites discussed in the volume. Site entries are modified by subject subentries.

The indexes cover text figures and tables but not core description forms ("barrel sheets") or core photographs. Also excluded are bibliographic references, names of individuals, and routine front and back matter.

For further information, and for information on electronic formats available, contact the Chief Production Editor, Ocean Drilling Program, 1000 Discovery Drive, College Station, Texas 77845-9547.

**SUBJECT INDEX**

Acoustic impedance, Atlantis Bank, seismic reflection profiling and, 246b  
 Acoustic properties  
     Atlantis Bank, 553b  
         hydraulic transmissivity and, 267b-268b  
 Actinolite, Atlantis II Fracture Zone: Site 732, replacement, 53a  
 Actinolite corona, Atlantis Bank, 133a, 135a  
 Actinolite, green, Atlantis Bank, 138a  
 Actinolitic hornblende-sodic plagioclase assemblage, Atlantis Bank, 160b  
 Alteration  
     Atlantis Bank  
         albitization, 208b  
         cataclastic deformation, 137a-138a  
         chemical effects, 546b-547b  
         deformation and styles of, 207b-208b, 212b  
         isotopic effects, high- vs. low-temperature, 138b  
         late-stage oxidation, 546b  
         layer 3, 543b-548b  
         low-temperature, 114b  
         magnetic properties and degree of, 294b  
             296b, 300b, 303b  
         near late intrusives, 163b, 165b  
         physical property effects of, 233b  
         porosity and, 261b-262b, 263b, 266b-267b  
         pressure of, 211b-212b, 213b  
         role of seawater in, 208b-209b  
         styles of, 113b-114b, 543b-546b  
         temperature of, 209b-211b, 213b  
 Atlantis II Fracture Zone: Site 732,  
     low-temperature, 53a  
     layer 3, dynamic model, 547b-548b  
     *See also* Greenschist-facies alteration  
 Alteration, hydrothermal  
     Atlantis Bank  
         amphibolite-facies, 488b  
         contribution to marine magnetic anomalies, 302b-303b  
         in iron-titanium oxide gabbro, 59b  
         isotopic changes with, 136b-138b, 141b  
         low-temperature, 489b  
         relation to plastic deformation, 106a  
         sulfur and secondary sulfides, 120b-121b  
 Atlantis II Fracture Zone: Site 732  
     moderate temperature, 53a  
     transform plate boundary, 43a  
 Alteration, hydrous  
     Atlantis Bank  
         deformation and, 174b  
         oxygen isotope depletion, 173b-174b  
         relation to synkinematic cracks, 163b  
 Alteration index, Atlantis Bank, vein density correlation, 171b  
 Alteration, oxidative, Atlantis Bank, 138a-139a, 209a  
 Alteration, static  
     Atlantis Bank, 90a, 209a, 489b  
         olivine replacement, 544b  
         origin and distribution, 544b  
         in undeformed gabbro, 138a  
     Atlantis II Fracture Zone  
         amphiboles, 68a  
         calcium silicate veins and, 86a  
         gabbro, 51a  
 Aluminum  
     Atlantis Bank  
         in amphiboles, 198b  
         geochemical log, 175a, 178a

    Atlantis II Fracture Zone, in ultramafic rocks, 13a  
 Aluminum activation clay tool (AACT), Atlantis Bank, 271b  
 Aluminum oxide  
     Atlantis Bank  
         in amphiboles, 56b  
         in clinopyroxene, 51b, 65b  
         geochemical logging data, 279b  
         in orthopyroxene, 53b, 67b  
     Atlantis II Fracture Zone, 150b  
 Aluminum/titanium ratio, Atlantis Bank, clinopyroxene, 140b  
 American-Antarctic Ridge  
     Atlantis II Fracture Zone and, 361b  
     basalt, geochemistry, transform vs. ridge axis, 472b  
 Amphibole  
     Atlantis Bank  
         anisotropy and orientation of, 249b, 251b  
         brown to green, 160b  
         deformation features, 130a  
         distribution of calcic, 166b  
         ferric vs. ferrous iron, 190b, 197b-198b, 199b  
         formation temperature, 132a  
         gabbro replacement, 137a  
         from metagabbros and late magmatic intrusions (LMIs), 165b  
         metamorphic vs. hydrothermal, 208b  
         mineral chemistry, 190b-198b  
         minor elements, 56b  
         orientation in deformed metagabbros, 240b, 241b  
         oxygen isotopes, parental magma, 136b  
         secondary matrix, 504b, 506b  
         strontium isotopes, 136b  
         temperature of alteration, 209b, 210b  
         vein type vs. red-brown type, 136a  
 Atlantis II Fracture Zone: Site 733, deformation and synkinematic growth, 68a  
 Amphibole, actinolitic, Atlantis Bank, oscillatory zonation, 198b  
 Amphibole, blue-green, Atlantis Bank, 133a  
 Amphibole, brown  
     Atlantis Bank  
         with gabbro deformation, 131a  
         magmatic origin, 186b  
         in olivine and ferrogabbros, 126a-127a  
         oxide association during deformation, 133a-134a  
 Atlantis II Fracture Zone: Site 733, 66a  
     replacement pattern, 69a  
 Amphibole, green  
     Atlantis Bank  
         oxygen isotopes and formation of, 174b  
         secondary sulfides in recrystallized, 117b  
         sulfide minerals and, 127a  
 Amphibole, iron-magnesium, Atlantis Bank, in reaction coronas, 198b  
 Amphibole, magnesian, Atlantis Bank, volume increase with depth, 138a  
 Amphibole, secondary, Atlantis Bank, strontium isotopes, 132b-133b  
 Amphibolite  
     Atlantis II Fracture fine: Site 733  
         foliation, 63a  
         plastic deformation, 60a  
         primary mineralogy, 65a  
 Amphibolite, foliated, Atlantis II Fracture Zone: Site 733, plastic deformation, 75a  
 Amphibolite mylonite  
     Atlantis II Fracture Zone: Site 734, 84a  
         metamorphism, 86a  
         petrography, 85a  
 Amphibolite-facies metamorphism  
     Atlantis Bank, 497b, 545b  
         conditions, 422b  
         dislocation slip, 407b, 421b-422b  
         Skaergaard Intrusion compared, 505b-506b  
         subsolidus deformation, 503b  
 Amphibolitization  
     Atlantis Bank  
         in recrystallized gabbros, 488b-489b  
         second stage, 132a  
 Andesite, Atlantis Bank, crystallization of Fe-Ti oxides and sulfide segregation, 93b-94b  
 Anisotropy  
     Atlantis Bank, 231b  
         in deformed gabbros, 234b, 236b  
         effect of mineral orientation on, 248b-249b  
         foliated gabbros and metagabbros, 157a, 159a  
         mineralogical effects, 236b  
         mylonitic metagabbros, 163a, 209a-210a  
         preferred crack orientation and, 231b  
         vs. pressure, oriented amphiboles, 240b  
         vs. pressure, oriented plagioclase, 240b, 241b  
 Anna de Koenig Seamount, plutonic rocks, shallow depth, 77b  
 Anorthite  
     Atlantis Bank, 99a  
         in plagioclase, 399b  
     in plagioclase, 12b, 18b, 42b, 49b, 62b, 64b, 69b  
 Anthophyllite, Atlantis Bank, in porphyroblast cores, 137a  
 Apatite  
     Atlantis Bank  
         fluid inclusions, 201b  
         in iron-titanium oxide gabbro, 116a  
         magmatic and metamorphic phases, 167b  
         primary origin, 117a  
         saturation, 99b, 100b  
 Atlantis II Fracture Zone: Site 732, 50a  
 Aragonite, Atlantis II Fracture Zone: Site 734, 87a  
 Atlantic Ocean, basalts, radiogenic isotopes, 131b  
 Atlantic Ocean N  
     gabbro, braided brittle-ductile deformation, 501b  
     seismic stratigraphy, low-dipping reflectors, 507b  
 Atlantis Bank  
     alteration  
         layer 3, 543b-548b  
         stratigraphy, 503b-505b  
     association, late magmatic intrusions, 159b  
     bathymetry, 22b, 76b  
     borehole televiewer, 205a, 262b, 264b  
     borehole temperature profile, 266b  
     deformation  
         brittle, 105a-106a, 159b  
         ilmenite-magnetite characteristics, 132a-134a  
         plastic, 103a-105a, 155b, 159b  
         reorientation of structures, 409b-414b  
     drill-string packer experiments, 333b-339b  
     dynamothermal metamorphism, 129a-132a  
     electrical conductivity, oceanic gabbros, 323b-329b  
     electrical properties, 555b  
     fluid inclusions, 200b-206b  
     gabbro  
         iron-titanium oxide vs. olivine, 56b, 59b  
         parental magma composition, 139b-141b

## SUBJECT INDEX

permeability, 333b-339b  
 plastic deformation, 399b-408b  
 tectonic evolution, 415b-429b  
 genetic relationship, principal rock types, 33b-35b  
 geochemical logging, 271b-272b  
 geochemistry, 4b-5b, 139a-150a  
     alteration effects, 136b-138b  
 geological setting, 21b, 44b, 90a-91a, 113b-114b, 127b-128b, 182b, 347b  
 glass composition  
     oxides, 86b-87b  
     potassium oxide and phosphate, 88b  
     sulfur, 88b-89b  
 hard-rock drilling  
     core recovery and penetration rate, 31a-33a  
     diamond bits, 31a-32a, 93a  
     hard-rock guidebase (HRGB), 31a  
     navidrill coring system, 36a-37a  
     positive displacement coring system, 31a-33a  
 hydraulic testing, 266b  
 hydrogen isotopes, 133b, 135b-136b  
 igneous sulfide distribution, 119b-120b  
 interval velocities, 222b-223b  
 intrusive events, 507b-509b  
 liquid line of descent, 82b, 95b-101b  
 lithologic summary, 77b-79b  
 lithology, schematic log, 183b  
 lithostratigraphy, 22b, 113b, 232b, 272b, 324b, 410b, 416b, 417b, 424b, 541b-542b  
 geochemical correlation, 147a  
 magnetostratigraphic correlation, 310b-313b  
 petrographic correlation, 186b  
 principal rock types, 23b  
 revised, 154b  
 unit contacts, 3b-4b  
 Unit I, 89a, 97a, 128b, 208a, 311b, 542b  
 Unit II, 89a, 97a, 99a-100a, 128b, 208a-209a, 311b, 542b  
 Unit II/Unit III boundary, 99a, 100a-101a  
 Unit III, 90a, 100a-101a, 128b, 208a-209a, 312b, 542b  
 Unit IV, 102a, 128b-129b, 208a-209a, 312b-313b, 542b-543b  
 Unit V, 102a, 129b-130b, 208a-209a, 313b, 543b  
 Unit V/Unit VI boundary, 103a  
 Unit VI 102a-103a, 208a-209a, 313b, 543b  
 location, 89a, 90a, 114b, 182b, 207a, 262b, 272b, 324b, 334b, 348b  
 logging, 165a, 209a-210a, 555b-556b  
     azimuth, 170a  
     borehole conditions, 169a  
     caliper, 170a  
     deviation, 170a  
     dual induction-sonic-caliper-natural gamma, 168a  
     dual laterolog resistivity, 169a, 177a  
     geochemical, 169a, 173a, 177a-178a  
     geophysical, 265b  
     induction resistivity-caliper, 169a, 177a  
     litho-density, 169a  
     magnetometer, 171a  
     natural gamma radiation, 173a  
     operations, 96a-97a  
     physical properties, 169a, 172b-177a  
     temperature, 171a  
 magmatic differentiation, deforming crust, 103b-104b  
 magmatic stratigraphy, 154b-155b  
 magnetic properties, 285b-305b, 554b-555b  
     alternating field demagnetization, 153a-154a  
     discrete samples vs. logging data, 309b-310b  
     inclination, 154a-156a

Koenigsberger Ratio, 153a  
 oceanic gabbros, 318b-320b  
 remanent magnetization, 151a  
 susceptibility, 151a-153a  
 thermal demagnetization, 154a  
 magnetometer logs, 191a-193a, 197a  
 melanogranophyres, 102b-103b  
 metamorphic minerals, 171b-174b  
 mineral chemistry, 189b-200b  
     minor elements, 51b, 53b, 56b  
 mineralogy, 90b-95b  
     ambiguous origin, 166b-170b  
     between rock types, 27b, 29b, 49b  
     covariation, 12b-13b  
     iron-titanium oxide gabbro, 49b  
     late-stage differentiation, 101b-102b  
     layer 3, 542b-543b  
     normative vs. microprobe compositions, 5b  
     oxide, 122a-129a  
     post-kinematic, 163b, 165b-166b  
     principal rock types, 23b-27b  
     sulfide, 122a-129a  
     synkinematic, 160b, 163b  
     within rock types, 29b-31b, 33b  
 model for development, 35b  
 multichannel sonic logs, 264b, 266b  
     operations, 178a, 180a-182a  
 navigation, 363b  
 operations, television/sonar survey, 92a, 94a  
 oxygen isotopes, 133b, 135b-136b, 170b-171b  
     trondhjemite, 138b-139b  
 parent magmas, diversity of, 83b-86b  
 permeability, 341b-342b  
 petrography, 44b, 47b-48b, 132b-133b, 186b-189b  
     basalt, 117a  
     felsic rocks, 117a-119a  
     iron-titanium oxide gabbro, 112a-117a  
     layer 3, 542b-543b  
     layering, 121a-122a  
     lithostratigraphic correlation, 48b, 186b  
     olivine-bearing and olivine gabbros, 107a-111a  
     porphyroclastic gabbro, 107a  
     textural variations, 119a-121a  
     troctolites, 111a-112a  
 petrology, 77b-79b  
     opaque minerals, 114b-117b  
 physical properties, 158a-162a, 227b-237b  
 radiogenic isotopes, 131b-135b  
 residual melt porosity, 79b-82b  
 rotary coring bit  
     operations, 32a-33a, 93a, 95a-96a  
     performance, 33a-34a  
 seismic properties, 553b-554b  
 seismic stratigraphy, reflection profiling, 223b-225b  
 stable isotopes, 206b-207b  
 stress and seawater circulation, 505b-507b  
 structural stratigraphy, 410b, 417b, 424b  
 subrift magma chamber, 14b  
 sulfur  
     carbon content, 117b  
     distribution, 119b-120b  
     hydrothermal alteration, 120b-121b  
     isotopes, 117b, 121b-123b  
 summary logs, 214a-222a  
 synthetic seismogram, 245b-247b  
 tectonic history, 90a-91a  
     effect of rotation on magnetic properties, 319b  
     three-step scenario, 425b-426b, 427b-428b  
 tectonic setting, 396b, 415b-116b, 541b  
 tectonic stratigraphy, 501b-503b  
 texture of principal rock types, 23b-27b

thermal conductivity, 163a-165a  
 thermal modeling, 347b-354b  
 topographic relief, 207a-208a  
     transverse ridge, 91a  
 trace elements in silicate minerals, 5b-6b, 12b  
 ultrasonic attenuation, 253b-257b  
 vertical seismic profile, 182a-191a, 217b-220b  
 Atlantis II Fracture Zone  
     basement description conventions, 10a-11a  
     bathymetry, 6a, 91a, 364b, 410b, 416b, 440b  
     magnetic anomalies, 378b  
     southern ridge-transform intersection, 365b  
 biostratigraphy, 19a  
 deformation and alteration, 488b-501b  
 formation, 361b  
 gabbro  
     geochemistry, 470b-488b  
     mineralogy, 470b-488b  
 geological setting, 3a-5a, 361b, 440b-441b  
 gravels, 431b-438b  
 hard-rock drilling  
     hard-rock guidebase (HRGB), 28a-30a  
     positive displacement coring motor (PDCM), 25a-28a  
 igneous rock, classification, 444b-148b  
 igneous stratigraphy, 448b-470b  
     discrete primary subintervals, 457b  
     Unit I, 449b, 450b, 456b-457b  
     Unit II, 451b, 457b-459b  
     Unit II/III contact, 458b-459b  
     Unit III, 452b, 459b-462b  
     Unit IV contact, 462b, 463b  
     Unit IV, 453b, 462b, 464b-465b  
     Unit IV/Unit V contact, 465b, 466b  
     Unit V, 453b-454b, 465b-466b  
     Unit V/Unit VI contact, 467b  
     Unit VI, 454b-455b, 467b-470b  
 inactive valley  
     bathymetry, 375b  
     morphology, nontransform vs. paleotransform walk, 373b  
 layer 3, lithostratigraphic evolution, 439b-513b  
 lithology, 4a  
     association of rock types, 377b, 380b  
     distribution, 377b  
     transform vs. nontransform regions, 360b, 380b  
 magnetic properties, 21a, 382b-384b, 388b-389b  
 median tectonic ridge, formation, 360b-361b  
 nodal basin, Southwest Indian Ridge  
     intersection, 360b  
 physical properties, 19a  
 physiography, 359b-360b  
     SeaBeam survey, 363b-364b  
 ridge axis, right-lateral offsets, 392b  
 ridge-transform intersection  
     mass-wasting features, 431b  
     physiography, 366b-368b  
 rift valleys, physiography, 364b-366b  
 sedimentology of fracture zone walls, 431b  
 spreading direction change, 360b, 376b-377b  
 spreading rate, asymmetry, 360b  
 structural model, 87a  
 tectonic evolution, 359b-396b  
 tectonic history, 60a, 79a, 359b-396b  
     median tectonic ridge, 43a  
     transtensional transform basins, 389b-392b  
 tectonic setting, 76b-77b, 361b-363b  
 topography, 145b  
 transform tectonic zone, 378b  
     dredge survey, 377b-382b  
     median tectonic ridge, 4a  
 physiography, 368b-377b

Atlantis II Fracture Zone: Site 732  
 bathymetry, 22b, 45a, 76b  
 transform valley, 43a  
 biostratigraphy, 48a-49a  
 borehole televiewer survey, median tectonic ridges, 373b  
 clay mineralogy, 146b-147b  
 deformation  
   cataclastic, 51a-52a  
   plastic, 51a  
 geochemistry, 54a, 147b-149b  
 geologic setting, 43a  
 hard-rock drilling rate, 47a  
 lithologic sequence, 56a  
 lithology, 42a, 50a, 145b  
 lithostratigraphy, 49a, 55a  
 location, 42a, 43a, 145b, 146b  
 navigation, 363b  
 operations, 43a-48a  
   television/sonar survey, 42a-44a  
 petrography, 132b-133b  
   basalt, 49a  
   diabase, 49a-50a  
   gabbro, 50a  
 rare earth elements, 150b  
 rotary coring bit, bottom-hole assembly, 46a-47a  
 sedimentology, 48a  
 Atlantis II Fracture Zone: Site 733  
 bathymetry, 22b, 61a, 76b  
 geochemistry, gabbro, 71a-73a  
 geologic setting, 60a  
 hard-rock drilling, beacon placement, 63a  
 lithostratigraphy, 63a-65a  
 location, 59a  
 magnetic properties, 73a  
 metamorphism and deformation, 67a-71a  
 navigation, 64a, 363b  
 petrography, 132b-133b  
 physical properties, 73a-75a  
 primary mineralogy, 65a-67a  
 Atlantis II Fracture Zone: Site 734  
 bathymetry, 22b, 76b, 79a  
 biostratigraphy, 85a  
 clay mineralogy, 146b-147b  
 geochemistry, 86a-87a, 147b-149b  
 geologic setting, 79a  
 gravel stratification, 432b-433b  
 lithology, 78a, 146b  
 lithostratigraphy, 85a  
 location, 78a, 79a, 145b, 146b  
 metamorphosis, 86a  
 navigation, 363b  
 petrography, 85a, 87a, 132b-133b  
 plastic deformation, serpentinized peridotites, 85a  
 rare earth elements, 150b  
 sedimentary sequence, 87a  
 sedimentology, 84a, 433b-434b  
 static alteration, moderate temperatures, 86a  
 television/sonar survey, 80a, 81a, 82a, 83a  
 Attenuation  
 Atlantis Bank  
   alteration/deformation effects, 256b  
   errors in measurement, 254b-255b  
   experimental setup, 253b-257b  
   pressure effect, 255b-256b  
   vs. velocity, 258b, 259b  
 Augen, chromite, Atlantis II Fracture Zone: Site 732, 55a  
 Augen gneissic texture, Atlantis Bank, in deformed gabbros, 103a  
 Augen, rounded, Atlantis II Fracture Zone: Site 732, 54a

Augite  
 Atlantis Bank  
   oxygen isotopes, 206b  
   velocity and orientation of, 236b  
 Basalt  
 Atlantis Bank  
   contribution to marine magnetic anomalies, 300b-305b  
   petrography, 117a  
   texture, 117a  
 Atlantis II Fracture Zone  
   classification, 10a  
   geochemistry, dredged vs. drilled samples, 471b-472b  
   hydrothermal alteration, 4a  
   layer 2, 377b, 380b  
   liquidus trends with proximity to mantle hot spots, 380b-381b  
   magma chamber eruption vs. crystallization at depth, 472b  
   MORB characteristics, 361b  
   valley vs. transform walls, 380b, 381b  
 Atlantis II Fracture Zone: Site 732, 48a  
 alteration, 49a  
 calcium vs. magnesium number, 55a  
 geochemistry, 54a  
 incompatible elements, 54a  
 as mineralogical source, 149b  
 petrography, 49a  
 quench crystallization textures, 49a, 50a  
 weight loss on ignition, 54a  
 Atlantis II Fracture Zone: Site 733  
 cumulus phase assemblage, 72a  
 geochemistry, 71a-73a  
 metamorphism, 67a-71a  
 petrography, 65a-67a  
 trapped liquid, 72a  
 Atlantis II Fracture Zone: Site 734, as  
   mineralogical source, 149b  
 Southwest Indian Ridge, depleted-N-type, 54a  
 thermal conductivity, 20a-21a  
 Basalt, holocrystalline, Atlantis II Fracture Zone: Site 732, 53a  
 Basalt, pillow, Atlantis II Fracture Zone, hooked ridges, 376b  
 Basement  
 Atlantis Bank, 89a  
   lithostratigraphy, 98a  
 Atlantis II Fracture Zone, alteration, 11a  
 Atlantis II Fracture Zone: Site 734, 79a, 87a  
 Bay of Islands, ophiolites, magma mixing and fractionation of, 6b  
 Biotite  
 Atlantis Bank  
   in felsic rock, 117a  
   magnetite, association with, 167b  
   mineral chemistry, 200b  
   red-green association, 167b  
 Biotite, green, Atlantis Bank, synkinematic minerals, 180b  
 Blake Spur Fracture Zone, Moho sequence, seismic stratigraphy, 219b, 225b  
 Borehole televiewer (BHTV), Atlantis II Fracture Zone, 205a-207a, 264b, 373b  
 Bornite, Atlantis Bank, in oxide-rich ferrogabbros, 125a  
 Bottom-hole assembly  
 Atlantis II Fracture Zone  
   with navidrill, 37a, 38a  
   with positive displacement coring motor (PDCM), 25a  
 Bouvet Fracture Zone, peridotite, 361b

Breccia  
 Atlantis Bank  
   carbonate vein relationship, 173b, 175b  
   cataclastic deformation, 106a  
   hydrothermal alteration, 159b, 175b  
   leucocratic diopside-plagioclase veins, 207b  
   low-temperature metamorphism, 172b  
   magmatic/hydrothermal zones, mineralogy, 165b-166b, 175b  
   magmatic/hydrothermal zones, texture, 159b  
   mineralogy, alteration effects, 171b  
   oxidation, 139a  
   oxygen isotope depletion, 171b, 175b  
   seawater hydration, 209b  
   secondary minerals, 208b  
   structure relation to late magmatic intrusions (LMIs), 172b-173b  
   zones, 186b  
 Atlantis II Fracture Zone, in mylonite zone, 465b  
 Atlantis II Fracture Zone: Site 732, 56a  
   origin, 49a  
   felsic intrusion, Atlantis Bank, 102a  
 Breccia, carbonate-cemented, Atlantis II Fracture Zone: Site 734, 85a  
 Breccia, cataclastic  
 Atlantis Bank, 419b  
   origin, 427b  
 Breccia, hydrothermal  
 Atlantis Bank, 502b  
   clinopyroxene composition, 169b, 170b  
   formation, 489b  
   greenschist-facies alteration, 505b  
   secondary minerals, 179b  
   zircon in, 172b  
 Breccia, trondhjemite  
 Atlantis Bank, 27b  
   crosscutting late-magmatic foliation, 503b, 535b  
 Atlantis II Fracture Zone, 499b  
   petrography, 447b-448b  
 Brevard fault zone, mylonites, velocity and silicon dioxide content, 247b  
 Bushveld Complex  
   apatite, chlorine-rich, 167b  
   biotite-magnetite association, 167b  
   fluid inclusions, methane-bearing, 209b  
   hortonolite pipes, 486b  
   hydrothermal veins, 208b  
   mineralogy, incompatible element-rich mineral association, 166b-167b  
 Calcareous nannofossils  
 Atlantis II Fracture Zone  
   biostratigraphy, 19a  
   zonation, 20a  
 Atlantis II Fracture Zone: Site 732, Zone NN20, 49a  
 Atlantis II Fracture Zone: Site 734  
   Zone NN19, 84a  
   Zone NN21, 84a  
 Calcite, Atlantis Bank, strontium isotopes, 200b  
 Calcium  
 Atlantis Bank  
   diabase, 477b  
   gabbros, 477b  
   geochemical log, 178a  
 Atlantis II Fracture Zone, in ultramafic rocks, 13a  
 Calcium oxide  
 Atlantis Bank, geochemical logging data, 279b  
 Atlantis II Fracture Zone, 150b  
 Calcium oxide/aluminum oxide ratio, Atlantis Bank, 13b-14b

## SUBJECT INDEX

Calcium/magnesium ratio, Atlantis Bank, amphibole precipitation and, 506b  
 Canyon-fan system, basin analysis models, 432b  
 Carbonate minerals, Atlantis II Fracture Zone, source provinces, 147b  
 Carbonate, pelagic, Atlantis II Fracture Zone, 431b  
 Cataclastic rock  
   Atlantis Bank  
 association, late magmatic intrusions (LMIs), 159b  
 mineralogy, 178b  
 Cation exchange capacity  
   Atlantis Bank, 325b, 326b-327b  
     vs. electrical resistivity, 327b  
 Cayman Trough  
   greenschist-facies alteration, 489b  
   mica, in gabbro, 168b  
 Central Indian Ridge, fracture zones, crystallization of gabbro, 35b  
 Chalcocpyrite  
   Atlantis Bank, 125a  
     in globular sulfides, 93b, 94b  
     in green amphibole, 117b  
     in secondary sulfides, 115b-116b  
 Charlie Gibbs Fracture Zone, density-velocity relationship, 225b  
 Chloride, Atlantis Bank, geochemical log, 174a  
 Chlorine  
   Atlantis Bank  
     in amphiboles, 208b, 212b  
     geochemical log, 178a  
     in hornblende, 198b  
 Chlorite  
   Atlantis Bank  
     in corona structures, 172b  
     in felsic veins, 551b  
     influence of water/rock ratio on, 174b  
     magnesian composition, 138a  
     metamorphosed gabbros and late magmatic intrusions (LMIs), 735b  
     plagioclase replacement, 138a  
     seawater downflow vs. upflow and composition of, 208b  
     temperature of alteration, 211b  
   Atlantis II Fracture Zone  
     plagioclase replacement, 53a  
     source provinces, 147b  
 Chrome, Atlantis Bank, in oxide-bearing gabbros, 483b, 486b  
 Chrome diopside, Atlantis II Fracture Zone, in troctolitic rocks, 447b  
 Chromium  
   Atlantis Bank, 136b  
     cyclical variations, 5b  
     diabase, 480b  
     gabbros, 480b  
 Chromium oxide  
   Atlantis Bank  
     in amphiboles, 56b  
     in clinopyroxene, 6b, 53b, 65b, 140b  
     in orthopyroxene, 53b  
 Chromian-spinel, Atlantis II Fracture Zone, in troctolite, 441b, 447b, 470b  
 Circulation, oceanic  
   Atlantis Bank  
     brittle-ductile shear zone formation, 511b  
     *in-situ* stress state and, 505b-507b  
 Clasts, igneous, Atlantis II Fracture Zone: Site 732, 49a, 55a  
 Clasts, metamorphic, Atlantis II Fracture Zone: Site 732, 49a, 55a  
 Clasts, sedimentary, Atlantis II Fracture Zone: Site 732, 49a

Clay mineralogy  
   Atlantis II Fracture Zone, 146b-147b  
   Atlantis II Fracture Zone: Site 732, gabbros, 53a  
   Atlantis II Fracture Zone: Site 734, mineral sources, 147b  
 Clinopyroxene  
   Atlantis Bank, 82b, 180b  
     actinolite replacement, 138a  
     aluminum vs. titanium, 17b  
     augen, 107a  
     chemical composition, 57b-59b, 69b  
     chromium oxide vs. magnesium number, 17b  
     chromium oxide-enriched, 6b  
     deformation features, 130a  
     exsolution lamellae, 4b  
     in felsic rock, 117a  
     gabbro compositions, 25b  
     inverted pigeonite in, 47b, 49b, 71b  
     in iron-titanium oxide gabbro, 113a  
     iron-titanium oxide vs. olivine gabbro, 44b  
     manganese oxide vs. iron content, 91b  
     metamorphic, 160b  
     Mg#, 62b  
     Mg#, vs. anorthite in plagioclase, 18b  
     mineral chemistry, 14b, 15b  
     mineral zoning, 5b  
     mineralogy, late differentiation, 97b  
     minor elements, 51b, 53b, 65b  
     minor oxide abundances, 97b  
     modal proportion, 116a  
     oikocrysts, 109a, 121a, 124a  
     in olivine gabbro, 126a  
     orientation of porphyroclasts, 104a  
     orthopyroxene intergrowths, 44b  
     plutonic rocks, 10b-11b  
     replacement by amphibole, 171b  
     replacement by hornblende, 136a-137a  
     seawater availability and recrystallization of, 174b  
     sodium oxide vs. grain size, 51b  
     strontium isotopes, 131b-132b  
     texture, 116a  
     titanium oxide, 6b, 80b  
     titanium vs. aluminum, 12b  
     titanium vs. chromium, 12b, 17b  
     zirconium in cumulus, 79b-80b  
     zonation, olivine gabbros, 25b  
 Atlantis II Fracture Zone, in troctolitic rocks, 447b  
 Atlantis II Fracture Zone: Site 733  
   porphyroclasts, 70a  
   recrystallization and mechanical twins, 69a  
   replacement pattern, 65a-66a, 69a  
 Atlantis II Fracture Zone: Site 734  
   relic minerals, 85a  
   replacement pattern, 78a  
 Cobalt, Atlantis Bank, in igneous sulfides, 115b  
*Coccoilithus pelagicus*, Atlantis II Fracture Zone: Site 734, glacial/interglacial cycles, 84a  
 Collapse caldera, Atlantis II Fracture Zone, transform volcano, 366b  
 Copper  
   Atlantis Bank  
     stratigraphic variation, 147a  
     vs. iron, in igneous sulfides, 119b-120b, 123b  
 Corona structure  
   Atlantis Bank, 114b, 136a  
     in gabbronorite, 44b, 47b  
     igneous phase replacement, 179b  
     interstitial olivine-bearing titanium oxide gabbro, 47b  
     iron-magnesium amphiboles in, 198b  
     mineralogy, 177b  
     olivine, 118b, 546b

phlogopite content, 168b  
 secondary sulfides, 115b  
 in undeformed gabbros, 138a  
 Covellite, Atlantis Bank, in oxide-rich ferrogabbros, 125a  
 Crack porosity  
   Atlantis Bank  
     effect on velocity, 231b, 233b, 234b, 235b, 247b  
     radiogenic isotopes and, 132b  
 Cracking front  
   Atlantis Bank  
     elastic strain available for, 506b  
     in seawater penetration, 505b  
 Crozier Basin, smectite source, 147b  
 Crust, oceanic  
   Arctic Ridge, magma supply and thickness of, 361b  
   Atlantis Bank  
     age, 415b-416b  
     excluded element concentration, 86b  
     magnetic properties, 318b  
     seismic stratigraphy, 219b, 225b  
 Atlantis II Fracture Zone  
   asymmetry across fracture zone, 392b  
   plate readjustment and thinning of, 392b  
   thickness of, 363b  
   formation, slow-spreading ridges, 513b  
   fracture zones, 90a  
     high vs. low offsets, 225b  
     thickness, 42a  
   hydrothermal circulation, black smokers, 136b-137b  
   Southwest Indian Ridge, magma supply and thickness of, 361b  
 Crust, plutonic, Atlantis II Fracture Zone, as magnetic anomaly source, 360b  
 Crustal weld  
   Atlantis II Fracture Zone  
     formation, 392b  
     ridge-transform intersections, 394b, 441b  
 Crystal layering, Atlantis Bank, 106a  
 Crystal mush  
   Atlantis Bank  
     formation, 509b  
     magmatic foliation, 427b  
   Atlantis II Fracture Zone  
     formation, 442b  
     intrusive blebs, 458b  
     syntectonic differentiation, 488b  
 Crystallization  
   Atlantis Bank  
     hydrous magmatic, 208b  
     lunar approximation model, 101b, 102b  
     metamorphic styles, 160b, 171b  
   Atlantis II Fracture Zone, of discrete magma body, 486b  
 Cummingtonite  
   Atlantis Bank, 163b  
     in porphyroblast cores, 137a  
   Atlantis II Fracture Zone: Site 733, 67a, 69a, 71a  
 Cumulates  
   Atlantis Bank, 4b  
     temperature of formation, 13b  
 Cyprus, pillow basalts, 431b  
 Deformation  
   Atlantis Bank  
     correlation with structural domains, 159b  
     effect on mineral chemistry, 88b  
     hydration and degree of, 186b  
     melt porosity and, 80b  
     olivine vs. titanium oxide gabbro, 56b  
     oxide formation, 90b

- oxide olivine gabbro, 26b-27b  
penetrative, 502b  
porphyroclastic/gneissic, 78b  
reorientation of structures, 409b-414b  
temperature, 319b  
textural types produced, 231b  
Zones 1-5, 501b-503b
- Atlantis II Fracture Zone  
braided, 457b  
brittle-ductile transition, 488b  
with melt present, 461b  
nonlinear progression, 495b  
brittle-ductile transition, slow-spreading vs.  
fast-spreading ridges, 513b
- Deformation, brittle  
Atlantis Bank, 105a-106a, 209a  
gabbros, 426b-427b  
intrusion of oxide gabbros and initiation of,  
27b  
mineralogical changes, 114b  
post-kinematic, 159b  
sulfur depletion, 121b  
synkinematic, 159b  
Atlantis II Fracture Zone, 499b  
macroscopic evidence, 491b
- Deformation, brittle-ductile  
Atlantis Bank  
braiding, 501b  
initiation, 511b  
late-magmatic, 501b-502b  
Atlantis II Fracture Zone  
amphibolite facies, 497b  
late-magmatic, 496b-497b
- Deformation, brittle-ductile, subsolidus, Atlantis  
Bank, 488b, 501b, 511b
- Deformation, cataclastic  
Atlantis Bank  
alteration with, 136a-138a  
metagabbro breccia, 106a  
Atlantis II Fracture Zone: Site 732,  
medium-temperature metamorphism and,  
51a-52a
- Deformation, ductile  
Atlantis Bank, 159b  
low-temperature, 428b  
Lower Structural Domain, 426b  
magmatic, 418b  
magmatic differentiation, 103b-104b  
Middle Structural Domain, 413b, 425b-426b  
migration of intercumulus melt, 488b, 491b  
mineral assemblages associated with, 160b  
residual melt expulsion during, 82b  
seawater availability, 174b, 175b  
Upper Structural Domain, 411b, 413b, 422b-  
423b, 425b
- Atlantis II Fracture Zone  
classification, 492b-493b  
extent and location of, 492b-496b  
two phases, 456b
- Deformation, ductile, solid-state  
Atlantis Bank, 418b-422b, 428b  
synkinematic metamorphic conditions, 420b-  
421b
- Deformation, ductile, subsolidus  
Atlantis Bank  
gabbronorite, 536b  
olivine gabbro, 536b
- Deformation, hypersolidus, Atlantis Bank, vein  
and breccia formation, 488b
- Deformation, magmatic  
Atlantis Bank, textural characteristics, 537b  
Atlantis II Fracture Zone, disseminated oxide  
olivine gabbros, 461b
- Deformation, mylonitic, late-stage subsolidus, At-  
lantis II Fracture Zone, oxide-bearing layers,  
458b
- Deformation, plastic  
Atlantis Bank, 103a-105a  
foliated vs. unfoliated specimens, 103a  
ilmenite-magnetite enrichment and, 105a  
petrofabric studies, 159b  
relation to hydrothermal alteration, 106a  
stratigraphy, 156b-157b  
synkinematic veins, 172b  
textural types, 155b, 171b  
Atlantis II Fracture Zone  
high-temperature, 51a, 469b  
serpentinized peridotite, 85a
- Deformation, plastic, solid-state  
Atlantis Bank  
gabbros with large recrystallized grains, 403b-  
405b, 406-407b  
gabbros with small recrystallized grains,  
403b-405b, 407b  
recrystallization mechanisms, 404b, 405b,  
406b  
strain localization, 406b-407b  
strain-softening effects, 405b, 406b, 407b,  
408b  
temperature of, 139a, 399b, 407b
- Deformation structures  
Atlantis Bank  
dip of foliation, 105a-106a, 107a  
intrusive contacts, 106a  
olivine gabbros, 108a  
textural distribution, 107a
- Deformation, subsolidus  
Atlantis Bank  
amphibolite-facies, 503b  
iron-titanium oxide gabbro, 44b  
Atlantis II Fracture Zone, water/rock ratio,  
497b-498b
- Density  
Atlantis Bank, 230b-231b, 324b-326b  
alteration effects, 233b  
crystalline rock, 157a  
metagabbro, 157a  
Unit IV/Unit V boundary, 507b  
variations between rock types, 507b  
velocity and, 231b-232b, 238b, 239b  
vs. electrical resistivity, 326b-327b
- Atlantis II Fracture Zone: Site 733, 73a-74a  
metagabbro, 60a, 76a  
*See also* Melt density
- Diabase  
Atlantis Bank, magnetization intensity values,  
302b
- Atlantis II Fracture Zone  
crosscutting relationships, 441b  
recrystallized foliated, 449b
- Atlantis II Fracture Zone: Site 732, 54a  
alteration, 50a  
petrography, 49a-50a
- Atlantis II Fracture Zone: Site 733, primary  
mineralogy, 66a
- Differentiation, late-stage  
Atlantis Bank  
conditions of, 101b-102b  
melt densities, 98b  
mineralogical effect, 96b-97b
- Diffusion rate, Atlantis Bank, olivine vs. iron-tita-  
nium oxide gabbro, 59b, 70b
- Dike, basalt  
Atlantis Bank, 100a  
geochemistry, 147a  
magnetic properties, 150a  
petrography, 117a
- Atlantis II Fracture Zone: Site 732,  
geochemistry, correlation, Site 735, 147a
- Dike, diabase  
Atlantis Bank, 504b  
Atlantis II Fracture Zone, 458b
- Dike, metabasalt, Atlantis Bank, 27b
- Dike, microgabbro, Atlantis Bank, 25b-26b
- Dike, trondhjemite, Atlantis Bank, 117a
- Dikelet, basaltic, Atlantis II Fracture Zone, 49a-  
50a, 68a
- Dikelet, leucocratic, Atlantis Bank, 426b
- Dikelet, trondhjemitic, Atlantis Bank, mineral-  
ogy, 90b, 104b
- Diopside  
Atlantis Bank  
episodic growth, 159b  
in felsic veins, 545b  
fluid inclusions, 201b  
hydrothermal origin, 200b  
in late magmatic intrusions (LMIs), 168b  
magmatic vs. hydrothermal, 168b, 169-170b  
origin, 168b  
oxygen isotopes, 206b  
sodic plagioclase association, 168b  
temperature of formation, 168b  
Atlantis II Fracture Zone, in troctolite, 470b
- Duluth layered intrusion, post-cumulus re-equili-  
bration, 5b
- DuToit Fracture zone, transverse ridges, 441b
- Earth's geomagnetic field, Atlantis Bank, 198a,  
200a, 210a
- East Pacific Rise  
basalt, lead isotopes, 136b, 137b, 138b  
strontium, basalt glasses, 84b
- Elasticity, Atlantis Bank, velocity and formation  
of, 236b, 242b-243b, 264b
- Electrical conductivity  
Atlantis Bank  
iron-titanium oxide contribution, 324b  
mechanisms, 327b, 329b  
salinity and, 327b-328b, 329b, 330b  
vs. resistivity, 328b
- Electrical resistivity  
Atlantis Bank, 324b-326b  
vs. cation exchange capacity, 327b  
vs. conductivity, 323b-324b, 327b  
vs. density, 326b-327b  
vs. porosity, 328b, 331b, 555b
- Emiliana huxleyi*, Atlantis II Fracture Zone: Site  
732, 49a
- Emiliana huxleyi* Acme Zone, Atlantis II Frac-  
ture Zone: Site 734, 84a
- Epidote  
Atlantis Bank, 165b, 168b  
iron content, 211b  
mineral chemistry, 200b  
in plagioclase-diopside veins, 551b  
Atlantis II Fracture Zone: Site 732, 53a
- Estimated Fractionation Effect, Atlantis Bank, 85b
- Faulting, detachment, Atlantis II Fracture Zone,  
393b-394b, 441b, 511b
- Faulting, ductile, Atlantis Bank, 107a, 511b
- Faulting, listric  
Atlantis Bank, 428b  
high-angle, 77b
- Faulting, listric normal  
Atlantis Bank, 547b  
as seawater conduits, 206b
- Faulting, normal  
Atlantis Bank, ridge-transform intersection,  
428b  
Atlantis II Fracture Zone

## SUBJECT INDEX

- paleotransform wall scarp, 376b  
transform wall, 396b  
in transverse ridge evolution, 392b, 395b  
Atlantis II Fracture Zone: Site 734, 79a
- Feldspar  
    Atlantis Bank  
        mineral chemistry, 198b-199b, 201b  
        strontium concentration, 85b  
    Indian Ocean, strontium concentration, 84b
- Felsic rock  
    Atlantis Bank  
        mixing with gabbroic rock, 119a  
        petrography, 117a, 119a
- Ferroandesite, Atlantis Bank, crystallization of Fe-Ti oxides and sulfide segregation, 93b-94b
- Ferrobasalt  
    Atlantis Bank, intrusions, 103b  
    Atlantis II Fracture Zone, absence of, 509b
- Ferrogabbro  
    Atlantis Bank  
        ilmenite/magnetite ratio, 124a  
        iron-titanium oxides, relation to silicate phases, 126a  
        magnesian, 5b  
        petrology, 155b  
    Atlantis II Fracture Zone: Site 732, static metamorphism, 53a, 56a
- Fluid inclusions  
    Atlantis Bank  
        clathrate melting temperature, 204b, 206b  
        fluid types, 208b-209b  
        in gabbro, 200b-206b  
        homogenization temperature, 202b, 203b-204b, 205b, 206b  
        ice-melting temperature, 203b, 205b  
        laser Raman spectroscopy, 205b-206b  
        leucocratic diopside-bearing veins, 204b-205b  
        lithostratigraphic variation, 208b  
        methane-bearing, 212b  
        opaque hexagonal platelet daughter crystal, 206b  
        plagioclase vs. diopside, 205b  
        trondjemite, 204b  
        veined metagabbro, 202b-204b  
        veined mylonitic metagabbro, 202b
- Foliation  
    Atlantis Bank  
        fanning, 502b  
        microshears, 425b  
    Middle Structural Domain, 425b  
    shear bands, 423b  
    Atlantis II Fracture Zone  
        deformation intensity and inclination, 490P-491b  
        east-west strike, 441b  
        symmetry, brittle-ductile shear zones, 500b
- Foliation, gneissic  
    Atlantis Bank  
        crosscutting relationships, 504b  
        development, 106a  
        dip, 501b
- Foliation, late magmatic  
    Atlantis Bank  
        dip orientation, 502b  
        overprinting, 501b
- Foliation, magmatic  
    Atlantis Bank, 159b, 419b  
        crosscutting, 425b  
        dip orientation, 413b  
        overprinting, 418b  
        oxide gabbros, 427b  
        with solid-state deformation, 427b-428b  
    Atlantis II Fracture Zone, crosscutting, 499b
- Foliation, magmatic flow, Atlantis Bank, oxide gabbros, 27b  
Foliation, mylonitic  
    Atlantis Bank  
        crosscutting relationships, 504b  
        dip orientation, 425b, 501b
- Foliation, penetrative  
    Atlantis II Fracture Zone  
        oxide olivine gabbro, 464b, 469b  
        oxide-bearing gabbros, 460b
- Foliation, solid-state, Atlantis Bank, dip orientation, 413b
- Forsterite, Atlantis Bank, 62b, 68b, 99a
- Fractionation  
    Atlantis Bank  
        across nickel-nickel oxide (NNO) buffer, 101b, 102b  
        excluded elements contributed, 86b  
        isotopic variation with, 131b, 132b, 141b  
        magma mixing and, ophiolitic sequences, 12b  
        mineral bimodality and, 34b  
        syndeformation magmatic, 155b
- Fracture zones  
    Atlantis Bank  
        seismic structure, 183a, 185a  
        semblance velocity logs, 183a  
        crustal thickness, 42a  
        crustal thinning around, 3a  
        models, 42a, 90a  
        plutonic rock, spreading rate and, 3a  
        Southwest Indian Ridge, oceanic nature, 361b  
        spreading ridge segmentation and, 3a  
        transverse ridges, high vs. low offsets, 225b
- Fractures  
    Atlantis Bank  
        apparent apertures, 264b, 268b  
        compressional velocity and shear semblance decreases, 206a  
        density, 105a, 337b  
        dip, 206a, 207a, 208a, 209a, 269b  
        late subvertical, 505b  
        negative temperature gradient anomalies, 266b, 267b  
        permeability and density of, 266b-267b, 339b  
        post-kinematic, 159b  
        smectite-lined, 546b  
        vertical distribution, 556b
- Gabbro  
    Atlantis Bank  
        alteration, 113b-114b, 261b-262b  
        brittle deformation, 136a, 426b-427b  
        chromium-rich, 130b  
        contribution to marine magnetic anomalies, 320b  
        crystal intergrowths, 110a  
        crystallization, location of, 396b  
        deformation and hydrous alteration, 174b-175b  
        deformation textures, 231b, 400b, 403b, 418b-419b, 422b-426b  
        deformed, with large recrystallized grains, 403b-405b, 406b-407b  
        deformed, with small recrystallized grains, 405b-407b  
        equigranular, 109a  
        finite strain intensity and texture of, 418b-419b  
        formation, location of, 441b  
        fractionation trends, 12b-13b  
        fracture density, 105a  
        granular, 109a-110a, 117a  
        hydration source, 209b  
        interaction with crustal fluids, 186b
- intrinsic rock properties, 556b  
layering, 121a-122a  
magmatic evolution, 6b, 12b  
magnetic properties, 283b-303b, 318b-319b  
magnetic susceptibility, 197a, 209a  
metasomatized, 119a  
Mg#, Indian ocean sites compared, 82b  
Mg#, vs. normative anorthite, 77b, 78b-79b, 80b  
Mg#, vs. normative magnetite, 77b  
microgabbro contact, 122a  
mineral covariation, 12b-13b  
mineralogy, metamorphosed rock, 167b  
oxide and sulfide mineralogy, 122a-129a  
parental magma, 85b, 86b, 98b-101b, 139b-141b  
permeability, 200a-205a, 333b-339b  
petrography, 143b  
petrology, 154b-155b  
phase assemblage vs. mineral changes, 26b, 33b  
phase layering, 128a  
physical properties, 90a, 209a-210a, 323b  
plagioclase An vs. mafic mineral content, 42b, 49b, 50b-51b  
plagioclase neoblasts, 131a  
plastic alteration, 129a-132a  
poikilitic, 110a, 116a  
porosity, 261b-262b  
primitive and evolved, mixed occurrence, 44b  
pyroxene porphyroclasts and neoblasts deformed, 161b-162b  
recrystallization, 104a, 108a  
seawater interaction, isotopic effects, 170b, 174b  
static alteration of undeformed, 138a  
tectonic evolution, 427b-428b  
textural types I-V, 418b, 421b  
textural variation, 119a-121a  
timing of magnetization, 300b-301b  
upward evolution, 130b-131b
- Atlantis II Fracture Zone  
    abundance of, 361b  
    cumulate, 473b  
    dredged vs. drilled collections, 381b, 385b  
    from transform volcano, 366b  
    geochemistry, 470b-488b  
    geochemistry, crust vs. ridge axis, 472b  
    laminations, mineral lineation, 460b  
    late intrusives, 445b  
    magmatic extension cracks, 460b  
    mineralogy, 470b-488b  
    oxide-bearing vs. oxide-free, 456b  
    petrography, 445b  
    petrography of late-magmatic, 448b  
    primary, 445b  
    recrystallization, 441b  
    reintrusion of migrating melt, 442b-443b  
    vs. other Southwest Indian Ridge transforms, 361b
- Atlantis II Fracture Zone: Site 732  
    alteration and deformation, 56a  
    cataclastic deformation, 51a-52a  
    metamorphic reaction rim, 53a  
    metamorphism, 51a-52a  
    petrography, 50a  
    plastic deformation, 51a  
    replacement pattern, 51a-52a  
    serpentinitized porphyroclasts, 51a
- Atlantis II Fracture Zone: Site 733  
    cumulus phase assemblage, 72a  
    gabbronorite and, 71a  
    geochemistry, 71a-73a  
    magnetic properties, 73a

- metamorphism, 67a-71a  
petrography, 65a-67a  
physical properties, 73a-75a  
protolith, 67a  
replacement minerals, 65a-66a  
trapped liquid, 72a  
Atlantis II Fracture Zone: Site 734, petrography, 85a  
slow-spreading centers, cooling and hydrothermal alteration model, 175b
- Gabbro**, augen gneissic  
    Atlantis Bank, 130a, 132a  
        anisotropy and shear-wave splitting of, 236b
- Gabbro**, diopsidic olivine, Atlantis II Fracture Zone, petrography, 447b
- Gabbro**, disseminated iron-titanium oxide, Atlantis Bank, petrography, 47b
- Gabbro**, disseminated iron-titanium oxide olivine, Atlantis Bank, mineralogy, 71b
- Gabbro**, disseminated oxide, Atlantis Bank, mineralogy, 90b
- Gabbro**, disseminated oxide olivine  
    Atlantis Bank, 32b  
        crystallization model, 35b, 37b  
        low-calcium pyroxene formation, 33b  
        mineralogy, 26b  
        modality, 31b  
        texture, 26b, 38b
- Atlantis II Fracture Zone  
        contacts, 459b, 461b  
        origin, 461b  
        petrography, 448b  
        texture, 459b-460b  
        vs. olivine gabbro, 459b
- Gabbro**, evolved oxide-bearing, Atlantis Bank, mineral chemistry, 5b
- Gabbro**, foliated  
    Atlantis Bank  
        during viscous flow, 401b-403b  
        orientation, 396b  
        physical properties, 157a  
        seismic reflection profiling, 224b
- Gabbro**, ilmenite, Atlantis Bank, ilmenite replacement, 135a
- Gabbro**, interstitial olivine-bearing iron-titanium oxide, Atlantis Bank, mineralogy, 72b
- Gabbro**, interstitial olivine-bearing titanium oxide  
    Atlantis Bank  
        inverted pigeonite in clinopyroxene cores, 49b, 72b  
        petrography, 47b
- Gabbro**, iron- and titanium-rich, Atlantis Bank, permeability, 205a
- Gabbro**, iron-titanium oxide  
    Atlantis Bank, 100a, 102a, 209a  
        crosscutting relationships, 83b, 98b  
        deformation, 90b, 121a  
        density, 232b  
        distribution, 113a-115a  
        electrical conductivity, 330b  
        euhedral, 44b  
        geochemical logging data, 276b  
        geochemistry, 146a-147a  
        grain size variation, 48b  
        in-situ vs. observed NRM values, 293b  
        intensity of magnetization, 315b  
        layer 3, 319b  
        logging, 178a  
        magmatic foliation, 121a  
        magmatic lamination, 44b, 48b  
        magnetic properties, 288b-289b, 292b, 312b, 313b, 318b  
        metamorphism and alteration, 116a  
        mineralogy, 41b, 44b, 49b, 71b, 72b
- origin of variations, 59b, 61b-62b  
    unimodal distribution, 61b
- modal abundance, 48b  
    olivine gabbro contact, 56b, 59b, 73b, 119a  
    petrography, 43b, 112a-117a  
    plagioclase An vs. mafic mineral content, 64b  
    pyroxene-rich layer, 116a  
    secondary component of magnetization, 293b, 312b, 314b  
    seismic reflection profiling, 224b  
    texture, 41b, 44b  
    velocity and alteration of, 233b-234b  
    velocity-density relationship, 238b  
    Atlantis II Fracture Zone, 448b
- Gabbro**, magnesian, Atlantis Bank, normative vs. microprobe mineralogy, 5b
- Gabbro**, magnetite-ilmenite  
    Atlantis Bank  
        demagnetization, 154a  
        magnetic susceptibility, 151a-152a  
        magnetometer logging, 198a  
        oxygen isotopes, 139b
- Gabbro**, mylonitic  
    Atlantis Bank, 130a, 132a  
        deformation structures, 103a  
        density, 232b  
        physical properties, 157a  
        plagioclase composition, 199b
- Gabbro**, olivine  
    Atlantis Bank, 89a, 90a, 100a, 102a, 208a, 209a  
        bimodal distribution, 103b-104b  
        contact relationships, 28b  
        crystallization model, 35b, 37b  
        deformation, 3b, 501b-502b, 534b  
        elevated oxide percentages, 31a, 33a  
        formation, 110a-111a  
        gabbronorite and, 509b-510b  
        geochemical logging data, 276b  
        geochemistry, 146a  
        grain size transition, 29b  
        hydrodynamic crystal sorting, 29b  
        ilmenite-magnetite ratio, 124a  
        intercumulus phases, 111a  
        iron-titanium oxide gabbro contact, 56b, 59b, 73b, 119a  
        lithologic composition and structure, 34b  
        magnetic properties, 287b-288b, 289b-290b, 311b, 312b, 313b  
        mineralogy, 25b, 30b, 41b  
        modality, 29b, 31b  
        nickel-rich, 6b  
        oxide-bearing gabbro contact, 27b, 34b, 79b  
        permeability vs. log porosity, 267b  
        petrography, 107a-111a  
        poikilitic, 109a  
        size grading, 129a  
        solid-state deformation, 426b  
        sulfides in magnesian, 90b  
        texture, 25b, 38b, 41b, 121a  
        titanium oxide content of clinopyroxene, 51b  
        troctolites and, 27b  
        velocity and alteration of, 233b  
        velocity-density relationship, 238b
- Atlantis II Fracture Zone  
        crosscutting veins and lenses, 457b  
        folded contacts, 469b  
        geochemistry, vs. olivine-free gabbros, 459b  
        late-magmatic deformation, 457b, 467b  
        microgabbro contacts, 457b-158b  
        modal and grain-size layering, 442b  
        oxide gabbro contacts, 457b, 459b  
        oxide-rich intervals, 465b, 466b  
        patchy olivine gabbro relationship, 458b  
        pegmatitic patches, 468b
- petrogenesis, 459b  
    petrography, 445b  
    Unit V, 465b  
    Unit VI, 467b
- Gabbro**, olivine iron-titanium oxide, Atlantis Bank, petrography, 48b
- Gabbro**, olivine oxide  
    Atlantis Bank  
        fine- and coarse-grained contacts, 32b  
        modality, 33b
- Gabbro**, olivine pigeonite  
    Atlantis Bank  
        crystallization temperature, 49b  
        petrography, 47b
- Gabbro**, oxide  
    Atlantis Bank  
        chrome content, 483b, 486b  
        copper content, 120b, 123b  
        emplacement, 33b  
        igneous sulfides, 114b  
        lithologic composition and structure, 34b  
        magmatic foliation, 427b  
        mineralogy, 90b  
        nickel content, 120b, 123b  
        olivine gabbro contact, 27b, 31b, 34b, 79b  
        recrystallization of oxide minerals, 91b  
        sulfur content, 120b, 123b  
        troctolites and, 27b, 29b  
        trondhjemetic intrusion breccia, 503b
- Atlantis II Fracture Zone  
        absence of olivine, 448b  
        in brittle-ductile deformation zones, 469b  
        contact relationships, oxide-free gabbros, 480b, 482b, 488b  
        contact relationships, undeformed olivine gabbros, 470b  
        cross-foliated contacts, 460b  
        deformation and distribution of, 486b  
        formation, 486b  
        geochemistry, 456b, 479b-480b, 482b-483b  
        mineralogy, 479b-480b, 482b-483b  
        penetrative foliation, 460b  
        petrography, 448b  
        protolith, 469b-470b  
        textural relationships, 461b, 498b
- Gabbro**, oxide, foliated, Atlantis II Fracture Zone, soda content, 498b-499b
- Gabbro**, oxide olivine  
    Atlantis Bank  
        clothespin structure, 538b  
        crystallization model, 35b, 37b  
        mineralogy, 26b-27b  
        texture, 26b-27b
- Atlantis II Fracture Zone  
        contact relations, 462b, 464b  
        origin, 464b-465b  
        oxide content, 462b  
        penetrative deformation, 469b  
        texture, 464b, 475b  
        in weakly ductily deformed shear zones, 468b
- Gabbro**, oxide-free  
    Atlantis II Fracture Zone  
        deformation resistance, 462b  
        geochemistry, 475b-479b  
        mineralogy, 476b-477b  
        origin, 473b  
        texture, hypersolidus vs. subsolidus, 498b
- Gabbro**, patchy olivine  
    Atlantis Bank, texture, 507b-508b  
    Atlantis II Fracture Zone, 443b, 444b-445b  
        intrusive contacts, 462b  
        petrogenesis, 458b  
        petrography, 447b  
        protolith, 470b

## SUBJECT INDEX

texture, 467b-468b  
**Gabbro**, patchy oxide olivine, Atlantis II Fracture Zone, petrography, 448b  
**Gabbro**, pigeonite  
 Atlantis Bank, 116a, 120a  
 crystallization temperature, 49b  
 petrography, 47b  
**Gabbro**, pigeonite-oxide, Atlantis Bank, 32b  
**Gabbro**, porphyroclastic  
 Atlantis Bank, 130a  
 amphibole replacement, 134b  
 foliation, 131a, 134b  
 ilmenite-magnetite intergrowths, 134a, 135a  
 primary petrography, 107a  
 protoliths, 107a  
**Gabbro**, pyroxene  
 Atlantis II Fracture Zone: Site 732,  
 geochemistry, 54a  
 Atlantis II Fracture Zone: Site 733,  
 deformation, 63a  
**Gabbro**, troctolitic  
 Atlantis Bank, 90a, 209a  
 Atlantis II Fracture Zone  
 geochemistry, 478b  
 petrography, 447b  
 undeformed, 469b  
 Unit VI, 468b, 469b  
**Gabbro**, troctolitic oxide, Atlantis II Fracture Zone, petrography, 448b  
**Gabbro**, veined mylonitic  
 Atlantis Bank  
 hydrous and anhydrous, 187b  
 petrography, 215b  
**Gabbronorite**  
 Atlantis Bank, 155b  
 mineralogy, 25b-27b, 71b  
 olivine gabbro contact, 71b  
 petrography, 44b  
 texture, 25b-27b  
 Atlantis II Fracture Zone, 449b, 456b-457b  
 bimodal distribution of plagioclase, 477b  
 olivine gabbro contact, 456b, 458b  
 petrography, 445b  
 Atlantis II Fracture Zone: Site 733  
 gabbros and, genetic link, 71a  
 protoliths, 66a  
**Gabbronorite**, olivine  
 Atlantis Bank  
 mineralogy, 71b  
 petrography, 47b  
**Gabbronorite**, oxide, Atlantis II Fracture Zone, petrography, 448b  
**Gabbronorite**, oxide-free, Atlantis II Fracture Zone, geochemistry, 478b-479b  
**Gabbronorite**, poikilitic, Atlantis Bank, 44b  
**Gadolinium**, Atlantis Bank, geochemical logging data, 175a, 178a, 280b  
**Galapagos Rift**  
 basalt, fractionation, 29b  
 ferroandesite, ilmenite-magnetite proportions, 31b  
 ferrogabbro, ilmenite-magnetite proportions, 31b  
**Galapagos Spreading Center**  
 Atlantis Bank, crystallization intervals, 90b  
 liquid evolution, vs. Skaergaard Intrusion, 100b-101b  
 mineralogy, during differentiation, 91b, 93b  
 oxygen fugacity, 102b  
 strontium, basalt glasses, 84b  
**Garrett Fracture Zone**  
 clinopyroxene, chromium oxide-enriched, 6b  
 gabbro  
 high temperature deformation, 51a  
 mineral covariation, 12b

nickel-rich olivine, 6b  
**Geochemical logs**  
 Atlantis Bank, 169a, 174a, 175a, 177a-178a  
 comparison, XRF analyses, 178a, 179a  
 lithostratigraphic correlation, 276b-280b  
**Geochemical logging tool (GLT)**  
 Atlantis Bank, 271b  
 centroid compositions, 277b  
 chemostratigraphic Zones P1-P7, 277b-280b  
 chemostratigraphic Zones Z1-Z6, 276b-277b  
 data analysis, 272b-281b  
 INCA analysis, 273b-280b, 281b, 282b  
 principal component analysis, 273b, 283b  
**Geochemistry**  
 Atlantis Bank  
 alteration effects, 136b-138b, 546b-547b  
 chemical stratigraphy, 147a-150a  
 loss-on-ignition values, 139a  
 methods, 139a  
 modal gap, 480b, 482b, 488b  
 permeability and, 502b  
 stratigraphic logs, 13b  
 Unit IV/Unit V boundary, 147a  
 Zone IV, 77b-78b  
 Zones I-III, 77b  
 Atlantis II Fracture Zone  
 active ridge sediments compared, 148b  
 between rock types, 474b  
 bimodal distribution, 473b, 475b  
 bulk-rock composition, diabase correlation, 483b-485b  
 dredged vs. drilled gabbros, 471b  
 lithologic variability, 471b  
 MORB-like composition, 485b-486b  
 oceanic pelagic sediments compared, 147b  
 oxide-bearing vs. oxide-free gabbros, 486b  
 post-cumulus exchange, 487b  
 transform vs. ridge axis samples, 472b  
 Atlantis II Fracture Zone: Site 732  
 igneous vs. biogenic components, 147b  
 sand- vs. clay-sized fraction, 148b-149b  
 source provinces, 149b  
 Atlantis II Fracture Zone: Site 733  
 gabbro, 71a  
 trace elements, gabbros vs. basalts, 72a  
 Atlantis II Fracture Zone: Site 734  
 gabbros, 86a  
 igneous vs. biogenic components, 147b  
 sand- vs. clay-sized fraction, 148b-149b  
 source provinces, 149b  
**Geochemistry**, whole-rock  
 Atlantis Bank, 527b-530b  
 cyclic variations, 5b  
 lithostratigraphic correlation, 4b-5b  
**Glass**, Atlantis II Fracture Zone, mineralogy, 382b  
**Glass**, basalt  
 Atlantis Bank  
 major oxides, 86b-87b  
 phosphate, 88b  
 phosphate vs. magnesium oxide, 103b  
 potassium oxide, 88b  
 sulfur content, 88b-89b, 99b  
 Type 2, 86b-87b  
 Type 3, 87b  
 Atlantis II Fracture Zone  
 calcium vs. magnesium number, 384b  
 mineral composition, east-west discontinuity, 88b  
**Gneiss**, Atlantis Bank, alteration, 179b  
**Gneiss**, amphibole  
 Atlantis Bank  
 foliated, 549b  
 mineralogy, 543b

**Gneiss**, foliated, Atlantis II Fracture Zone, 51a, 52a, 449b  
**Gorringe Bank**  
 gabbro, braided brittle-ductile deformation, 501b  
 greenschist-facies alteration, absence of, 489b  
**Grain size**  
 Atlantis Bank, 124a  
 basalt, 117a  
 with deformation, 130a  
 iron-titanium oxide gabbro, 48b  
 magmatic origin and variation in, 3b  
 olivine gabbro, 107a  
 in plastically deformed gabbros, 400b, 403 b-406b  
 Atlantis II Fracture Zone  
 disseminated oxide olivine gabbro, 459b  
 oxide olivine gabbros, 464b  
 Atlantis II Fracture Zone: Site 734, 84a  
**Granulite facies** deformation, Atlantis Bank, temperature-dependent diffusion, 407b, 421b  
**Gravel**  
 Atlantis II Fracture Zone, 431b-438b  
 gravity-driven transport, 431b-432b  
 Atlantis II Fracture Zone: Site 732, Pleistocene, 373b  
 Atlantis II Fracture Zone: Site 734  
 depositional mechanism, 436b  
 graded beds, 432b-433b  
 high-temperature metamorphism, 435b  
 length measurements, 435b  
 origin, 434b-435b  
 shape, 437b  
 shape, transport behavior and, 434b  
 size distribution, 436b  
 sorting, 432b-433b  
**Greenschist**  
 Atlantis Bank, 165b  
 in felsic veins, 504b  
**Greenschist-facies** alteration  
 Atlantis Bank, 172b  
 absence of, 319b, 489b  
 in hydrothermal breccias, 505b  
 Atlantis II Fracture Zone, dredged vs. drilled gabbros, 381b  
 retrograde metamorphic association, 489b, 491b  
**Halogen**, Atlantis Bank, in hydrated minerals, 208b  
**Hard-rock guidebase (HRGB)**  
 Atlantis II Fracture Zone, assembly and deployment, 28a-30a  
 mechanical design evacuation, 30a-31a  
**Harzburgite**, Atlantis II Fracture Zone: Site 732, serpentinization, 53a-54a  
**Hawaiian Kileau Rift**, constructional volcanism, 366b  
**Heat flux**  
 Atlantis Bank, 350b-351b  
 ocean temperature warming and surface, 354b  
**Heliosphaera sellii**, Atlantis II Fracture Zone: Site 734, 94a  
**Hornblende**  
 Atlantis Bank  
 chlorine content, 427b  
 clinopyroxene replacement, 198b  
 episodic growth, 159b  
 fluid inclusions, 201b, 202b  
 mineral chemistry, 212b  
 oscillatory zonation, 188b  
 oxygen isotopes, 206b  
 vein to foliation-plane mineral phase, 163b  
 Atlantis II Fracture Zone: Site 733, velocity, 74a  
**Hornblende**, actinolitic  
 Atlantis Bank

in amphibolite gneisses, 544b  
temperature of formation, 172b

Hornblende, brown  
Atlantis Bank  
clinopyroxene replacement, 136a  
recrystallized, 403b, 404b-405b  
secondary parasitic, 160b  
Atlantis II Fracture Zone: Site 733, anhedral relicts and blebs, 69a

Hornblende, brownish-green  
Atlantis II Fracture Zone: Site 732, 51a  
Atlantis II Fracture Zone: Site 733, 67a

Hornblende, green  
Atlantis Bank, 160b  
Atlantis II Fracture Zone, 67a  
crosscutting veins, 71a  
porphyroclasts, 52a

Hornblende, parasitic, Atlantis Bank, magmatic origin, 170b

Hornblende, reddish-brown  
Atlantis Bank, texture, 117a  
Atlantis II Fracture Zone, 50a, 67a

Hornblende-plagioclase assemblage, Atlantis Bank, 160b

Horsts  
Atlantis Bank, formation, 153b  
Atlantis II Fracture Zone  
formation, 441b  
irregularly shaped, 368b

*huxleyi*, *Emiliania*, Atlantis II Fracture Zone: Site 732, 49a

Hydration  
Atlantis Bank  
degree of deformation and, 186b  
seawater source, 209b, 213b

Hydrogen  
Atlantis Bank  
geochemical log, 174a, 178a  
porosity and content of, 262b

Hydrogen isotopes  
Atlantis Bank, 133b, 135b-136b  
gabbro, 129b-131b

Hydrothermal fluid, methane-bearing, Atlantis Bank, role in deep crustal processes, 209b

Icelandic Ridge, constructional volcanism, 366b

Igneous lamination, Atlantis Bank, in olivine gabbro, 125a

Igneous rock  
Atlantis Bank  
mineralogy, 23b-27b  
primary lithologies, 509b-510b, 516b-526b  
primary stratigraphy, 22b  
texture, 23b-27b

Atlantis II Fracture Zone  
classification, 444b-448b  
distribution, 444b  
intrusion model, 512b-513b  
petrogenesis, 441b-444b  
petrography, 65a-67a

Illite, Atlantis II Fracture Zone, source provinces, 147b

Ilmenite  
Atlantis Bank  
in deformed gabbros, 131a, 133a-134a, 155b  
derivation, 160b  
effect on oxide concentrations, 91b  
host-rock foliation and, 134a  
magnesium oxide vs. manganese oxide, 99b  
magnetite intergrowths, 124a-125a  
 $Mg\#$ , 93b  
minor elements, 56b  
mobilization, 135a  
in orthopyroxene-bearing gabbro, 135a

oxidative alteration, 127a  
parental magma concentration, 93b  
titanium oxide and, 5b

Atlantis II Fracture Zone  
grain size, in oxide olivine gabbros, 464b  
ore bodies, 444b  
Atlantis II Fracture Zone: Site 732, 51a

Ilmenite, cumulus, Atlantis II Fracture Zone: Site 732, 53a

Ilmenite, primary igneous, Atlantis II Fracture Zone: Site 732, 52a

Ilmenite/magnetite ratio  
Atlantis Bank  
in deformed gabbro, 132a-134a  
in ferrogabbros, 124a  
in iron-titanium oxide gabbro, 115a  
in undeformed olivine gabbros and troctolites, 124a

Impedance, Atlantis Bank, changes causing seismic reflection, 190a

Indian Ocean  
basalt, lead isotopes, 136b, 137b, 138b  
mica, in silicic dikelets, 168b  
strontium, basalt glasses, 84b

Indian Ocean E, geochemistry, 148b

Indian Ocean S  
bathymetry, 4a  
geochemistry, 148b

Indian Ocean Triple Junction, 361b

Intrusion, assimilative, Atlantis II Fracture Zone, olivine gabbros, 467b

Intrusion, late, Atlantis Bank, gabbro contact relationships, 507b-509b

Intrusion, microgabbro  
Atlantis Bank  
crosscutting olivine gabbro, 507b, 509b, 511b  
podiform, 511b

Intrusion, troctolitic  
Atlantis Bank, 511b  
Atlantis II Fracture Zone, 442b  
thermal gradient at emplacement, 470b

Intrusive events  
Atlantis Bank, 56b, 59b  
deformation and, 85b, 103b-104b  
during deformation, 27b  
sequence, 85b  
trondhjemite breccia, 27b

*See also* Late magmatic intrusions (LMIs)

Iron  
Atlantis Bank  
geochemical log, 175a  
mafic phases, 30b

Atlantis II Fracture Zone, dredged vs. drilled gabbros, 381b

Atlantis II Fracture Zone: Site 732, xenocrysts, 49a

Iron oxide  
Atlantis Bank, 135b  
geochemical logging data, 276b, 278b  
velocity and percentage of, 247b-248b, 250b

Iron oxide/magnesium oxide ratio  
Atlantis Bank  
Atlantic vs. Pacific lavas, 13b  
clinopyroxene, coexisting liquid correlation, 19b  
liquids in equilibrium with cumulus olivine, 13b, 20b  
olivine, coexisting liquid correlation, 19b

Iron oxyhydroxides, Atlantis Bank, 125a

Iron-enrichment differentiation  
Atlantis Bank  
liquid immiscibility and, 96b  
melt densities produced, 98b  
pyroxenes, 91b

Iron-titanium oxides  
Atlantis Bank, 99a  
iron-titanium mobility during metamorphism, 127a  
plastic deformation association, 105a  
relation to silicate phases, 126a-128a  
secondary, 107a

Iterative nonhierarchical cluster analysis (INCA), Atlantis Bank, 273b-275b

Jasper Seamount, abyssal-gabbro xenolith, mineralogy during differentiation, 91b, 93b

Juan de Fuca Ridge, xenoliths, in ferrobasalt, 4b

Kane Fracture Zone  
detachment fault, 441b  
gabbro, 361b  
magnetic susceptibility, 209a  
geometry, 366b  
greenschist-facies alteration, 489b  
hooked ridge, pillow lava flows, 376b  
magnetic properties, Koenigsberger Ratio, 153a  
metagabbro, velocity, 74a  
neovolcanic zones, 376b  
normal faulting, ridge-transform intersection, 428b  
northern active transform wall, comparison, Atlantis II Fracture Zone, 376b  
oblique normal faults, 501b  
seismic stratigraphy, 219b  
seismic structure, 183a, 185a  
talus ramps, 431b  
transform valley, intersection with Mid-Atlantic Ridge, 394b  
transform walls  
dredging survey, 377b  
vs. nontransform walls, 392b  
transverse ridge, interval velocity, 225b  
velocity, 190a

Kaolinite, Atlantis II Fracture Zone, source provinces, 147b

Kiglapait Intrusion, melt porosity, 81b-82b

Labradorite  
Atlantis Bank  
in feldspar, 199b  
fluid inclusions, 187b, 202b

*lacunosa*, *Pseudoemiliania*  
Atlantis II Fracture Zone: Site 732, 49a  
Atlantis II Fracture Zone: Site 734, 94a

Lamellae, exsolution, Atlantis Bank, pyroxenes, 4b

Lamination, igneous, Atlantis Bank, 101a, 106a-107a, 121a

Lamination, magmatic, Atlantis Bank, iron-titanium oxide gabbro, 44b, 48b

Late magmatic intrusions (LMIs)  
Atlantis Bank, 155b  
alteration near, 163b, 165b  
assimilation, 177b-178b  
cataclastic association, 159b  
magmatic pyroxene, 161b-162b  
mineralogy, 167b, 168b  
structures not related to, 172b-173b  
structures related to, 172b

Atlantis II Fracture Zone, contact relationships, 445b

Layer 3  
Atlantis Bank  
alteration, 543b-548b  
attenuation, 253b-257b  
contribution to marine magnetic anomalies, 300b-302b, 305b, 320b  
fluid-rock interaction, 186b

## SUBJECT INDEX

- fluids, 181b  
 formation, 513b  
 geological evolution, 509b, 510b  
 logging data, 325b  
 magnetic properties, 320b  
 mineralogy, 542b-543b  
 petrography, 542b-543b  
 physical properties, 323b  
 seismic reflection profiling, 245b  
 sulfur depletion and seawater interaction, 123b, 124b  
**Atlantis II Fracture Zone**, lithostratigraphic evolution, 439b-513b  
 development, slow-spreading ridges, 21b  
 Southwest Indian Ridge, transforms, 361b  
**Layering**  
 Atlantis Bank, 100a, 130b  
 olivine gabbro, 47b  
**Lead isotopes**  
 Atlantis Bank  
   comparison, MORB and Southwest Indian Ridge (SWIR) basalts, 131b  
 gabbro, 129b-133b, 137b  
   systematic variation, 132b  
   vs. neodymium isotopes, 138b  
**Atlantis II Fracture Zone**, vs. neodymium isotopes, 138b  
**Leucogabbronorite**, Atlantis II Fracture Zone: Site 733, 66a  
**Liquid line of descent**  
 Atlantis Bank, 95b-101b  
   attributes, 82b-83b  
   in gabbro formation, 75b  
   liquid immiscibility, 96b-98b, 104b, 119b  
   melt density, 98b  
   mineral composition and, 27b  
   petrography, 90b  
**Madagascar Basin**  
   kaolinite source, 147b  
   quartz source, 147b  
**Mafic rock**  
   Atlantis Bank, sulfur isotopes, 121b, 122b  
   Atlantis II Fracture Zone, petrography, 85a  
**Magma chamber**  
   Atlantis Bank, distinctive magma introduction through time, 140b, 141b  
   fast-spreading ridges, 509b  
   spreading ridge segmentation and, 3a  
   subrift, gabbros generated by, 14b  
**Magma chamber process**, slow-spreading ridges, boundary-layer diffusion, 86b  
**Magma mixing**  
 Atlantis Bank, 85b  
**Magma, parental**  
 Atlantis Bank  
   heterogeneity of, 83b-86b, 139b-141b  
   lithostratigraphic correlation, 85b  
   oxide content, 98b-101b  
   signature retention, 85b-86b  
   sodic, 86b  
   sulfur content, 101b  
**Magmatic differentiation**, Atlantis Bank, deforming crust, 103b-104b  
**Magmatic events**, Atlantis Bank, downward intrusion and evolution, 62b  
**Magmatic fluid**, Atlantis II Fracture Zone, source, 487b-488b  
**Magmatic layering**, Atlantis Bank, as hydrothermal conduit, 139a  
**Magnesio-hornblende**, Atlantis Bank, chlorine content, 198b, 543b  
**Magnesium**  
 Atlantis Bank  
   normative vs. measured, 5b  
   vs. sodium, 272b  
**Atlantis II Fracture Zone**, in ultramafic rocks, 13a  
**Magnesium number**  
 Atlantis Bank  
   diabase, 476b  
   gabbros, 100a, 476b  
   liquid vs. glass group averages, 82b  
   oxide vs. olivine gabbros, 83b  
**Magnesium oxide**  
 Atlantis Bank, 136b  
   geochemical logging data, 279b  
   in ilmenite, 56b  
**Magnesium/calcium ratio**, Atlantis Bank, oxide-free gabbros, 475b-476b  
**Magnetic anomalies**  
 Atlantis Bank, 555b  
   correlation, reversed magnetization, 292b  
   magnetic properties and, 320b  
   models, 307b  
   oceanic intrusive rocks and origin of, 300b-305b  
   Vine-Matthews model, 150a, 309b  
**Atlantis II Fracture Zone**  
   anomalies 1-5, 360b, 382b-384b, 386b  
   central anomaly, 382b  
   eastern vs. western rift valley, 388b  
   negative mass anomaly, 395b  
   transform slip, anomalies, 1-2A, 392b  
   truncated magnetic-anomaly source layer, 384b, 388b-389b  
**Magnetic properties**  
 Atlantis Bank, 151a-157a, 209a  
   alternating-field demagnetization, 153a-154a, 155a, 156a, 291b-292b, 293b-294b, 302b  
   degree of metamorphism/alteration and, 294b-296b, 300b  
   directional changes, 291b  
   directional coherence, 319b-320b  
   discrete samples vs. logging data, 309b-310b, 311b, 313b-318b  
   drilling-induced remanent magnetization, 294b  
   gabbros, 197a-198a, 287b-290b, 318b-319b  
   inclination, 154a-156a, 286b, 291b, 292b, 293b, 296b, 297b-300b, 301b, 302b, 315b-317b  
   intensity, 313b-315b, 318b  
   Koenigsberger Ratio, 153a, 154a, 290b-291b, 292b, 293b, 296b, 300b, 305b  
   lithostratigraphic correlation, 310b-313b  
   logging errors, 317b-318b  
   median demagnetizing field, 153a-154a, 155a, 291b, 292b, 300b, 305b  
   NRM intensity, 286b, 554b-555b  
   oxide minerals, 78b  
   remanent magnetization, 151a, 153a, 156a  
   secondary component of magnetization, 293b-294b, 314b, 317b, 320b  
   stable inclination, 157a  
   susceptibility, 151a-153a, 154a, 286b, 291b, 314b, 315b, 316b, 318b  
   thermal demagnetization, 154a, 156a, 291b-292b, 296b  
   Unit II/Unit III boundary, 313b  
   vertical magnetic field, 316b, 317b  
   viscous remanent magnetization, 294b  
**Atlantis II Fracture Zone**, 21a  
**Atlantis II Fracture Zone: Site 733**, 73a, 75a  
   vs. physical properties, 76a  
*See also* Magnetic anomalies; Stable remanent magnetic vector  
**Magnetite**  
 Atlantis Bank  
   alteration, 91b  
   in amphibole/talc reaction zones, 127a-128a  
   in deformed gabbro, 133a-134a, 155b  
   ilmenite intergrowths, 124a-125a  
   NRM intensity effect, 293b  
   oxidative alteration, 127a  
   parental magma concentration, 93b  
   Unit IV gabbros, 312b  
**Atlantis II Fracture Zone: Site 732**, parallel to foliation, 54a  
**Magnetometer log**  
 Atlantis Bank, 210a  
   azimuthal reference readings, 192a  
   conductivity signal, 192a  
   equipment, 191a-193a, 197a  
   gravitational orientation of elements, 191a, 201a  
   parameters, 192a, 193a, 197a, 202a  
   susceptibility, 192a, 197a-198a, 201a, 554b  
   three-component magnetometer probe, 191a, 200a  
   USGS magnetic-susceptibility probe, 191a-192a, 200a  
   variability of magnetic north, 198a  
**Manganese oxide**  
 Atlantis Bank  
   in amphiboles, 56b  
   in clinopyroxene, 51b, 53b, 65b  
   in olivine, 56b  
   in orthopyroxene, 53b, 67b  
**Mantle rock**, fracture zones, ridge spreading rate and, 42a  
**Marie Celeste Fracture Zone**, geochemistry, 148b  
**Marine magnetic anomalies**. *See* Magnetic anomalies  
**Mass-wasting**  
 Atlantis II Fracture Zone  
   alteration minerals, 435b  
   ridge-transform intersection, 431b  
**Median tectonic ridge**  
 Atlantis II Fracture Zone  
   origin, northern ridge, 371b  
   physiography, 371b-373b  
   seismic stratigraphy, 370b  
   transtensional spreading and formation of, 392b  
**Melanogranoophyses**, Atlantis Bank, inhibited evolution, 102b-103b  
**Melt density**, Atlantis Bank, effect of iron-rich differentiation on, 98b  
**Melt**, leucocratic  
 Atlantis Bank, 425b  
   dikelet, 426b  
**Melt migration**  
 Atlantis Bank, during deformation, 488b, 507b  
 Atlantis II Fracture Zone, permeability and pressure controls, 487b  
**Melt porosity**  
 Atlantis Bank  
   deformation and, 81b  
   determining residual, 79b-82b, 106b-107b  
   phosphate content and, 81b  
   shear zones, 82b  
   vs. Skaergaard Intrusion, 81b  
   zirconium content and, 80b, 81b, 83b  
**Melt, tholeiitic**, Atlantis II Fracture Zone, crystallization models, 487b-488b  
**Melt, trapped**  
 Atlantis Bank  
   expelled during deformation, 82b  
   in olivine gabbro, 59b  
   phosphate/zirconium ratio and, 79b  
**Melville Bank**, plutonic rocks, shallow depth, 77b

- Metabasalt**
- Atlantis II Fracture Zone: Site 733
    - petrography, 67a
    - primary mineralogy, 65a
    - replacement pattern, 70a–71a, 75a
  - Atlantis II Fracture Zone: Site 734,
    - metamorphism, 86a, 435b
  - Metabasalt, foliated, Atlantis II Fracture Zone: Site 734, phosphate/titanium oxide composition, 88b
  - Metabasalt, granoblastic, Atlantis II Fracture Zone: Site 734, geochemistry, 86a–87a
  - Metagabbro
    - Atlantis Bank
      - with amphibole veins, 136a
      - anisotropy and preferred plagioclase orientation, 234b, 236b
      - fluid inclusions in plagioclase, 190b
      - hornblende replacing clinopyroxene, 198b
      - physical properties, 157a
      - plagioclase composition, 199b
    - Atlantis II Fracture Zone: Site 732, 50a
    - Atlantis II Fracture Zone: Site 733
      - foliation, 67a–68a
      - magnetic properties, 60a
      - physical properties, 60a, 75a, 76a
      - plastic deformation, 60a
      - protoliths, 60a
    - Atlantis II Fracture Zone: Site 734, 78a
      - incompatible trace elements, 86a
      - metamorphism, 86a, 435b  - Metagabbro, brown
    - Atlantis II Fracture Zone: Site 733
      - metamorphism, 68a–69a
      - protolith, 75a
      - replacement pattern, 69a, 70a, 71a  - Metagabbro, foliated
    - Atlantis Bank, 89a
      - magnetic properties, 287b
      - magnetic properties, logged, 311b
      - physical properties, 157a
      - pyroxene porphyroblast, 549b
      - sulfide and oxide mineralogy, 125a
      - sulfur depletion, 121b, 123b
    - Atlantis II Fracture Zone: Site 733
      - plastic deformation, 75a
      - primary mineralogy, 65a  - Metagabbro, green
    - Atlantis II Fracture Zone: Site 733
      - gneissic foliation and lineation, 69a
      - metamorphism, 67a–68a
      - protolith, 75a  - Metagabbro, mylonitic
    - Atlantis Bank
      - physical properties, 157a
      - velocity, 163a  - Metagabbro, mylonitic, veined, Atlantis Bank, fluid inclusions, 202b
  - Metagabbro, porphyroclastic
    - Atlantis II Fracture Zone: Site 734
      - geochemistry, 86a
      - phosphate/titanium oxide composition, 88b  - Metagabbro, veined
    - Atlantis Bank
      - fluid inclusions, 202b–204b
      - modality and extent of deformation, 186b–187b
      - petrography, 186b–187b, 214b–215b  - Metagabbronorite, Atlantis Bank, ductile deformation, 423b
  - Metamorphic rock
    - Atlantis Bank, 130a
      - stratigraphy, 156b–157b, 172b, 175b
- Metamorphism**
- Atlantis Bank, 209a
    - contribution to marine magnetic anomalies, 303b–304b
    - deformation and, 139a, 171b–172b
    - extent, 153b–154b
    - low-temperature, breccia zones, 172b
    - magnetic properties and degree of, 294b–296b, 300b, 303b
    - mineral assemblages associated with, 159b–170b
    - mineralogical effect, 171b–172b
    - oxide-sulfide relationship, 134a–136a
    - seawater penetration and, 153b, 174b, 175b
  - Atlantis II Fracture Zone
    - dredged vs. drilled gabbros, 381b
    - mafic rocks, 86a
  - Metamorphism, dynamothermal
    - Atlantis Bank, 90a
      - gabbro textures, 129a–132a
      - mineralogical changes, 113b–114b
      - sequence of events, 129a
      - sulfur depletion, 121b
  - Metamorphism, high-temperature, Atlantis Bank, mineral assemblages, 543b–544b
  - Metamorphism, hydrothermal, Atlantis Bank, alteration of gabbros, 208b
  - Metamorphism, medium-temperature, Atlantis II Fracture Zone: Site 732, cataclastic deformation and, 51a–53a
  - Metamorphism, static
    - Atlantis Bank
      - pseudomorphous and coronitic replacements, 163b
      - sulfur depletion, 121b
  - Metamorphism, synkinematic, Atlantis Bank, assemblages, 420b–421b, 423b, 426b
  - Methane
    - Atlantis Bank
      - in plagioclase inclusions, 204b, 206b
      - reduction of seawater carbonate to, 209b
      - source, 209b
      - in veined metagabbros and trondjemite, 182b
  - Mica
    - Atlantis Bank, 168b
      - metamorphosed gabbros and late magmatic intrusions (LMIs), 735b
      - mineral chemistry, 200b
  - Mica, brown, Atlantis Bank, 119a
  - Mica, green, Atlantis Bank, 119a
  - Microcracks
    - Atlantis Bank
      - opaque mineral petrology, 116b
      - syokinetic, 159b
  - Microfaults, Atlantis II Fracture Zone, crosscutting late-magmatic foliation, 499b
  - Microgabbro
    - Atlantis Bank, 118a
      - amphibolitized, 117a
      - gabbro contacts, 122a
      - textural variation, 130a
    - Atlantis II Fracture Zone
      - crosscutting olivine gabbro, 443b
      - mineralogy, late intrusive, 474b
      - origin, 474b
  - Microgabbro, olivine
    - Atlantis Bank, 102a
      - contact relationships, 28b
      - mineralogy, 25b
      - modal proportions, 110a
      - texture, 25b
    - Atlantis II Fracture Zone
      - intrusive into olivine gabbro, 468b
      - olivine gabbro contacts, 457b
  - petrography, 445b, 447b
  - texture, 467b–468b
  - Microgabbro, olivine iron-titanium oxide, Atlantis Bank, petrography, 48b
  - Microgabbro, oxide olivine
    - Atlantis II Fracture Zone
      - contact relations, dip orientation, 464b
      - petrography, 448b
      - Unit IV, 462b
    - Microgabbro, troctolitic
      - Atlantis Bank
        - mineralogy, 24b–25b
        - texture, 24b–25b
      - Atlantis II Fracture Zone
        - disseminated oxide olivine gabbro contact, 460b
        - geochemistry, 458b
        - olivine gabbro contacts, 457b–458b
        - petrography, 447b
        - undeformed, 469b
        - Unit VI, 468b, 469b
    - Microgabbronorite, oxide
      - Atlantis Bank
        - olivine gabbro contacts, 533b
        - texture, 38b
      - Atlantis II Fracture Zone
        - geochemistry, 467b
        - olivine gabbro contacts, 467b
        - petrography, 445b
    - Mid-Atlantic Ridge
      - basalt, lead isotopes, 136b, 137b, 138b
      - eastern rift valley wall, Kane Fracture Zone
        - transform valley intersection, 394b
      - gabbro, 361b
        - magnetic properties, 73a, 76a
        - seamounts, ridge-transform intersection, 365b
    - Mid-Cayman Rise
      - amphibole gneiss, 543b
      - calcium oxide/aluminum oxide ratio, 13b–14b
      - gabbro
        - aluminum oxide, crystallization effect, 140b
        - clinopyroxene, chemical evolution, 6b
        - mineral covariation, 12b
        - nickel-rich olivine, 6b
    - Mid-Indian Ocean Ridge
      - geochemistry, 148b
      - uplift blocks, 91a
    - Mid-ocean-ridge basalt (MORB)
      - Atlantis II Fracture Zone
        - mineralogy, 361b
        - soda content, 380b
        - oxygen isotopes, 170b
        - sulfur isotopes, correlation, Atlantis Bank, 118b–119b, 121b
      - Mid-ocean ridges (MOR), crystallization, magma interaction and, 127b
      - Millerite, Atlantis Bank, in green amphibole, 117b
    - Mineralogy
      - Atlantis Bank, 5b, 99a, 108b–115b, 189b–200b
        - ambiguous origin, 166b–170b
        - amphiboles, 190b, 197b–198b
        - between rock types, 49b
        - bimodality, 27b, 33b, 34b, 78b, 86b
        - covariation, 12b–13b
        - crystallization model, 35b
        - cumulus phase changes, 49b
        - degree of deformation and, 128b, 129b
        - distribution of metamorphic, 171b–172b
        - elements excluded during fractionation, 86b
        - feldspar, 198b–199b
        - gabbros, 481b
        - genetic relationship between rock types, 33b–35b
        - high water/rock deformation assemblage, 177b

## SUBJECT INDEX

- igneous lithologies, 26b  
 immiscible separated siliceous vs. iron-rich liquids, 101b  
 incompatible element-rich mineral association, 166b-167b  
 iron-titanium oxide gabbros, 115a-116a  
     origin of variations, 59b  
     vs. olivine gabbro and troctolites, 49b  
 late differentiation, 97b  
 lithostratigraphic correlation, 77b, 184b-185b  
 low-water/rock deformation assemblage, 176b-177b  
 mafic mineral variations, 45b  
 mafic mineral variations, iron-titanium oxide gabbro, 46b, 64b  
 metamorphic assemblages, 420b  
 mineral orientation and seismic velocity, 248b-249b  
 minor elements in constituent minerals, 51b, 53b, 56b  
 moderate-temperature, high-water/rock deformation assemblage, 177b  
 normative vs. microprobe compositions, 5b  
 olivine vs. iron-titanium oxide gabbro, 59b  
 opaque phases, 101a  
 preferred orientation, 505b  
 secondary minerals, 199b-200b, 202b-203b  
 secondary minerals, temperature of alteration, 209b, 211b  
 Skaergaard Intrusion compared, 78b-79b, 95b, 102b  
 synkinematic, 160b, 163b  
 tholeiitic high-iron differentiation, 95b  
 unimodal distribution, 61b  
 Unit II/Unit III boundary, 208b  
 Unit III/Unit IV contact, 33b  
 variation between rock types, 27b, 29b  
 variation within rock types, 23b, 29b-31b  
 velocity and, 233b  
 zoning, 5b, 545b
- Atlantis II Fracture Zone  
 between rock types, 474b-475b  
 bimodal distribution, 444b  
 gabbros, 484b-485b  
 oxide-bearing vs. oxide-free gabbros, 475b, 486b  
 oxide-free gabbros, 476b-477b  
 source provinces, 149b
- Atlantis II Fracture Zone: Site 733  
 igneous rocks, 65a-67a  
 opaque phases, 66a  
 phase layering, 66a
- Moho discontinuity, serpentinization, physical properties, 163a
- Moho sequence, seismic stratigraphy, 219b, 225b
- Multichannel sonic (MCS) logging  
 Atlantis Bank, 264b, 266b  
     compressional and shear velocity correlation, 181a, 184a  
     fracture zones, 206a  
 digital waveforms, 180a-181a, 182a, 213a  
 MCS vs. LSS tools, 178a, 181a  
 operations, 178a, 180a-182a  
 semblance velocity logs, 181a-182a, 555b-556b
- Muskox Intrusion, reverse fractionation, 61b
- Mylonite  
 Atlantis Bank, 179b, 416b  
     anisotropy and orientation of amphiboles, 236b  
     crosscutting foliated oxide olivine gabbro, 532b  
     deformation mechanisms, 406b-407b  
     horizontal foliation, 425b
- oxygen isotope values, 174b  
 synkinematic assemblages, 423b  
 zonation, 159b
- Atlantis II Fracture Zone  
 albitization, 465b  
 amphibolite-facies deformation, 460b  
 foliation plane, 449b  
 texture, hypersolidus vs. subsolidus, 498b  
*See also* Amphibolite mylonite
- Mylonite, iron-titanium oxide-rich  
 Atlantis Bank, 117a, 402b, 406b  
     origin of oxides in, 407b
- Mylonite, serpentinized peridotite, Atlantis II Fracture Zone: Site 732, 53a, 54a
- Mylonite, veined  
 Atlantis Bank  
     modality and extent of deformation, 186b-187b  
     petrography, 186b-187b
- Mylonitic band, Atlantis II Fracture Zone: Site 733, geometry, 68a
- Mylonitic zones  
 Atlantis Bank  
     deformation structures, 103a  
     frequency of occurrence, 114a
- Nannofossils, calcareous. *See* Calcareous nannofossils
- Navdrill coring system  
 Atlantis II Fracture Zone, 6a  
     assembly, 34a-35a  
     deployment, 36a-37a  
     mechanical design evaluation, 37a-38a  
     testing, 35a-36a, 96a  
     flow rate, 34a, 36a, 38a
- Neodymium isotopes  
 Atlantis Bank  
     gabbro, 129b-133b  
     vs. strontium isotopes, 140b  
 Atlantis II Fracture Zone  
     vs. lead isotopes, 138b  
     vs. strontium isotopes, 136b
- Neovolcanic zone  
 Atlantis II Fracture Zone  
     median tectonic ridge, 371b  
     nontransform walls, 376b  
     rift valley, 365b-366b
- Nickel  
 Atlantis Bank, 136b  
     cyclical variations, 5b  
     in igneous sulfides, 115b, 119b-120b, 123b
- Nickel oxide, Atlantis Bank, in olivine, 56b
- Nodal basins  
 Atlantis II Fracture Zone  
     bisected and infilled, 373b  
     formation, 392b, 395b  
     physiography, 366b
- Oceanographer Fracture Zone  
 metagabbro, velocity, 74a  
 neovolcanic zones, 376b  
 refraction data, 225b  
 ridge axis, right-lateral offsets, 392b
- Oceanographer Ridge, seamounts, ridge-transform intersection, 365b
- Olivine  
 Atlantis Bank, 89a, 99a, 107a  
     chemical composition, 52b-53b, 69b  
     crystallization, 49b, 90b, 95b  
     crystallographic orientation, 403b, 404b, 407b  
     deformation features, 130a  
     dislocation slip, 404b, 406b  
     dynamic recrystallization, 401b, 406b  
     ferrosilite, vs. anorthite in plagioclase, 18b
- in foliated gabbro, 402b  
 forsterite, 62b, 68b  
 fractured, 402b  
 growth rate, 29b  
 magnesium number, 481b-483b  
 in metagabbro, 173b  
 mineral chemistry, 14b, 15b  
 mineralogy, olivine vs. oxide gabbros, 90b  
 minor elements, 56b  
 modal abundance, 82b, 116a  
 modality, 31b  
 neoblasts, 401b, 403b  
 nickel vs. forsterite content, 17b  
 plastic deformation, 104a  
 plutonic rocks, 6b-7b  
 pseudomorphed, 550b  
     with reaction corona, 137a  
 replacement with Iddingsite, 163b  
 replacement by talc and magnetite, 138a  
 texture, 117a  
 Unit V/Unit VI boundary, 4b
- Atlantis II Fracture Zone  
 phenocrysts, 49a  
 relict minerals, 85a
- olivine, cumulus, Atlantis Bank, in olivine iron-titanium oxide gabbro, 72b
- Olivine, magnesian, Atlantis Bank, formation of low-calcium pyroxene from, 33b
- Oman ophiolite  
 deformed gabbro, textures, 465b  
 gabbro  
     crosscutting relationship, 499b  
     undeformed igneous texture, 461b
- Ooze, calcareous, Atlantis II Fracture Zone, as mineralogical source, 149b
- Ooze, siliceous, Atlantis II Fracture Zone: Site 732, 42a, 48a, 49a
- Opaque minerals  
 Atlantis Bank, 119a  
     petrology, 114b-117b  
     texture, 117a
- Ophiolite  
 Atlantis Bank, magma mixing and fractionation of, 6b  
*See also* Oman ophiolite; Samail ophiolite; Sarmiento ophiolite; Troodos ophiolite
- Orthopyroxene  
 Atlantis Bank, 99a  
     augen, 107a  
     chemical composition in gabbro, 60b-61b  
     clinopyroxene intergrowths, 44b, 122a  
     deformation features, 131a  
     exsolution lamellae, 4b  
     metamorphic, 160b  
     minor elements, 53b, 67b  
     modality, 31b  
     oikocrysts, 121a  
     plutonic rocks, 12b  
     replacement with iddingsite, 163b  
     texture, 117a
- Atlantis II Fracture Zone  
 late fractures, 69a-70a  
 relict minerals, 85a  
 replacement pattern, 66a, 71a
- Atlantis II Fracture Zone: Site 733, recrystallization and mechanical twins, 69a
- Owen Fracture Zone, lithology, 91a
- Oxidation, late-stage, Atlantis Bank, 546b
- Oxide, iron-titanium  
 Atlantis Bank  
     distribution, 123a-124a  
     effect on Mg#, 82b-83b  
     mobilization, 135a  
     sulfide minerals and, 125a

- Atlantis II Fracture Zone, 464b  
crystallization, effect on liquid line of descent, 101b
- Oxide minerals  
Atlantis Bank, 94b  
ductile deformation and formation of concentrated, 104b  
magnetic susceptibility, 78b  
principal component analysis, 273b, 274b  
temperature vs. plagioclase composition, 91b
- Oxides  
Atlantis Bank  
activation of strain-softening mechanisms, 407b  
distribution, 122a–125a  
magmatic origin, 155b  
noncoticetic proportions, 31b  
relation to metamorphism, 134a–136a, 547b
- Oxygen fugacity  
Atlantis Bank  
crustal buffering, 101b–102b  
crystallization controls, 91b  
troilite and variations in, 119b, 121b–122b
- Oxygen isotopes  
Atlantis Bank, 135b–136b, 158b  
alteration index vs. whole-rock values, 173b  
clinopyroxene vs. plagioclase, 139b, 140b  
closed vs. open system conditions, 133b  
correlation, central Indian Ridge, 133b  
depletion zone distribution, 171b  
disequilibrium, pyroxene vs. plagioclase pairs, 141b  
felsic veins, 546b  
gabbro, 129b–131b  
mineral separates, 206b–207b, 212b  
modal composition and variation in, 170b  
open-system characteristics, 135b, 140b  
seawater interaction effects, 170b, 174b, 206b  
vein density and depletion of, 173b–174b  
vs. normal MORB values, 133b  
water/rock ratio and composition of, 135b–136b  
whole-rock powders, 170b–171b, 206b
- Parental Feldspar (FS) Array, Atlantis Bank, abyssal tholeiites, 84b–85b
- Pargasite  
Atlantis Bank, 163b  
temperature of formation, 172b
- pelagicus*, *Coccolithus*, Atlantis II Fracture Zone: Site 734, glacial/interglacial cycles, 84a
- Penetration distance, Atlantis Bank, olivine vs. iron-titanium oxide gabbro, 59b, 70b
- Pentlandite  
Atlantis Bank  
in green amphibole, 117b  
in igneous sulfides, 115b  
igneous vs. secondary, 117b  
in secondary sulfides, 115b–116b
- Peridotite  
Atlantis Bank, physical properties, 157a  
Atlantis II Fracture Zone  
Bouvet Fracture Zone compared, 361b  
classification, 10a  
melting point, 361b, 363b  
petrography, 381b  
Atlantis II Fracture Zone: Site 734, 78a  
East Pacific Rise, 42a  
Southwest Indian Ridge, 42a  
vs. East Pacific Rise, 3a
- Peridotite, foliated serpentinized, Atlantis II Fracture Zone, 84a
- Peridotite, mantle, Atlantis II Fracture Zone, mineralogy, 361b
- Peridotite, serpentinized  
Atlantis II Fracture Zone, 4a  
geochemistry, 87a  
petrography, 85a  
plastic deformation, 85a
- Peridotite/mylonite, serpentinized, Atlantis II Fracture Zone: Site 732, 50a
- Permeability  
Atlantis Bank, 90a, 210a  
comparison, Sites 504 and 395, 347b  
constant-rate injection tests, 202a  
drill-string packer measurements, 201a, 202a–203a, 333b–336b, 337b–339b  
effect on physical and chemical properties, 200a  
foliation fanning and, 502b  
fracture density and, 266b–267b, 339b, 554b  
in high-conductivity layers, 352b  
horizontal advection, 351b–352b  
*in situ*, 201a, 336b  
interval, 339b, 345b  
lithostratigraphic correlation, 205a, 266b, 346b  
logging indicators, 267b–268b  
model-fitting procedures, 336b  
operations, 203a–205a  
porosity and, 351b  
pressure-time records, 204a–205a  
pressurized fluid properties and, 336b–337b  
slug tests, 202a, 341b–342b  
TAM packer, 202a–203a  
and temperature anomalies, 349b  
vertical advection, 351b–352b, 556b
- Atlantis II Fracture Zone,  
deformation-enhanced, 443b–444b
- Philippines, serpentinized ultramafic rock, methane-rich gas seeps, 209b
- Phlogopite, Atlantis Bank, in metamorphic coronas, 168b
- Phosphate  
Atlantis Bank  
in basalt glasses, 88b  
during silicate mineral fractionation, 99b–101b  
expected vs. measured, 80b  
melt porosity and, 81b
- Phosphate/magnesium oxide ratio, Atlantis Bank, apatite saturation relationship, 99b
- Phosphate/zirconium ratio, Atlantis Bank, trapped melt composition and, 79b
- Phosphorus, Atlantis II Fracture Zone, bulk-rock vs. diabase concentrations, 484b–485b
- Photoelectric factor (PEF), Atlantis Bank, 271b–272b
- Physical properties  
Atlantis Bank, 158a–162a, 227b–237b  
crystalline rocks, 158a–159a, 160a–161a  
logging, 169a, 177a  
Unit I vs. Unit IV, 162a  
Atlantis II Fracture Zone: Site 733  
gabbros, 73a–75a  
vs. magnetic properties, 76a
- Pigeonite  
Atlantis Bank, 4b  
in iron-titanium oxide gabbro, 113a  
texture, 117a
- Plagioclase  
Atlantis Bank, 107a  
anisotropy and orientation of, 249b, 251b  
anorthite, 12b, 18b, 42b, 49b, 62b, 64b, 68b, 69b  
boudinaged porphyroblast, 402b  
chemical composition, 25b, 54b–56b, 59b, 69b, 100a, 102a  
crystallographic orientation, 403b, 404b, 405b, 407b
- deformation features, 131a  
diffusion rate, 59b, 70b  
dislocation slip, 405b, 407b  
in disseminated oxide olivine gabbro, 31b  
dynamic recrystallization, 400b, 401b, 403b, 407b–408b, 423b, 426b  
in felsic rock, 117a  
fluid inclusions, 190b, 200b–201b, 202b, 204b  
in foliated gabbro, 402b  
formed from iron-rich liquids, 97b  
growth rate, 29b  
igneous, 180b  
in late magmatic intrusions (LMIs), 164b  
lead isotopes, 132b  
magmatic vs. recrystallized, 399b  
mechanical twins, 400b, 403b  
in metagabbro, 164b, 189b  
metamorphic vs. hydrothermal, 208b  
mineral chemistry, 14b, 15b, 199b  
mineral zoning, 5b  
mineralogy, olivine vs. oxide gabbros, 90b  
minor elements, 53b, 56b  
in mixed gabbro-microgabbro interval, 127a  
modal abundance, 116a  
neodymium isotopes, 132b  
in olivine gabbro, 30b  
orientation in deformed metagabbros, 239b  
orientation of porphyroclasts, 103a–104a  
oxygen isotopes, 174b, 206b–207b, 212b  
plastic deformation, 418b, 419b  
plutonic rocks, 8b–9b  
recrystallized grain size with deformation changes, 421b–422b  
replacement, 138a, 163b  
reverse zonation, 27b  
theological effects of deformation, 399b  
temperature of alteration, 209b, 210b, 211b  
temperature of formation, synkinematic veins, 172b  
texture, 117a  
troctolitic olivine/iron-titanium oxide gabbro contact, 68b  
zonation, olivine gabbros, 25b
- Atlantis II Fracture Zone  
augen, 449b, 456b  
bimodal distribution, 477b–478b  
in olivine gabbro, 465b  
zonation, oxide-bearing gabbros, 486b–487b
- Atlantis II Fracture Zone: Site 732  
phenocrysts, 49a  
recrystallized, 52a  
xenocrysts, 49a
- Atlantis II Fracture Zone: Site 733  
porphyroclasts, 70a  
recrystallization and mechanical twins, 69a  
replacement pattern, 65a
- Atlantis II Fracture Zone: Site 734, replacement pattern, 78a
- central Indian Ocean, oxygen isotopes, 137b
- Plagioclase, calcic, Atlantis Bank, magmatic liquid surrounding xenoeyysts, 179b
- Plagioclase, sodic  
Atlantis Bank, 136a  
diopside association, leucocratic veins, 168b  
in veins and breccias, 172b
- Plagioclase-clinopyroxene pairs, Atlantis Bank, oxygen isotopes, 133b, 135b–136b
- Plate tectonics  
Atlantis II Fracture Zone, 5a  
magnetic anomalies, 44a, 45a  
transverse ridges, 77b
- Porosity  
Atlantis Bank, 230b–231b, 324b–326b

## SUBJECT INDEX

- alteration effects, 150a, 261b-262b, 263b, 266b-267b  
 electrical resistivity and, 555b  
 formation factor and, 328b-329b, 331b  
 hydrogen content and, 261b-262b  
*in-situ* measurements, 323b  
 permeability and, 351b  
 unsaturated, 554b  
 vs. electrical resistivity, 328b, 331b  
 vs. velocity, 159a, 161a-162a  
**Atlantis II Fracture Zone**, 73a-74a  
 dredged vs. drilled gabbros, 381b  
*See also* Crack porosity; Melt porosity  
**Porphyroclasts**, Atlantis Bank, microcracks, 159b  
**Porphyroclasts**, augen-type, Atlantis Bank, deformation features, 130a  
**Positive displacement coring motor (PDCM)**  
**Atlantis II Fracture Zone**  
 mechanical evaluation, 26a-28a, 82a  
 operating characteristics, 25a-26a  
 recommended design modifications, 28a  
 unsupported spudding, 25a-28a  
**Atlantis II Fracture Zone: Site 734**, 80a  
**Post-kinematic minerals**  
**Atlantis Bank**  
 alteration near late intrusive, 163b, 165b  
 magmatic/hydrothermal breccias, 165b-166b  
 pseudomorphs and coronas, 163b  
 veins, 166b  
**Potassium**  
 Atlantis Bank, geochemical log, 178a  
 Atlantis II Fracture Zone: Site 732, basalt, 54a  
**Potassium oxide**  
**Atlantis Bank**  
 in amphiboles, 56b  
 in basalt glasses, 88b  
 geochemical logging data, 280b  
 in plagioclase, 56b  
**Potassium oxide/phosphate ratio**, Atlantis Bank, east-west fracture zone discontinuity, 88b  
**Prehnite**  
 Atlantis Bank, iron content, 211b  
 Atlantis II Fracture Zone: Site 732, 53a  
**Pseudoemiliana lacunosa**  
 Atlantis II Fracture Zone: Site 732, 49a  
 Atlantis II Fracture Zone: Site 734, 94a  
**Pyrite**  
**Atlantis Bank**  
 in green amphibole, 117b  
 in secondary sulfides, 116b  
 sulfur isotopes, 119b, 122b  
**Pyrite**, secondary, Atlantis Bank, sulfur isotopes, igneous vs. basaltic sulfides, 123b  
**Pyrite-marcasite pseudomorphs**, Atlantis Bank, iron oxyhydroxide association, 117b  
**Pyroxene**  
**Atlantis Bank**  
 from hydrothermal breccias, 169b, 170b  
 from late magmatic intrusives, 161b-162b  
 growth rate, 29b  
 iron enrichment, 91b  
 late-stage oxidation, 546b  
 mineralogy, high- vs. low-calcium, 90b  
 mineralogy, late magmatic intrusions (LMIs) vs. iron-rich neoblast mineralogy, 168b-169b  
 morphology, oxide-free vs. oxide-bearing gabbros, 531b  
 porphyroclasts and neoblasts, deformed gabbros, 161b-162b  
 in silicate minerals, 92b-93b  
**Atlantis II Fracture Zone**  
 morphology, 461b  
 in olivine gabbro, 465b  
 in troctolitic rocks, 447b  
**Pyroxene augen**, Atlantis II Fracture Zone, plagioclase and, 461b  
**Pyroxene**, brown, Atlantis Bank, 102a  
**Pyroxene**, calcium-poor  
 Atlantis Bank, 115a  
 texture, 117a  
**Pyroxene**, green, Atlantis Bank, 102a  
**Pyroxene-hornblende-plagioclase assemblage**, Atlantis Bank, 160b  
**Pyrrhotite**  
 Atlantis Bank, 125a  
 in green amphibole, 117b  
 igneous vs. secondary, 115b-116b, 117b, 123b  
 in sulfides, 115b-116b  
**Quartz**  
**Atlantis Bank**  
 in felsic rock, 117a  
 fluid inclusions, 197b, 201b, 204b  
 oxygen isotopes and stabilization of, 138b-139b  
**Atlantis II Fracture Zone**, source provinces, 147b  
**Radiogenic isotopes**, Atlantis Bank, high- vs. low-temperature alteration effects, 138b  
**Rare earth elements**, Atlantis II Fracture Zone, 148b, 150b  
**Recrystallization**  
**Atlantis Bank**  
 oxide minerals, 91b-93b  
 oxygen fugacity, 91b  
 temperature, 91b  
**Recrystallization**, subsolidus  
**Atlantis Bank**  
 pyrite-pyrrhotite-chalcopyrite assemblages, 94b-95b  
 temperatures, 94b-95b  
**Recrystallization**, syntectonic, Atlantis Bank, to relieve elastic strain, 506b  
**Resistivity**, Atlantic Bank, 90a, 169a, 176a, 177a, 210a  
**Rhum Layered Intrusion**, troctolites, picritic intrusion, 487b  
**Ridge dynamics**, models, 3a  
**Ridge-transform intersection**  
**Atlantis Bank**, tectonic evolution of gabbros, 428b  
**Atlantis II Fracture Zone**  
 bathymetry, 367b  
 inside corner highs, 368b  
 lithology, 441b  
 mass-wasting features, 431b  
 nodal basins, 366b  
 outside corners, 368b  
 topographic asymmetry, 368b  
 transform volcano, 366b, 368b  
**Rift valley**  
**Atlantis II Fracture Zone**  
 flanking deeps, 364b, 366b  
 northern, 366b  
 southern, 364b-366b  
 wall heights, 364b, 366b  
**Southwest Indian Ridge**, bathymetry, present-day, 396b  
**Romanche transform zone**, gabbro, high temperature deformation, 51a  
**Rotary-Coring System (RCB)**  
 Atlantis Bank, 93a, 95a-96a  
 summarized, 6a  
**Rubidium**  
 Atlantis Bank, 135b  
 stratigraphic variation, 147a, 150a  
**Atlantis II Fracture Zone: Site 732**, basalt, 54a  
**St. Paul Fracture Zone**, tectonic setting, 91a  
**St. Paul's Rocks**, foliated peridotites, orientation, 396b, 441b  
**St. Peter Fracture Zone**, tectonic setting, 91a  
**Salinity**, Atlantis Bank, electrical conductivity and, 327b-328b, 329b, 330b  
**Samail ophiolite**  
 magma mixing and fractionation of, 6b  
 plagiogranite, oxygen isotopes, 139b  
 seismic reflection profiling, 246b  
**Sand**, Atlantis II Fracture Zone: Site 732, Pleistocene, 373b  
**Sand**, foraminiferal, Atlantis II Fracture Zone: Site 734, stratification, 433b  
**Sand**, igneous, Atlantis II Fracture Zone: Site 734, 435b  
**Sand**, unconsolidated, Atlantis II Fracture Zone: Site 732, 42a  
**Sand**, volcanicogenic, Atlantis II Fracture Zone: Site 732, 48a  
**Sandstone**, Atlantis II Fracture Zone: Site 732, 49a  
**Sandstone**, indurated, Atlantis II Fracture Zone: Site 732, 42a  
**Sandstone**, lithic  
 Atlantis II Fracture Zone: Site 732, 51a  
 petrography, 50a  
**Sarmiento ophiolite**  
 magnetite-ilmenite crystals, 139b  
 metagabbros, oxygen isotopes, 135b  
**Sea-surface temperature**, Atlantis Bank, climatic effects, 353b  
**Seamounts**, Atlantis II Fracture Zone, ridge-transform intersection, 365b  
**Seawater/rock interaction**  
**Atlantis Bank**  
 deformation-enhanced, 153b, 174b, 175b  
 in deformed zones, 544b  
 isotopic effects, 137b, 170b, 174b, 206b  
 in magnesio-hornblende formation, 543b-544b  
 oxide elevation, 546b  
 sulfur depletion, 123b, 124b  
**Sediment ponds**  
**Atlantis II Fracture Zone**  
 crust under, 392b  
 transform valley floor, 371b, 372b  
**Seismic anisotropy**  
**Atlantis Bank**  
 in deformed gabbros, 405b  
 parallel vs. perpendicular to foliation, 554b  
**Seismic stratigraphy**  
**Atlantis Bank**  
 air gun vs. water gun results, 224b  
 brittle-ductile deformation zones, 507b  
 coherence between seismograms, 223b-224b  
 correlation, physical properties, 163a  
 factors affecting reflectivity, 247b-249b  
 layer 3 gabbroic sequence, 245b-247b  
 lithologic correlation, 224b  
 lithostratigraphic correlation, 246b  
 reflection profiling, 91a-92a, 223b-225b, 554b  
 synthetic seismogram, 245b-247b, 248b  
**Atlantis II Fracture Zone**  
 across southern median tectonic ridge, 372b-373b  
 reflection profiling, 43a, 46a, 62a, 63a, 81a  
 transverse ridges, 219b  
*See also* Vertical seismic profiles (VSP)  
**Seismic velocity**, Atlantis Bank, 553b-554b  
**sellii**, *Helicosphaera*, Atlantis II Fracture Zone: Site 734, 94a

- Semblance anomalies, 266b  
 Sense of shear  
   Atlantis Bank, indicators, 419b-420b, 423b  
   Atlantis II Fracture Zone, amphibole vein linings, 499b  
 Serpentinite, rotated, Atlantis II Fracture Zone: Site 732, 55a  
 Serpentinite  
   Atlantis II Fracture Zone, classification, 10a-11a  
   Atlantis II Fracture Zone: Site 732, 48a  
     foliation, 54a  
 Serpentization, Atlantis Bank, of ultramafic rock, 209b  
 Shear, mylonitic, Atlantis Bank, in textural type IV gabbro, 421b  
 Shear polyhedra, oxide-free, Atlantis II Fracture Zone, in deformation zones, 488b, 501b  
 Shear strain  
   Atlantis Bank, 104a  
     textural types and, 103a  
   Unit III/Unit IV boundary, 107a  
 Shear zone  
   Atlantis Bank, 159b  
     foliation, 425b  
   Atlantis II Fracture Zone  
     formation in solidifying pluton, 487b  
     morphology of late-magmatic, 497b  
   *See also* Sense of shear  
 Shear zone, amphibolite facies, Atlantis Bank, dip orientation, 414b  
 Shear zone, brittle-ductile, Atlantis II Fracture Zone, permeability, 443b-444b  
 Shear zone, ductile, Atlantis Bank, dip orientation, 428b  
 Shear zone, granulite-facies, Atlantis Bank, dip orientation, 414b  
 Shear zone, mylonitic  
   Atlantis Bank, 416b  
     foliation, 418b  
     foliation, dip orientation, 423b, 425b  
     foliation and lineation orientation, 411b  
     as hydrous fluid pathway, 427b, 428b  
   Lower Structural Domain, 426b  
   Middle Structural Domain, 425b  
   Upper Structural Domain, 423b  
 Shear zone, normal, Atlantis Bank, foliation and lineation orientation, 413b  
 Shear-wave splitting, Atlantis Bank, anisotropy and, 236b, 241b  
 Silica  
   Atlantis Bank  
     geochemical log, 174a, 178a  
     geochemical logging data, 276b, 278b  
     in iron-rich liquids, 97b  
     velocity and percentage of, 247b, 249b  
 Silicates  
   Atlantis Bank, 108b-111b  
     analogy to lava lakes, 30b  
     grain size variations, 25b  
     mineral composition, 24b  
     nucleation rate and grain size variation, 30b  
     post-cumulus homogenization, 29b  
     pyroxenes, 92b-93b  
     in recrystallized gabbro, 406b-407b  
     trace elements, 5b-6b, 12b  
 Atlantis II Fracture Zone  
   intragranular diffusion rates, 486b  
   mineral composition, 473b  
   modal variation, 465b  
 Silicon, Atlantis II Fracture Zone, in ultramafic rocks, 13a  
 Skaergaard Intrusion  
   clinopyroxene, hydrothermal, 168b  
   cooling history, vein mineralization and, 175b
- crystallization intervals, 61b, 90b  
 diopside, temperature of formation, 168b  
 diopside-bearing veins, 504b  
 gabbro  
   geochemistry, 470b-471b  
   mineral composition and phase assemblages, 33b  
   plagioclase An vs. Mg# of mafic minerals, 49b  
   strontium concentration, 85b  
 hydrothermal veins, 208b  
   vertically oriented crack network, 505b  
 iron-titanium-rich gabbro pegmatites, magmatic origin, 155b  
 liquid evolution, 82b  
   vs. Galapagos Spreading Center, 100b-101b  
 melt porosity, 81b  
 Layer Zone and Marginal Border Series vs.  
   Upper Border Series, 81b  
 mineralogy  
   comparison, Atlantis Bank, 95b  
   cumulus phase changes, 49b  
   olivine, 48b  
     crystallization, temporary cessation of, 49b  
   pyroxene-oxide veins, magmatic origin, 155b  
 recrystallization, oxygen fugacity, 91b, 102b  
 trondhjemite, Tinden granophyre, 79b  
 Upper Zone melanogranophyres, 102b  
 Skaergaard Layered Series, crystallization, pyroxene composition, 91b  
 Skinner Cove Sequence, ophiolites, fracture zone rocks, 431b  
 Slip component, Atlantis II Fracture Zone, strike-slip vs. dip-slip, 501b  
 Slow-spreading ridges  
   characteristic features, 361b  
   lithology, transform edge effect, 360b  
   ophiolite formation, 509b  
   plutonic crust, heterogeneity with fault zones, 473b  
 Slump scars, Atlantis II Fracture Zone, 48a, 371b  
 Smectite  
   Atlantis Bank, lining fractures, 546b  
   Atlantis II Fracture Zone, source provinces, 147b  
 Sodium metasomatism, Atlantis Bank, of vein structures, 136a  
 Sodium oxide  
   Atlantis Bank, 135b  
     in amphiboles, 56b  
     basalt glasses, 86b-87b  
     in diabase, 478b  
     in gabbros, 478b  
     vs. Mg# of clinopyroxene, 51b, 65b  
 Southwest Indian Ridge, 361b  
 basalt, geochemistry, transform vs. ridge axis, 472b  
 gabbro, trapped intercumulus melt, 472b  
 lithology, American-Antarctic Ridge transforms compared, 377b  
 rift valley, 512b  
 spreading rate, 361b  
 tectonic setting, 362b  
 uplift, 441b  
 Sphalerite, Atlantis Bank, in oxide-rich ferrogabbros, 125a  
 Sphe ne  
   Atlantis Bank  
     in felsic veins, 549b  
     in hydrothermal veins, 180b  
 Sphe ne, vermicular pink, Atlantis Bank, ilmenite association, 133a  
 Spinel  
   Atlantis Bank, distribution, 122a-123a  
   Atlantis II Fracture Zone, source, 447b
- See also* Chromium-spinel  
 Spreading rate  
   Atlantis II Fracture Zone  
     across transform, 390b  
     asymmetry, 389b  
     magnetic anomaly correlation, 382b, 387b  
 Stable isotopes, Atlantis Bank, temperature of formation and, 211b  
 Stable remanent magnetic vector (SRMV)  
   Atlantis Bank, 409b-411b  
     declination, Lower Structural Domain, 413b-414b  
     declination, Middle Structural Domain, 413b  
     declination, Upper Structural Domain, 411b, 413b  
 Stillwater complex  
   apatite, chlorine-rich, 167b  
   magmatic biotite, phlogopite association, 167b  
   mineralogy, incompatible element-rich mineral association, 166b-167b  
   pargasite, 170b  
 Strain, elastic, Atlantis Bank, available for cracking, 506b  
 Stress  
   Atlantis Bank, 556b  
     seawater circulation and, 505b  
 Strontium  
   Atlantis Bank  
     in calcite veins, 200b  
     in parental magma, 83b, 107b  
     stratigraphic variation, 147a  
   Atlantis II Fracture Zone: Site 734, 87a  
 Strontium isotopes  
   Atlantis Bank, 139b  
     amphiboles, 136b  
     gabbro, 129b-133b  
     magmatic pile buffer, 137b-138b  
     seawater interaction effects, 137b  
     systematic variation, 132b  
     vs. neodymium isotopes, 136b, 140b  
   Atlantis II Fracture Zone, vs. neodymium isotopes, 136b  
 Strontium/aluminum ratio, Atlantis Bank, parental magma composition and, 84b, 85b  
 Sulfate, Atlantis Bank, incorporation of seawater, 122b-123b  
 Sulfate, soluble, Atlantis Bank, sulfur isotopes, 119b  
 Sulfide  
   Atlantis Bank  
     comparison, Skaergaard Intrusion, 101b  
     distribution, 122a-125a  
     in metamorphosed gabbros, 125a-126a, 134a-136a  
     mineralogy, 95b  
     oxide mineral association, 125a  
     recrystallization changes, 93b-95b  
     relation to silicate phases, 126a-128a  
     saturation, 101b  
     segregation, during recrystallization, 93b-94b  
     segregation, mineralogical changes, 120b  
     in undeformed gabbros, 125a  
 Sulfide, globular, Atlantis Bank, in oxide concentrates, 93b, 104b  
 Sulfide, igneous  
   Atlantis Bank  
     altered, 118b  
     distribution, 119b-120b  
     globular interstitial, 119b  
     immiscible, 119b  
     opaque mineral petrology, 116b  
     petrology, 114b-115b  
     in silicates, 119b

## SUBJECT INDEX

Sulfide, primary, Atlantis Bank, in iron-titanium oxide gabbro, 115a–116a  
 Sulfide, secondary  
     Atlantis Bank  
         in altered olivine, 121b  
         hydrothermal alteration, 120b–121b  
         opaque mineral petrology, 115b–117b, 118b  
 Sulfur  
     Atlantis Bank  
         in basalt glasses, 88b–89b  
         carbon content, 117b  
         distribution, 119b–120b  
         dynamothermal metamorphism and depletion of, 121b  
         east-west fracture zone discontinuity, 88b  
         geochemical logging data, 279b  
         hydrothermal alteration, 120b–121b, 123b  
         low-temperature alteration, 121b  
         mantle source properties affecting, 89b  
         modeling in submarine hydrothermal systems, 122b  
         MORB-type source, 121b, 123b  
         in oxide gabbros, 120b, 123b  
         Rayleigh fractionation, 99b  
         saturation, 99b–100b  
         troilite and variations in fugacity, 119b, 121b–122b  
 Sulfur isotopes  
     Atlantis Bank, 118b–119b  
         correlation, MORB glasses, 121b  
         pyrite vs. monosulfide sulfur, 122b  
         sulfur and oxygen fugacity and variations in, 121b–122b, 123b  
 Sulfur, monosulfide, Atlantis Bank, sulfur isotopes, 122b  
 Synkinematic differentiation, Atlantis II Fracture Zone, 442b, 444b, 486b–487b, 511b  
 Synkinematic minerals  
     Atlantis Bank  
         host rock, 160b  
         textural type correlation, 171b  
 Syntectonic differentiation, Atlantis II Fracture Zone, 488b  
 Talc, Atlantis Bank, metamorphosed gabbros and late magmatic intrusions (LMIs), 735b  
 Talus  
     Atlantis II Fracture Zone, 75a  
         ridge-transform intersection, 431b  
 Tectonites, metagabbro, Atlantis Bank, velocity, 159a  
 Television/sonar system  
     Atlantis Bank, 94a  
     use with positive displacement coring motors (PDCM), 25a, 42a–44a, 80a, 81a, 82a, 92a  
 Temperature  
     Atlantis Bank, 90a  
         anomalies, 349b  
         corrected for drilling disturbance, 347b, 349b  
         of deformation, 131a–132a, 139a  
         equilibrium profile, 167a–168a, 352b–353b, 354b  
         ilmenite-magnetite pairs vs. plagioclase, 98b  
         logging data, 166a–168a, 169a, 171a, 266b, 347b, 555b  
         logging equipment, 165a–166a  
         negative gradient, 168a, 266b, 267b, 268b  
         positive anomalies, 167a  
 Texture  
     Atlantis Bank  
         in ductilely deformed gabbros, 422b–426b  
         gabbro, types I–IV, 418b, 421b  
         intrusive rocks vs. patchy olivine gabbros, 507b–508b, 537b

of plastically deformed gabbros, 400b, 403b  
     shear bands, 425b  
 Atlantis II Fracture Zone  
     deformation, 461b  
     hypersolidus vs. subsolidus gabbros, 498b  
     oxide-bearing gabbros, 461b  
 Thermal conductivity  
     Atlantis Bank, 167a  
         F factors, silica glass vs. Macor standard, 164a, 166a  
         horizontal seawater advection and, 351b  
         measuring techniques, 163a–164a  
 Atlantis II Fracture Zone, 19a–21a  
     F factors, silica glass vs. Macor standard, 21a, 23a  
     gabbro, 60a  
     silica glass vs. Macor standard, 21a  
         variation with probe orientation, 74a, 76a  
         calibration tests, 20a–21a  
 Thermal modeling  
     Atlantis Bank  
         with enhanced thermal interaction, 351b  
         linear vs. step change, 353b  
         one-dimensional conduction from transient boundary temperatures, 352b–354b  
         single-and double-layer models, 351b  
         two-dimensional conduction simulation of advection, 349b–352b  
         uniformly conducting models, 350b, 351b  
 Thermocline, Atlantis Bank, thickness, seasonal variation, 353b  
 Thingmuli volcano, mineralogy, during differentiation, 93b  
 Tholeiite, abyssal  
     Atlantis Bank  
         crystallization history, 85b  
         Parental Feldspar Array, 84b–85b  
         sulfur content, 88b  
         liquid line of descent, iron-titanium oxide crystallization effects, 101b  
 Thomsonite, Atlantis Bank, in hydrothermal veins, 180b  
 Thorium, Atlantis Bank, geochemical logging data, 178a, 280b  
 Titanite, Atlantis Bank, mineral chemistry, 200b  
 Titanium  
     Atlantis Bank, geochemical log, 175a  
     Atlantis II Fracture Zone, in troctolitic spinel, 447b  
 Titanium oxide  
     Atlantis Bank, 135b  
         basalt glasses, 86b  
         in clinopyroxene, 80b  
         in diabase, 479b–480b  
         in gabbros, 479b–480b  
         geochemical logging data, 276b, 278b  
         ilmenite-rich layers, 5b  
         in orthopyroxene, 53b, 67b  
         parental magmas, 91b  
         vs. ferrosilite, 17b  
         vs. Mg#, 51b, 99b  
         vs. Mg#, of amphiboles, 56b  
         vs. Mg#, of clinopyroxene, 65b  
 Atlantis II Fracture Zone  
     basalt vs. oxide-free gabbros, 473b  
     vs. magnesium, dredged vs. drilled samples, 473b  
     xenocrysts, 49a  
 Titanium-pargasite, Atlantis Bank, magmatic red-brown, 160b  
 Titanomagnetite  
     Atlantis Bank  
         crystallization, 101b–102b, 104b  
         magnesium oxide vs. aluminum oxide, 99b  
 Trace elements  
     Atlantis Bank  
         in iron-titanium oxide gabbro, 146a–147a  
         magma mixing and variation in, 6b  
         olivine gabbro, 146a  
         in silicate minerals, 5b–6b, 12b  
         stratigraphic variation, 147a, 150a  
         in troctolite, 146a  
 Transform tectonic zone  
     Atlantis II Fracture Zone  
         hooked ridges, 373b, 375b  
         nontransform walls, 373b, 376b  
         paleotransform walls, active vs. inactive, 376b  
         present-day transform slip zone, 373b  
         lithology, 380b  
 Transform valley, Atlantis II Fracture Zone, physiography, 371b  
 Transform walls  
     Atlantis II Fracture Zone  
         cirque-shaped scars, 371b  
         physiography, eastern vs. western slope, 371b, 390b, 392b  
 Transmissivity  
     Atlantis Bank  
         constant-rate injection tests, 343b, 344b, 345b  
         interval, 345b  
         slug tests, 341b–342b, 343b, 345b  
 Transtensional spreading, Atlantis II Fracture Zone, 360b  
 Transtensional transform basin  
     Atlantis II Fracture Zone  
         crustal thinning during formation of, 392b  
         formation model, 389b–391b  
         spreading direction shift, 389b  
 Transverse ridges  
     Atlantis II Fracture Zone  
         bathymetry, 370b, 372b  
         crustal thickness under, 368b  
         elevation above normal seafloor, 368b  
         evolution, 392b–395b  
         excessive subsidence, 395b  
         saddle morphology, 368b, 371b  
         uplift, 511b  
         seismic stratigraphy, 219b, 225b  
 Tremolite, Atlantis Bank, 138a  
 Troctolite  
     Atlantis Bank, 90a, 118a, 129b–130b, 209a, 427b  
         crystallization model, 35b, 37b  
         equigranular, 112a  
         geochemistry, 146a  
         ilmenite-magnetite ratio, 124a  
         magma composition, 13b  
         magnetic properties, 289b–290b, 313b  
         mineralogy, 24b–25b, 41b  
         olivine gabbro and, 27b  
         olivine-liquid relationship, 82b  
         oxide gabbro and, 27b, 29b  
         permeability vs. log porosity, 267b  
         petrography, 111a–112a  
         poikilitic, 111a–112a  
         porosity and zirconium content, 80b  
         texture, 24b–25b, 38b, 41b  
 Atlantis II Fracture Zone  
     geochemistry, 478b  
     magmatic sedimentary layering, 470b  
     olivine gabbro contact, 470b  
     petrography, 447b  
     Unit VI, 468b  
 Troctolite, olivine-rich, Atlantis Bank, 102a  
 Troilite, Atlantis Bank, in igneous sulfides, 115b, 119b  
 Troilite-pyrrhotite exsolution, Atlantis Bank, oxygen fugacity, 119b

Trondhjemite  
 Atlantis Bank  
   alteration stratigraphy, 503b  
   clinopyroxene and MnO content of, 91b  
   crystallization model, 35b, 37b  
   fluid inclusions, 197b, 204b  
   formation, 511b  
   geochemical logging, 178a  
   geochemistry, 147a  
   granophytic texture, 190b  
   intrusion, 155b, 208b  
   iron-titanium gabbro contact, 119a  
   methane signal, 206b  
   oxygen isotopes, 133b, 138b-139b, 206b  
   penetration of deformed rock, 90b  
   petrography, 215b  
   temperature of alteration, 209b  
 Atlantis II Fracture Zone  
   formation, 461b  
   oxide-bearing vs. oxide-free gabbros, 461b  
   petrography, 447b-448b  
 Troodos ophiolite  
   formation, 509b  
   magnetic properties, 73a  
   Koenigsberger Ratio, 153a  
 Turbidity currents, Atlantis II Fracture Zone: Site 734, gravel deposition, 436b

Ultramafic rock  
 Atlantis Bank, serpentinization and reduction of seawater carbonate, 209b  
 Atlantis II Fracture Zone  
   classification, 12a  
   as mineralogical source, 149b  
 Atlantis II Fracture Zone: Site 732  
   alteration and deformation, 53a-54a  
   petrography, 50a

Uplift  
 Atlantis Bank  
   blocks, 91a, 207a-208a  
   transform wall, 264b  
 Atlantis II Fracture Zone: Site 732, median ridge, 56a  
 Uranium, Atlantis Bank, geochemical logging data, 178a, 280b

Vanadium, Atlantis Bank, stratigraphic variation, 147a

Veinlet, andesine, Atlantis Bank, fluid inclusions, 189b

Veinlet, plagioclase, Atlantis Bank, 188b

Veins  
 Atlantis Bank  
   alteration index and density of, 171b, 173b-174b  
   dip, 209a  
   dip, vs. foliation strike, 555b  
   distribution, 506b  
   miarolitic mineralization, 136a  
   post-kinematic minerals, 166b  
   structure relation to late magmatic intrusions (LMIs), 172b-173b  
   synkinematic minerals, 160b, 163b, 171b  
 Atlantis II Fracture Zone  
   density, 494b-496b  
   dip, 492b-493b, 499b, 511b

Veins, amphibole  
 Atlantis Bank  
   crosscutting relationship, 187b, 504b  
   in deformed metagabbro, 136a  
   density, 186b  
   mineral chemistry, 208b, 543b  
   monomineralic, 545b

orientation, deformed vs. undeformed zones, 505b-506b  
 subsolidus ductile deformation and, 504b  
 Atlantis II Fracture Zone, 499b  
   oblique slip, 497b  
   slip direction, bimodal distribution, 501b  
 Veins, analcime, Atlantis Bank, actinolite inclusions, 166b  
 Veins, calcium silicate, Atlantis II Fracture Zone: Site 734, static alteration and, 86a  
 Veins, carbonate  
 Atlantis Bank, 186b  
   breccia zone relationship, 173b, 175b  
   crosscutting hornblende-Plagioclase veins, 552b  
 Veins, conjugate, Atlantis Bank, 106a  
 Veins, diopside  
 Atlantis Bank, 166b, 180b, 504b  
   fluid inclusions, 197b  
   temperature of formation, 545b  
 Atlantis II Fracture Zone, in olivine gabbro, 466b  
 Veins, diopside-bearing  
 Atlantis Bank  
   dip orientation, 502b  
   hydrothermal origin, 546b  
 Veins, diopside-hornblende-plagioclase, Atlantis Bank, origin, 175b  
 Veins, diopside-Plagioclase  
 Atlantis Bank, 504b  
   oxygen isotopes, 171b  
   temperature of formation, 506b  
 Veins, felsic  
 Atlantis Bank  
   alteration stratigraphy, 504b-505b  
   dip orientation, 502b  
   origin, 119a, 546b  
   paragenesis, 503b  
   trondhjemetic-appearing, 504b  
   zoned mineral assemblages, 545b  
 Atlantis II Fracture Zone, 448b  
   in olivine gabbro, 466b  
 Veins, green amphibole, Atlantis Bank, 105a  
 Veins, hornblende  
 Atlantis Bank, 136a, 549b  
   dip orientation, 414b, 427b  
   ductile deformation and density of, 171b  
   mineralogy, 177b  
 Veins, hornblende-andesine, Atlantis Bank, pressure of alteration, 212b  
 Veins, hornblende-plagioclase, Atlantis Bank, mineralogy, 114b  
 Veins, hydrothermal  
 Atlantis Bank, 105a, 502b  
   formation of, 208b, 212b  
   inclination, 545b  
   non-synkinematic, 427b  
 Veins, hydrothermal-plagiogranite, Atlantis Bank, 138a  
 Veins, late magmatic, Atlantis Bank, inclination, 545b  
 Veins, leucocratic  
 Atlantis Bank  
   diopside-plagioclase association, 168b  
   mineralogy, 155b  
   replacement minerals, 172b-173b  
   seawater hydration, 209b  
   zircon in, 155b, 163b, 165b, 172b  
 Veins, leucocratic diopside-bearing, 206b  
 Atlantis Bank, 186b, 198b  
   fluid inclusions, 204b-205b  
   petrography, 188b-189b, 215b  
   temperature of mineral formation, 211b, 212b

Veins, leucocratic diopside-plagioclase, Atlantis Bank, isotopic composition, 207b  
 Veins, monomineralic, Atlantis Bank, pyroxene composition, 169b  
 Veins, oxide gabbro, Atlantis II Fracture Zone, cutting oxide-free olivine gabbro, 462b, 464b  
 Veins, plagioclase, Atlantis Bank, 504b  
 Veins, plagioclase-amphibole  
 Atlantis Bank, 551b  
   crosscutting relationship, 187b  
   density, 186b  
 Veins, quartz, Atlantis II Fracture Zone: Site 732, 53a  
 Veins, smectite-carbonate, Atlantis Bank, 505b, 508b  
 Veins, synkinematic  
 Atlantis Bank  
   plastic deformation, 172b  
   temperature of formation, 172b  
 Veins, thomsonite, Atlantis Bank, 179b  
 Veins, trondhjemitic  
 Atlantis Bank, 27b, 35b, 106a  
   dip orientation, 502b  
   formation, 488b  
   isotopic composition of plagioclase, 207b  
   origin, 503b, 545b  
   in oxide-bearing gabbros, 503b  
   temperature of formation, 545b  
 Atlantis II Fracture Zone  
   crosscutting mylonite, 465b  
   petrography, 447b-448b  
 Veins, zeolite, Atlantis Bank, crosscutting felsic veins, 552b

Velocity  
 Atlantis Bank, 164a, 227b-231b  
   alteration effects, 233b  
   chemical composition effect, 232b  
   crack porosity and, 233b, 234b, 235b, 247b  
   crystalline rock, 157a  
   density and, 231b-232b, 238b, 239b  
   elastic constants and, 236b, 242b-243b  
   foliated gabbros and metagabbros, 157a-159a  
   gabbro, 209a  
   isotropic gabbro, 244b  
   logging data, 246b  
   metagabbro, 157a  
   method, 227b, 229b-230b, 231b  
   oceanic crust variations, 227b  
   seismic stratigraphy correlated, 163a  
   semblance calculation, 266b  
   vertical vs. horizontal layers, 162a-163a  
   vs. attenuation, 258b, 259b  
   vs. porosity, 159a, 161a-162a  
   vs. sonic and vertical seismic profile, 159a-161a  
 Atlantis II Fracture Zone, 19a, 73a-74a  
   metagabbro, 60a

Velocity, interval  
 Atlantis Bank, 222b-223b, 225b  
   deformation zones, 223b

Vema Fracture Zone  
 amphibole gneiss, 543b  
 gabbro, high temperature deformation, 51a  
 geometry, 366b  
 greenschist-facies alteration, 489b  
 neovolcanic zones, 376b  
 normal faulting, ridge-transform intersection, 428b  
 transverse ridge, excessive subsidence, 395b

Vertical seismic profiles (VSP)  
 Atlantis Bank, 553b  
   correlation, hand sample velocities, 189a  
   data processing, 221b-222b  
   deconvolved traces, 221b-222b, 223b

## SITE INDEX

geologic setting, 183a, 185a  
 gun mooring, 190a  
 gun signatures, 191a  
 logging cable attenuation, 186a  
 methods, 219b-221b  
 navigation and weather, 187a  
 objectives, 182a  
 operations, 187a-188a  
 origin, 225b  
 previous oceanic seismology, 182a-183a  
 processing, 186a-187a, 188a  
 receivers, 185a, 192a  
 recording systems, 185a, 188a  
 reflector, 190a  
 seismic sources, 186a, 187a, 192a  
 sources of error, 189a  
 stacked seismograms, 190a, 194a, 195a, 196a, 199a  
 testing and calibration, 185a  
 three-component seismometer, 185a, 197a  
 timing signal flow, 189a  
 traveltimes distances, 189a, 197a  
 vertical-component seismometer, 185a, 197a  
 water-gun record, 222b  
 Vine-Matthews-Morley type initial magnetization, Atlantis Bank, oceanic gabbros, 301b, 304b, 309b  
 Viscous flow  
 Atlantis Bank  
 critical melt fraction, 403b  
 magmatic foliation produced by, 427b

Site 334  
 gabbro, strontium/aluminum values, 84b  
 greenschist-facies alteration, 489b

Site 395  
 basalt, porosity-hydrogen relationship, 262b  
 basement, permeability, 347b  
 drill-string packer experiments, 335b  
 oceanic crust, depth of drilling, 323b  
 permeability, 339b

Site 417  
 oceanic crest, depth of drilling, 323b  
 physical erosion, 435b

Site 418  
 basalt, porosity-hydrogen relationship, 262b  
 electrical conductivity, 328b  
 oceanic crust, depth of drilling, 323b

Site 504  
 basalt  
 layer 2, 127b  
 porosity-hydrogen relationship, 262b  
 sulfur isotopes, 122b  
 basement, permeability, 347b  
 depth of drilling, 323b  
 drill-string packer experiments, 335b  
 electrical conductivity, 328b  
 gabbro mineralogy, INCA study, 276b  
 magnetic properties  
 logging data, 309b  
 lower oceanic crust, 285b  
 permeability, 339b  
 titanium oxide, 272b

in solidifying gabbroic magma, 400b, 401b, 403b  
 Voight-Reuss-Hill average, Atlantis Bank, iron-titanium oxide gabbros, 233b-234b  
 Volcanism, constructional, Atlantis II Fracture Zone, 366b, 376b  
 Volcano, transform, Atlantis II Fracture Zone, 366b, 368b  
 Water content, Atlantis Bank, 230b-231b  
 Water/rock ratio  
 Atlantis Bank  
 chlorine contents and, 208b  
 control on vein assemblages, 504b, 506b  
 ductile deformation, 423b, 544b  
 isotopic composition and, 135b-136b, 170b, 174b, 175b, 206b  
 plagioclase-water fractionation and, 174b  
 pyroxene replacement, 544b  
 Weathering, biogenic, Atlantis II Fracture Zone: Site 734, 435b-436b  
 Wharton Basin, geochemistry, 148b  
 Woods Hole Oceanographic Institution vertical seismic profile tool, Atlantis II Fracture Zone, 96a

Yttrium  
 Atlantis Bank, stratigraphic variation, 150a  
 Atlantis II Fracture Zone, bulk-rock vs. diabase concentrations, 484b-485b

Yttrium/niobium ratio, Atlantis II Fracture Zone: Site 732, basalt, 54a, 56a  
 Yttrium/zirconium ratio  
 Atlantis Bank, 5b  
 trapped melt, 83b  
 Atlantis II Fracture Zone, veins, 486b  
 Zeolite, Atlantis Bank, 168b  
 Zeolite grade alteration, Atlantis Bank, 173b  
 Zeolite-facies alteration, Atlantis Bank, in hydrothermal breccias, 505b  
 Zinc, Atlantis Bank, stratigraphic variation, 147a  
 Zircon  
 Atlantis Bank  
 in leucocratic veins, 155b, 163b, 165b, 172b  
 stratigraphic distribution, 167b  
 Zirconium  
 Atlantis Bank, 135b  
 in clinopyroxene, 80b  
 stratigraphic variation, 150a  
 Atlantis II Fracture Zone  
 bulk-rock vs. diabase concentrations, 484b-485b  
 in trondhjemites, 486b  
 Zirconium/triobium ratio  
 Atlantis Bank, trapped melt, 83b  
 Atlantis II Fracture Zone: Site 732, basalt, 54a, 56a  
 Zirconium/yttrium ratio, Atlantis Bank, 5b, 83b

## SITE INDEX

Site 556, greenschist-facies alteration, 491b  
 Site 642, basalt, porosity-hydrogen relationship, 262b

Site 732  
 alteration  
 low-temperature, 53a  
 static hydrothermal, 53a  
 basalt  
 geochemistry, 54a, 55a, 56a, 380b-381b  
 petrography, 49a  
 petrology, 380b-381b  
 basement description conventions, 7a-11a, 12a  
 bathymetry, 4a, 6a, 45a, 61a, 76b, 79a, 91a, 364b  
 inactive northern trace, 374b-375b  
 biostratigraphy, 19a, 48a-49a  
 cataclastic deformation, 51a-53a  
 clinopyroxene  
 chemical composition, 10b-11b, 15b, 17b  
 iron oxide/magnesium oxide ratio, 19b  
 coring summary, 48a  
 detachment faulting, 392b-393b  
 diabase, petrography, 49a-50a  
 dredge survey, 377b, 379b-381b  
 gabbro  
 geochemistry, 381b  
 petrography, 50a, 132b-133b  
 petrology, 381b  
 plastic deformation, 50a  
 geochemistry, 54a, 147b-149b, 382b  
 trace elements, 5b-6b, 14a, 17b  
 whole-rock, 4b-5b  
 geological setting, 3a-5a, 43a  
 glass, 382b

gneiss, foliated gabbro, 52a  
 hard-rock guidebase (HRGB)  
 assembly and deployment, 28a-29a  
 mechanical design evaluation, 30a-31a  
 hornblende, 52a  
 ilmenite, primary igneous, 52a  
 lead isotopes, 137b  
 lead/neodymium isotope ratio, 138b  
 lithic sandstone, 50a  
 lithology, 42a, 149b  
 distribution and associations, 377b, 380b  
 lithostratigraphy, 49a  
 location, 42a, 145b, 334b, 432b  
 magma chamber, 14b  
 magnetic anomalies  
 bathymetry, 378b  
 identification, 382b-384b  
 near transform, 384b-389b  
 magnetic properties, hard rock, 21a  
 median tectonic ridges, 371b-373b  
 tectonic evolution, 392b  
 metamorphism, medium-temperature, 51a-53a  
 mineral chemistry, 5b  
 mineralogy, 5b, 146b-147b  
 covariation, 12b-13b, 18b  
 mylonite, serpentinized peridotite, 54a  
 navidrill coring system, 34a-36a  
 normal faulting, ridge-transform intersection, 394b, 396b  
 objectives, 3a, 42a-43a  
 olivine  
 chemical composition, 6b-7b, 15b, 20b  
 iron oxide/magnesium oxide ratio, 19b  
 operations, 43a-48a  
 television/sonar survey, 43a-44a, 47a  
 orthopyroxene, chemical composition, 12b

oxides, 14a  
 peridotite, 381b  
 physical properties, 19a  
 physiography, 359b-361b  
   SeaBeam survey, 363b-364b  
 plagioclase  
   chemical composition, 8b-9b, 15b  
   recrystallized, 52a  
 plastic deformation, 50a, 51a  
 plate tectonics, 44a, 45a  
 positive displacement coring motor (PDCM)  
   mechanical evaluation, 26a-28a  
   operating pressures, 25a-26a  
   unsupported spudding, 25a  
 rare earth elements, 148b, 150b  
 ridge-transform intersection, 366b  
   bathymetry, 365b, 367b  
   inside corner highs, 368b  
   nodal basins, 366b  
   outside corners, 368b  
   topographic asymmetry, 368b  
   transform volcano, 366b, 368b  
 rift valleys, physiography, 364b-366b  
 sediment ponds, seismic reflection profiling, 370b  
 sedimentology, 48a  
 seismic stratigraphy, reflection profiling, 46a  
 serpentine, with iron chromite augen, 55a  
 silicates, 5b-6b  
 source provinces, 149b  
 strontium/neodymium ratio, 136b  
 tectonic evolution, 43a, 359b-396b  
 thermal conductivity, 19a-21a, 23a  
 transform tectonic zone  
   inactive fracture zone valley, 373b  
   nontransform walls, 373b, 376b  
   paleotransform walls, 376b-377b  
   present-day transform slip zone, 373b  
   transform walls, 371b  
 transtensional transform basins, tectonic evolution, 389b-392b  
 transverse ridges, 368b, 369b, 371b  
   tectonic evolution, 392b-396b  
 ultramafic rock  
   petrography, 50a  
   serpentinitization, 53a-54a

Site 733  
 basalt  
   geochemistry, 380b-381b  
   petrology, 380b-381b  
 basement description conventions, 7a-11a, 12a  
 bathymetry, 4a, 6a, 61a, 76b, 79a, 91a, 364b  
   inactive northern trace, 374b-375b  
 biostratigraphy, 19a  
 clinopyroxene  
   chemical composition, 10b-11b, 15b, 17b  
   iron oxide/magnesium oxide ratio, 19b  
 coring summary, 64a  
 deformation, 67a-71a  
 detachment faulting, 392b-393b  
 dredge survey, 377b, 379b-381b  
 gabbro  
   geochemistry, 71a-73a, 381b  
   magnetic properties, 73a  
   metamorphism and deformation, 67a-71a  
   petrographic summary, 132b-133b  
   petrology, 381b  
   physical properties, 73a-75a  
 geochemistry, 71a-73a, 382b  
   trace elements, 12b, 14a, 17b  
   whole-rock, 4b-5b  
 geological setting, 3a-5a, 60a  
 glasses, 382b

hard-rock guidebase (HRGB)  
   assembly and deployment, 28a-29a  
   mechanical design evaluation, 30a-31a  
 lead isotopes, 137b  
 lead/neodymium isotope ratio, 138b  
 lithology, 60a  
   distribution and associations, 377b, 380b  
 lithostratigraphy, 63a, 65a  
 location, 59a, 334b, 432b  
 magma chamber, 14b  
 magnetic anomalies  
   bathymetry, 378b  
   identification, 382b-384b  
   near transform, 384b-389b  
 magnetic properties, 60a, 73a, 75a  
   hard rock, 21a  
 median tectonic ridges, 371b-373b  
   tectonic evolution, 392b  
 metabasalt, 71a  
 metagabbro  
   brown, 70a, 71a  
   foliated, 65a  
   green, 68a, 69a  
   physical properties, 76a  
 metamorphism, 67a-71a  
 mineral chemistry, 5b  
 mineralogy, 5b  
   covariation, 12b-13b, 18b  
 navidrill coring system, 34a-36a  
 navigation, 64a  
 normal faulting, ridge-transform intersection, 394b, 396b  
 objectives, 3a, 59a, 60a  
 olivine  
   chemical composition, 6b-7b, 15b, 20b  
   iron oxide/magnesium oxide ratio, 19b  
 operations, 60a, 62a-63a  
   television/sonar survey, 60a, 62a-63a  
 orthopyroxene, chemical composition, 12b  
 oxides, 14a  
 peridotite, 381b  
 physical properties, 19a, 73a-75a, 76a  
 physiography, 359b-361b  
   SeaBeam survey, 363b-364b  
 plagioclase, chemical composition, 8b-9b, 15b  
 positive displacement coring motor (PDCM)  
   mechanical evaluation, 26a-28a  
   operating pressures, 25a-26a  
   unsupported spudding, 25a  
 primary mineralogy, 65a-67a  
 ridge-transform intersection, 366b  
   bathymetry, 365b, 367b  
   inside corner highs, 368b  
   nodal basins, 366b  
   outside corners, 368b  
   topographic asymmetry, 368b  
   transform volcano, 366b, 368b  
 rift valleys, physiography, 364b-366b  
 sediment ponds, seismic reflection profiling, 370b  
 sedimentary rock  
   classification, 13a-15a  
   description conventions, 15a-18a  
 seismic stratigraphy, reflection profiling, 62a  
 silicates, 12b  
 strontium/neodymium ratio, 136b  
 tectonic evolution, 60a, 359b-396b  
 thermal conductivity, 19a-21a, 23a, 60a, 74a-75a  
 transform tectonic zone  
   inactive fracture zone valley, 373b  
   nontransform walls, 373b, 376b  
   paleotransform walls, 376b-377b  
   present-day transform slip zone, 373b

transform walls, 371b  
 transtensional transform basins, tectonic evolution, 389b-392b  
 transverse ridges, 368b, 371b  
   tectonic evolution, 392b-396b  
 troctolite, chemical composition, 13b

Site 734  
 basalt  
   geochemistry, 380b-381b  
   petrology, 380b-381b  
 basement description conventions, 7a-11a, 12a  
 bathymetry, 4a, 6a, 61a, 76b, 79a, 91a, 364b  
   inactive northern trace, 374b-375b  
 biostratigraphy, 19a, 84a  
 clinopyroxene  
   chemical composition, 10b-11b, 15b, 17b  
   iron oxide/magnesium oxide ratio, 19b  
 coring summary, 83a  
 detachment faulting, 392b-393b  
 dredge survey, 377b, 379b-381b  
 gabbro  
   geochemistry, 381b  
   petrographic summary, 132b-133b  
   petrology, 381b  
 geochemistry, 86a-87a, 147b-149b, 382b  
   trace elements, 12b, 14a, 17b  
   whole-rock, 4b-5b  
 geological setting, 3a-5a, 79a  
 glasses, 382b  
 gravel, 432b-438b  
   depositional mechanisms, 436b, 438b  
   grain sorting, 432b-433b  
   proximal source, 434b-435b  
   sedimentology, 433b-434b  
 hard-rock guidebase (HRGB)  
   assembly and deployment, 28a-29a  
   mechanical design evaluation, 30a-31a  
 lead isotopes, 137b  
 lead/neodymium isotope ratio, 138b  
 lithologic sequence, 78a  
 lithology, 149b  
   distribution and associations, 377b, 380b  
 lithostratigraphy, 85a  
 location, 78a, 145b, 335b, 432b  
 mafic rock  
   metamorphosis, 86a  
   petrography, 85a  
 magma chamber, 14b  
 magnetic anomalies  
   bathymetry, 378b  
   identification, 382b-384b  
   near transform, 384b-389b  
 magnetic properties, hard rock, 21a  
 mass-wasting, 435b  
 median tectonic ridges, 371b-373b  
   tectonic evolution, 392b  
 metamorphism, 85a-86a  
 mineral chemistry, 5b  
 mineralogy, 5b, 146b-147b  
   covariation, 12b-13b, 18b  
 navidrill coring system, 34a-36a  
 navigation, 82a, 83a  
 normal faulting, ridge-transform intersection, 394b, 396b  
 objectives, 3a, 78a-79a  
 olivine  
   chemical composition, 6b-7b, 15b, 20b  
   iron oxide/magnesium oxide ratio, 19b  
 operations, 79a-84a  
   television/sonar survey, 80a, 81a, 82a, 83a  
 orthopyroxene, chemical composition, 12b  
 oxides, 14a  
 peridotite, 381b

## SITE INDEX

petrography, 85a  
 physical properties, 19a  
 physiography, 359b-361b  
   SeaBeam survey, 363b-364b  
 plagioclase, chemical composition, 8b-9b, 15b  
 positive displacement coring motor (PDCM)  
   mechanical evaluation, 26a-28a  
   operating pressures, 25a-26a  
   unsupported spudding, 25a, 80a  
 rare earth elements, 148b, 150b  
 ridge-transform intersection, 366b  
   bathymetry, 365b, 367b  
   inside corner highs, 368b  
   nodal basins, 366b  
   outside corners, 368b  
   topographic asymmetry, 368b  
   transform volcano, 366b, 368b  
 rift valleys, physiography, 364b-366b  
 sediment ponds, seismic reflection profiling, 370b  
 sedimentology, 84a  
 seismic stratigraphy, 80a  
   reflection profiling, 81a  
 serpentized peridotite  
   deformation, 85a  
   petrography, 85a  
 silicates, 12b  
 source provinces, 149b  
 static alteration, 86a  
 strontium/neodymium ratio, 136b  
 tectonic evolution, 79a, 359b-396b  
 thermal conductivity, 19a-21a, 23a  
 transform tectonic zone  
   inactive fracture zone valley, 373b  
   nontransform walls, 373b, 376b  
   paleotransform walls, 376b-377b  
   present-day transform slip zone, 373b  
   transform walls, 371b  
 transtensional transform basins, tectonic evolution, 389b-392b  
 transverse ridges, 368b, 371b  
   tectonic evolution, 392b-396b  
 troctolite, chemical composition, 13b  
 veins, calcium silicate, 86a

## Site 735

acoustic borehole televiewer, 262b, 264b  
 acoustic log, 553b  
 alteration, 488b-505b  
   cataclastic, 138a  
   chemical effects, 546b-547b  
   dynamic model for layer 3, 547b-548b  
   hydrothermal, 120b-121b, 136b-138b  
   late-stage oxidation, 546b  
   lithostratigraphic correlation, 266b-267b  
   oxidative, 138a-139a  
   porosity and, 261b-262b, 263b  
   pressure of, 211b-212b  
   static, 138a, 544b  
   stratigraphy, 503b-505b  
   styles of, 207b-209b, 543b-546b  
   temperature of, 209b-211b  
   in undeformed gabbro, 138a  
   vein density and, 233b  
 alteration index, 171b, 173b  
 aluminum oxide/titanium oxide ratio, 140b  
 amphibole  
   in metagabbros and late magmatic intrusions (LMIs), 164b  
   mineral chemistry, 190b-198b  
   orientation in metagabbro, 241b  
   replacement, 134b  
   secondary matrix, 506b  
   trace elements, 56b

amphibole gneiss, foliated, 549b  
 amphibole veins, 545b  
   alteration stratigraphy, 504b  
   dip orientation, 499b  
   oblique slip, 497b  
   slip direction, 501b  
 amphibole-plagioclase veins, 551b  
 andesine veinlet, 189b  
 anisotropy, 234b, 236b  
   foliated gabbro and metagabbro, 157a, 159a  
   vs. pressure, 240b  
 apatite, 167b  
 basalt  
   geochemistry, 380b-381b  
   petrography, 117a  
   petrology, 380b-381b  
 basalt dike, geochemistry, 147a  
 basement, 89a  
   description conventions, 7a-11a, 12a  
   lithostratigraphy, 98a  
   permeability, 347b  
 bathymetry, 4a, 6a, 61a, 76b, 79a, 91a, 364b,  
   400b, 410b, 416b  
   inactive northern trace, 374b-375b  
 biostratigraphy, 19a  
 biotite, 167b-168b, 180b  
 borehole televiewer, 205a-207a  
 breccia  
   cataclastic, 419b  
   hydrothermal, 179b  
   magmatic/hydrothermal, 165b-166b  
   mineralogy, 165b-166b, 172b-173b, 179b  
   trondhjemite intrusion, 535b  
 calcium number, 477b  
 calcium/magnesium ratio, 475b  
 carbon, 117b  
 cation exchange capacity, 325b, 326b  
   vs. electrical resistivity, 327b  
 chlorite, 167b  
 chromium, 136b, 480b  
 clinopyroxene  
   chemical composition, 10b-11b, 15b, 17b, 69b  
   contact relationships, 69b  
   iron oxide/magnesium oxide ratio, 19b  
   in iron-titanium oxide gabbro, 64b  
   oxide minerals, 65b, 66b, 97b  
   trace elements, 51b, 53b, 57b-59b  
   constant-rate injection tests, 336b, 344b  
   corona structure, 118b, 163b, 165b  
   crack porosity, 231b, 233b  
   crystalline rock, physical properties, 157a,  
     158a-159a, 160a-161a  
 deformation, 488b-505b  
   alteration with, 136a-138a  
   braided zone, 457b  
   brittle, 90a, 105a, 159b, 426b-427b, 499b,  
     501b  
   brittle-ductile, 491b-499b, 500b  
   cataclastic, 106a, 136a-138a  
   classification, 492b-493b  
   distribution, 411b  
   Lower Structural Domain, 413b-114b  
   Middle Structural Domain, 413b  
   partial reorientation, 409b-414b  
   petrofabric studies, 159b  
   post-kinematic, 159b  
   stratigraphy, 156b-157b  
   symmetry of foliation, 500b  
   textural types, 155b, 171b  
   timing and temperature, 172b  
 Upper Structural Domain, 411b, 413b  
 See also Ductile deformation; Plastic deformation

deformation structures, 209a  
 density, 231b-232b, 325b-326b, 327b  
   seismic stratigraphy and, 507b  
   vs. electrical resistivity, 326b-327b  
 detachment faulting, 392b-393b  
 diopside, 180b  
   in veins, 551b  
 diopside olivine gabbro, 447b  
 disseminated iron-titanium oxide gabbro,  
   petrography, 47b  
 disseminated oxide olivine gabbro, 448b  
   contact relationships, 459b-460b  
   mineralogy, 26b  
   origin, 461b-462b  
 dredge survey, 377b, 379b-381b  
 ductile deformation, 163b  
   Lower Structural Domain, 426b  
   magmatic flow, 418b  
   Middle Structural Domain, 425b-426b  
   solid-state, 418b-422b  
   subsolidus, 536b  
   Upper Structural Domain, 423b  
 elasticity, 236b  
 electrical conductivity  
   mechanisms, 329b  
   salinity and, 327b-328b, 329b  
 electrical resistivity, 325b-326b, 555b  
 epidote, 168b  
   in plagioclase-diopside veins, 551b  
 feldspar, mineral chemistry, 198b-199b, 201b  
 felsic rock, 117a-119a  
 felsic veins, 545b-546b  
   alteration stratigraphy, 504b-505b  
   spheire crystals, 549b  
 ferrogabbro, 155b  
 fluid inclusions, 200b-206b  
   homogenization temperature, 203b, 204b,  
     205b, 206b  
   ice-melting temperature, 203b, 204b, 205b  
   laser Raman spectroscopy, 205b-206b  
   in plagioclase, 190b  
   in quartz, 197b  
   in vein diopside, 197b  
 foliation, dip orientation, 107a  
 fracturing  
   dip vectors, 269b  
   late subvertical, 505b  
   lithostratigraphic correlation, 266b-267b  
   smectite lining, 546b  
   vertical distribution, 556b  
 gabbro  
   augen gneissic, 132a, 133a  
   brittle deformation, 426b-427b  
   crystallization model, 35b, 36b  
   deformed textures, 103a, 129a-132a, 497b  
   ductile deformation, 418b-426b  
   electrical conductivity, 323b-330b  
   geochemistry, 25b, 381b  
   intrinsic rock properties, 556b  
   mafic minerals, 42b, 45b, 50b-51b  
   magnetic properties, 285b-305b, 309b-320b  
   mineral chemistry, 484b-485b  
   petrography, 132b-133b, 154b-155b, 445b  
   petrology, 381b  
   plastic deformation, 399b-408b  
   poorly foliated, 131a  
   porphyroclastic, 131a, 133a, 134a, 134b  
   radiogenic isotopes, 131b-133b  
   small-scale textures, 121a  
   stable isotopes, 129b-133b  
   strontium/aluminum values, 84b  
   textural variation, 119a-121a  
   ultrasonic attenuation, 253b-259b  
 gabbronorite, 445b, 449b

mineralogy, 25b-26b  
petrography, 44b, 47b  
geochemical log  
statistical analysis, 272b-281b  
Zones P1-P7, 277b-280b  
geochemistry, 139a-150a  
bulk hole composition, 483b-486b  
chemical stratigraphy, 147a-150a  
lithologic variability, 470b-488b  
trace elements, 14a, 17b  
whole-rock, 4b-5b, 527b-530b  
geological setting, 3a-5a, 21b-22b, 44b,  
90a-91a, 113b-114b, 127b-128b, 182b,  
272b, 285b, 347b, 440b-441b  
geomagnetic field, 198a  
geophysical log, 265b  
glasses  
geochemistry, 382b  
major oxides, 86b-89b  
gneiss, alteration, 179b  
grain-size variations, 124a, 129a  
gravel  
shape factors, 437b  
size distribution, 436b  
greenschist-facies alteration, 489b, 491b  
hard-rock guidebase (HRGB)  
assembly and deployment, 28a-29a  
mechanical design evaluation, 30a-31a  
hornblende, 549b  
oscillatory zonation, 188b  
hydraulic testing, 266b  
hydrogen isotopes, 129b-131b, 133b,  
135b-136b  
igneous lamination, 106a-107a  
igneous rock  
classification, 444b  
lithology, 441b-444b  
petrogenesis, 441b-444b  
primary lithologies, 516b-526b  
igneous stratigraphy, 22b  
Unit I, 449b, 450b, 456b-457b  
Unit II, 451b, 457b-459b  
Unit III, 459b-462b  
Unit III/Unit IV contact, 462b, 463b, 532b  
Unit IV, 453b, 462b, 464b-465b  
Unit IV/Unit V contact, 465b, 466b  
Unit V, 453b, 465b-466b  
Unit V/Unit VI contact, 467b, 533b  
Unit VI, 454b-455b, 467b-470b  
ilmenite  
magnesium oxide/manganese oxide content,  
99b  
trace elements, 56b  
ilmenite-magnetite, in deformed gabbros,  
132a-134a, 135a  
impedance, 190a  
intrusive contacts, 106a  
interstitial olivine-bearing iron-titanium oxide  
gabbro, petrography, 47b-48b  
iron oxide, 135b  
velocity and, 250b  
iron-titanium oxide gabbro, 121a, 464b  
geochemistry, 146a-147a  
magnetic properties, 288b-289b  
mineralogy, 41b, 44b, 46b, 49b, 71b, 72b  
mineralogy, origin of variation, 59b, 61b-62b  
olivine gabbro contact, 56b, 59b, 73b  
petrography, 43b, 112a-117a  
undeformed, 123a  
iron-titanium oxide-rich mylonite  
origin of oxides in, 407b  
plastic deformation, 402b, 406b  
strain localization, 406b-407b

iterative nonhierarchical cluster analysis  
(INCA), 273b-275b  
model 1, 275b-277b, 281b  
model 2, 277b-280b, 281b, 282b  
layer 3  
lithostratigraphic evolution, 439b-513b  
logging data, 325b  
metamorphism and alteration, 541b-548b  
seismic reflection profiling, 245b-250b  
ultrasonic attenuation, 253b-259b  
layering, 121a-122a, 128a  
lead isotopes, 129b-133b, 137b, 139b  
lead/neodymium isotope ratio, 138b  
leucocratic diopside-bearing veins, 198b  
fluid inclusions, 204b-205b  
petrography, 188b-189b  
leucocratic rock, 155b  
liquid immiscibility, 96b-98b  
liquid line of descent, 82b-84b, 90b, 95b-96b,  
98b-101b  
lithology  
distribution and associations, 377b, 380b  
logging data, 183b  
variability, 444b  
lithostratigraphy, 4b, 22b, 77b-78b, 113b-114b,  
128b-131b, 232b, 324b, 411b, 415b-417b,  
424b, 443b  
Unit I, 89a, 97a, 208b, 311b, 542b  
Unit II, 89a, 97a, 99a-100a, 208a-209a, 311b,  
542b  
Unit III, 90a, 100a-101a, 209a, 312b, 542b  
Unit IV, 90a, 102a, 209a, 312b-313b, 542b-  
543b  
Unit V, 90a, 102a, 209a, 313b, 543b  
Unit VI, 90a, 102a-103a, 209a, 313b, 543b  
location, 89a, 90a, 114b, 182b, 262b, 272b,  
324b, 336b, 410b, 416b, 432b  
logging, 209a-210a  
azimuth, deviation, and caliper, 170a  
dual induction-sonic-caliper-natural gamma,  
168a, 173a  
dual laterolog resistivity, 169a, 176a, 177a  
geochemical, 169a, 174a, 175a, 177a-178a  
geochemical, vs. X-ray fluorescence, 179a  
induction-resistivity-caliper, 169a, 176a, 177a  
litho-density, 169a  
magnetometer, 171a  
physical properties, 169a, 172a, 177a  
summary logs, 214a-222a  
temperature, 171a  
logging comparisons, 555b-556b  
macroscopic features, 143b  
magma  
liquid source, 487b-488b  
viscous flow in solidifying, 400b, 403b, 404b  
magma chamber, 14b  
magma mixing, 508b  
magma, parental  
composition, 83b-86b  
excluded elements, 86b  
Types A-D, 85b  
magmatic differentiation  
conditions, 101b-102b  
in deforming crust, 103b-104b  
magmatic foliation, 419b, 499b  
magmatic intrusions, 445b, 507b-509b  
amphibole composition, 165b  
mineralogy, 167b  
plagioclase composition, 164b  
pyroxene composition, 164b  
structures not related to, 173b  
structures related to, 172b  
magmatic stratigraphy, 154b-155b  
magnesium number, 476b

magnesium oxide, 136b  
magnetic anomalies, 307b  
bathymetry, 378b  
identification, 382b-384b  
intrusive rocks and origin of, 300b-302b  
layer 3 magnetic properties and, 320b  
near transform, 384b-389b  
magnetic properties, 90a, 150a-157a, 209a  
alternating field demagnetization, 153a-154a,  
155a, 156a, 291b-292b, 294b-295b, 302b  
degree of metamorphism/alteration and, 294b-  
296b, 300b, 303b  
directional coherence, 319b-320b  
discrete samples vs. magnetic logs, 313b-318b  
hard rock, 21a  
inclinations, 154a-156a, 157a  
intensity of magnetization vs. magnetic field,  
313b-315b  
Koenigsberger ratio, 153a, 154a, 290b-301b,  
305b  
lithostratigraphic correlation, 310b-313b  
logging data, 309b-320b, 554b-555b  
median demagnetizing field, 305b  
natural remanent magnetization, 151a, 153a,  
156a, 286b, 291b, 304b, 317b  
secondary component, 293b-294b  
stable inclination, 292b-293b, 296b, 297b-  
300b, 301b, 302b, 315b-317b  
susceptibility, 151a-153a, 154a, 286b, 291b,  
304b, 314b, 315b, 316b  
thermal demagnetization, 154a, 291b-292b,  
296b  
vertical magnetic field, 316b, 317b  
magnetometer logs, 191a-193a, 200a, 201a,  
202a, 203a, 210a  
median tectonic ridges, 371b-373b  
evolution, 392b  
melanogranophyre absence, 102b-103b  
melt density, 98b  
metagabbro  
amphibole composition, 165b  
deformation, 136a  
foliated, 287b  
magnetic properties, 287b  
physical properties, 157a  
metagabbro, veined  
fluid inclusions, 202b-204b  
petrography, 186b-187b  
metal content, 120b  
metamorphism  
dynamothermal, 90a, 129a-132a  
high-temperature, 543b-544b  
mineral assemblages, 159b-160b, 163b, 165b-  
170b  
mineral distribution, 171b-172b  
stratigraphy, 156b-157b  
styles of, 171b-172b  
sulfide-oxide relationship, 134a-136a  
synkinematic conditions, 420b-421b  
mica, 167b  
microgabbro  
contact relationships, 458b  
location, 118a  
textural variation, 130a  
mineral chemistry, 5b, 49b-56b, 154b,  
189b-200b  
variations between rock types, 49b  
mineralogy, 5b, 90b-96b, 99a, 184b-185b  
of ambiguous origin, 166b-170b  
covariation, 12b-13b, 18b, 78b-79b, 80b  
genetic relationships, 33b-35b  
igneous rock, 23b-27b  
incompatible elements, 166b-167b  
iron-titanium oxides, 123a-125a

## SITE INDEX

lithologic variability, 470b-488b  
 lithostratigraphic correlation, 542b-543b  
 of metamorphosed gabbro, 125a-126a, 167b  
 post-kinematic, 160b, 163b, 165b-166b  
 primary, 108a-115a  
 secondary, 199b-200b, 202b-203b  
 silicate phase relationship, 126a-129a  
 sulfide, 125a-126a  
 in undeformed gabbro, 125a  
 variations between rock types, 27b, 29b-31b,  
   33b-35b  
 velocity and, 233b-234b  
 multichannel sonic logs, 213a, 264b, 266b  
 operations, 178a, 180a-182a, 183a, 184a  
 tools, 178a  
 mylonite  
   shear, 421b  
   veined, 186b-187b  
 mylonitic gabbro, 132a  
 mylonitic metagabbro, veined, fluid inclusions,  
   202b  
 navidrill coring system, 96a  
   assembly, 34a-35a  
   deck test, 35a-36a  
   deployment, 36a-37a  
   mechanical design evaluation, 37a-38a  
   unsupported spudding, 37a  
 neodymium isotopes, 129b-133b, 139b  
 nickel, 136b  
 nickel/aluminum ratio, 120b  
 niobium, 135b  
 normal faulting, ridge-transform intersection,  
   394b, 396b  
 objectives, 3a, 90a, 127b  
 olivine  
   chemical composition, 6b-7b, 15b, 20b, 23b,  
   52b-53b  
   chemical composition, at contacts, 69b  
   dynamic recrystallization, 401b  
   iron oxide/magnesium oxide ratio, 19b  
   in metagabbro, 173b  
   mineralogy, 90b, 481b-483b  
   plastic deformation, 402b  
   pseudomorphed, 550b  
   trace elements, 56b, 62b  
 olivine gabbro, 125a, 445b, 449b, 465b-466b,  
   467b, 468b  
 clothespin structure, 538b  
 contact relationships, 28b, 32b, 457b-459b  
 ductile deformation, 534b  
 geochemistry, 146a  
 magnetic properties, 287b-288b, 289b-290b  
 mineralogy, 25b, 41b  
 oxide-bearing gabbro contact, 79b  
 petrography, 107a-111a  
 olivine gabbronorite, petrography, 47b  
 olivine iron-titanium oxide microgabbro, 48b  
 olivine microgabbro, 445b, 447b, 467b-468b  
   mineralogy, 25b  
 olivine pigeonite gabbro, 47b  
 olivine-bearing gabbro  
   magnetic properties, 287b-288b  
   mineralogy, 41b  
   petrography, 107a-111a  
 olivine-rich gabbro, magnetic properties, 290b  
 olivine-titanium oxide gabbro, 48b  
 opaque minerals, 101a  
   petrology, 114b-117b  
 operations, 91a-97a, 324b, 334b-335b  
   logging, 96a-97a  
   television/sonar survey, 92a, 94a  
 orthopyroxene  
   chemical composition, 12b, 53b, 60b-61b  
   in clinopyroxene, 122a

oxide minerals, 67b  
 oxide gabbro, 448b  
   geochemistry, 479b-480b, 482b-483b  
   mineralogy, 479b-480b  
 oxide gabbronorite, 448b  
 oxide microgabbro, 445b, 467b  
 oxide minerals, 69b, 91b, 94b, 102b  
   in orthopyroxene, 67b  
 oxide olivine gabbro, 26b-27b, 448b, 468b  
   contact relationships, 462b, 464b  
   modality, 33b  
   origin, 464b-465b  
   penetrative deformation, 469b  
 oxide olivine microgabbro, 448b  
 oxide-bearing gabbro  
   in brittle-ductile deformation zones, 469b  
   penetrative foliation, 460b  
   textural relationships, 461b  
 oxide-free gabbro  
   geochemistry, 475b-479b  
   mineralogy, 475b-479b  
 oxide-rich gabbro, magmatic extension cracks,  
   460b  
 oxides, 14a  
 oxygen isotopes, 129b-131b, 133b, 135b-136b,  
   154b, 158b, 170b-171b  
   controls on, 173b-174b  
   plagioclase vs. clinopyroxene, 139b, 140b  
 patchy olivine gabbro, 447b, 467b-468b  
   protolith, 470b  
 patchy oxide olivine gabbro, 448b  
 peridotite, 381b  
 permeability, 90a, 200a-205a, 210a  
   along crosscutting fractures, 554b  
   constant-rate injection vs. slug tests, 345b  
   drill-string packer experiments, 336b-339b  
   *in-situ* vs. calculated, 336b  
   lithostratigraphic correlation, 266b-267b,  
   346b  
   logging data, 267b-268b  
   transmissivity and, 341b-342b, 343b, 345b  
   vertical distribution, 556b  
 petrography, 90b, 107a-122a  
   lithostratigraphic correlation, 48b, 186b,  
   542b-543b  
 petrology, Zones I-IV, 77b-78b  
 physical properties, 19a, 90a, 157a-163a, 227b,  
   229b-231b  
 physiography, 359b-361b  
   SeaBeam survey, 363b-364b  
 pigeonite gabbro, 47b, 120a  
 plagioclase  
   chemical composition, 8b-9b, 15b, 53b-56b,  
   69b  
   contact relationships, 69b  
   in disseminated oxide olivine gabbro, 31b  
   dynamic recrystallization, 401b, 419b  
   in iron-titanium oxide gabbro, 62b, 64b, 69b  
   in late magmatic intrusions, 164b  
   in metagabbro, 164b, 189b, 239b  
   mineralogy, 90b-91b  
   in olivine gabbro, 30b  
   orientation, 239b, 251b  
   in oxide olivine gabbro, 34b  
   plastic deformation, 402b  
   recrystallized grain size, 421b-422b  
   trace elements, 52b-56b, 62b  
   veinlet, 188b  
   xenocrysts, 179b  
 plastic deformation, 103a-105a, 155b, 159b,  
   172b  
 gabbros with large recrystallized grains, 403b-  
   405b, 406b-407b

gabbros with small recrystallized grains,  
   405b-406b, 407b  
 porosity, 258b, 325b-326b, 327b  
 alteration and, 261b-262b, 263b  
 formation factor and, 328b, 331b  
   vs. resistivity, 328b, 331b  
 porphyroclastic gabbro, petrography, 107a  
 positive displacement coring motor (PDCM)  
 mechanical evaluation, 26a-28a  
 operating pressures, 25a-26a  
 unsupported spudding, 25a, 31a-32a  
 pressure, permeability and, 337b-339b,  
   341b-342b, 344b  
 principal component analysis, 273b, 274b, 283b  
 pseudomorphs, 163b, 165b  
 pyroxene  
   breccia and veins, 162b  
   in clinopyroxene, 169b  
   in deformed gabbros and late magmatic intru-  
   sions (LMIs), 161b  
   in foliated metagabbro, 549b  
   mineralogy, 90b-91b  
   olivine vs. oxide-bearing olivine gabbro, 531b  
 reaction corona, 135a, 137a  
 residual melt porosity, 79b-82b, 106b-107b  
 lithostratigraphic correlation, 81b-82b  
 ridge-transform intersection, 366b  
   bathymetry, 365b, 367b  
   inside corner highs, 368b  
   nodal basins, 366b  
   oblique slip, 498b  
   outside corners, 368b  
   topographic asymmetry, 368b  
   transform volcano, 366b, 368b  
 rift valleys, physiography, 364b-366b  
 rotary coring bit, 93a, 95a-96a  
   operations, 32a-33a  
   performance evacuation, 33a-34a  
 rubidium, 135b  
 salinity and, 330b  
 sediment ponds, seismic reflection profiling,  
   370b  
 seismic anisotropy, 554b  
 seismic stratigraphy  
   data processing, 221b-222b  
   methods, 219b-221b  
   properties controlling reflectivity, 247b-249b  
   reflection profiling, 93a, 223b-225b  
 seismic velocity, 553b  
 shear-wave splitting, 241b  
 silica, velocity and, 247b  
 silicates  
   in igneous rock, 24b  
   mineral fractionation, 99b-100b  
   mineralogy, 92b-93b, 108b-111b  
   trace elements, 12b  
   slug tests, 335b-336b, 341b-342b  
   sodium oxide, 135b, 478b  
   stable isotopes, 206b-207b  
   stable remanent magnetic vector, 409b-410b  
 stress, 556b  
   seawater circulation and, 505b-507b  
 strontium isotopes, 129b-131b, 139b  
   vs. oxygen isotopes, 140b  
 strontium in parental magma, 107b  
 strontium/neodymium ratio, 136b  
 structural stratigraphy, 411b, 417b, 424b  
 sulfide  
   igneous, 114b-115b, 116b, 119b-120b  
   recrystallization changes, 93b-95b  
   secondary, 115b-117b, 118b, 120b-121b  
 sulfur, 117b  
   in basaltic glasses, 88b-89b  
   distribution, 119b-120b

evolution, 99b, 101b  
 hydrothermal alteration, 120b-121b, 123b  
 sulfur isotopes, 118b-119b, 121b-123b  
 surface conductivity vs. resistivity, 328b  
 synkinematic differentiation, 486b-487b  
 synkinematic late-magmatic gabbro, 448b  
 synkinematic minerals, 160b, 165b-166b  
   textural type correlation, 171b  
 synthetic seismogram, 245b-247b, 248b, 251b  
 talc, 167b  
 tectonic evolution, 90a-91a, 359b-396b  
   three-step scenario, 427b-428b  
 tectonic setting, 76b-77b, 415b-416b, 541b  
 tectonic stratigraphy, Zones 1-5, 501b-503b  
 temperature, 98b  
   borehole profile, 266b  
   equilibrium, 352b-353b, 354b  
   logging data, 165a-168a, 171a, 347b, 349b  
 texture  
   deformed gabbros, 403b, 421b, 508b  
   distribution by type, 107a  
   igneous rock, 23b-27b, 38b-39b, 41b-44b  
   late-magmatic deformation, 537b  
 thermal conductivity, 19a-21a, 23a, 163a-165a,  
   166a, 167a  
 thermal modeling, 347b-354b  
   one-dimensional conduction from transient  
     boundary temperatures, 352b-354b  
   two-dimensional conduction simulation of ad-  
     vection, 349b-352b  
   uniformly conducting model, 350b  
 titanium oxide, 135b, 479b  
 titanium oxide/calcium ratio, 474b  
 titanium oxide/magnesium ratio, 473b  
 titanomagnetite, magnesium oxide/aluminum  
   oxide content, 99b

trace elements, 138a  
 trace minerals, 147a, 150a  
 transform tectonic zone  
   inactive fracture zone valley, 373b  
   nontransform walls, 373b, 376b  
   paleotransform walls, 376b-377b  
   present-day transform slip zone, 373b  
   transform walls, 371b  
 transtensional transform basins, tectonic  
   evolution, 389b-392b  
 transverse ridges, 368b, 369b, 371b  
   tectonic evolution, 392b-396b  
 troctolite, 427b, 447b, 468b  
   chemical composition, 13b  
   geochemistry, 146a  
   location, 118a  
   magnetic properties, 290b  
   mineralogy, 24b-25b, 41b  
   petrography, 111a-112a  
 troctolitic gabbro, 447b, 468b-469b  
 troctolitic microgabbro, 24b-25b, 447b, 468b  
 troctolitic oxide gabbro, 448b  
 trondjemite, 447b-448b, 461b  
   alteration stratigraphy, 503b  
   fluid inclusions, 204b  
   intrusion breccia, 535b  
   oxygen isotope values and, 138b-139b  
   petrography, 187b  
   veins, 545b  
 trondjemite, geochemistry, 147a  
 ultrasonic attenuation  
   errors in measurement, 254b-255b  
   experimental methods, 253b-254b  
   measurements under pressure, 255b-256b  
 uplifted blocks, 207a-208a  
 veins

amphibole, 105a, 136a  
 carbonate, 104a, 552b  
 density, 171b, 494b-496b  
 dip orientation, 492b-493b  
 distribution, 506b  
 felsic, 119a  
 filling of inclined, 545b-546b  
 hornblende, 549b  
 hydrothermal-plagiogranite, 138a  
 mineral assemblages, 159b-160b, 163b, 165b-  
   170b  
 mineralogy, 172b-173b  
 synkinematic minerals, 165b-166b  
 thomsonite, 179b  
 velocity, 164a, 222b-223b, 228b-229b, 231b  
 attenuation and, 258b  
 crack porosity and, 233b, 234b  
 density relationship, 162a, 238b, 239b  
 for isotropic gabbros, 244b  
 logging data, 246b  
   vs. elasticity, 242b-243b  
   vs. porosity, 163a  
 vertical seismic profile, 182a-191a, 192a, 193a,  
   197a, 225b, 553b-554b  
 geological setting, 183a, 185a  
 methods, 185a-187a  
 navigation and weather, 187a  
 objectives, 182a  
 operations, 187a-188a  
 previous borehole seismology, 182a-183a  
 stacked seismograms, 194a-196a  
 water/rock ratio, 174b  
 zeolite, 168b  
   veins, 552b  
 zirconium, 135b