

INDEX TO VOLUME 119

This index provides coverage for both the *Initial Reports* and *Scientific Results* portions of Volume 119 of the *Proceedings of the Ocean Drilling Program*. Index entries with the suffix *a* refer to pages in the *Initial Reports*, and those with *b*, to pages in the *Scientific Results* (this book).

The index is presented in three parts: (1) a Subject Index, (2) a Site Index, and (3) a Paleontological Index.

The index was prepared by Wm. J. Richardson Associates, Inc., under subcontract to the Ocean Drilling Program. It expands the concept developed by the Deep Sea Drilling Project at Scripps Institution of Oceanography for a comprehensive, cumulative index of DSDP volumes. Both of these indexes are based on a hierarchy of entries: (1) a main entry, defined as a key word or concept followed by a reference to the page on which that word or concept appears; (2) a subentry, defined as a further elaboration on the main entry followed by a page reference; and (3) a sub-subentry, defined as an even further elaboration on the main entry or subentry followed by a page reference.

The Subject Index follows a standard format. Geographic and individual names are referenced in the index only if they are subjects of discussion. This index also includes broad fossil groups, such as foraminifers and radiolarians, which also appear in the Paleontological Index.

The Site Index is structured to contain entries for the sites discussed in the volume. Site entries are modified by subject subentries.

The Paleontological Index is an index relating to significant findings and/or substantive discussions, not of species names *per se*. This index covers three varieties of information: (1) broad fossil groups, including individual genera and species that have been erected or emended formally; (2) biostratigraphic zones; and (3) fossils depicted in illustrations. Taxonomic entries consisting of both genera and species are listed alphabetically *by species*; biostratigraphic zones, however, are listed alphabetically *by genus*.

The indexes cover text figures and tables but not core description forms ("barrel sheets") or core photographs. Also excluded are bibliographic references, names of individuals, and routine front and back matter.

For further information, and for information on electronic formats available, contact the Chief Production Editor, Ocean Drilling Program, 1000 Discovery Drive, College Station, Texas 77845-9547.

SUBJECT INDEX

Acoustic basement
 Kerguelen-Heard Plateau N: Site 737, sonobuoy velocity, 160a–161a
 Kerguelen-Heard Plateau S: Site 738, seismic reflection profile, 231a–232a
 Prydz Bay, 9a, 370a, 372a
 Acoustic impedance
 Kerguelen-Heard Plateau, reflection amplitude changes, 212a, 274a–275a
 Prydz Bay, 331a–332a, 438a
 Acoustic stratigraphy
 Prydz Bay
 angular unconformity, 9b, 11b
 areal distribution, 13b–14b, 16b–17b
 lithostratigraphic correlation, 99b–101b
 strata thickness, 18b
 Unit PS.1, 7b–8b, 14b, 18b–19b, 27b, 29b, 33b–34b, 36b, 99b–100b
 Unit PS.1/Unit PS.4 unconformity, 12b
 Unit PS.2A, 8b, 9b, 11b, 16b–17b, 18b–19b, 29b, 30b, 36b, 100b–101b, 876b
 Unit PS.2A/PS.2B unconformity, 8b, 20b
 Unit PS.2B, 8b, 11b–12b, 13b, 18b, 22b, 36b, 100b–101b
 Unit PS.4, 12b, 18b, 36b
 Unit PS.4/Unit PS.2B unconformity, 12b, 14b, 15b, 32b
 Unit PS.4/Unit PS.5 unconformity, 18b, 20b, 22b
 Units PS.5–6, 12b–13b, 36b
 vertical reflection changes, 18b
 Adelie Coast, Antarctic Bottom Water, 232b
 Air/sea interface
 Kerguelen Plateau S: Site 744, physical characteristics, 493a
 Kerguelen Sediment Ridge: Site 745, physical characteristics, 519a
 Alkalinity
 Kerguelen Plateau S: Site 744, 491a
 pore waters, 367b
 Kerguelen Sediment Ridge, 517a–518a, 545a
 pore waters, 369b, 373b
 Kerguelen-Heard Plateau N, 139a–140a, 186a
 carbonate precipitation and, 359b, 372b
 sulfate and calcium correlation, 362b
 Kerguelen-Heard Plateau S: Site 738, 256a–257a
 pore waters, 364b, 366b
 Prydz Bay, 312a, 362a, 374a, 418a
 calcium correlation, 380b, 385b
 measured vs. predicted values, 389b
 phosphate and ammonium correlation, 383b
 sediment/water interface, 385b
 sulfate reduction and, 388b
 n- Alkanes
 Prydz Bay, 410b
 odd carbon number predominance, 419b, 423b
 source, 408b, 409b
 Alteration, cold seawater, Kerguelen-Heard Plateau S: Site 738, basalts, 307b, 314b–315b
 Alteration, Kerguelen-Heard Plateau S: Site 738, basalts, 313b–315b
 Alteration, low-temperature
 Kerguelen-Heard Plateau S: Site 738
 basalts, 307b
 basement rocks, 314b
 Aluminum, Kerguelen-Heard Plateau N: Site 736, 356b
 Amery Depression, Holocene sediment accumulation, 747b
 Amery Formation

alnoite sill and mafic dikes, 795b, 804b
 kaolinitic matrix, 247b
 organic carbon content, 113b
 red beds, 797b
 Amery Group
 Lambert Graben, correlation, Prydz Bay: Site 740, 346a
 Permian sedimentary sequence, 53b, 66b
 correlation with Beacon Supergroup, 795b
 Amery Ice Shelf
 basal melting, 62b, 64b
 flow lines, seaward divergence, 62b, 64b
 freeze-on ice, 64b
 Permian strata, 9a
 position changes, 64b
 Prydz Bay, 7a
 regional geology, 5b
 shear and thinning of, 62b
 sources, 290a, 338a
 surface sediments, 749b
 Amino acids
 Kerguelen Plateau S: Site 744, 114a–115a
 Prydz Bay, 114a–115a, 116a, 117a
 Ammonia
 Kerguelen Plateau, high-resolution vs. regular sampling results, 395b
 pore water gradient, sedimentation rate and, 402b–403b
 Prydz Bay: Site 739, high-resolution vs. regular sampling results, 395b
 sulfate gradient and, relation to carbon flux, 401b, 403b
 Ammonium
 Kerguelen Plateau S: Site 744, 491a
 Kerguelen Sediment Ridge, 518a, 544a–545a
 pore waters, 369b, 373b
 Kerguelen-Heard Plateau N: Site 736, 137a–138a, 140a, 156a
 Kerguelen-Heard Plateau N: Site 737, 187a, 359b
 Kerguelen-Heard Plateau S: Site 738, 257a
 Prydz Bay: Site 739, 313a
 sulfate anticorrelation, 380b
 Prydz Bay: Site 740, 362a–363a, 374a
 phosphate and alkalinity correlation, 383b
 Prydz Bay: Site 741, 385a
 vs. phosphate, pore waters, 384b
 Prydz Bay: Site 742, 429a
 pore waters, 386b
 sulfate anticorrelation, 385b
 Prydz Bay: Site 743, 466a
 Amphibolite
 Prydz Bay, 138b, 453a
 provenance, 139b
 Amygdule, Kerguelen-Heard Plateau S: Site 738, 241a
 Anorthite, Kerguelen-Heard Plateau S: Site 738, 301b
 Antarctic Bottom Water
 deep-sea hiatus, Atlantic and Indian oceans, 874b
 formation, 246b, 694b
 glacial/interglacial cycles, 232b
 Oligocene hiatus association, Indian and Pacific Oceans, 713b–714b
 sediment composition effects, 506a
 sediment transport, from Antarctic shelf, 551a
 Antarctic Circumpolar Current, progressive polar cooling and, 124a
 Antarctic Convergence

Antarctic Surface Water and, 185b
 calcareous-siliceous transition, 7a, 124a
 movement through time, 124a
 Antarctic Divergence, Circumpolar-Coastal Current separation, 185b, 187b, 246b
 Antarctic Ice Sheet
 development, 290a, 303a–304a
 flow lines, 69b
 growth model, 873b
 initiation, 340a
 Antarctic shelf
 Prydz Bay: Site 739, reverse deepening, 336a
 sediment transport, by Antarctic Bottom Water, 551a
 Antarctica, 417b
 alluvial plains, Permian development, 797b
 biosiliceous sediments, latitudinal migration, 874b
 circum-Antarctic current, thermal isolation role, 201b, 246b
 continental shelf, progradation, Oligocene–Miocene, 115b, 122b
 deep waters
 high-latitude cooling, Eocene, 882b–883b
 Tethyan source, 713b
 deglaciation, Pliocene, 10a, 13a
 glaciation
 Cenozoic, 869b–875b
 climatic changes, 122b
 continental, 79b–80b, 693b–694b
 deep-sea record, 78b
 Eocene, 882b–883b
 ice sheet timing and initiation, 115b–118b
 isotopic evidence, 715b
 Oligocene, 122b, 693b–694b, 715b, 874b–875b
 Pliocene deglaciation event, 120b, 888b–889b
 pollen record, 873b–874b
 sea level, 122b
 sediment record, 870b–873b
 glaciomarine sediments, reworking process, 103b
 iceberg calving, Oligocene, 872b
 intermediate waters, Tethyan source, 713b
 paleogeographic reconstruction, 804b, 806b
 paleotemperature, Oligocene, 872b
 polar-wander paths, 767b
 preglacial conditions, 200b–201b
 surface water temperatures, Paleogene decline, 693b
 weathering change, Eocene/Oligocene boundary, 199b
 Antarctica, E
 alluvial plains, drainage and deposition, 53b
 geology, Lambert Glacier–Prydz Bay region, 137b
 glaciation
 Eocene, 875b
 onset, 201b
 stratigraphic record, 118b–120b
 ice erosion and sediment transport, 133b
 India separation, rift structures, 9a
 See also East Antarctic Craton; East Antarctic Ice Sheet
 Antarctica, W, glaciation, Paleocene–Eocene, 79b
 Antarctica-India separation, Kerguelen Plateau origin, 5a–6a
 Antarctica/Australia landmass, Eocene separation, low pressure centers, 873b

SUBJECT INDEX

Antimony, Kerguelen-Heard Plateau S: Site 738, Cretaceous/Tertiary boundary, 724b-725b
 Aragonite
 Prydz Bay: Site 739
 interstitial waters, 390b
 strontium isotopes, 741b, 745b
 Arsenic, Kerguelen-Heard Plateau S: Site 738, Cretaceous/Tertiary boundary, 724b-725b
 Aseismic ridges, subsidence curve, 280a, 282a
 Ash, basanitic, Kerguelen-Heard Plateau, Miocene–Quaternary, 326b, 333b
 Ash layers
 Kerguelen Sediment Ridge, 325b, 329b
 basaltic shards, 326b
 Kerguelen-Heard Plateau N, 325b, 326b, 328b-329b
 Ash, rhyolitic, Kerguelen-Heard Plateau, 326b, 328b
 Ash, volcanic
 Kerguelen Sediment Ridge: Site 745, 513a
 origin, 514a
 Kerguelen-Heard Plateau N, 130a, 168a
 Atlantic Ocean, Eocene cooling trend, 201b
 Atlantic Ocean S
 calcium compensation depth, Pliocene shoaling, 195b
 carbonate deposition, Eocene/Oligocene boundary, 714b
 deep waters, low-latitude origin, 694b, 713b
 intermediate waters, Antarctic origin, 694b
 lysocline, Pliocene shoaling, 195b
 Atterberg limits, Prydz Bay, 150b-151b, 152b
 Australian-Antarctic Basin, paleoceanographic reconstruction, 244b, 246b-247b
 Bacteria, living, Antarctic sediments, density changes, 687b-689b
 Baffin Bay, smectite-kaolinite correlation, 198b
 Bainmedart Coat Measures, Amery Group, 797b
 Barents Sea, slumping, 468a
 Barite
 Kerguelen Plateau S: Site 744, 216b-217b
 biogenic opal correlation, 218b
 Kerguelen-Heard Plateau S: Site 738, 216b-217b
 hydrothermal association, 217b
 Barium
 Kerguelen-Heard Plateau S: Site 738
 in altered basalts, 313b-314b
 comparison, other MORB-type basalts, 318b
 Cretaceous/Tertiary boundary, 725b
 Barium/niobium ratio, Kerguelen-Heard Plateau S: Site 738, 318b
 Barium/tantalum ratio, Kerguelen-Heard Plateau S: Site 738, basalts, 318b
 Basalt melting, Amery Ice Shelf, 62b, 64b
 Basalt
 Kerguelen-Heard Plateau N: Site 736, 130a
 Kerguelen-Heard Plateau S: Site 738
 alteration, 241a, 243a
 depositional environment, 279a
 petrology, 240a-241a
 secondary minerals, 364b
 Basalt, alkali, Kerguelen Sediment Ridge, dispersed ashes, 326b
 Basalt, aphyric, Kerguelen-Heard Plateau S: Site 738, plagioclase microphenocrysts, 295b
 Basalt, tholeiitic
 continental margins, niobium content, 294b
 Kerguelen-Heard Plateau S: Site 738, 302b-303b, 307b
 alteration, 307b, 313b-315b
 comparison, continental margin basalts, 296b
 continental flood, 318b

geochemistry, 294b, 312b
 immobile element concentrations, 317b
 isotopic data, 294b-295b
 mineralogy, 293b-294b
 petrography, 300b-301b
 plume-component bearing, 317b
 source, 296b
 subduction-related arc, 319b
 Basalt, trachytic, Kerguelen Sediment Ridge, dispersed ashes, 326b
 Basement
 Kerguelen Plateau
 pre-Turonian, 277a
 thickness, 7a
 Kerguelen-Heard Plateau N: Site 737, geochemistry, 187a-188a
 Kerguelen-Heard Plateau S: Site 738
 alteration, 314b, 315b
 geochemistry, 309b-311b
 igneous petrology, 240a
 structure, 300b
 Basement, metamorphic
 Prydz Bay
 acoustic Unit PS.5, 13b
 erosion with rift onset, 20b
 Precambrian age, 33b, 36b-37b
 Beacon Supergroup, Antarctica, correlation with Permian Amery Group, 795b
 Beaver Lake
 kaolinite, in glacial tills, 114b
 sedimentary strata, 65b-66b
 Phanerozoic, 7b
See also Amery Formation
 Bedrock, Prydz Bay, Precambrian igneous-metamorphic, 8a
 Biogenic sediments
 Kerguelen-Heard Plateau N, detrital components, 285b
 Prydz Bay, 112b-113b
 Southern Ocean, high-latitude cooling, 874b
 Biomimicry, Kerguelen-Heard Plateau S: Site 738, 238a
 Biotite, Prydz Bay: Site 740, petrography, 50b
 Biscuit structure, Kerguelen-Heard Plateau S: Site 738, 237a
 Bottom current
 Kerguelen Sediment Ridge: Site 745, sedimentation effects, 525a, 526a
 Kerguelen Sediment Ridge: Site 746
 detrital transport, 551a
 silt layers and, 542a
 Bottom water, formation, mechanisms, 232b
 Boulders, crystalline, Prydz Bay: Site 740, 93b, 125b
 Braided stream deposits
 Prydz Bay: Site 740
 fining-upward sequences, 47b, 53b
 red beds, 797b
 Bransfield Basin, siliceous surface sediments, 112b
 Bransfield Strait, organic matter, recycling, 417b
 Breccia
 Kerguelen-Heard Plateau S: Site 738
 clast types, 301b
 provenance, 307b
 water/rock ratio, 313b
 Breccia, amphibolite gneiss, Prydz Bay: Site 740, 383b
 Breccia, basaltic, Kerguelen-Heard Plateau S: Site 738, 238a-240a
 Breccia, igneous, Prydz Bay: Site 740, 383b
 Breccia, volcanic, Kerguelen-Heard Plateau S: Site 738, physical properties, 269a
 Broken Ridge

rifting from Kerguelen-Heard Plateau, 218a
 tectonic history, 5a
 Brunhes Chron
 Kerguelen Plateau S: Site 744, 490a, 782b, 820b
 Kerguelen Sediment Ridge: Site 745, 818b
 Brunhes/Matuyama Chron boundary, Kerguelen Sediment Ridge: Site 745, 752b
 Bunbury basalts, isotopic data, 296b
 Calcareous-siliceous transition
 Antarctic Convergence, 7a, 124a
 Kerguelen Plateau S: Site 744, 503a
 Kerguelen-Heard Plateau S: Site 738
 logging, 271a
 physical property changes, 269a
 Calcite
 Kerguelen-Heard Plateau S: Site 738
 in altered basalts, 314b-315b
 negative scandium correlation, 722b-723b
 Prydz Bay: Site 739, interstitial waters, 390b
 Calcite, geopetal, Kerguelen-Heard Plateau S: Site 738, 240a
 Calcite, micritic, Kerguelen-Heard Plateau N: Site 737, 357b
 Calcite, sparry, Kerguelen-Heard Plateau S: Site 738, 239a, 240a-241a
 Calcium
 high-resolution vs. regular sampling results, 397b-398b
 Kerguelen Plateau S: Site 744, 491a
 pore waters, 930b-931b
 vertical distribution gradient, 367b, 372b
 vs. magnesium, 493a
 Kerguelen Sediment Ridge, 516a-517a, 544a
 pore waters, 930b-931b
 vertical distribution gradient, 373b
 vs. magnesium, 522a
 Kerguelen-Heard Plateau N, 139a
 chemical gradient, 359b
 pore waters, 929b-930b
 sulfate and alkalinity correlation, 362b
 vertical distribution gradient, 363b, 372b
 Kerguelen-Heard Plateau N: Site 737, 185a-186a
 discontinuities, 187a
 vs. magnesium concentration, 193a
 Kerguelen-Heard Plateau S: Site 738, 255a
 in sheet silicate formation, 313b
 vertical distribution gradient, 366b, 372b
 vs. magnesium concentration, 259a
 Prydz Bay: Site 739, 312a
 pore waters, 380b, 930b-931b
 shallow-level source, 312a
 vertical distribution gradient, 380b-381b, 391b
 Prydz Bay: Site 740, 360a, 362a
 pore waters, 380b
 vertical distribution gradient, 388b, 391b
 Prydz Bay: Site 741, 385a
 pore waters, 380b
 vertical distribution gradient, 385b, 391b
 Prydz Bay: Site 742, 418a
 pore waters, 380b, 385b
 vertical distribution gradient, 386b, 391b
 vs. alkalinity, 385b
 vs. magnesium, 420a
 Prydz Bay: Site 743, 466a
 pore waters, 380b
 vertical distribution gradient, 391b
 Calcium carbonate
 Kerguelen Plateau S: Site 744, 367b, 898b-899b
 Kerguelen-Heard Plateau N: Site 736, 142a, 357b
 Kerguelen-Heard Plateau N: Site 737, 168a, 193a-194a, 196a, 359b

- alkalinity and, 186a
temperature and solubility of, 359b
- Kerguelen-Heard Plateau S: Site 738, 261a
364b
alkalinity and precipitation of, 257a
- Prydz Bay, 314a, 378b, 385b
deposition, 112b, 113b
site correlations, 113b
strontium release during saturation, 391b
vs. magnesium, 362a
- Calcium/magnesium ratio, Kerguelen Sediment Ridge, 371b, 373b
- Calcrete formation, Prydz Bay: Site 740, 50b
- California, Gulf of, barite dissolution, 217b
- Carbon, preservation, oxygen requirement, 401b
- Carbon, carbonate
Kerguelen Plateau S: Site 744, 493a-494a
Kerguelen Sediment Ridge, 522a, 548a
Kerguelen-Heard Plateau N: Site 736, 142a
Kerguelen-Heard Plateau S: Site 738, 259a-260a
Prydz Bay, 386a, 423a, 424a, 467a
- Carbon, combustible
Kerguelen Plateau S: Site 744, 114a-115a
Prydz Bay, 114a-115a
- Carbon-14 dating, Prydz Bay: Site 740, 748b
- Carbon dioxide, Kerguelen-Heard Plateau S: Site 738, in altered basalts, 313b
- Carbon, inorganic
Kerguelen Plateau S: Site 744, 491a, 493a-494a, 906b
Kerguelen Sediment Ridge, 522a, 545a, 548a, 906b
Kerguelen-Heard Plateau N, 142a, 189a, 193a-194a, 196a, 904b-905b
- Kerguelen-Heard Plateau S: Site 738, 259a-260a, 261a, 905b
Prydz Bay, 313a-314a, 363a, 385a, 386a, 422a, 423a, 905b-906b
- Carbon isotopes
Atlantic Ocean S vs. equatorial Pacific, Oligocene, 713b
Atlantic Ocean S vs. Pacific deep water, Miocene gradient, 713b
deep waters, vs. carbon dioxide record, 704b
- Eocene sediments, magnetostratigraphic correlation, 695b
- Holocene planktonic foraminifers, dissolved carbon dioxide disequilibrium, 697b
- Kerguelen Plateau S: Site 744, 696b, 698b-699b, 705b
climatic changes, Eocene/Oligocene boundary, 709b
enrichment, Eocene/Oligocene boundary, 704b
hiatus correlation, 936b
intersite comparison, 704b, 713b
Miocene, 935b-939b
Monterey Excursion, 937b
Oligocene enrichment, 709b-710b
paleoceanographic implications, 714b-715b
Paleocene/Eocene boundary, 697b, 704b stratigraphy, 935b
- Kerguelen-Heard Plateau S: Site 738, 700b-701b, 702b-703b, 705b
Cretaceous/Tertiary boundary, 859b
Eocene/Oligocene boundary, 709b
intersite comparison, 704b, 706b-707b, 713b
Oligocene enrichment, 709b-710b
paleoceanographic implications, 714b-715b
Paleocene/Eocene boundary, 697b, 704b, 706b surface to intermediate water gradients, 709b-710b, 714b
- planktonic and benthic foraminifers
interspecific differences, 697b
Paleogene decline, 695b
- systematic, 696b-697b
Prydz Bay, 408b, 410b, 742b
- Carbon, organic
Kerguelen Plateau S: Site 744, 190b, 355b, 367b, 906b
Kerguelen Sediment Ridge, 355b, 373b, 522a, 523a, 530a, 545a, 548a, 551a, 906b
glacial/interglacial cycles, 240b
- Kerguelen-Heard Plateau N: Site 736, 140a, 142a, 354b, 357b, 904b
- Kerguelen-Heard Plateau N: Site 737, 189a, 193a-194a, 195a, 354b, 359b, 904b-905b
abundance of and phosphate and ammonium content, 187b
sulfate concentration and, 186a-187a
- Kerguelen-Heard Plateau S: Site 738, 259a-260a, 354b-355b, 364b, 905b
- Prydz Bay
glaciation effects, 388b
sources, 113b
- Prydz Bay: Site 739, 81b-82b, 85b, 378b, 905b-906b
erosion of preglacial sediments and, 113b
recycled, 411b
terrestrial origin, 409b-410b, 411b
- Prydz Bay: Site 740, 374a-375a, 378b
- Prydz Bay: Site 741, 378b, 386a, 418b-419b
terrestrial origin, 410b, 412b
- Prydz Bay: Site 742, 81b-82b, 84b, 378b, 385b, 412a, 413a, 418b-419b, 423a, 424a
erosion of preglacial sediments and, 113b
magnetic susceptibility and, 113b
- Prydz Bay: Site 743, 378b, 467a
- Carbon, pyrolyzed, Prydz Bay: Site 741, 388a
- Carbon, total
Kerguelen Sediment Ridge, 522a, 548a
Kerguelen-Heard Plateau N: Site 736, 140a, 142a
Kerguelen-Heard Plateau N: Site 737, 193a-194a, 195a
Kerguelen-Heard Plateau S: Site 738, 257a, 259a-260a
- Prydz Bay: Site 739, 313a-314a
Prydz Bay: Site 740, 362a, 363a
Prydz Bay: Site 741, 385a, 386a
Prydz Bay: Site 742, 423a
- Carbon, total organic
Kerguelen-Heard Plateau N: Site 736, 140a
Kerguelen-Heard Plateau N: Site 737, 189a
Kerguelen-Heard Plateau S: Site 738, 257a
Prydz Bay: Site 739, 313a-314a, 408b
Prydz Bay: Site 740, 363a
Prydz Bay: Site 741, 385a-386a, 387a
Prydz Bay: Site 742, 422a, 424a
Prydz Bay: Site 743, 467a
- Carbonate
Kerguelen Plateau, ice-rafting and decrease in, 120b
- Kerguelen Plateau S: Site 744, 187b, 189b, 215b, 355b, 482a, 491a, 503a, 713b
Eocene/Oligocene boundary, 711b
ice-rafting association, 194b, 201b, 202b
preservation, 935b
- Kerguelen Sediment Ridge: Site 745, 227b, 355b, 519a
glacial/interglacial cycles, 240b
- Kerguelen Sediment Ridge: Site 746, 227b, 355b, 545a, 551a
- Kerguelen-Heard Plateau N: Site 736, 354b
- Kerguelen-Heard Plateau N: Site 737, 173a, 354b
diagenesis, 173a
vs. clinoptilolite content, 218b
- Kerguelen-Heard Plateau S: Site 738, 187b, 208b, 216b, 277b, 354b-355b
gray clay layer, 854b
ice-rafting association, 194b, 201b, 202b
recrystallization, 364b
vs. clinoptilolite content, 218b
oxygen isotopes, modern vs. Oligocene deep waters, 713b
- Prydz Bay
equilibria, thermodynamic modeling, 389b-391b
source, 113b
- Prydz Bay: Site 739, 82b, 85b
opal correlation, 113b
- Prydz Bay: Site 741, 386a, 395a
physical properties and, 392a
- Prydz Bay: Site 742, 82b, 84b, 422a
clotted matrix structure, 135b
- Carbonate compensation depth (CCD)
Atlantic Ocean, S, lowering, Eocene/Oligocene boundary, 714b
- Kerguelen Plateau S: Site 744, 194b
Pliocene shoaling, 195b
- Kerguelen Sediment Ridge, 230b, 818b
- Kerguelen-Heard Plateau N: Site 737, 357b
- Kerguelen-Heard Plateau S: Site 738, 194b
Pliocene shoaling, 195b
- Pacific Ocean, equatorial, lowering, Eocene/Oligocene boundary, 714b
- Carbonate critical depth (CCrD)
Kerguelen Plateau S: Site 744, Pliocene shoaling, 195b
- Kerguelen Sediment Ridge, 230b
- Kerguelen-Heard Plateau S: Site 738, Pliocene shoaling, 195b
- Carbonate, pedogenic, Prydz Bay: Site 740, lack of, 50b
- Carbonate-cemented layers, Prydz Bay: Site 739, 128b
- Carbonized plant protein
Prydz Bay: Site 741, 381a, 395a
physical properties, 391a
seismic stratigraphy, 393a
- Caribbean Core P6304, iridium accumulation rate, 855b
- Celadonite
Kerguelen-Heard Plateau S: Site 738, 301b
conditions of formation, 307b, 313b
- Cementation
Kerguelen-Heard Plateau N: Site 737, velocity and degree of, 197a
Prydz Bay: Site 739, physical properties, 326a
- Cementation, carbonate, Prydz Bay, 303a, 381a, 394a
- Cerium, Kerguelen-Heard Plateau S: Site 738, negative anomaly, 726b
- Cerium/lanthanum ratio, Kerguelen-Heard Plateau S: Site 738, Cretaceous/Tertiary boundary, 726b
- Cesium
Kerguelen-Heard Plateau S: Site 738
calcite-free data, 728b-729b
scandium-normalized distribution, 725b
- chalk
Kerguelen-Heard Plateau S: Site 738
iridium content, 854b
Maestrichtian sedimentation rate, 852b-853b
- Chalk, calcareous, Kerguelen-Heard Plateau S: Site 738, magnetic properties, 758b
- Chalk, nannofossil
Kerguelen Plateau N, sedimentation rate, 353b
- Kerguelen Plateau S: Site 744, chalk-to-ooze transition, 501a
- Kerguelen-Heard Plateau S: Site 738, 237a

SUBJECT INDEX

- chalk-to-ooze transition, 269a, 271a
- Channel sand bodies
Prydz Bay: Site 740
depositional environment, 47b-49b, 53b
original bedform identification, 49b
- Channel shifting, Prydz Bay, 47b-49b, 53b, 57b
- Charnockite, Kerguelen Sediment Ridge, Precambrian, 247b
- Charybdis Glacier, origin, 62b
- Chelnok glaciation, 749b
- Chert layers
Kerguelen-Heard Plateau N: Site 737, 171a, 207a
mineralogy, 214b-215b
- Kerguelen-Heard Plateau S: Site 738, 237a, 363b-364b
mineralogy, 215b-216b, 217b, 218b-219b
photoelectric value, 279b
physical properties, 266a
silica diagenesis, 372b
silica precipitation and, 254a-255a, 256a
- Chloride
Kerguelen Plateau S: Site 744, 491a
interstitial waters, 354b-355b
- Kerguelen Sediment Ridge, 516a, 544a
interstitial waters, 354b-355b
- Kerguelen-Heard Plateau N, 139a, 185a
interstitial waters, 354b
- Kerguelen-Heard Plateau S, 255a
- Prydz Bay, 310a-312a, 360a, 418a, 465a-466a
interstitial waters, 378b, 380b, 383b, 385b
- Chlorite
in high-latitude marine sediments, 198b
- Kerguelen Plateau S: Site 744, 194b, 199b
peak development, 199b
- Kerguelen Sediment Ridge, 244b
downslope transport, 231b
negative illite correlation, 247b
- Kerguelen-Heard Plateau S: Site 738, 194b
- Prydz Bay, 86b
clay matrix, 50b
sources, 113b-114b
- Chromium
Kerguelen-Heard Plateau S: Site 738
in basalt, 303b, 317b
Cretaceous/Tertiary boundary, 724b
scandium-normalized distribution, 724b
- Chromium/iridium ratio, Kerguelen-Heard Plateau S: Site 738, 724b
- Circumpolar Deep Water
deep-sea hiatus role, 201b, 874b
erosion, 478a
- Kerguelen-Heard Plateau, erosive periods, 230a
rafting, 478a
- Clasts
Kerguelen Plateau, 7a
lithology, 195b
- Kerguelen Sediment Ridge: Site 746, 539a, 550a
Antarctic basement source, 541a, 551a
- Kerguelen-Heard Plateau N: Site 736, 139a
- Kerguelen-Heard Plateau S: Site 738, lithology, 195b
- Prydz Bay
composition, 303a
grain size, 81b
metamorphic groups V-XV, 137b-138b
orientation, 107b, 299a, 300a
sedimentary groups I-IV, 134b-137b
shapes, 90b-91b, 97b, 98b
See also Gravel clasts
- Clasts, basaltic, Kerguelen-Heard Plateau S: Site 738, 238a, 301b
- Clasts, macrofossil, Kerguelen-Heard Plateau S: Site 738, 238a
- Clasts, manganese-oxide-coated, Kerguelen Plateau S: Site 744, 482a-483a
- Clasts, metamorphic
Kerguelen Sediment Ridge: Site 745, 508a
origin, 530a
- Clasts, sandstone, Kerguelen Plateau S: Site 744, Beacon Supergroup strata, 195b
- Clasts, tachylitic, Kerguelen-Heard Plateau S: Site 738, 301b
- Clay
Kerguelen Sediment Ridge: Site 745, origin, 530a
Prydz Bay, 385b
distribution, 149b
- Clay, diatomaceous
Kerguelen Sediment Ridge, 249b, 510a, 513a, 541a
facies intervals, 252b-253b
glacial/interglacial cycles, 226b-232b, 233b
ice-rafting maximums, 248b, 249b
origin, 514a
- Clay, iridium-rich
Kerguelen-Heard Plateau S: Site 738
calcite-free data, 728b-729b
formation, 727b
vs. clay-rich laminations, elemental ratios, 727b-728b
- Clay matrix, Prydz Bay: Site 740, mineralogy, 50b
- Clay mineralogy
East Antarctica, chemical weathering effects, 874b
- Kerguelen Plateau, correlation, Prydz Bay, 114b
- Kerguelen Plateau S: Site 744
chemical-to-physical weathering transition, 884b
as climatic indicator, 198b-200b
detrital components, 199b
Eocene, 199b, 882b
glaciation effects, Eocene, 201b
multiple source areas, 199b
Oligocene warming trend, 199b
- Kerguelen Sediment Ridge, 242b, 244b
facies, 228b-229b
source, 247b
- Kerguelen-Heard Plateau S: Site 738, 194b, 208b-209b
as climatic indicator, 198b-200b
Cretaceous/Tertiary boundary, 854b
detrital components, 199b
Eocene-Oligocene weathering effects, 199b, 201b, 882b
glaciation effects, Eocene, 201b
magnetic properties, 758b
multiple source areas, 199b
Oligocene warming trend, 199b
- Prydz Bay, 84b, 86b, 87b
correlation, Kerguelen Plateau, 114b
Pliocene compositional shift, 114b
sources, 113b-114b
- Clay, silty
Kerguelen Sediment Ridge: Site 746, 539a
Prydz Bay: Site 740
depositional environment, 749b
sedimentation rate, 749b
- Prydz Bay: Site 743, 462a
depositional environment, 463a
- Clay, terrigenous
Kerguelen Plateau S: Site 744, 197b, 209b-210b
- Kerguelen Sediment Ridge, 230b, 245b, 246b
mineralogy, 237b, 259b-260b
- Kerguelen-Heard Plateau S: Site 738, 196b, 208b-209b
- Claystone
Kerguelen-Heard Plateau S: Site 738, 168a
diagenetic greenish laminations, 170a
- Claystone, diatomaceous, Prydz Bay: Site 739, 300a
- Claystone, glauconite-bearing, Kerguelen Sediment Ridge: Site 745, 513a
- Claystone, glauconitic calcareous, Kerguelen-Heard Plateau N: Site 737, 159a
- Claystone, silty
Kerguelen-Heard Plateau N: Site 737, 168a
Prydz Bay: Site 742, 409a, 410a
depositional environment, 413a
- Claystone/clay layer, Western Interior of North America, geochemistry, 726b-727b
- Claystone/limestone boundary, Kerguelen-Heard Plateau N: Site 737, reflection peak, 214a
- Clinoptilolite
Kerguelen-Heard Plateau, 215b
precipitation, Eocene-Oligocene, 218b
- Clinopyroxene, Kerguelen-Heard Plateau S: Site 738, chemical composition, 304b, 317b
- Coal-bearing sediment
Prydz Bay: Site 741, 57b-60b, 836b-837b
Prydz Bay: Site 742, Permian Amery Group, 795b
- Coarsening-upward sequences
Prydz Bay: Site 740, 47b, 50b
sandstone, 45b
Prydz Bay: Site 741, relation to fining-upward sequences, 57b
- Cobalt
Kerguelen-Heard Plateau S: Site 738
anoxic depositional environment, 724b
Cretaceous/Tertiary boundary, 724b
scandium-normalized distribution, 724b
- Cobb Mountain Event
Kerguelen Sediment Ridge: Site 745, 753b
Kerguelen-Heard Plateau N: Site 737, 754b
- Cochiti Subchron, Kerguelen Sediment Ridge: Site 745, 753b, 818b
- Color banding
Kerguelen Sediment Ridge, 369b, 539a-540a
- Kerguelen-Heard Plateau N: Site 737, 170a-171a, 357b
- Prydz Bay: Site 740
diagenetic red and green, 46b
grain size variation with, 49b
- Conductivity-temperature-depth profile
Kerguelen-Heard Plateau S: Site 738, 109a
Prydz Bay, 109a
- Conglomerate, polymictic, Prydz Bay: Site 740, 351a
- Conglomerate, Prydz Bay: Site 741, 381a
- Consolidation tests
Prydz Bay, 172b, 175b
analytic methods, 170b-172b
diamictite, 174b, 884b
diamictons, 152b, 154b, 174b
siliciclastic rocks, 174b
See also Terzaghi consolidation theory
- Coral Sea, Oligocene hiatuses, 874b
- Creep rate, Prydz Bay, 176b, 179b
- Cretaceous/Tertiary boundary
Kerguelen-Heard Plateau S: Site 738, 724b, 724b-725b
carbon isotopes, 859b

- cerium/lanthanum ratio, 726b
 clay mineralogy, 854b
 claystone, 279a
 element stratigraphy, 719b-726b
 gray clay layer, 850b, 862b, 864b
 iridium, 720b, 722b, 723b, 851b, 854b-856b
 iron, 723b-724b
 laminations, clay-rich, 727b-728b
 magnetic properties, 253a, 757b-758b, 760b, 764b, 765b
 magnetobiochronology, 825b
 magnetostratigraphy, 757b-759b, 851b-853b
 oxygen isotopes, 859b
 planktonic foraminifers, 455b-460b
 sedimentation rate, 852b-853b
- Crevasses, Lambert Glacier-Amery Ice Shelf system, orientation, 71b
 Cross-bedding, planar, Prydz Bay: Site 740, 49b
 Cross-bedding, trough, Prydz Bay: Site 740, sandstones, 46b
 Crust, oceanic
 Kerguelen Plateau, 7a
 tectonic history, 5a-6a
 Kerguelen-Heard Plateau, 7a
- Debris flow, Kerguelen-Heard Plateau N: Site 736, deposition, 156a
 Deep-water circulation
 Antarctica
 glacial/interglacial cycles, 246b
 isotopic correlation, 712b-714b
 Kerguelen Sediment Ridge, sedimentary cycles, 232b
 Deformation
 Prydz Bay: Site 742, 413a
 soft-sediment, 411a-412a
 Density
 Kerguelen Plateau S: Site 744, 495a
 with silica variations, 499a
 Kerguelen Sediment Ridge: Site 745, 520a, 522a-524a
 Kerguelen Sediment Ridge: Site 746, 546a-547a
 Kerguelen-Heard Plateau N: Site 736, 148a, 153a
 grain-bulk correlation, 144a-145a
 Kerguelen-Heard Plateau N: Site 737
 discontinuities, 202a-203a
 synthetic seismogram, 212a
 Kerguelen-Heard Plateau S: Site 738, 262a, 264a, 266a
 scatter, 267a-268a
 Prydz Bay: Site 739, 145b, 148b, 319a, 325a
 Prydz Bay: Site 740, 365a, 369a
 Prydz Bay: Site 742, 146b, 428a, 434a
 logging vs. laboratory measurements, 435a
 Prydz Bay: Site 743, 146b, 148b, 469a-470a
 Density flow, Kerguelen-Heard Plateau N: Site 736, origin, 131a
 Density-porosity relationship
 Kerguelen-Heard Plateau N: Site 737, calcareous sequences, 272b, 275b
 Kerguelen-Heard Plateau S: Site 738, calcareous sequences, 277b, 281b
 Prydz Bay, glaciomarine sequences, 280b-281b, 285b
 Diagenesis
 Kerguelen Sediment Ridge: Site 745, 369b
 sedimentary structures, 513a
 Kerguelen-Heard Plateau N: Site 737, 173a, 357b
 Prydz Bay: Site 739, 303a
 physical property effects of, 326a
 Diagenetic front
 Kerguelen-Heard Plateau N: Site 737
- physical property correlation, 363b
 upward movement, 372b
- Diamictite
 definition, 88b
 grain size, meltwater activity and, 120b-121b
 Kerguelen-Heard Plateau S: Site 738, lithologic Unit V, 838b
 Prydz Bay
 intersite correlation, 442a
 McMurdo Sound sediments compared, 121b
 overconsolidation, 174b
 petrography, gravel claws, 452a-454a
 physical properties, 157b
 shear strength, soft vs. overconsolidated, 154b-155b
 silica content, 388b
 weathered source rocks, 876b
 Prydz Bay: Site 739, 290a, 295a, 296a, 299a, 300a-301a, 302a
 clast lithology, 302a
 clay-rich, 286b-287b
 correlation, Prydz Bay: Site 742, 442a
 depocenter, 884b
 distal to proximal glaciomarine transition, 107b
 gravel content, 81b
 massive, 301a, 302a
 mollusk shells, 739b
 Oligocene, 880b, 883b-884b, 888b
 organic matter, 411b
 physical properties, 147b-149b, 328a, 329a-330a
 Pliocene-Quaternary, 882b
 source, 74b
 stratified, 301a, 302a
 textural classification, 90b
 Prydz Bay: Site 740, 383b
 Prydz Bay: Site 741, mixing with preglacial sediments, 118b
 Prydz Bay: Site 742, 280b, 397a, 403a-405a, 410a-411a, 412a
 clay content, 281b
 density, 437a
 Eocene-Oligocene, 447a, 888b
 gravel content, 81b
 magnetic properties, 760b
 mixing with preglacial sediments, 118b
 physical properties, 147b-149b, 385b, 428a
 physical properties, changes with cementation, 434a
 Pliocene-Quaternary, 882b
 porosity and clay content, 285b
 reworking, 132b
 seismic profile, 443a, 444a
 source, 74b
 stratified, 408a
 textural classification, 90b
 velocity, 441a-442a
 Prydz Bay: Site 743, source, 74b
- Diamictite, massive
 Prydz Bay, 88b, 93b, 95b, 126b, 127b, 129b
 depositional environment, 285b
 genetic classification, 103b, 107b
 grain size, 90b
 gravel content, 127b, 128b
 velocity, 31b
- Diamictite, weakly stratified
 Prydz Bay, 88b-89b, 95b, 127b, 129b
 deposition, 107b
 mottling, 129b
- Diamictite, well-stratified
 Prydz Bay, 89b, 127b, 129b
 deposition, 107b
- Diamictite-siliceous ooze, interbedded, Prydz Bay: Site 740, Holocene, 882b
 Diamicton
 definition, 88b
 Prydz Bay, 88b, 296a, 403a, 462a
 flat-lying sequence, 107b
 glacial deposition, 463a
 physical properties, 147b-158b, 325a
 shear strength, 156b
 stratified, 303a
 Diamicton, massive
 Prydz Bay, 93b, 127b, 128b
 genetic classification, 103b
 gravel content, 126b
 Diamicton, unconsolidated, Prydz Bay, Pleistocene, 94b
 Diamicton, underconsolidated, Prydz Bay: Site 743, consolidation tests, 174b
 Diamicts
 definition, 88b
 formation, distal to proximal glaciomarine environment, 107b
 Prydz Bay
 grain fabrics, 91b-92b
 organic carbon content, 113b
 textural classification, 88b
 Diamicts, massive, Prydz Bay, as debris-flow deposits, 107b
 Diatomite, Prydz Bay, 89b, 130b, 405a, 409a, 412a
 Dike, basalt
 Kerguelen Plateau, classification, 24a, 44a
 Prydz Bay, classification, 24a, 44a
 Dike, clastic, Kerguelen-Heard Plateau N: Site 737, 170a, 171a, 173a
 Dome Argus, ice cover, thin areas, 64b
 Drake Passage, opening, 200b
 Dropstones
 Kerguelen Plateau S: Site 744, 479a
 Prydz Bay: Site 739, 301a
 in stratified mudstone, 89b, 107b, 127b, 130b
 Prydz Bay: Site 742, 409a, 410a, 413a
- East Antarctic craton, as clay mineral source, 198b
 East Antarctic Ice Sheet
 drainage, Lambert Glacier, 20b
 growth model, Pliocene-Pleistocene, 888b
 Oligocene, 201b
 Eh/pH conditions, Kerguelen-Heard Plateau S: Site 738, during alteration, 314b
 Elan Bank, geographic setting, 5a
 Electrical resistivity
 Kerguelen-Heard Plateau S: Site 738, 366b
 Prydz Bay, 379b-380b
 ease of diffusion and, 385b
- Eocene/Oligocene boundary
 Antarctica, 199b
 carbonate compensation depth (CCD), Pacific Ocean, equatorial, 714b
 Kerguelen Plateau S, ice rafting, 710b-711b
 Kerguelen Plateau S: Site 744
 carbon isotopes, 704b
 carbonate, 711b
 Kerguelen-Heard Plateau S: Site 738, carbon isotopes, 709b
 oxygen isotopes
 deep-sea benthic foraminifers, 694b
 Kerguelen Plateau S: Site 744, 704b, 708b, 710b, 711b-712b
 Kerguelen-Heard Plateau S: Site 738, 704b, 708b, 714b
 Prydz Bay, 10a
 Pacific Ocean, equatorial, carbonate deposition, 714b

SUBJECTINDEX

- Erosion
 Kerguelen Sediment Ridge: Site 745, 526a
 Prydz Bay, 462a-463a
 Oligocene–Miocene hiatus, 885b
- Erosion, glacial
 Prydz Bay
 deformable till bed scenario, 178b
 ice thickness and, 178b-179b
 Oligocene–Miocene unconformities, 118b
 overconsolidation effects, 156b, 173b-174b, 178b
 Pliocene style change, 114b
 sulfur and organic carbon changes with, 113b
- Ethane, Prydz Bay: Site 742, 420a, 422a, 423a
- Europium/lanthanum ratio, Kerguelen-Heard Plateau S: Site 738, Cretaceous/Tertiary boundary, 726b
- Eustatic sea level
 Kerguelen Plateau S: Site 744, Antarctic glacial expansion and, 885b
 Prydz Bay, overconsolidation effects, 174b-175b, 181b
 Ross Sea, Oligocene shallowing event, 872b
- Extractable organic matter
 Prydz Bay
 analytical methods, 407b
 gas chromatograms, 412b, 413b, 414b
 maturity, 408b-411b, 412b
 thermal histories, 409b
- Extrusive
 Kerguelen Plateau, classification, 24a, 44a
 Prydz Bay, classification, 24a, 44a
- Faeroe Islands, tholeiitic basalt, Cretaceous niobium spike, 294b
- Falkland Plateau, microfossil assemblages, Eocene–Oligocene diversity decrease, 874b
- Feldspar
 Kerguelen Plateau, northern vs. southern plateau, 217b
 Kerguelen Plateau S: Site 744, 216b
 quartz correlation, 217b
 Kerguelen Sediment Ridge: Site 745, 237b, 510a
 origin, 514a
 Kerguelen-Heard Plateau S: Site 738, 216b
 primary and secondary, 302b-303b
 Prydz Bay: Site 739
 alteration to smectite, 391b
 fluvial reworking, 115b
 Prydz Bay: Site 740, 46b
 orthoclase, 52b
 petrography, 50b
- Ferrar Fjord, glaciogenic sediments, Pliocene, 119b
- Ferrar Glacier, advances and retreats, Pliocene-Quaternary, 79b
- Fining-upward sequences
 Prydz Bay: Site 740
 braided river systems and, 47b, 53b
 claystones, 46b
 mechanism, 52b-53b
 sandstone, 45b, 48b
 siltstones, 46b
 Prydz Bay: Site 741, relation to coarsening-upward sequences, 57b
- Fisher Glacier
 longitudinal foliation, 75b
 surging, 62b, 73b
See also Lambert Glacier
- Flagstone Bench Formation, 797b
 Prydz Bay red bed correlation, 54b
- Flood plain deposits, Prydz Bay: Site 740, 49b-50b, 53b
- Fluvial sediments, Prydz Bay: Site 741, 376b
- Fluvio-lacustrine sediments, Prydz Bay: Site 741, lithology, 59b, 60b
- Four Ladies Bank
 depth, 80b
 development, ice-shelf stability and, 115b
 glacial tills, 290a
 topography, 64b
- Fractures, Kerguelen-Heard Plateau N: Site 737, 170a
- Fram Bank, development, ice-shelf stability and, 115b
- Freeze-on ice
 Amery Ice Shelf, 64b, 73b
 Kerguelen Sediment Ridge, 247b
- Gamburtsev Subglacial Mountains
 glaciation, 74b, 118b
 interior basin, drainage, 80b
 topography, 64b
- Garnet
 Prydz Bay: Site 740, 56b
 petrography, 51b, 52b
- Gauss Chron
 Kerguelen Plateau S: Site 744, 782b, 820b
 Kerguelen Sediment Ridge: Site 745, 753b
- Gaussberg (Antarctica), leucites, correlation, Prydz Bay basalts, 295b, 296b
- Geochemistry
 Kerguelen Islands, tephra vs. whole-rock, 335b
 Kerguelen Plateau, 926b-927b, 928b
 Kerguelen-Heard Plateau S: Site 738, 301b-302b
 neutron activation analysis, 721b
 with water as alteration index, 307b
 with zirconium as differentiation index, 303b, 313b
 Prydz Bay: Site 739, 927b-928b
- Geochemistry, inorganic, 156a
 diagenetic features and, 254a-255a
 Kerguelen Plateau S: Site 744, 35a, 490a-491a, 493a
 Kerguelen Sediment Ridge, 35a, 516a-518a, 544a-545a
 Kerguelen-Heard Plateau N, 35a, 137a-140a, 185a-188a, 219a
 Kerguelen-Heard Plateau S: Site 738, 35a, 254a-257a
 Prydz Bay: Site 739, 35a, 310a-313a
 Prydz Bay: Site 740, 35a, 53b, 360a-362a, 374a
 low-salinity water source, 360a
 Prydz Bay: Site 741, 35a, 385a, 395a
 Prydz Bay: Site 742, 35a, 415a, 418a-420a
 Prydz Bay: Site 743, 35a, 465a-466a, 475a
- Geochemistry, organic
 Kerguelen Plateau S: Site 744, 34a-35a, 491a
 Kerguelen Sediment Ridge, 34a-35a
 518a-519a, 545a
 Kerguelen-Heard Plateau N, 34a-35a, 140a, 156a, 188a-191a, 219a
 Kerguelen-Heard Plateau S: Site 738, 34a-35a, 257a
 Prydz Bay: Site 739, 34a-35a, 313a
 Prydz Bay: Site 740, 34a-35a, 362a, 374a-375a
 Prydz Bay: Site 741, 34a-35a, 385a-387a
 Prydz Bay: Site 742, 34a-35a, 420a-422a, 838b
 Prydz Bay: Site 743, 34a-35a, 466a, 475a
- Rock-Eval pyrolysis
 Kerguelen Sediment Ridge: Site 745, 519a, 523a, 524a
 Kerguelen-Heard Plateau N: Site 736, 140a-143a
 Kerguelen-Heard Plateau N: Site 737, 189a-190a, 196a, 197a, 198a
 Prydz Bay: Site 739, 409b
- Prydz Bay: Site 741, 386a-387a, 409b
 type III kerogen, 386a-387a, 395a
 Prydz Bay: Site 742, 424a-425a
 type III kerogen, 113b, 422a
 Prydz Bay: Site 743, 466a, 467a, 468a
 type III kerogen, 113b
- George V Coast
 Beacon Supergroup, kaolinitic sediments, 247b
 siliceous surface sediments, 112b
- George V Land, siltstone, Cretaceous, 60b
- Gilbert Chron
 Kerguelen Sediment Ridge: Site 745, 818b
 Kerguelen Sediment Ridge: Site 746, 544a
 Kerguelen-Heard Plateau N: Site 737, 754b
- Gilsa Subchron
 Kerguelen Sediment Ridge: Site 745, 753b
 Kerguelen-Heard Plateau N: Site 737, 757b
- Glacial debris
 Kerguelen Plateau S: Site 744, in oxygen isotope enrichment zone, 712b
 Lambert Glacier-Amery Ice Shelf system surging effects, 71b, 73b
 transport paths, 71b
- Prydz Bay
 sources, 73b-74b
 transport, 115b
- Glacial sediments
 Prydz Bay, 881b, 885b
 vertical solute diffusion, 387b-388b
- Prydz Bay: Site 739
 Neogene warming trend, 886b
 physical properties, 326a
- Glacial stratigraphy, Prydz Bay, seismically defined units, 92b-101b
- Glacial/interglacial cycles
 Kerguelen Sediment Ridge, 226b-232b, 240b-244b, 551a
 downslope transport, 230b-231b, 232b
 productivity, 230b
- Norwegian Sea, 514a
- Prydz Bay, 118b, 122b
 physical properties, 326a
 prograding sequences, 19b
- Glaciation
 Antarctica
 continental shelf and terrestrial record, 79b-80b
 deep-sea record, 78b
 ice build-up, Eocene/Oligocene boundary, 78b
 onset unconformity, 22b
 Prydz Bay-Kerguelen Plateau hiatuses with Oligocene expansion, 885b
 sea level, Cretaceous, 200b
 terrestrial record, 10a
- Kerguelen Plateau S
 continental vs. mountain, 201b
 Eocene, 882b
 ice rafting with intensification of, 202b
 Oligocene, 883b-884b
 onset, 120b
- Kerguelen Sediment Ridge, 247b-249b
- Kerguelen-Heard Plateau N
 erosion and gravel transport, 131a
 sea level changes with, 216a
- Kerguelen-Heard Plateau S, 201b-202b, 248b-249b
 continental vs. mountain, 201b
 ice rafting with intensification of, 202b
 onset, 120b
- Paleogene, Prydz Bay and Kerguelen Plateau S compared, 885b
- Prydz Bay, 79b-80b, 180b
 advances and retreats, 118b, 875b
 biogeochemical effects of, 388b

- continental shelf development and, stages of, 116b-118b
depositional record, 340a
Eocene, 882b-883b
Holocene readvance, 749b
ice expansion events, 99b
marine conditions and onset of, 397a
Miocene–Pliocene transition, 248b-249b
Miocene–Quaternary advances and retreats, 122b
Neogene, 885b-889b
Oligocene, 883b-885b
Oligocene–Miocene hiatus, 268b
onset, 290a
Pliocene recession, 119b
Glaciation, continental, Oligocene, oxygen isotope record, 201b
Glacier, tidewater, ground-line positions, 339a
Glaciogenic deposition, Prydz Bay, front geometry, 19b
Glaciogenic sediment
genetic classification, 101b, 103b
Prydz Bay
consolidation, 151b-152b
distribution mode, 14b, 16b
erosion and overconsolidation of, 178b-179b
by lithofacies types, 104b, 105b, 107b
permeability vs. preconsolidation stress, 155b
plasticity, 154b
source area, 14b, 16b
transport and depositional environment, 114b-115b
Glaciomarine deposition, continental shelves, high-latitude, 80b
Glaciomarine sediment
Kerguelen Sediment Ridge, facies, 240b
Prydz Bay
erosional unconformity, 797b
Miocene deposition, 118b
pebbly muds, 749b
proximal vs. distal, 103b, 107b, 248b-249b, 882b, 888b
Glaciotectonic deformation, Prydz Bay, facies, 112b
Glass
Kerguelen Plateau, textures, 325b
Kerguelen Sediment Ridge
basalt-andesitic to andesitic, 328b-329b
geochemistry, 337b-343b
Kerguelen-Heard Plateau N, 218b
alteration, 344b
shard morphology, 344b
Glaucousites, Kerguelen-Heard Plateau N: Site 737, 165a, 167a, 173a
Glaucous minerals, Kerguelen-Heard Plateau S: Site 738, laminated intervals, 724b
Gneiss
Prydz Bay
extension of Proterozoic and Archaean, 53b-54b
provenance, 52b
Gneiss, biotite-garnet-orthopyroxene, Prydz Bay, 453a
Gneiss, biotite-quartz-feldspar, Prydz Bay, 247b
Gneiss, charnockitic granite, Prydz Bay, 138b
Gneiss, clinopyroxene-amphibole-garnet, Prydz Bay, 453a
Gneiss, cordierite-sillimanite, Prydz Bay, 454a
Gneiss, garnet-bearing quartzofeldspathic, Prydz Bay, 453a-454a
Gneiss, granite, Prydz Bay, 138b
Gneiss, mafic, Prydz Bay, 138b, 453a
Gneiss, orthopyroxene, Prydz Bay, 453a
Gneiss, quartzofeldspathic, Prydz Bay, 454a
Godovari rift graben, deposition, 21b
Godovari Valley, 9a
Gondwana, rift graben systems, 20b
Gondwana sequence, 9a
Gondwanaland
paleogeographic reconstruction, 804b, 806b, 807b
red beds, Devonian through Triassic, 795b
Grain orientation, Prydz Bay, 99b, 100b, 105b, 107b
Grain size
Kerguelen Plateau S, 195b
nonbiogenic sediment, 191b, 193b, 207b-208b
Kerguelen Sediment Ridge
facies A and B compared, 227b-228b
tephra layers, 328b
terrigenous component, 236b, 240b, 242b, 243b, 244b
Kerguelen-Heard Plateau N, tephra layers, 328b
Kerguelen-Heard Plateau S: Site 738, nonbiogenic sediment, 190b-191b, 206b-207b
Prydz Bay, 81b, 405a
detrital quartz and, fluvial environments, 50b, 52b
diamictite, 296a, 299a
by facies type, 89b-90b, 91b, 94b
permeability and, 154b
physical properties and, 149b-150b, 156b-157b, 392a, 431a, 434a
porosity and, 282b, 385b
sorting, 90b
source of clay minerals and, 114b
transport distance and, 121b, 388b
Granite/gneiss rock, Prydz Bay: Site 740, in sandstones, 50b
Granulite, Prydz Bay, 453a-454a
GRAPE density
Kerguelen Plateau S: Site 744, 37a, 495a
Kerguelen Sediment Ridge, 37a, 520a, 522a-524a
Kerguelen-Heard Plateau N, 37a, 145a, 202a
Kerguelen-Heard Plateau S: Site 738, 37a, 262a, 264a, 265a, 267a
Prydz Bay, 37a, 319a, 365a, 369a
Gravel
Kerguelen Plateau, origin, 131a
Kerguelen Plateau S: Site 744, grain size distribution, 191b, 192b
Kerguelen Sediment Ridge
ice rafting association, 231b, 248b, 249b
sea level control, 249b
Kerguelen-Heard Plateau N: Site 736, temperature, 152a
Kerguelen-Heard Plateau S: Site 738
grain size distribution, 190b
ice rafted origin, 201b
Prydz Bay, 405a
composition, 299a, 300a
distribution, 149b
grain size variation, 81b, 82b, 83b
Gravel clasts
Prydz Bay
shape, 90b-91b
source of, 114b
weathering rinds, 114b
Gravity flow, Prydz Bay, diamictites, 112b
Gray clay layer
Kerguelen-Heard Plateau S: Site 738
Cretaceous/Tertiary boundary, 850b, 862b, 864b
iridium content, 854b
magnetic properties, 852b
mineralogy, 854b
sedimentation rate, 852b-853b
SEM analysis, 857b
stable isotopes, 859b
Green clay layer, Kerguelen-Heard Plateau S: Site 738, microfossil remains, 866b, 867b
Greenschist, Prydz Bay: Site 739, 453a
Grove Mountains
alpine glaciation, initiation, 74b
ice cover, 64b
Hafnium
Kerguelen-Heard Plateau S: Site 738
calcite-free data, 728b-729b
scandium-normalized distribution, 725b, 726b
Halo structures
Kerguelen Plateau S: Site 744, 484a, 771b-772b
Kerguelen Sediment Ridge: Site 745, 510a
Heat flow
Kerguelen Plateau S: Site 744, 499a
Kerguelen-Heard Plateau N: Site 736, 153a, 156a
Kerguelen-Heard Plateau N: Site 737, 198a
Hopane/moretane ratio, Prydz Bay: Site 739, 408b
Hydrocarbon gases
Kerguelen Plateau S: Site 744, 34a, 491a
Kerguelen Sediment Ridge, 34a, 519a, 545a
Kerguelen-Heard Plateau, 34a, 140a, 189a, 257a, 259a
Prydz Bay, 34a, 313a, 363a, 385a-386a, 420a-421a, 466a
Hydrocarbons, aliphatic
Prydz Bay, 411b, 412b, 413b, 415b, 418b
gas chromatograms, 420b
odd/even carbon number predominance, 419b
Hydrocarbons, aromatic, Prydz Bay: Site 739, 409b
Hydrocarbons, steroid
Prydz Bay
mass fragmentograms, 422b
source and maturity, 422b-423b
Tm/Ts and C₃₁hopane epimer ratios, 419b
Hydrocarbons, triterpenoid
Prydz Bay
mass fragmentograms, 421b
maturity level, 419b, 422b
Tm/Ts ratio, 419b
Hydrogen index
Kerguelen-Heard Plateau N, 144a, 145a, 198a, 199a
Prydz Bay, 388a, 408b, 410b, 426a, 469a
Hydrogen isotopes
Prydz Bay, 377b
depletion in meteoric waters, 388b
Prydz Bay: Site 739, weathering effects, 381b, 383b
Prydz Bay: Site 740
Antarctic Bottom Water compared, 383b
interstitial waters, 384b
Prydz Bay: Site 742
below sediment/water interface, 385b
pore waters, 386b-387b
Ice flow, Prydz Bay, source, 71b, 115b
Ice grounding
depositional model, 19b
Prydz Bay
Oligocene expansion, 180b
physical property effects, 157b
position fluctuation, 115b
sediment transport, 16b
stability, 115b
Prydz Bay: Site 739
line fluctuations, 326a
seismic reflection profiling, 19b-20b

SUBJECT INDEX

- Prydz Bay: Site 742, 446a, 448a
initiation, 452a
sediment mixing from, 412a
Prydz Bay: Site 743, 465a
Ice loading
Prydz Bay, 378a, 412a
event chronology, 176b, 178b, 179b-181b,
884b-886b
intersite differences, 179b
overconsolidation effects, 156b, 157b, 175b-
176b, 178b
physical property changes, 339a, 368a, 431a
Pliocene ice sheet growth and, 886b
unconformities and, 179b-181b
velocity indicators, 36b
Ice rafting
Antarctica, Oligocene, 78b
Kerguelen Plateau, Oligocene-Holocene
sediment record, 217b
Kerguelen Plateau S
comparison, Southern Indian Ocean, 120b
glacial events and intensity of, 248b
Miocene-Holocene maximum, 120b
Oligocene decrease, 201b, 202b
temporal variation, 195b, 197b-198b
Kerguelen Plateau S: Site 744, isotopic
correlation, Eocene/Oligocene boundary,
710b-711b
Kerguelen Sediment Ridge, 505a, 508a, 514a,
530a, 541a, 550a
glacial/interglacial cycles, 231b, 234b
grounding line limitations, 232b
iceberg density and, 247b
with sea level changes, 249b
Kerguelen-Heard Plateau N: Site 736, 123a,
131a
Kerguelen-Heard Plateau N: Site 737, 172a
Kerguelen-Heard Plateau S: Site 738
comparison, Southern Indian Ocean, 120b
glacial events and intensity of, 248b
Miocene-Holocene maximum, 120b
Oligocene decrease, 201b, 202b
Prydz Bay, 450a-451a
influence on biogenic diatom ooze, 381a
sea level control, 249b
Ice sheet
Pleistocene, meltwater association, 121b
Prydz Bay
past thickness profiles, 178b
thermal regime, 120b-121b
Ice Shelf Water, formation, 232b
Ice volume, deep waters, oxygen isotopic evi-
dence, 695b, 715b
Ice-raftered debris
Antarctica, Cenozoic record, 873b
Kerguelen Plateau S: Site 744
Antarctic glacial expansion and, 885b
clay mineralogy with, 875b
magnetic anomaly 13, 101b
negative carbonate correlation, 194b
Neogene, 886b
Oligocene, 198b, 884b
origin, 195b
Kerguelen Sediment Ridge
clean vs. dirty snow, 247b
grain-size distribution, 231b-232b
ice shelf fluctuation correlation, 248b
Neogene, 886b, 887b
origin, 247b
transport, 247b
Kerguelen-Heard Plateau S: Site 738, 198b
Eocene, 201b, 882b
negative carbonate correlation, 194b
Oligocene, 884b
origin, 195b
ice-rafted detritus, Kerguelen Plateau S: Site 744,
502a-503a
Iceberg calving
Antarctica
isotopic correlation, 712b, 714b
sea level, 873b
ice streams and outlet glaciers
debris transport, 247b
vs. ice shelf calving, 248b, 249b
Prydz Bay, sea level rise and increase in, 249b
Igneous petrology, Kerguelen-Heard Plateau S:
Site 738, 234a-244a
Igneous rock, alkaline mafic, Prydz Bay, Creta-
ceous-Eocene, 1b
Illite
in high-latitude marine sediments, 198b
Kerguelen Plateau S: Site 744
negative smectite correlation, 194b, 199b
peak development, 199b
Kerguelen Sediment Ridge, 244b
downslope transport, 231b
negative chlorite correlation, 247b
vs. smectite, 249b
Kerguelen-Heard Plateau S: Site 738, negative
smectite correlation, 194b
Prydz Bay, 86b
potassium decrease and formation of, 391b
sources, 113b-114b
Ilmenite, Prydz Bay: Site 740, in sandstones, 51b,
56b
Impact-ejecta fallout layer, Kerguelen-Heard Pla-
teau S: Site 738, iridium anomaly association,
727b-728b
Indian Ocean
Eocene cooling trend, 201b
intermediate waters, Antarctic origin, 694b,
704b, 707b
Inertinite, Prydz Bay: Site 739, 408b
Intermediate waters
Antarctic-Australian opening, isotopic
correlation, 713b
Indian Ocean, 694b, 704b, 707b
Interstitial-water chemistry
charge imbalance, 350b
contamination sources, 350b-351b, 353b-354b
high-resolution sampling technique, 348b, 372b,
376b, 393b-399b, 903b
Kerguelen Plateau N, alteration effects, 372b
Kerguelen Plateau S, solute distribution, 368b
Kerguelen Plateau S: Site 744, 367b-369b,
490a-491a, 493a
Kerguelen Sediment Ridge, 369b-371b,
516a-518a, 520a, 521a
diffusion coefficient variations, 373b
solute distribution, 370b
Kerguelen-Heard Plateau N, 137a-140a, 141a,
185a-188a, 191a, 192a
biogeochemical processes, 357b, 359b
chemical gradients, 359b, 361b
diffusion coefficient variations, 362b-363b
inorganic processes, 359b, 362b-363b
solute distribution, 357b, 358b, 360b
Kerguelen-Heard Plateau S: Site 738,
254a-257a, 258a, 279a-280a, 364b-367b
diffusion coefficient variations, 366b
dissolved vs. dissolved magnesium concentra-
tion, 255a
solute distribution, 365b
major cation concentrations, 351b
Prydz Bay
biogeochemical processes, 359b
charge imbalance, 378b
Fick's law relationship, 362b
origin data, 379b
solute distribution, 382b, 384b, 387b
weathering effects, 391b
Prydz Bay: Site 739, 310a-313a, 380b-383b
Prydz Bay: Site 740, 358a, 360a-362a
ocean-terrestrial ground water mixing, 383b
Prydz Bay: Site 741, 384b-385b, 385a, 386a
bacterial activity below sediment/water inter-
face, 384b
Prydz Bay: Site 742, 385b, 415a, 418a-420a,
451a
Prydz Bay: Site 743, 385b-387b, 465a-466a
temperature change effects, 351b-352b
Intraclasts, claystone
Prydz Bay: Site 740, 46b, 48b
floodplain deposition, 50b
Intraclasts, siltstone
Prydz Bay: Site 740, 46b
floodplain deposition, 50b
Intrusions, igneous
Lambert Glacier-Amery Ice Shelf system, 65b
Prydz Bay: Site 740, 54b
Iridium
in the biosphere, 856b
in the hydrosphere, 855b-856b
Kerguelen-Heard Plateau S: Site 738
accumulation rates, 855b
analytical methods, 719b-720b
anomaly, calcite carbon isotopes and, 725b
anomaly, origin, 726b-729b
Cretaceous/Tertiary boundary, 720b, 722b,
723b, 851b, 854b-856b
enrichment event, 858b-859b
magnetic susceptibility, 758b
red clay vs. carbonate environment, 855b
scandium-normalized distribution, 720b,
722b, 723b
Iron
Kerguelen-Heard Plateau N: Site 736, 356b
Kerguelen-Heard Plateau S: Site 738
Cretaceous/Tertiary boundary, 723b-724b
excess iron/excess iridium ratio, 724b
scandium-normalized distribution, 724b
in sheet silicate formation, 313b
Iron, banded
Prydz Bay, 138b
microfolded formations, 453a
provenance, 139b
James Ross Island, oxygen isotopes, Cretaceous,
200b
Jaramillo Subchron
Kerguelen Plateau S: Site 744, 820b
Kerguelen Sediment Ridge: Site 745, 752b
Kaena Subchron
Kerguelen Plateau S: Site 744, 820b
Kerguelen Sediment Ridge: Site 745, 753b
Kaolinite
climatic variation, 114b, 198b
Kerguelen Plateau S: Site 744, 194b
detrital source, 198b
peak development, 199b
Kerguelen Sediment Ridge, 244b
source, 247b
Kerguelen-Heard Plateau S: Site 738, 194b
detrital source, 198b
Prydz Bay, 86b, 198b
Eocene, 882b
Kerguelen Islands, petrology, 325b
Kerguelen Plateau
arching, Maestrichtian-Campanian
unconformity and, 280a
bathymetry, 6a, 349b

- geographic setting, 5a
 geological setting, 7a, 353b
 ice rafting, Oligocene, 78b
 magnetic properties, seafloor-spreading anomalies, 5a
 magnetobiochronology, 817b-836b
 Miocene–Eocene hiatus, 7a
 Neogene unconformity, 7a
 Oligocene–Neogene hiatuses, climatic deterioration and, 120b
 tectonic history, 5a
Kerguelen Plateau S
 crustal structure, 7a
 volcano-tectonic evolution, 299b
Kerguelen Plateau S: Site 744
 bathymetry, 5a, 45a
 biology and oceanography, 35a-37a, 493a-495a
 biostratigraphy, 484a-488a, 824b, 825b
 benthic foraminifers, Eocene–Holocene, 614b-615b
 calcareous nannofossils, 474b-476b
 datum levels, 826b-828b
 diatoms, 559b-573b
 Neogene events, 829b
 paleogene events, 830b
 planktonic foraminifers, Paleogene–Neogene, 435b-437b
 radiolarians, 519b
 silicoflagellate, 933b-934b
 zonation, 24a, 26a-33a
 calcareous-biosiliceous transition, 820b
 Circumpolar Deep Water, erosional hiatuses associated with, 201b
 clay mineralogy, 194b
 depositional environment, 477a, 484a evolution, 217b-219b
 drifting sediment trap deployment, 111a
 drilling deformation, 15a, 480a
 geochemistry, organic, 34a-35a, 491a
 geophysics, 478a
 glaciation
 Neogene, 885b-888b
 Oligocene–Holocene, 201b-202b
 Paleogene, 882b-885b
 lithology, 187b, 215b
 lithostratigraphy, 479a-484a
 biogenic vs. nonbiogenic components, 482a
 coarse fraction source, 484a
 diatom-nannofossil ooze transition, 367b, 888b
 lithologic Unit I, 213b, 480a, 482a, 502a, 771b
 lithologic Unit I/Unit II contact, 820b
 lithologic Unit II, 213b, 482a, 503a, 771b-772b
 magnetobiostratigraphic correlation, 696b
 nannofossil-diatom transition, 202b
 nannofossil-diatomaceous ooze transition, 477a-478a, 484a, 503a
 seismic stratigraphic correlation, 501a
 summary log, 481a
 living bacteria, density changes, 687b-689b
 location, 144b, 169b, 186b, 212b, 300b, 324b, 350b, 376b, 477a, 694b, 714b, 731b, 751b, 814b
 logging, magnetic susceptibility, 895b
 magnetic properties, 33a-34a, 45a, 488a-490a, 772b-782b
 magnetobiochronology, 819b-822b
 magnetostratigraphy
 biostratigraphic correlation, 781b-782b
 reversal sequence, 772b, 781b
 mineralogy, 215b-217b
 Miocene unconformity, 477a, 478a, 502a
 nannofossil-diatom ooze transition, 190b
 navigation, 45a, 47a, 48a-85a, 87a, 89a
 Oligocene unconformity, 477a, 478a, 502a, 885b
 Oligocene–Miocene hiatus, 736b, 782b
 particulate fluxes, 109a-110a, 114a
 physical properties, 37a-39a, 495a-500a
 consolidation and, 151b-152b
 preglacial conditions, 200b-201b
 sediment composition, 188b, 204b-205b
 environmental implications, 194b-200b
 nonbiogenic sediment, 190b-193b
 sedimentary structures, 20a
 sedimentation rate, 490a
 seismic stratigraphy, 45a, 102a, 500a-501a
 digital magnetic tape parameters, 106a
 system geometry, 106a
 stable isotopes
 Eocene–Miocene, 708b-710b
 Eocene/Oligocene transition, 710b-712b
 paleoceanographic implications, 712b-714b
 Paleocene–Eocene, 697b-708b
 stratigraphy, 935b-939b
 strontium isotopes, mixed planktonic foraminifers, 731b-736b
Kerguelen Ridge E
 morphology, 225b
 sediment drift, Antarctic Bottom Water responsibility, 232b
Kerguelen Sediment Ridge
 ash layers
 geochemistry, 325b
 petrography, 325b
 bathymetry, 5a, 45a
 biology and oceanography, 35a-37a, 519a-520a
 clasts, 257b
 glaciation
 Miocene, 248b
 Miocene–Pliocene transition, 248b-249b
 Neogene, 249b, 885b-888b
 Paleogene, 882b-885b
 sedimentary record, 240b-244b
 ice rafting, origin, 247b
 interstitial-water chemistry, 369b-371b
 high-resolution vs. regular sampling method, 393b-399b
 lithostratigraphy, 327b, 369b, 509a-514a
 location, 144b, 169b, 226b, 239b, 300b, 324b, 350b, 376b, 505a, 751b, 814b
 navigation, 45a, 47a, 48a-85a, 89a
 paleoceanographic reconstruction, 244b, 246b-247b
 sediment composition, 235b, 254b-256b
 sedimentary cycles
 grain size and composition, 226b-229b
 origin, 229b-232b
 sedimentary structures, 20a
 seismic stratigraphy, 45a, 103a-104a, 525a-526a, 548a
 system geometry, 106a
Kerguelen Sediment Ridge: Site 745
 biostratigraphy, 514a-516a, 818b
 diatoms, 573b-575b
 radiolarians, 519b-526b
 zonation, 24a, 26a-33a
 depositional environment, 513a-514a
 drilling deformation, 15a, 508a-509a
 geochemistry, inorganic, 35a, 516a-518a
 geochemistry, organic, 34a-35a, 518a-519a
 geophysics, 506a
 lithostratigraphy
 lithologic Unit I, 505a, 509a-510a, 512a, 530a
 mixed pelagic-terrigenous component, 505a, 508a
 seismic stratigraphic correlation, 525a
 summary, 511a
 logging, magnetic susceptibility, 895b, 896b-897b, 898b, 900b-901b
 magnetic properties, 33a-34a, 45a, 516a
 magnetobiochronology, 817b-818b
 magnetostratigraphy, 752b-753b
 physical properties, 37a-39a, 520a, 522a-525a
 sedimentation rate, 516a
Kerguelen Sediment Ridge: Site 746
 biostratigraphy, 542a-544a, 821b
 calcareous nannofossils, 476b
 diatoms, 575b-577b
 radiolarians, 526b-527b
 zonation, 26a-33a
 depositional environment, 541a-542a
 depression, sediment transport role, 542a, 551a
 drilling deformation, 15a, 539a
 geochemistry, inorganic, 35a, 544a-545a
 geochemistry, organic, 34a-35a, 545a
 geophysics, 537a-538a
 interstitial-water chemistry, 369b-371b
 lithostratigraphy
 lithologic Unit I, 537a, 539a-541a, 550a
 lithologic Unit II, 537a, 541a, 550a
 summary, 540a
 magnetic properties, 33a-34a, 45a, 544a-545a
 magnetobiochronology, 818b-819b
 magnetostratigraphy, 753b-754b
 physical properties, 37a-39a, 545a-548a
 sedimentation rate, 544a
Kerguelen Sedimentary Ridge, E, topography, 525a
Kerguelen-Heard Plateau
 crustal structure, 7a
 geographic setting, 5a
 geological evolution, 8a, 124a
 glaciation, 131a
 rifting from Broken Ridge, 218a
Kerguelen-Heard Plateau N
 ash layers
 geochemistry, 326b-333b
 petrography, 325b
 bathymetry, 5a, 45a
 biology and oceanography, 35a-37a, 140a, 142a-144a, 191a, 193a-194a
 glaciation
 Neogene, 885b-888b
 Paleogene, 882b-885b
 paleoceanography, 124a
 paleoclimatology, 156a
 regional geology, 124a
 seismic stratigraphy, system geometry, 106a
Kerguelen-Heard Plateau N: Site 736
 biostratigraphy, 131a-135a, 156a, 836b
 benthic foraminifers, Eocene–Holocene, 615b-616b
 calcareous nannofossils, 469b
 datum levels, 837b
 diatoms, 557b-558b
 radiolarians, 514b-516b
 silicoflagellate, 933b-934b
 zonation, 24a, 26a-33a
 drilling deformation, 15a, 128a-129a
 Eocene–Miocene unconformity, 124a
 geochemistry, inorganic, 35a, 137a-140a
 geochemistry, organic, 34a-35a, 140a
 geophysics, 124a-125a
 interstitial-water chemistry, 353b-357b
 lithostratigraphy, 327b, 353b
 correlation, seismic stratigraphy, 156a
 depositional environment, 130a-131a
 fining-upward sequence, 130a
 lithologic Unit I, 123a, 129a-130a, 156a
 lithologic Unit I/Unit II contact, 828b
 lithologic Unit II, 123a, 130a, 156a

SUBJECT INDEX

summary, 129a
 living bacteria, density changes, 687b-689b
 location, 123a, 124a, 144b, 169b, 300b, 324b, 350b, 376b, 751b, 814b, 871b
 magnetic properties, 33a-34a, 45a, 135a-137a
 magnetobiochronology, 825b-828b
 navigation, 45a, 47a, 48a-85a, 87a
 physical properties, 37a-39a, 144a-153a
 sedimentary structures, 20a
 sedimentation rate, 137a
 seismic stratigraphy, 45a, 90a, 153a-155a
Kerguelen-Heard Plateau N: Site 737
 biostratigraphy, 173a-181a, 218a
 calcareous nannofossils, 470b-471b
 datum levels, 841b
 diatoms, 558b-559b
 Neogene, 839b, 842b
 Paleogene, 840b
 radiolarians, 516b-518b
 trace fossils, 639b
 zonation, 24a, 26a-33a
 depositional environment, 173a-174a
 evolution, 217b-219b
 transition, Unit IV, 173a
 drilling deformation, 15a, 165a
 Eocene hiatus, 173a
 geochemistry, inorganic, 35a, 185a-188a
 geochemistry, organic, 34a-35a, 188a-191a
 geophysical log, 265b, 275b
 geophysics, 160a-162a
 interstitial-water chemistry, 357b-363b
 lithology, 266b
 summary, 359b
 lithology, log-inferred, 285b
 calcareous sequences, 272b, 274b-277b
 lithostratigraphy, 159a, 172a, 217a-218a, 327b
 calcareous-siliceous transition, 357b
 lithologic Unit I, 165a, 212b
 lithologic Unit I/Unit II contact, 165a, 167a, 829b
 lithologic Unit II, 167a-168a, 212b
 lithologic Unit III, 168a, 212b, 264b
 lithologic Unit IV, 168a, 212b-213b, 264b
 lithologic Unit V, 168a, 170a, 213b, 264b
 lithologic Unit V/Unit VI contact, 835b-836b
 lithologic Unit VI, 170a-171a, 213b, 264b
 nannofossil ooze-to-chalk-to-limestone transition, 357b, 888b
 summary, 165a
 location, 144b, 159a, 169b, 212b, 264b, 300b, 324b, 350b, 376b, 814b, 871b
 logging, 207a, 210a, 264b, 275b-277b
 lithoporosity combination, 39a-40a, 208a
 lithostratigraphic correlation, 275b-277b
 physical properties, 205a-206a
 quality, 205a
 seismic stratigraphic combination, 39a, 208a
 summary, 221a-227a
 synthetic seismogram, 40a-41a, 212a-214a, 215a
 magnetic properties, 33a-34a, 45a, 181a-182a
 magnetobiochronology, 828b-836b
 magnetostratigraphy, 754b-757b
 mineralogy, 214b-215b
 navigation, 45a, 47a, 48a-85a, 87a, 88a
 Oligocene-Miocene hiatus, 162a, 169a, 833b
 physical properties, 37a-39a, 194a-203a
 sedimentary structures, 20a
 sedimentation rate, 183a, 185a
 seismic stratigraphy, 45a, 91a-92a, 214a
Kerguelen-Heard Plateau S: Site 738
 basalt
 alteration, 307b, 313b-315b
 geochemistry, 294b, 301b-307b, 308b-312b

isotopic data, 294b-295b
 mineralogy, 293b-294b
 petrography, 300b-301b
 bathymetry, 5a, 45a
 biology and oceanography, 35a-37a, 257a, 260a-261a, 279a
 biostratigraphy, 244a-252a, 279a, 831b, 832b
 benthic foraminifers, 613b-615b, 857b-859b
 calcareous nannofossils, 472b-474b, 856b-857b
 Danian, 495b-505b
 datum levels, 833b
 dinoflagellates, 631b-633b
 Maestrichtian-Eocene events, 835b
 planktonic foraminifers, 428b-435b, 452b-460b, 857b-859b
 radiolarians, 518b, 527b-529b
 trace fossils, 637b-639b
 zonation, 24a, 26a-33a
 calcareous ooze-to-chalk transition, 288b
 calcareous-biosiliceous transition, 822b
 Circumpolar Deep Water, erosional hiatuses
 associated with, 201b
 clay mineralogy, 194b
 Cretaceous/Tertiary boundary, 854b
 color changes, lithologic Unit I/Unit II
 boundary, 243a
 conductivity-temperature-depth profile, 109a
 Cretaceous/Tertiary boundary
 sampling strategy, 849b, 851b
 sediment composition, 852b
 depositional environment, 238a, 240a
 evolution, 217b-219b
 drilling deformation, 15a
 element stratigraphy, Cretaceous/Tertiary
 boundary, 719b-726b
 geochemistry, inorganic, 35a, 254a-257a
 geochemistry, organic, 34a-35a, 257a
 geological evolution, 229a-230a, 280a
 geophysical log, 267b, 277b
 geophysics, 230a-232a
 glaciation
 Neogene, 885b-888b
 Oligocene-Holocene, 201b-202b
 Paleogene, 882b-885b
 interstitial-water chemistry, 364b-367b
 iridium, Cretaceous/Tertiary boundary clay,
 726b-729b, 854b-856b
 lithology, 187b, 216b, 268b
 lithology, log-inferred, 285b
 calcareous sequences, 277b-279b
 lithostratigraphy, 229a, 234a-244a, 307b, 849b
 correlation, physical properties, 267a
 lithologic Unit I, 213b, 235a, 279a
 lithologic Unit I/Unit II contact, 822b
 lithologic Unit II, 213b, 235a, 237a, 279a
 lithologic Unit III, 213b, 237a, 264b, 279a
 lithologic Unit IV, 213b, 237a, 264b, 279a
 lithologic Unit V, 213b-214b, 237a-238a,
 264b, 279a, 838b
 lithologic Unit VI, 214b, 238a, 264b, 279a,
 838b, 843b
 lithologic Unit VII, 214b, 238a-240a, 300b
 lithologic Unit VIII, 214b, 240a-243a, 300b
 Maestrichtian-Turonian sediments, 825b
 magnetic susceptibility and, 243a, 246a
 nannofossil ooze-to-chalk-to-limestone transition, 363b-364b
 nannofossil-diatom ooze transition, 202b
 summary, 237a
 location, 7a, 144b, 169b, 212b, 230a, 264b,
 293b, 300b, 324b, 350b, 376b, 693b, 694b,
 714b, 751b, 814b, 871b
 logging, 264b-265b

lithoporosity combination, 39a-40a, 268a
 lithostratigraphic correlation, 277b-279b
 nannofossil ooze-to-chalk transition, 271a
 physical properties, 272a
 seismic stratigraphic combination, 39a, 270a
 summary, 283a-288a
 synthetic seismogram, 40a-41a, 273a-276a
 Unit 1, 270a-271a, 277b-279b, 280a
 Unit 2, 271a, 279b, 280a
 Unit 3, 272a, 279b, 280a
 Unit 4, 272a
Maestrichtian-Eocene hiatus, 244a
 magnetic properties, 33a-34a, 45a, 252a-253a
 magnetobiochronology, 822b-825b
 magnetostratigraphy, 757b-759b
 Cretaceous/Tertiary boundary, 851b-853b
 mineralogy, 215b-217b
 Miocene-Pliocene hiatus, 243a
 nannofossil-diatom ooze transition, Polar Front
 migration and, 195b
 navigation, 45a, 47a, 48a-85a, 87a
 Oligocene-Miocene hiatus, 202b, 243a-244a
 paleoceanography, 229a-230a, 279a
 paleolatitude, 759b
 petrogenesis, 315b-319b
 physical properties, 37a-39a, 261a-269a
 preglacial conditions, 200b-201b
 sediment composition, 188b, 204b
 environmental implications, 194b-200b
 nonbiogenic sediment, 190b-193b
 sedimentary structures, 20a
 sedimentation rate, 253a-254a
 Cretaceous/Tertiary boundary, 852b-853b
 sedimentology, 234a-244a
 seismic stratigraphy, 45a, 93a, 276a-279a
 stable isotopes, 859b
 Eocene-Miocene, 708b-710b
 Eocene/Oligocene transition, 710b-712b
 paleoceanographic implications, 712b-714b
 Paleocene-Eocene, 697b-708b
Kerogen
 Prydz Bay
 chemical composition, 410b
 unreactive type III, 380b, 388b
 Prydz Bay: Site 739
 depositional environment, 408b, 411b
 lithologic Unit II vs. Unit V, 408b
 Prydz Bay: Site 741, highly altered Type IV,
 410b
 Prydz Bay: Site 742, reworked type III, 385b
 Kerogen, amorphous, Prydz Bay: Site 739, allochthonous origin, 48b
 King George Island, glaciation, 79b, 201b
 Lag deposit, gravel, Prydz Bay: Site 739, 129b
 Lambert Glacier-Amery Ice Shelf system
 bedrock topography, 64b, 68b
 debris transport, 71b, 73b-74b
 drainage basin, 65b, 66b
 flow lines, 65b
 grounding line
 Miocene-Pliocene fluctuation, 248b
 Oligocene, 875b
 grounding past present ice sheet edge, isotopic
 evidence, 714b-715b
 ice dynamics, 62b, 64b
 ice flow, 62b, 70b-71b, 72b, 73b
 ice thickness, 64b
 interior basin, mass flux, 62b
 rift structure, 9a
 structural glaciology, 63b, 68b-71b
 Landsat imagery, 68b-69b
 surging, model, 62b
 Wisconsin glacial maximum, retreat from, 749b

Lambert Glacier
ablation, 62b
channel geometry, 70b
drainage basin, 7a
glacial deposits, Cenozoic, 7b
grounding line, 339a
ice drainage basin, 290a
ice thickness, 64b
lobate prograding apron, 20b
longitudinal foliation, confluence region, Mellor and Fisher glaciers, 75b
origin, 62b
regional geology, 5b
Lambert Graben
age, 7a
basement depressions, 67b
development, timing of, 67b-68b
draining basin, 338a
formation, 346a
geology, 66b-67b
ice cover, thickness of, 64b
nonmagnetic rocks, Permian, 80b
northerly extension, 67b-68b
Prydz Bay development, 377a
regional geology, 5b
sediment transport, 80b
structural separation, Prydz Bay basin, 18b
subglacial floor depth, 80b
subsea-level feeder branches, 64b
topography, subglacial, 64b
Laminations
Kerguelen Sediment Ridge: Site 745, 510a
Kerguelen-Heard Plateau S: Site 738
calcareous chalk, 237a-238a
sedimentation rate, 853b
Prydz Bay: Site 740, siltstones and claystones, 46b
Laminations, clay-rich, Kerguelen-Heard Plateau S: Site 738, Cretaceous/Tertiary boundary, 723b, 727b-728b
Laminations, diagenetic greenish, Kerguelen-Heard Plateau N: Site 737, in calcareous claystone, 170a
Laminations, foreset, Prydz Bay: Site 740, sigmoidal, 46b
Laminations, ripple
Prydz Bay: Site 740, 46b
siltstones, 49b
Laminations, white ooze, Kerguelen-Heard Plateau S: Site 738, microfossil remains, 865b
Lanthanum, Kerguelen-Heard Plateau S: Site 738, scandium-normalized distribution, 725b
Lead isotopes, Kerguelen-Heard Plateau S: Site 738, basalts, 294b-295b, 298b
Legru Bay Group, glacial sediments, Oligocene, 871b
Limestone
Kerguelen-Heard Plateau S: Site 738, 238a
bioturbated, 239a
magnetic properties, 758b-759b
Limestone, calciclastic, Kerguelen-Heard Plateau S: Site 738, 238a, 239a
Limestone, clayey, Kerguelen-Heard Plateau N: Site 737, sedimentary structures, 172a
Limestone, sandy
Prydz Bay: Site 742, 135b, 141b
Jurassic-Cretaceous deposition, 138b
Lithofacies
Prydz Bay
diamict, 88b-89b
grain-size distribution, 89b-90b, 94b
sand/mud ratio by type, 95b, 96b
thicknesses and recovered percentages, 88b
Lonestone

Kerguelen Sediment Ridge: Site 745, metamorphic basement origin, 508a, 530a
Kerguelen-Heard Plateau N: Site 736, 131a
McMurdo Sound
diamictites, grain size and meltwater activity, 120b-121b
glaciation, Oligocene-Miocene, 79b, 118b, 872b
Magma
Kerguelen-Heard Plateau S: Site 738
comparison, Mesozoic Southern Hemisphere basalts, 315b, 317b
evolution, 317b
source regions, 317b, 318b, 319b
Magma, basaltic, Kerguelen Plateau, formation, 299b
Magnesium
high-resolution vs. regular sampling results, 395b, 397b
Kerguelen Plateau S: Site 744, 491a
pore waters, 930b-931b
vertical distribution gradient, 367b, 372b
vs. calcium, 493a
Kerguelen Sediment Ridge
pore waters, 930b-931b
vertical distribution gradient, 371b, 373b
Kerguelen Sediment Ridge: Site 745, 516a-517a
vs. calcium, 522a
Kerguelen Sediment Ridge: Site 746, 544a
Kerguelen-Heard Plateau N, pore waters, 929b-930b
Kerguelen-Heard Plateau N: Site 736, 139a, 356b
Kerguelen-Heard Plateau N: Site 737, 185a-186a
chemical gradient, 359b
discontinuities, 187a
vertical distribution gradient, 363b
vs. calcium concentration, 193a
Kerguelen-Heard Plateau S: Site 738, 255a
vertical distribution gradient, 366b, 372b
vs. calcium concentration, 259a
Prydz Bay, 312a, 360a, 385a, 418a, 466a
pore waters, 380b, 930b-931b
silicate formation, 312a
source, 362a
vertical distribution gradient, 385b, 386b, 388b, 391b
vs. calcium, 420a
Prydz Bay: Site 739, vertical distribution gradient, 380b-381b
Magnetic minerals, Kerguelen Sediment Ridge: Site 745, during enhanced glaciation, 757b
Magnetic properties
Kerguelen Plateau S: Site 744, 45a, 488a-490a, 503a, 783b-794b
alternating-field demagnetization, 772b, 773b-777b
direction changes, 775b
logging data, 895b, 896b-897b
multicomponent magnetization, 772b, 774b, 776b, 782b
NRM intensity, 778b, 779b, 780b
paleomagnetic directions, 776b
reversal stratigraphy, 489a-490a
shore-based vs. shipboard measurements, 772b
susceptibility, 34a, 821b
Kerguelen Sediment Ridge: Site 745, 45a, 516a, 517a, 518a
alternating-field demagnetization, 753b, 754b
logging data, 895b, 896b-897b, 900b-901b
mean destructive field, 757b, 761b
NRM intensity, 752b, 757b, 761b
stable inclinations, 754b, 756b
susceptibility, 34a, 757b, 761b
Kerguelen Sediment Ridge: Site 746, 45a, 544a, 545a, 546a, 547a
alternating-field demagnetization, 753b, 756b
NRM intensity, 753b
stable inclinations, 755b, 757b
Kerguelen-Heard Plateau N: Site 736, 45a, 135a-137a
alternating-field demagnetization, 135a, 138a
susceptibility, 34a, 137a, 139a
Kerguelen-Heard Plateau N: Site 737, 45a, 185a, 189a
alternating-field demagnetization, 181a-183a, 184a, 188a, 754b
intensity-susceptibility correlation, 181a, 182a-183a
NRM intensity, 754b
stable inclinations, 759b
Kerguelen-Heard Plateau S: Site 738, 45a, 252a-253a, 254a
alternating-field demagnetization, 758b, 763b
Cretaceous/Tertiary boundary, 253a, 757b, 758b, 760b, 764b, 765b
flux jumps, 253a
lithostratigraphic correlation, 243a, 246a
NRM intensity, 758b, 766b
stable inclinations, 766b
susceptibility, 34a, 758b
Prydz Bay, 45a, 465a
Prydz Bay: Site 739
alternating-field demagnetization, 305a, 310a
intensity, long-term oscillations, 310a
susceptibility, 310a
Prydz Bay: Site 740, 359a, 374a
alternating-field demagnetization, 357a, 358a, 798b
high-temperature magnetization, 803b-804b
lithological correlation, 804b
multicomponent magnetization, 798b, 802b-804b
NRM intensity, 357a, 804b
paleomagnetic directions, 797b-798b
red beds, 357a-358a, 797b-804b
remagnetization, 798b, 800b, 801b, 806b
thermal demagnetization, 357a, 358a, 798b-804b
univectorial magnetization, 798b, 799b
Prydz Bay: Site 742, 415a, 416a, 417a, 451a, 769b
anhydrous remanent magnetization, 415a
NRM intensity, 759b-760b, 768b
reversed polarity zones, 415a
stable inclinations, 770b
susceptibility, lithologic correlation, 759b-760b, 768b
susceptibility, organic carbon and, inverse correlation, 113b
Magnetobiochronology
Kerguelen Plateau S: Site 744, 820b-822b
Kerguelen Sediment Ridge: Site 745, 817b-818b, 820b
Kerguelen Sediment Ridge: Site 746, 818b-819b, 822b, 823b
Kerguelen-Heard Plateau N: Site 736, 825b, 828b, 838b
Kerguelen-Heard Plateau N: Site 737, 828b-829b, 831b, 833b-836b, 843b
Kerguelen-Heard Plateau S: Site 738, 822b, 834b
Cretaceous/Tertiary boundary, 825b
Prydz Bay, 836b-838b, 843b, 846b
Magnetostratigraphy
Kerguelen Plateau S: Site 744, 824b, 829b, 830b

SUBJECT INDEX

anomalies 7A, 8, 11–13, and 15, 782b
 biostratigraphic correlation, 781b–782b
 geomagnetic time scale correlation, 782b
 hiatuses, 821b
 polarity logs, 819b–820b
 reversed polarity, 772b, 781b
 Kerguelen Sediment Ridge, 760b
 Kerguelen Sediment Ridge: Site 745, 752b, 756b, 818b
 biostratigraphic correlation, 752b–753b
 Kerguelen Sediment Ridge: Site 746, 550a–551a, 755b, 818b–819b, 821b
 biostratigraphic correlation, 753b–754b
 Kerguelen–Heard Plateau N: Site 737, 183a, 190a, 218a, 757b, 828b
 biostratigraphic correlation, 754b
 Neogene, 760b, 839b, 842b
 subchron boundaries, 841b
 Kerguelen–Heard Plateau S: Site 738, 822b, 832b, 835b, 851b, 852b
 Cretaceous/Tertiary boundary, 757b–759b
 reversed polarity, 758b
 subchron boundaries, 833b
 Tertiary anomalies, 279a
 Prydz Bay, 16b
 anomaly, acoustic Unit PS.6, 13b
 Prydz Bay: Site 742, 451a, 759b–760b, 843b, 845b, 846b, 881b, 882b
 reversed polarity, 760b
 Mahanadi Basin, siltstone, Cretaceous, 60b
 Mahanadi rift graben, deposition, 21b
 Mahanadi Valley, 9a
 Malaspina Glacier, ice flow, geometry, 71b
 Mammoth Subchron
 Kerguelen Plateau S: Site 744, 820b
 Kerguelen Sediment Ridge: Site 745, 753b
 Manganese
 Kerguelen Plateau S: Site 744, 482a, 483a–484a
 Kerguelen–Heard Plateau N: Site 736, 356b
 Manop Site C, iridium accumulation rate, 855b
 Marie Byrd Land, hyaloclastite deposits, Cenozoic, 10a
 Marl, diatom-bearing ferruginous
 Prydz Bay: Site 739, 134b, 140b
 provenance, 138b
 Matuyama Chron
 Kerguelen Plateau S: Site 744, 490a
 Kerguelen–Heard Plateau N: Site 737, 757b
 Maud Rise
 carbonate, strontium isotopes, 741b
 carbonate–biosiliceous facies transition, 195b, 874b
 clay mineralogy, chemical weathering effects, 874b
 deep waters, low-latitude origin, Eocene, 707b
 oxygen isotopes, Paleogene record, 704b, 885b
 Mawson Escarpment, 80b
 basement depressions, 67b
 topography, 64b
 Mellor Glacier
 longitudinal foliation, confluence region,
 Lambert and Fisher glaciers, 75b
 See also Lambert Glacier
 Mertz–Ninnis Trough, Holocene sediment accumulation, 747b, 748b
 Metals
 Kerguelen Plateau S: Site 744, geochemical logs, 903b, 911b, 925b
 Kerguelen Sediment Ridge, geochemical logs, 903b, 912b, 914b, 916b, 918b, 922b, 925b
 Kerguelen–Heard Plateau N, geochemical logs, 903b, 908b–909b, 913b–914b, 915b–916b, 917b–918b, 920b–921b, 923b–924b

Kerguelen–Heard Plateau S: Site 738, geochemical logs, 903b, 909b–910b, 924b
 Prydz Bay: Site 739, geochemical logs, 903b, 910b–911b, 914b, 916b, 918b, 921b–922b, 924b–925b
 Metamorphic rock, Prydz Bay, 8a, 380a
 Meteoric waters, Prydz Bay, effect on interstitial water chemistry, 388b
 Methane
 Kerguelen–Heard Plateau S: Site 738, interstitial waters, 366b
 Prydz Bay, 385a, 420a, 422a
 Methane/ethane ratio, Prydz Bay: Site 742, 421a, 422a
 Mica, Prydz Bay: Site 740, petrography, 50b
 Mica schist, Prydz Bay, 138b, 453a
 Micrite
 Kerguelen Plateau S: Site 744, 480a, 482a, 503a, 771b
 Kerguelen–Heard Plateau N: Site 737, 168a, 357b
 clotted structures, 173a
 Prydz Bay, 302a–303a, 409a
 Microfaults, Kerguelen–Heard Plateau N: Site 737, 171a
 Microfractures
 Kerguelen Sediment Ridge: Site 746, 542a
 Kerguelen–Heard Plateau S: Site 738, 238a
 Microsparite, Prydz Bay: Site 739, 302a
 Mid-Ocean Ridge basalts (MORB), Southern Ocean, barium concentration, 315b–316b
 Mineralogy
 Kerguelen–Heard Plateau N: Site 737
 detrital, 168a
 lithological correlation, 276b–277b, 278b–279b
 tephras, 325b
 Kerguelen–Heard Plateau S: Site 738
 lithological correlation, 280b–281b
 secondary, basalts, 316b
 Prydz Bay: Site 742, lithological correlation, 282b–283b, 284b–285b, 286b–287b
 Miocene/Pliocene boundary, Polar Front, northward migration, 484a
 Montmorillonite, cryptocrystalline, Kerguelen–Heard Plateau S: Site 738, 301b
 Montmorillonite–beidellite, Kerguelen–Heard Plateau S: Site 738, 301b
 Mount Rubin, kaolinite, in glacial tills, 114b
 Mud, diatomaceous
 Kerguelen Plateau S: Site 744, organic carbon content, 113b
 Kerguelen–Heard Plateau N: Site 736, organic carbon content, 113b
 Prydz Bay
 organic carbon content, 113b
 terrigenous component, 107b
 Mud, pebbly, Prydz Bay: Site 740, depositional environment, 749b
 Mud, sandy, Prydz Bay: Site 739, 93b, 127b
 Mud, terrigenous, Prydz Bay, modern deposition, 112b
 Mudstone
 Prydz Bay: Site 739, with dropstone, 130b
 Prydz Bay: Site 742, sedimentological log, 111b
 Mudstone, diatomaceous
 Prydz Bay, 89b
 sorting, 90b
 terrigenous component, 107b
 Mudstone, sandy
 Prydz Bay, 89b, 93b, 126b
 sorting, 90b
 terrigenous component, 107b
 Mudstone, stratified

Prydz Bay, 89b, 127b, 130b
 lacustrine deposition, 107b
 Muscovite, Prydz Bay: Site 740, 50b
 Mylonite, Prydz Bay, 138b
 Neodymium isotopes, Kerguelen–Heard Plateau S: Site 738, basalts, 294b–295b, 297b, 298b
 Nickel, Kerguelen–Heard Plateau S: Site 738, in basalt, 303b
 Nitrogen, combustible
 Kerguelen Plateau S: Site 744, 114a–115a
 Prydz Bay, 114a–115a
 isotopic fractionation, 115a, 117a
 North Atlantic Deep Water, carbon isotope enrichment, Oligocene bicarbonates, 713b
 Northern Ice Limit, seasonal ice cover, 109a–110a, 112a–113a
 Norwegian Sea, Gilbert Chronozone, 753b
 Nova Scotia, smectite–kaolinite correlation, 198b
 Nunivak Event, Kerguelen Sediment Ridge: Site 745, 818b
 Olduvai Subchron, Kerguelen Sediment Ridge: Site 745, 752b–753b
 Olivine, Kerguelen–Heard Plateau S: Site 738, pseudomorphs, 301b–302b
 Ooze, calcareous
 Kerguelen Plateau S: Site 744, 480a, 482a
 Kerguelen–Heard Plateau N: Site 737, 168a
 Ooze, calcareous nannofossil, Kerguelen–Heard Plateau S: Site 738, 235a, 237a
 Ooze, clay rich, Kerguelen–Heard Plateau N: Site 737, 276b
 Ooze, clayey diatom
 Kerguelen Sediment Ridge, 539a
 glacial/interglacial cycles, 226b–232b, 233b
 vs. diatomaceous clays, color differences, 226b
 Ooze, diatom
 Kerguelen Plateau, ice rafting and onset of, 120b
 Kerguelen Plateau N, sedimentation rate, 353b
 Kerguelen Plateau S: Site 744, 771b
 colors, 480a
 Miocene accumulation onset, 190b
 organic carbon content, 113b
 sedimentation, 490a
 Kerguelen Sediment Ridge: Site 745, 508a
 colors, 509a–510a
 magnetic properties, 752b
 Kerguelen–Heard Plateau N: Site 736, 129a–130a
 depositional environment, 130a, 155a
 organic carbon content, 113b
 physical properties, 144a, 199a, 202a
 Kerguelen–Heard Plateau N: Site 737, 165a
 compaction rate and porosity of, 271a
 initiation of, 195b
 physical properties, 199a, 202a
 sedimentation rate, Pliocene, 185a
 seismic stratigraphy, 215a
 textural ranges, 167a–168a
 Kerguelen–Heard Plateau S: Site 738, 235a
 physical properties, 267a
 sedimentation rate, 253a
 Prydz Bay, 89b, 93b, 125b–126b, 372a, 374a, 380a–381a
 cold water characteristics, 112b
 enhanced productivity association, 107b
 organic carbon content, 113b
 Prydz Bay: Site 739, depositional environment, 112b
 Prydz Bay: Site 740
 geochemistry, inorganic, 358a
 physical properties, 365a, 383b

Prydz Bay: Site 741, bacterial activity, 384b
 Ooze, diatom nannofossil, Kerguelen-Heard Plateau N: Site 737, 168a
 Ooze, diatomaceous
 Kerguelen Plateau S: Site 744, 501a–502a
 physical properties, 495a
 Kerguelen Sediment Ridge: Site 745, physical properties, 520a
 Kerguelen-Heard Plateau N: Site 736, 825b
 magnetic properties, 135a
 Prydz Bay: Site 741, velocity, 34b
 Ooze, foraminiferal, Kerguelen Plateau S: Site 744, 480a
 Ooze, nannofossil
 Kerguelen Plateau S: Site 744, 483a, 502a,
 771b–772b
 chalk transition, 500a
 colors, 482a
 magnetic properties, 819b–820b
 physical properties, 495a
 sedimentation rate, 490a
 Kerguelen Sediment Ridge, 541a
 bioturbation, 246b–247b
 Kerguelen-Heard Plateau S: Site 738
 chalk transition, 269a, 271a
 compaction rate and porosity of, 271a
 sedimentation rate, 253a–254a
 Ooze, siliceous, Prydz Bay: Site 740, sedimentation rate, 748b–749b
 opal
 Kerguelen Plateau S: Site 744, 190b
 Kerguelen Sediment Ridge
 glacial/interglacial cycles, 240b
 inverse terrigenous correlation, 227b
 Kerguelen-Heard Plateau S: Site 738, 190b
 A to CT transition, 219b
 as productivity indicator, 201b
 Prydz Bay, 84b, 85b
 carbonate correlation, 113b
 Opal, biogenic
 Kerguelen Plateau S: Site 744, 217b
 Eocene, 882b
 Kerguelen-Heard Plateau N: Site 737, 215b
 Miocene, 218b
 Opal-CT
 Kerguelen-Heard Plateau N: Site 737, chert layers, 214b–215b
 Kerguelen-Heard Plateau S: Site 738, 217b, 220b
 chert layers, 218b
 formation, 219b
 Orkney Islands, S, modern bivalve, strontium isotopes, 741b
 Overconsolidated sediment
 Prydz Bay, 154b
 causes of, 174b–176b
 glacial advance and retreat effects, 157b
 ice loading effects, 884b
 physical properties, 31b, 46b, 150b
 transition to normally consolidated sediments, 181b
 Overconsolidation ratios (OCR)
 Prydz Bay, 152b
 sediment loading effects, 156b
 vs. shear strength, 155b–156b
 Oxidation, Kerguelen-Heard Plateau S: Site 738, altered basalts, 314b–315b
 Oxidation-reduction potential
 Kerguelen Plateau S: Site 744, sediment accumulation rate and, 367b
 Kerguelen Sediment Ridge, 371b, 373b
 Kerguelen-Heard Plateau S: Site 738, 364b
 Oxygen index

Kerguelen-Heard Plateau N: Site 736, 144a, 145a
 Kerguelen-Heard Plateau N: Site 737, 198a, 199a
 Prydz Bay: Site 739, 408b, 410b
 Prydz Bay: Site 741, 388a
 Prydz Bay: Site 742, 426a
 Prydz Bay: Site 743, 469a
 Oxygen isotopes
 Amery Ice Shelf, 64b
 Antarctica
 Cenozoic record, 869b–870b
 ice sheet documentation, 78b, 79b
 Atlantic Ocean S, Eocene enrichment, 882b–883b
 deep-sea benthic foraminifers
 Eocene/Oligocene boundary increase, 694b
 global warming and Tertiary decline, 694b
 magnetostratigraphic correlation, 694b, 695b
 depletion in meteoric waters, latitude correlation, 377b
 estimating ice volume from, 870b
 Holocene planktonic foraminifers, dissolved carbon dioxide equilibrium, 697b
 Kerguelen Plateau S: Site 744, 503a, 696b, 698b–699b, 826b–828b, 935b–939b
 carbonate minima and, 194b–195b
 cooling trend, 885b
 enrichment, Eocene/Oligocene boundary, 704b, 708b, 710b, 711b–712b
 Eocene–Miocene, 708b–709b, 882b, 884b, 935b–939b
 hiatus correlation, 936b
 intersite comparison, 713b
 Neogene, 887b
 Oligocene enrichment zone, glacial debris in, 712b
 Oligocene hiatus, 821b
 paleoceanographic implications, 714b–715b
 Paleocene/Eocene boundary, 697b, 704b
 planktonic foraminifers, 705b
 stratigraphy, 935b
 strontium isotope correlation, 734b
 Kerguelen Sediment Ridge, Neogene, 887b
 Kerguelen-Heard Plateau N, diatom species transition, Neogene, 887b–888b
 Kerguelen-Heard Plateau S: Site 738, 700b–701b, 702b–703b, 705b
 carbonate minima and, 194b–195b
 Cretaceous/Tertiary boundary, 859b
 enrichment, Eocene/Oligocene boundary, 704b, 708b, 714b
 Eocene–Miocene, 708b–709b
 intersite comparison, 713b
 latitudinal gradient, Eocene, 707b–708b, 709b
 Oligocene–Miocene record, 202b
 paleoceanographic implications, 714b–715b
 Paleocene/Eocene boundary, 697b, 704b, 706b
 surface to intermediate water gradients, 697b, 704b, 709b–710b, 714b
 Miocene–Pliocene cooling trend, 249b
 planktonic and benthic foraminifers
 interspecific differences, 697b
 systematic, 696b–697b
 Prydz Bay, 377b
 Eocene/Oligocene boundary, 10a
 glacial/interglacial cycles, 886b
 Miocene hiatus, 120b
 mollusk shells, 740b, 742b
 Oligocene cooling trend, 248b
 Prydz Bay: Site 739
 Neogene, 887b
 Oligocene sea surface temperature correlation, 745b

weathering effects, 381b
 Prydz Bay: Site 740
 correlation, Antarctic Bottom Water, 383b
 interstitial waters, 384b
 Prydz Bay: Site 742
 below sediment/water interface, 385b
 interstitial waters, 386b–387b
 seawater enrichment, ice volume association, 695b
 P- wave velocity
 Kerguelen Plateau S: Site 744, 37a–38a, 495a
 Kerguelen Sediment Ridge: Site 746, 37a–38a
 Kerguelen-Heard Plateau N, 37a–38a, 148a–149a, 197a
 Kerguelen-Heard Plateau S: Site 738, 37a–38a
 Prydz Bay, 37a–38a
 Pacific Ocean, Oligocene deep waters, low-latitude origin, 713b
 Pacific Ocean, equatorial
 carbon isotopes, comparison, Prydz Bay and Weddell Sea, 704b
 carbonate deposition, Eocene/Oligocene boundary, 714b
 Pacific Ocean S
 ice rafted debris, Eocene, 873b
 sand, glacially-derived, 201b
 Pack ice
 Kerguelen Plateau S: Site 744, 110a, 112a–113a
 Kerguelen-Heard Plateau S: Site 738, 110a, 112a–113a
 Prydz Bay, 110a, 112a–113a
 Paleocene/Eocene boundary
 carbon isotopes, Kerguelen Plateau S: Site 744, 697b, 704b
 oxygen isotopes
 Kerguelen Plateau S: Site 744, 697b, 704b
 Kerguelen-Heard Plateau S: Site 738, 697b, 706b
 Paragneiss, garnet-biotite-feldspar, Prydz Bay, 454a
 Paragneiss, garnet-biotite-sillimanite, Prydz Bay, 454a
 Paragneiss, granulitic, Prydz Bay, 138b
 Particulate flux
 Kerguelen Plateau S: Site 744
 collections, 109a–110a
 mass, 114a
 Prydz Bay
 collections, 109a–110a
 mass, 114a
 Pedogenesis, carbonate, climatic requirements, 374a
 Pedogenic sediment, Prydz Bay: Site 740, depositional environment, 356a
 Pegmatoid schlieren
 Kerguelen-Heard Plateau S: Site 738
 in basalts, 300b, 322b
 plagioclase composition, 303b
 Permeability, Prydz Bay, diamictons, 152b, 154b
 Permeability anisotropy, Prydz Bay: Site 740, 369a
 Permo-Triassic Beacon Supergroup, Prydz Bay, red beds, 54b
 Peru Margin
 ammonium, 138a
 phosphate, 138a
 pH
 Kerguelen Plateau S: Site 744, 491a
 Kerguelen Sediment Ridge: Site 745, 517a
 Kerguelen-Heard Plateau N, 139a–140a, 186a
 discontinuities, 187a
 Kerguelen-Heard Plateau S: Site 738, 256a
 Prydz Bay, 312a, 362a, 418a

SUBJECT INDEX

Phosphate
 Kerguelen Plateau S: Site 744, 491a
 Kerguelen Sediment Ridge, 518a, 545a
 pore waters, 369b, 371b, 373b
 Kerguelen-Heard Plateau N: Site 736,
 137a–138a, 140a, 156a
 Kerguelen-Heard Plateau N: Site 737, 187a,
 359b
 Kerguelen-Heard Plateau S: Site 738, 257a,
 301b
 Prydz Bay: Site 739, 313a
 Prydz Bay: Site 740, 362a–363a, 374a
 ammonium and alkalinity correlation, 383b
 Prydz Bay: Site 741, 385a
 vs. ammonium, pore waters, 384b
 Prydz Bay: Site 742, 420a
 pore waters, 386b
 sediment/water interface, 385b
 Prydz Bay: Site 743, 466a
 Photoelectric effect
 Kerguelen-Heard Plateau S: Site 738, 277b
 Prydz Bay: Site 742, diamictites, 281b
 Plagioclase, Kerguelen-Heard Plateau S: Site 738,
 in pegmatoid schlieren, 301b
 Plant pigments
 calculated fluxes, 120a
 degradation products, 118a–119a
 Plasticity, Prydz Bay, 150b, 151b
 Plutonic rock, Prydz Bay, 8a–9a
 Polar Front
 ice cover control, 230b
 Kerguelen Island, 244b, 246b
 northward migration, 195b
 Miocene/Pliocene boundary, 484a
 position of, climatic control, 229b
 Polar Frontal Zone
 biogenic sediments, present day, 874b
 biogeographic changes, 874b
 circumpolar current, 211b
 Polar Record Glacier, outflow, 133b
 Polonez Cove Formation, 79b
 glacially controlled deposits, 871b
 Porosity
 Kerguelen Plateau S: Site 744, 495a
 with silica variations, 499a
 Kerguelen Sediment Ridge: Site 745, 520a,
 522a–524a
 Kerguelen Sediment Ridge: Site 746, 546a–547a
 Kerguelen-Heard Plateau N: Site 736,
 144a–145a, 148a, 153a
 Kerguelen-Heard Plateau N: Site 737,
 calcareous ooze-limestone transition, 285b
 Kerguelen-Heard Plateau S: Site 738, 262a,
 264a, 266a
 Prydz Bay, grain size and lithological
 correlations, 388b
 Prydz Bay: Site 739, 145b, 148b, 325a, 327a
 electrical resistivity correlation, 379b
 magnesium and, 312a
 measured vs. calculated, 283b
 Raymer-Hunt equation, 269b
 sedimentation rate and, 330a
 silica and, 312a
 Prydz Bay: Site 740, diatom ooze, 383b
 Prydz Bay: Site 742, 146b, 428a, 434a
 grain size and, 385b
 massive vs. stratified diamictites, 281b–282b
 measured vs. calculated, 274b, 435a
 Prydz Bay: Site 743, 146b, 148b, 469a–470a
 Porosity-density relationship
 Kerguelen-Heard Plateau N: Site 737,
 calcareous sequences, 276b
 Kerguelen-Heard Plateau S: Site 738,
 compaction effect, 279b

Potassium
 Kerguelen Plateau S: Site 744
 pore waters, 930b–931b
 vertical distribution gradient, 367b, 372b
 Kerguelen Sediment Ridge
 pore waters, 930b–931b
 vertical distribution gradient, 371b, 373b
 Kerguelen-Heard Plateau N: Site 736, 356b
 pore waters, 929b
 Kerguelen-Heard Plateau N: Site 737
 chemical gradient, 359b
 pore waters, 929b–930b
 radioactivity relationship, 272b, 274b
 vertical distribution gradient, 363b
 Kerguelen-Heard Plateau S: Site 738, vertical
 distribution gradient, 366b, 372b
 Prydz Bay
 pore waters, 380b, 930b–931b
 vertical distribution gradient, 380b–381b, 391b
 Potassium-feldspar crystals, Kerguelen-Heard Plateau S: Site 738, in pegmatoid schlieren, 301b
 Potassium/rubidium ratio, Kerguelen-Heard Plateau S: Site 738, 302b
 Preconsolidation stress
 Prydz Bay, 152b, 158b
 vs. permeability, glaciogenic sediments, 155b
 Prince Charles Mountains
 alkali igneous intrusive, 22b, 295b
 alkaline mafic igneous rock,
 Cretaceous–Eocene, 7b
 alpine glaciation, initiation, 74b
 banded iron-formations, 65b
 clay mineralogy, 114b
 ice cover, thickness of, 64b
 ice-free areas, 62b
 Precambrian igneous and metamorphic rock,
 64b–65b
 red beds, 795b
 schematic-geologic map, 10a
 sedimentary rock, Phanerozoic, 7b
 topography, 64b
 uplift, 22b
 Production index, Prydz Bay, 408b, 410b
 Productivity
 Kerguelen Sediment Ridge
 facies A vs. facies B, 229b–230b
 glacial/interglacial cycles, 232b
 Kerguelen-Heard Plateau S: Site 738
 barite production and, 725b
 opal content indicators, 201b
 Prydz Bay: Site 741, 388a
 Prograding sequence
 Kerguelen-Heard Plateau N: Site 737, seismic
 reflection profile, 214a–215a, 217a
 Prydz Bay
 deposition, 118b, 459a
 depositional environment, 376b
 glacial units, 880b
 glacial/interglacial cycles, 19b
 ice-shelf stability, 115b
 inner shelf vs. outer, 876b
 lithostratigraphic correlation, 100b–101b
 overlying hiatus, 122b
 seismic stratigraphy, 81b, 336a–338a
 topset bed geometry, 19b
 upper and lower, 92b–93b
 Prydz Bay: Site 739
 age, 337a, 338a
 glacial topset beds, seaward thickening, 336a–
 337a
 gravel content, 81b
 Oligocene–Miocene, 885b
 thickness, 16b–17b
 uniformity, 339a

Prydz Bay: Site 742, seismic profile, 444a
 Prydz Bay: Site 743, 465a, 475a
 seismic stratigraphy, 474a
 Prydz Bay: Site 742, 420a, 422a, 423a
 Prydz Bay
 acoustic stratigraphy
 areal distribution, 13b–17b
 Unit PS.1, 7b–8b
 Unit PS.2, 8b–12b
 aliphatic hydrocarbons, 417b–423b
 bathymetry, 5a, 6b, 11a, 45a, 78b, 291a, 373a
 bedrock topography, 64b
 biogenic sedimentation, 112b–113b
 biology and oceanography, 35a–37a,
 313a–314a, 363a–365a, 387a–390a,
 422a–423a, 425a, 466a–468a
 biostratigraphy
 calcareous nannofossils, 477b, 743b
 diatoms, 645b–665b, 667b–673b, 743b–744b
 clasts
 metamorphic groups V–XIV, 137b–138b
 sedimentary group III, 135b, 137b
 clay mineralogy, 84b–86b
 conductivity-temperature-depth profile, 109a
 consolidation tests, 174b
 continental shelf, geology, 875b–876b
 depositional environment, 18b–19b
 diamictons, 154b
 drifting sediment trap deployment, 111a
 drilling deformation, 15a, 295a, 461a–462a
 evolutionary model, 20b–21b
 flat-lying sequence
 depositional mechanism, 115b
 ice fluctuation and deposition of, 122b
 lithology, 876b
 geographic setting, 80b–81b
 geologic setting, 5b, 6b, 58b, 80b–81b, 170b,
 375b–376b, 377b, 407b
 geological evolution, 7a–9a
 glaciation, 875b–882b
 continental shelf, 77b–122b
 continental shelf development and, stages of,
 116b–118b
 glacial-onset unconformity, 22b
 Neogene, 885b–888b
 Paleogene, 882b–885b
 glaciogenic sediment, genetic classification,
 22a–23a
 grain size distribution, 81b
 hinterland, 64b–68b, 69b
 ice loading, 176b–181b
 ice source, 20b
 lithostratigraphy
 intersite correlation, 413a
 post-breakup strata, 9a
 logging, geotechnical, 177b
 magnetobiochronology, 836b–846b
 morphology, 7a–8a, 290a
 navigation, 45a, 47a, 48a–85a, 87a, 88a
 overconsolidation, causes of, 174b–176b
 oxygen isotopes, and sea ice formation,
 Eocene/Oligocene boundary, 10a
 paleoclimatic evolution, 9a–10a
 particulate fluxes, 109a–110a
 mass, 114a
 physical properties, 37a–39a, 147b–158b,
 315a–328a, 365a–369a, 390a–392a,
 425a–434a, 468a–473a
 Atterberg limits, 150b–151b
 consolidation and, 155b–157b
 grain size and, 149b–150b
 permeability and, 152b–153b
 rift history, 20b

sedimentary structures, 20a
distribution, 17b
seismic stratigraphy, 9a, 333a–334a
geologic noise, 7b
outer shelf, 336a–338a
structured features, 17b–18b
sonobuoy seismic recording
seismic-refraction data, 28b–29b
velocity-depth study, 29b–30b, 30b–33b, 34b–35b, 35b
topography, subglacial, 80b
Prydz Bay: Site 739
biostratigraphy, 304a–305a, 339a, 844b
radiolarians, 518b–519b
zonation, 24a, 26a–33a
carbon isotopes, bivalve shells, 742b
depositional environment, 101b–115b,
303a–304a, 338a
glacial, 18b–20b
geochemistry, inorganic, 35a, 310a–313a
geochemistry, organic, 34a–35a, 313a
geophysical log, 269b, 272b
geophysics, 290a–292a
interstitial-water chemistry, 380b–383b
lithofacies, 86b, 88b–92b
lithology, 143b–145b, 273b, 379b–380b
lithology, log-inferred, 286b–287b
glaciomarine sequences, 283b–285b
lithostratigraphy, 295a–304a, 408b, 877b
acoustic unit correlation, 99b–101b
diagenesis, 303a
lithologic Unit I, 93b, 127b, 289a, 294a, 296a,
304a, 339a
lithologic Unit II, 93b, 127b, 269b, 289a,
294a, 296a, 299a, 304a, 339a
lithologic Unit III, 95b, 109b, 127b, 269b,
289a–290a, 294a–295a, 299a–301a, 304a,
339a
lithologic Unit IV, 95b, 127b–128b, 269b,
290a, 295a, 304a, 339a
lithologic Unit V, 95b, 128b, 269b, 290a,
302a–303a, 304a, 339a, 881b
logging summary, 297a, 381b
summary, 296a
living bacterial density changes, 687b–689b
location, 6b, 28b, 58b, 81b, 135b, 144b, 169b,
264b, 289a, 290a, 300b, 350b, 376b, 751b,
814b, 871b
logging, 269b
caved-noncaved zone alteration, 328a–329a
correlation, Prydz Bay: Site 742, 442a, 443a,
448a–449a
lithoporosity combination, 39a–40a, 328a–
329a
quality, 328a
seismic stratigraphic combination, 39a, 329a
stratigraphic, 177b
summary, 102b, 342a–344a
synthetic seismogram, 40a–41a, 331a–333a
units, 283b–284b, 329a–330a, 339a
magnetic properties, 33a–34a, 45a, 305a,
308a–310a
magnetobiochronology, 837b
Oligocene–Miocene unconformity, 97b, 118b,
330a, 884b, 888b
opal, 84b
organic matter, 81b–82b
source and maturity, 407b–410b
oxygen isotopes, bivalve shells, 742b
palynology, 942b, 944b–945b
preglacial-glacial transition, 837b
sedimentary succession, 106b
sedimentation
during continental shelf progradation, 115b

during nonprogradation of continental shelf,
115b
glacial processes, 303a
sedimentation rate, 310a
seismic stratigraphy, 45a, 94a, 333a–338a
strontium isotopes, mollusk shells, 739b–742b
Prydz Bay: Site 740
biostratigraphy, 356a–357a
radiolarians, 519b
zonation, 24a, 26a–33a
depositional environment, 101b–115b, 345a,
374a
fluvially influenced, 355a–356a
geochemistry, inorganic, 35a, 358a, 360a–362a
geochemistry, organic, 34a–35a, 363a
geophysics, 347a–349a
sedimentary section, 348a–349a
lithology, 134b, 143b–145b, 747b–748b
lithostratigraphy, 349a–356a, 383b, 836b
acoustic unit correlation, 99b–101b
coarsening-upward sequence, 356a, 374a
fining-upward sequence, 351a, 353a, 356a,
374a
lithologic Unit I, 93b, 125b, 345a, 350a–351a,
360a, 372a, 374a
lithologic Unit II, 345a, 350a, 351a, 374a
lithologic Unit III, 345a, 350a, 351a–355a,
374a
physical properties correlation, 369a
red bed sequence, 374a
summary, 351a
magnetic properties, 33a–34a, 45a, 357a–358a
magnetobiochronology, 836b
overbank environment, waterlogged, 49b–50b
palynology, 941b, 942b
Pleistocene–Holocene hiatus, 99b
red beds
correlation, Permian Amery Group, 797b
distribution, 806b
magnetic properties, 797b–806b
sedimentary succession, 47b–50b
fluvial deposition, 54b
sedimentation rate, 358a
seismic stratigraphy, 45a, 95a, 369a–372a
reflection profiling, 7b
unconsolidated sediments, Holocene
chronology, 747b–750b
Prydz Bay: Site 741
biostratigraphy, 382a–383a
radiolarians, 519b
zonation, 24a, 26a–33a
coal-bearing sediment, 57b–60b
angular unconformity, 57b
depositional environment, 101b–115b, 377a,
381a
acoustic Unit PS.2b, 22b
fluvially influenced, 394a–395a
nonmarine, 393a
preglacial, 18b
geochemistry, inorganic, 35a, 385a
geochemistry, organic, 34a–35a, 385a–387a
geological setting, 79b, 133b
geology, 170b
geophysics, 378a–380a
interstitial-water chemistry, 383b, 384b–385b
lithology, 134b, 143b–145b, 383b–384b
lithostratigraphy
acoustic unit correlation, 99b–101b
fining-upward sequence, 381a, 382a, 394a
lithologic Unit I, 93b, 125b–126b, 377a, 380a–
381a, 394a
lithologic Unit II, 93b, 126b, 377a, 380a–
381a, 394a

lithologic Unit III, 93b, 126b, 377a, 380a–
381a, 394a
lithologic Unit IV, 377a, 380a–381a, 394a
magnetic properties, 33a–34a, 45a, 384a–385a
magnetobiochronology, 836b–837b
organic matter, source and maturity, 410b–411b
palynology, 941b, 942b–944b
Pleistocene–Holocene hiatus, 99b
sedimentation rate, 385a
seismic stratigraphy, 45a, 96a, 392a–393a
Prydz Bay: Site 742, 110a
biostratigraphy, 413a–414a, 845b
radiolarians, 519b
zonation, 24a, 26a–33a
carbonate-biosiliceous facies transition
Neogene, 888b
Paleogene, 885b
depositional environment, 101b–115b,
412a–413a, 448a–451a
fluvial/lacustrine, 452a
glacial, 18b–20b
glaciomarine, 412a
lacustrine, 413a
nonmarine, 397a
preglacial, 18b
fossil remains, 409a–410a
geochemistry, inorganic, 354a, 415a, 418a–420a
geochemistry, organic, 34a–35a, 420a–422a
geophysical log, 270b
geophysics, 398a–400a, 447a
reflector EE, 397a
glacial-preglacial unconformity, 880b, 882b
hiatuses, flat-lying sequence, 97b, 99b
interstitial-water chemistry, 385b–387b
lithofacies, 86b, 88b–92b
lithology, 134b, 143b–145b, 271b, 285b
summary, 878b, 879b
lithology, log-inferred, 285b–286b
glaciomarine sequences, 279b–283b
lithostratigraphy, 385b, 877b
acoustic unit correlation, 99b–101b
biogenic/clastic facies mixing, 412a
carbonate-rich zones, 407a, 409a
coarsening-upward sequence, 407a
correlation, physical properties, 426a
glacial-preglacial sediment unconformity,
877b
lithologic Unit I, 93b, 126b, 402a, 403a, 412a,
448a
lithologic Unit II, 93b, 126b, 269b, 402a,
403a–404a, 412a, 448a
lithologic Unit III, 93b, 108b, 126b, 269b,
402a, 404a–405a, 412a, 448a
lithologic Unit IV, 93b, 126b, 269b, 402a,
405a, 412a, 448a–449a
lithologic Unit V, 93b, 126b, 269b, 402a,
405a, 407a, 409a, 412a, 449a–450a
lithologic Unit V/Unit VI contact, 97b, 127b,
130b
lithologic Unit VI, 93b, 110b, 126b–127b,
269b, 402a, 409a–412a, 413a, 450a–451a,
881b
preglacial sequence, 397a
seismic stratigraphic correlation, 444a
summary, 386b, 406a–407a
logging, 269b, 451a
compaction and dewatering effects, 442a
correlation, Prydz Bay: Site 739, 442a, 443a,
448a–449a
lithoposity, 39a–40a, 438a, 439a
low-velocity rock, 441a, 442a
seismic stratigraphic combination, 39a, 437a
stratigraphic, 177b
summary, 101b, 455a–458a

SUBJECT INDEX

- synthetic seismogram, 40a-41a, 437a-442a
unconformities, 442a
units, 281b-282b, 283b, 436a-437a, 838b, 882b
magnetic properties, 33a-34a, 45a, 415a
magnetobiochronology, 837b-846b
magnetostratigraphy, 759b-760b
opal, 84b
organic matter, 81b-82b
palynology, 941b-942b, 943b-944b
Pleistocene–Holocene hiatus, 99b
sedimentary succession, 105b
sedimentation, during continental shelf
 progradation, 115b
sedimentation rate, 415a
seismic stratigraphy, 45a, 97a–99a, 443a-447a
Prydz Bay: Site 743
 biostratigraphy, 465a
 radiolarians, 519b
 zonation, 24a, 26a–33a
 depositional environment, 101b-115b, 462a, 465a
 glacial, 18b-20b
 vs. paleodeposition, 473a
 depositional processes, 475a
 geochemistry, inorganic, 35a, 465a-466a
 geochemistry, organic, 34a-35a, 466a
 geophysics, 460a
 lithology, 134b, 143b-145b, 385b
 lithostratigraphy
 acoustic unit correlation, 99b-101b
 lithologic Unit I, 95b, 128b, 459a, 461a, 462a, 475a
 lithologic Unit II, 95b, 128b, 459a, 461a, 462a, 475a
 lithologic Unit III, 95b, 128b, 459a, 461a, 462a, 475a
 Pleistocene–Holocene hiatus, 99b
 summary, 465a
 magnetic properties, 33a-34a, 45a, 465a
 seismic stratigraphy, 45a, 100a-101a,
 473a-474a
Prydz Channel
 topography, 64b
 trough mouth fan, 80b
Pumice, Kerguelen–Heard Plateau N, 130a
Pyrite
 Kerguelen–Heard Plateau N, 359b
 Prydz Bay: Site 739, 134b, 303a
 Prydz Bay: Site 740, 45b
 Prydz Bay: Site 741, 381a, 395a
 Prydz Bay: Site 742, 409a
Pyroxene, Kerguelen–Heard Plateau S: Site 738,
 augitic, 301b

Quartz
 Kerguelen Plateau S: Site 744
 in ice-raftered sediments, 202b
 Oligocene peak, glaciation onset, 217b
 Kerguelen Sediment Ridge, 237b, 510a-511a
 glacial/interglacial cycles, 240b
 origin, 514a
 round grains, 229b, 231b
 Kerguelen–Heard Plateau S: Site 738, 215b,
 217b
 in ice-raftered sediments, 202b
 Prydz Bay: Site 739, fluvial reworking, 115b
 Prydz Bay: Site 740, 46b, 50b, 56b
Quartz, fuchsite, Prydz Bay, provenance, 139b
Quartz-micaceous-chlorite rock, Prydz Bay: Site
 740, 56b
Quartz-siltstone, metamorphosed, Prydz Bay: Site
 739, 138b

Quartz/feldspar ratio, Kerguelen Sediment Ridge,
 229b
Quartzite, Prydz Bay, 137b, 354a
Quartzite, fuchsite
 Prince Charles Mountains, 65b
 Prydz Bay, 138b, 453a

Radioactivity
 Kerguelen–Heard Plateau N: Site 737, as clay
 indicator, 274b, 275b
 Prydz Bay: Site 742, glaciomarine sediments,
 281b
Radok Conglomerate
 deposition, 54b
 Permian sediments, source, 797b
Radok Lake, Pagodroma Tillite, overlying Permian
 strata and Proterozoic basement, 888b
Raggatt Basin
 Cretaceous lavas, 293b
 seismic stratigraphy, Units A–F, 277a, 279a,
 280a
Rare earth elements
 Kerguelen–Heard Plateau S: Site 738
 basalt, 307b
 chondrite-normalized, 315b
 Cretaceous/Tertiary boundary, 726b
 neutron activation analysis, 722b
 vs. other Mesozoic Southern Hemisphere,
 317b
Red beds
 Prydz Bay, 53b
 Prydz Bay: Site 740, 376b, 383b
 blocking temperature, 803b-804b
 correlation, Flagstone Bench Formation, 54b
 correlation, Permian Amery Group, 797b,
 804b
 depositional environment, 797b
 intrusives, magnetic effect, 806b
 lithologic Unit III, 47b
 magnetic properties, 797b-804b
 paleolatitude, 795b, 806b
Resistivity, Kerguelen–Heard Plateau S: Site 738,
 logging data, 279b
Réunion Subchron
 Kerguelen Sediment Ridge: Site 745, 753b
 Kerguelen–Heard Plateau N: Site 737, 754b
Rhythmite
 Prydz Bay, 108b, 127b
 mudstone contact, 130b
Rift graben, Prydz Bay, deposition, 20b-21b
Rift graben system, continental margins, depositional
 environments, 18b
Rifting, Prydz Bay, rift-onset unconformity, 20b,
 22b
Rock flour, Prydz Bay, physical properties and
 content of, 157b
Ross continental shelf, glaciomarine sediments,
 Oligocene, 79b
Ross Ice Shelf, Ice Stream B, 115b
Ross Sea
 Antarctic Bottom Water, 232b
 continental glaciation, Oligocene, 201b
 diamictites, organic carbon content, 113b
 Dry Valley Drilling Project
 Miocene–Pliocene glaciation, 118b-119b
 Pliocene–Pleistocene record, 79b
 glacial sediment, Cenozoic, 10a, 340a, 872b
 glaciation, Oligocene, oxygen isotopic
 evidence, 695b
 ice sheets, temperate Oligocene climates, 121b
 ice shelf, expansion, 248b
 ice-raftered debris, Oligocene, 873b
 Miocene unconformity, ice loading event 2,
 885b
mudstone, ice-raftered detritus, 886b
organic matter, recycling, 417b
shallowing event, Oligocene unconformity,
 872b, 885b
siliceous surface sediments, 112b
Rubidium, Kerguelen–Heard Plateau S: Site 738,
 basalts, 315b

Salinity
 high altitudes, 870b
Kerguelen Plateau S: Site 744, 491a
 interstitial waters, 354b-355b
Kerguelen Sediment Ridge, 516a, 544a
 interstitial waters, 354b-355b
 sonobuoy stations, 107a
Kerguelen–Heard Plateau N, 139a, 185a
 interstitial waters, 354b-355b
 sonobuoy stations, 107a
Kerguelen–Heard Plateau S: Site 738, 110a, 255a
 interstitial waters, 354b-355b
 sonobuoy stations, 107a
Prydz Bay, 110a, 310a-312a, 360a, 418a
 interstitial waters, 378b, 383b
 sonobuoy stations, 107a
Samarium, Kerguelen–Heard Plateau S: Site 738,
 scandium-normalized distribution, 725b
Samarium/lanthanum ratio, Kerguelen–Heard Plateau
 S: Site 738, Cretaceous/Tertiary boundary, 726b
Sand
 Kerguelen Plateau S: Site 744, grain size
 distribution, 191b
 Kerguelen–Heard Plateau S: Site 738, grain size
 distribution, 191b
 Prydz Bay, 302a, 328a
 depositional environment, 475a
 distribution, 81b, 149b
 mineralogy, 410a
 seismic profile, 444a
Sand, black, Kerguelen–Heard Plateau N: Site
 737, 165a
Sand, diatomaceous, Prydz Bay: Site 743, 462a
Sand, greenish, Prydz Bay: Site 740, physical
 properties, 369a
Sand, terrigenous
 Kerguelen Plateau S: Site 744, grain-size
 maxima, 217b
 Kerguelen Sediment Ridge
 chemical composition, 229b
 facies A vs. facies B, 228b
 gravel content and, 242b, 249b
 ice-raftered transport, 249b
 sea level control, 249b
Sand, volcanic, Kerguelen–Heard Plateau N: Site
 737, 168a
Sand, well-sorted, Prydz Bay: Site 743, 128b
Sand-silt, diatomaceous, Prydz Bay: Site 742,
 93b, 126b, 403a
Sand-silt-claystone, Prydz Bay, 300a, 409a, 410a
Sand/mud ratio, Prydz Bay, by facies type, 95b,
 96b
Sandstone
 Prydz Bay: Site 740, 383b
 color and redox potential, 45b, 49b
 cross-bedding, 46b
 depositional environment, 49b
 fining-upward trend, 45b
 magnetic minerals, 51b
 mineralogy, 352a
 petrography, 50b-52b
 provenance, 51b-52b
 red bed sequence, 351a-352a, 354a, 356a,
 365a
 rippled, 49b

rock fragments in, 50b
 Prydz Bay: Site 741, 381a, 385a
 depositional environment, 57b
 fining-upward sequences, 57b
 petrography, 57b
 seismic stratigraphy, 393a, 394a
 Spreiten-type burrows, 384a
 Prydz Bay: Site 742, 89b, 410a
 Sandstone, compacted subarkosic, Prydz Bay, 137b
 Sandstone, muddy, Prydz Bay: Site 742, 89b
 Sandstone, subarkosic, Prydz Bay, provenance, 138b-139b
 Sandstone, subarkosic, plant-bearing
 Prydz Bay
 carbonate-altered, 135b, 137b, 141b
 micaceous, 137b
 Sandstone, well-sorted
 Prydz Bay
 carbonate-cemented, 137b
 origin, 107b-108b
 Sandwich Islands S, island-arc tholeiitic series, 328b
 Santa Barbara Basin, iridium accumulation rate, 855b
 Saponite, Kerguelen-Heard Plateau S: Site 738,
 conditions of formation, 307b, 313b
 Saponite, brown, Kerguelen-Heard Plateau S: Site 738, 301b
 Saponite, green, Kerguelen-Heard Plateau S: Site 738, 301b
 Scandium
 Kerguelen-Heard Plateau S: Site 738
 calcite-free data, 728b-729b
 Cretaceous/Tertiary boundary, 722b-723b, 724b
 Schist, garnet-biotite-magnetite, Prydz Bay: Site 739, 453a
 Sea level
 Pliocene deglaciation, enhanced iceberg calving, 249b
 See also Eustatic sea level
 Sea-surface temperature
 Kerguelen-Heard Plateau S: Site 738, Eocene
 vs. modern, 707b-708b
 Paleocene, surface-to-bottom thermal stratification, 714b
 Prydz Bay: Site 743, oxygen isotope correlation, Oligocene, 742b
 Southern Ocean, 551a-553a
 tropical waters, vs. bottom water temperatures, 870b
 Seafloor spreading, Indian-Antarctic continental separation, breakup unconformity, 22b
 Sediment accumulation rate
 Kerguelen Plateau S: Site 744,
 oxidation-reduction potential and, 367b
 Kerguelen Sediment Ridge, 373b
 Kerguelen-Heard Plateau S: Site 738, organic carbon correlation, 364b
 noncarbonate, Kerguelen-Heard Plateau S: Site 738, Cretaceous/Tertiary boundary, 854b
 Sedimentary cycles
 Kerguelen Sediment Ridge, clay mineralogy, 228b-229b
 Pliocene-Holocene, 249b
 Sedimentary structures
 Kerguelen Sediment Ridge: Site 745, diagenetic origin, 513a
 Prydz Bay: Site 740
 planar sets, 352a-353a, 355a
 trough sets, 352a-353a
 Prydz Bay: Site 741, 382a

Sedimentation, Kerguelen-Heard Plateau N: Site 736, post-compaction, 828b
 Sedimentation rate
 ammonia pore water gradient and, 402b-403b
 estimating, 403b-404b
 Kerguelen Plateau N, 353b
 thermal gradient and, 372b
 Kerguelen Plateau S: Site 744, 490a
 post-compaction, 820b-821b, 821b-822b
 Kerguelen Sediment Ridge, 516a, 544a
 post-compaction, 818b, 819b
 Kerguelen-Heard Plateau N: Site 736, 136a
 heat flow and, 153a
 Kerguelen-Heard Plateau N: Site 737, 183a,
 185a, 191a, 754b
 post-compaction, 818b-836b
 Kerguelen-Heard Plateau S: Site 738,
 253a-254a, 256a
 Cretaceous/Tertiary boundary, 852b-853b
 post-compaction, 822b, 825b
 thermal gradient and, 372b
 vs. compaction rate, 272a-273a
 organic carbon content and, 401b
 Prydz Bay, 310a, 358a, 385a, 415a
 organic carbon correlation, 376b
 Pleistocene-Holocene, 748b-750b
 post-compaction, 837b
 reservoir correction, 748b, 749b
 silty clay vs. siliceous ooze, 749b
 Site 739, seaward edge of grounded ice sheet, 744b
 sulfate pore water gradient and, 402b
 Seismic reflection profiles
 Kerguelen Plateau, 7a
 Kerguelen Plateau S: Site 744, 45a
 Kerguelen Sediment Ridge, 45a, 535a, 550a
 Kerguelen-Heard Plateau N: Site 736, 45a,
 124a, 126a, 127a, 154a
 Kerguelen-Heard Plateau N: Site 737, 45a,
 160a-161a
 Kerguelen-Heard Plateau S: Site 738, 45a,
 230a-232a, 233a
 Prydz Bay, 9b
 Prydz Bay: Site 739, 19b, 45a
 vs. logging data, 10b
 Prydz Bay: Site 740, 45a, 373a, 374a
 Prydz Bay: Site 741, 45a
 Prydz Bay: Site 742, 19b, 45a
 vs. logging data, 10b
 Prydz Bay: Site 743, 45a
 Seismic stratigraphy
 data processing and availability, 23b-24b
 digital magnetic tape parameters, 106a
 Kerguelen Plateau, discordance A, 7a
 Kerguelen Plateau S: Site 744, 102a
 correlation, Kerguelen-Heard Plateau S: Site 738, 501a
 lithostratigraphic correlation, 501a
 units, 500a
 Kerguelen Sediment Ridge, 103a-104a,
 525a-526a, 534a, 548a
 bottom current influence, 525a, 526a, 530a
 Kerguelen-Heard Plateau, units, 124a
 Kerguelen-Heard Plateau N: Site 736, 90a,
 153a-155a, 828b
 correlation, lithostratigraphy, 156a
 discordance A, 124a, 153a
 mounds, 154a-155a
 units, 154a-155a
 Kerguelen-Heard Plateau N: Site 737, 91a-92a
 Eocene-Miocene unconformities, 218a
 erosional unconformity, 215a
 mounds, 214a
 reflector A, 159a

synthetic vs. observed seismic-reflection traces, 214a, 274a, 277a
 units, 160a, 214a-216a
 Kerguelen-Heard Plateau S: Site 738, 93a,
 276a-279a
 basement arch, 276a-277a
 correlation, Kerguelen Plateau S: Site 744,
 501a
 units, 276a, 280a
 Prydz Bay, 9a, 11a, 12a
 basement high, 18b, 22b
 discontinuous reflections, midshelf swale, 11b
 flat-lying sequence, 876b
 geologic noise, 7b
 intersite correlations, 445a, 446a-447a, 450a,
 452a
 seismic-refraction profiles 5-7, 36b, 38b, 40b
 structured features, 17b-18b
 Prydz Bay: Site 739, 94a, 333a-338a
 dip changes, 9b, 18b
 Eocene, 882b
 glacial strata, 19b
 logging data, 127b-128b, 288b
 low-velocity layers, 336a
 noise sources, 334a
 random events, 334a
 synthetic vs. observed seismic-reflection traces, 332a
 units, 334a-336a, 338a, 339a-340a
 Prydz Bay: Site 740, 95a
 compacting sand/shale sequence, 370a
 dip orientation, 797b
 nonmarine sequence, 372a
 units, 346a, 369a, 370a
 Prydz Bay: Site 741, 96a
 Cretaceous unconformity, 393a
 geologic variations, 25b
 sedimentary section thickness, 393a, 395a
 units, 377a, 392a-393a
 Prydz Bay: Site 742, 97a-99a
 discontinuous reflections, 441a, 443a
 Eocene, 882b
 line PB-21, 398a
 lithostratigraphic correlation, 444a, 451a
 logging data, 126b-127b, 288b
 nonmarine sequence, 445a
 reflection peaks, 441a-442a
 seismically distorted unit, 877b, 880b
 structures, 446a
 synthetic vs. observed seismic-reflection traces, 439a-441a, 442a
 units, 443a-445a, 447a, 451a-452a
 Prydz Bay: Site 743, 100a-101a
 contemporary vs. paleodepositional environments, 473a-474a
 units, 473a, 475a
 system geometry, 106a
 Sericite, Prydz Bay: Site 740, clay matrix, 50b
 Shale, carbonaceous, Prydz Bay: Site 742, glaciectonic origin, 877b
 Shear strength
 Kerguelen Plateau S: Site 744, 38a, 495a, 499a
 Kerguelen Sediment Ridge, 38a, 520a,
 522a-524a, 547a-548a
 Kerguelen-Heard Plateau N: Site 736, 38a,
 145a, 148a, 149a, 153a, 197a
 Kerguelen-Heard Plateau N: Site 737, 38a,
 195a, 197a, 199a, 202a, 203a
 Kerguelen-Heard Plateau S: Site 738, 38a, 264a,
 267a, 268a
 Prydz Bay, 38a, 145b, 146b, 148b, 156b, 158b,
 365a, 428a, 469a-470a
 diamictites, 154b-155b
 negative pore pressures, 155b, 156b

SUBJECT INDEX

- Shetland Islands S, glaciomarine elastic deposits, Paleocene–Eocene, 870b
- Sidufjall event, Kerguelen Sediment Ridge: Site 745, 818b
- Silica
formation, controlling factors, 219b
high-resolution vs. regular sampling results, 398b
Kerguelen Plateau S: Site 744, 491a
Miocene cooling event, 218b
opal contents and, 217b
physical property correlation, 499a
- Kerguelen Sediment Ridge, 517a, 544a
- Kerguelen-Heard Plateau N: Site 736, 137a–138a, 139a, 156a
- Kerguelen-Heard Plateau N: Site 737, 186a
clinoptilolite and diagenesis of, 218b
discontinuities, 187a
Miocene cooling event, 218b
shear strength and, 197a
vertical concentration gradient, 363b
- Kerguelen-Heard Plateau S: Site 738, 255a–256a
chert and precipitation of, 254a
clinoptilolite and diagenesis of, 218b
diagenesis, 364b
Miocene cooling event, 218b
pore waters, 372b
- Prydz Bay: Site 739, 312a
- Prydz Bay: Site 740, 362a
- Prydz Bay: Site 741, 385a
- Prydz Bay: Site 742, 418a
pore waters, 386b
- Prydz Bay: Site 743, 466a, 475a
- Silica, aqueous
Kerguelen Plateau S: Site 744, pore waters, 367b, 369b
- Kerguelen Sediment Ridge, 371b
- Kerguelen-Heard Plateau S: Site 738,
pore-water chemistry anomaly, 366b–367b
- Prydz Bay
pore waters, 381b, 384b–385b
solute diffusion inhibition, 387b–388b
- Silica, authigenic, Prydz Bay: Site 740, lack of, 52b
- Silica, biogenic
Kerguelen Plateau S: Site 744
carbonate correlation, Oligocene, 884b
fluxes, 114a
- Kerguelen-Heard Plateau N: Site 737, 173a
- Prydz Bay, fluxes, 114a
- Silica/aluminum ratio, opal precipitation and, interstitial waters, 218b
- Silicate, authigenic, Southern Ocean, formation, 218b–219b
- Silicate, sheet
Kerguelen-Heard Plateau S: Site 738
brown sheet phases, 301b, 302b
chemical composition, 305b–306b, 308b, 313b
- Siliceous microflora, Prydz Bay, cold water characteristics, 112b
- Silicon, Kerguelen-Heard Plateau S: Site 738, in sheet silicate formation, 313b
- Silicon dioxide, Kerguelen-Heard Plateau S: Site 738, 301b
- Sillimanite needles
Prydz Bay: Site 740
in quartz grains, 56b
in sandstones, 50b–51b
- Silt
Kerguelen Plateau S: Site 744, grain size distribution, 191b, 193b
- Kerguelen Sediment Ridge, 513a
current activity and deposition of, 514a, 542a
turbiditic origin, 231b
- Kerguelen-Heard Plateau N: Site 737, turbiditic origin, 285b
- Kerguelen-Heard Plateau S: Site 738, grain size distribution, 191b
- Silt, clayey
Prydz Bay, 372a, 374a, 381a, 462a
laminations, 301a
origin, 475a
physical properties, 365a
- Silt, terrigenous, Kerguelen Plateau S: Site 744, grain-size maxima, 217b
- Siltstone
Prydz Bay: Site 740
depositional environment, 356a
laminated, 49b
mottled zones, 46b–47b, 53b
petrography, 50b–52b
sedimentary structures, 353a–354a
- Prydz Bay: Site 741, 381a
seismic stratigraphy, 393a
- Prydz Bay: Site 742, 410a
- Siltstone, clayey, Prydz Bay, 385a, 405a
- Siltstone, laminated, Prydz Bay: Site 739, 127b
- Siltstone, sandy, Kerguelen-Heard Plateau N: Site 737, 168a
- Siltstone, subarkosic, plant-bearing, Prydz Bay, carbonate-altered, 135b, 137b
- Siltstone-claystone, Prydz Bay: Site 742, 405a, 881b
- Sirius Formation, microfossils, Pliocene to Quaternary emplacement, 120b
- Slumping
Prydz Bay, 301a, 468a–469a
diamictite, 112b, 132b
- Smectite
from chemically weathered volcanics, 198b
- Kerguelen Plateau S, oxygen isotope correlation, 199b–200b
- Kerguelen Plateau S: Site 744
climatic variation, 202b
negative illite correlation, 194b, 199b
peak development, 199b
- Kerguelen Sediment Ridge, 244b
vs. illite, Pliocene–Holocene, 249b
- Kerguelen-Heard Plateau S: Site 738
climatic variation, 202b
negative illite correlation, 194b
- Prydz Bay, 86b
plasticity and water content relationship, 150b
- Sodium
Kerguelen Plateau S: Site 744
pore waters, 930b–931b
vertical distribution gradient, 367b
- Kerguelen Sediment Ridge, pore waters, 930b–931b
- Kerguelen-Heard Plateau N
chemical gradient, 359b
pore waters, 372b, 929b, 929b–930b
vertical distribution gradient, 363b
- Kerguelen-Heard Plateau S: Site 738, pore waters, 930b
- Prydz Bay
pore waters, 380b, 930b–931b
vertical distribution gradient, 380b–381b, 385b, 388b, 391b
- Sonobuoy seismic recording
equipment and parameters, 107a
- Kerguelen Sediment Ridge: Site 745, 506a, 507a, 508a, 509a
- Kerguelen-Heard Plateau N: Site 736, 125a, 126a
- Kerguelen-Heard Plateau N: Site 737, 160a–162a
- Kerguelen-Heard Plateau S: Site 738, 230a–232a, 233a
procedures, 45a–46a
- Prydz Bay
deployment sites, 28b
ray-trace model, 28b–35b, 37b, 39b, 41b–43b
sonobuoys 4–8, 29b–42b
- Prydz Bay: Site 739, 291a–292a, 293a, 294a
correlation, Sites 739 and 742, 31b
curved vs. linear refraction arrivals, 292a
- Prydz Bay: Site 740, 347a–348a
- Prydz Bay: Site 741, 378a–380a
noise levels, 378a
refraction arrivals, 378a–379a
secondary reflections, 378a
- Prydz Bay: Site 742, 398a–430a, 401a, 402a
calculated vs. observed values, 29b–30b
curved refraction arrivals, 398a, 400a
top-down modeling, 29b–30b, 33b
wide-angle reflections, 398a
- Sørdsal Glacier
ice cover, 749b
outflow, 133b
- South Shetland Islands, glacial sediment, Eocene, 340a
- Southern Hemisphere, surface-water masses, 872b
- Southern Ocean
Eocene cooling trend, 201b
ice cover, climatic fluctuations, 230b
kaolinite, continental origin, 198b
magnetobiochronology, 813b–817b
Oligocene cooling trend, oxygen isotope correlation, 194b–195b
- Sri Lanka, Jurassic dolerites, magnetic properties, 806b
- Stable isotopes
Kerguelen Plateau S: Site 744, 114a–115a
Prydz Bay, 114a–115a
- Steranes, Prydz Bay: Site 739, gas chromatography, 408b–409b
- Stevns Klint, boundary clay, origin, 854b
- Strontium isotopes
Antarctica, ice growth events, 870b
Kerguelen Plateau S: Site 744, 826b–828b
analytical methods, 732b
foraminifer preservation, 732b–733b
magnetobiostratigraphic correlation, 736b
mixed planktonic foraminifers, 733b–736b
Neogene, 829b
Oligocene hiatus, 821b
Oligocene/Miocene boundary unconformity, 736b
Paleogene, 830b
- Kerguelen-Heard Plateau S: Site 738, basalts, 294b–295b, 297b, 298b
- Prydz Bay, seawater composition, 741b–742b
- Prydz Bay: Site 739, 844b
Eocene–Oligocene, 881b
mollusk shells, 739b–742b, 837b
pore waters, 740b–741b
seawater vs. pore waters, 881b
sources, 391b
- Strontium/calcium ratio, Kerguelen Plateau S: Site 744, 733b
- Subsidence history, Kerguelen-Heard Plateau S: Site 738, 194b, 280a
- Subtropical Convergence, biogeographic changes, 874b
- Sulfate
ammonia gradient and, relation to carbon flux, 401b, 403b
high-resolution vs. regular sampling results, 398b–399b
- Kerguelen Plateau S: Site 744, 491a

- Kerguelen Sediment Ridge, 518a, 544a
pore waters, 369b
- Kerguelen-Heard Plateau N: Site 736, 137a, 139a–140a
- Kerguelen-Heard Plateau N: Site 737, 186a–187a
calcium and alkalinity correlation, 362b
organic carbon correlation, 359b
pore waters, 218b
- Kerguelen-Heard Plateau S: Site 738, 257a
pore waters, 364b
pore water gradient, sedimentation rate and, 402b
- Prydz Bay: Site 739, 312a–313a
ammonium anticorrelation, 380b
- Prydz Bay: Site 740, 362a, 374a
pore waters, 388b
- Prydz Bay: Site 741, 385a
pore waters, 384b
- Prydz Bay: Site 742, 418a, 420a
ammonium anticorrelation, 385b
- Prydz Bay: Site 743, 466a
- Sulfate reduction
Kerguelen Plateau S: Site 744, lack of organic matter and, 367b
- Kerguelen Sediment Ridge
bacterial, 371b
magnesium-potassium uptake and, 371b
- Kerguelen-Heard Plateau N, 359b
anaerobic, 355b
- Kerguelen-Heard Plateau S: Site 738, lack of organic matter and, 366b, 372b
- Prydz Bay
bacterial, 380b
effect on pore water alkalinity, 388b
pore waters, 385b
sedimentation rate and, 401b
- Sulfide, Prydz Bay: Site 740, anoxic conditions, 383b
- Sulfur
Prydz Bay, 82b, 84b, 85b
erosion of preglacial sediments and, 113b
- Surface-water mass, Southern Ocean, present-day location, 9a
- Surface-water temperature, Kerguelen Plateau S: Site 744, isotopic correlation, Oligocene cooling trend, 711b–712b, 714b–715b
- Surging
Lambert Glacier-Amery Ice Shelf system, 62b
effect on debris distribution, 71b, 73b
- Svenner Channel, topography, 64b
- Tantalum
Kerguelen-Heard Plateau S: Site 738
calcite-free data, 728b–729b
scandium-normalized distribution, 725b
- Tantalum/lanthanum ratio, Kerguelen-Heard Plateau S: Site 738, basalts, 318b
- Tasman Sea, Oligocene hiatuses, 874b
- Tectonic history, Kerguelen Plateau, crustal structure, 5a–6a
- Tectonic uplift, Prydz Bay: Site 740, of source area, 52b, 53b
- Temperature
Kerguelen Plateau S: Site 744, 495a, 499a, 500a, 502a, 503a
- Kerguelen Sediment Ridge: Site 745, 524a, 533a, 534a
- Kerguelen-Heard Plateau N: Site 736, 150a–153a
- Kerguelen-Heard Plateau N: Site 737, 198a, 206a, 207a
drift rate, 205a
- Kerguelen-Heard Plateau S: Site 738, 110a
- Prydz Bay, 110a
Southern Ocean, sonobuoy stations, 107a
- Temperature gradient
Kerguelen Plateau N, sediment thickness and, 353b
- Kerguelen-Heard Plateau N: Site 737, 357b
- Tephra
Kerguelen Islands
basaltic vs. trachytic to rhyolitic, 329b, 333b
ice-rafterd transport, 333b
- Kerguelen Plateau, mineralogy, temporal changes, 325b
- Kerguelen Sediment Ridge
discrete layers, 326b, 331b
dispersed, 326b
geochemistry, 331b–332b
geochronology, 336b
grain size and magnetic separating, 328b
provenance, 329b, 333b
- Kerguelen-Heard Plateau N
discrete layers, 326b, 329b
dispersed, 326b
geochemistry, 330b–331b
geochronology, 336b
grain size and magnetic separating, 328b
provenance, 329b
- Tephra, calc-alkaline, Kerguelen Islands, comparison, South Sandwich Islands, 335b
- Terbium/lanthanum ratio, Kerguelen-Heard Plateau S: Site 738, Cretaceous/Tertiary boundary, 726b
- Terrigenous sediment
classification, grain size, 24a
- Kerguelen Sediment Ridge
glacial transport, 249b, 530a
glacial/interglacial cycles, 231b
grain-size distribution, 228b, 236b, 240b, 242b, 258b
gravel content, 243b, 244b
inverse opal correlation, 227b, 240b
- Kerguelen-Heard Plateau S: Site 738
Cretaceous, 200b
Eocene, 197b
- Terzaghi consolidation theory, principles, 172b, 174b
- Thermal conductivity
Kerguelen Plateau S: Site 744, 38a, 499a, 500a
with silica variations, 499a
- Kerguelen Sediment Ridge, 38a, 520a, 522a–524a, 548a
- Kerguelen-Heard Plateau N: Site 736, 38a
bimodal distribution, 150a
temperature, 153a
- Kerguelen-Heard Plateau N: Site 737, 38a, 197a–198a, 205a
bimodal distribution, 197a
- Kerguelen-Heard Plateau S: Site 738, 38a
Prydz Bay, 38a, 428a, 431a, 470a, 473a
- Thorium
Kerguelen-Heard Plateau S: Site 738
calcite-free data, 728b–729b
Cretaceous/Tertiary boundary, 725b–726b
- Thvera Subchron, Kerguelen Sediment Ridge: Site 745, 753b, 818b
- Till, INQUA classification, 103b
- Till, basal
clast orientation, 107b
deposition, 103b
- Till, glacial, Four Ladies Bank, 290a
- Till, lodgement
deposition, 103b
- Prydz Bay, 326a, 339a, 356a, 463a, 465a
correlation, Oligocene prograding sequence, 115b
- diamict deposition as, 107b
- Till, melt-out, deposition, 103b
- Till, waterlain
Prydz Bay
deposition, 103b
depositional effects, 412a
Eocene, 882b
Oligocene deposition, 118b
- Tillite, grounded and marine, depositional environment, 287b
- Titanomagnetite, Prydz Bay: Site 740, in sandstones, 51b, 56b
- Total alkalis vs. silica (TAS)
Kerguelen Sediment Ridge, discrete vs. dispersed tephras, 326b, 333b, 334b
- Kerguelen-Heard Plateau N, discrete tephra layers, 326b, 333b
- Trace elements, Kerguelen-Heard Plateau S: Site 738, basalts, 317b
- Transantarctic Mountains
glaciation
East Antarctic Ice Sheet development and, 340a
ice sheet development, 118b
uplift history and onset, 289a
- Transantarctic Mountains, Beacon Supergroup, kaolinitic sediments, 247b
- Tsunami event, Kerguelen-Heard Plateau S: Site 738, Cretaceous/Tertiary boundary, 853b
- Tuff
Kerguelen-Heard Plateau N: Site 736, 130a
depositional environment, 131a
- Turbidite, Kerguelen-Heard Plateau N: Site 736, disordered, 131a
- Unconsolidated sediments, Prydz Bay: Site 740, Holocene chronology, 747b–750b
- Uplift history, Prydz Bay, mechanism, 22b
- Uranium
Kerguelen-Heard Plateau S: Site 738
calcite-free data, 728b–729b
Cretaceous/Tertiary boundary, 725b–726b
- Vein, calcite, Kerguelen-Heard Plateau S: Site 738, in volcaniclastic rock, 244a
- Velocity
Kerguelen Plateau S: Site 744, 499a
refraction, 46a
- Kerguelen Sediment Ridge: Site 745, 520a, 522a–524a
sonobuoy stations, 107a
- Kerguelen Sediment Ridge: Site 746, 548a
- Kerguelen-Heard Plateau N: Site 736, 150a, 153a
sedimentary rock, 125a
- Kerguelen-Heard Plateau N: Site 737, 204a
discontinuities, 162a, 202a–203a
oscillations, 214a
synthetic seismogram, 212a
- Kerguelen-Heard Plateau S: Site 738, 264a, 266a–267a, 268a–269a
oscillations, 275a
sonic vs. sonobuoy seismic logs, 276a
- Lambert Glacier-Amery Ice Shelf system, surface measurements, 62b
- measurement errors, 46a
- Prydz Bay
acoustic Units PS.1–PS.5, 36b
glacially overconsolidated sediment, 31b
- Prydz Bay: Site 739, 145b, 148b, 319a
inversion, 292a, 400a
laboratory vs. downhole measurements, 326a, 327a
logging data, 8b, 19b

SUBJECT INDEX

sonobuoy seismic study, 30b-31b, 35b
 Prydz Bay: Site 740, 365a, 369a
 high-velocity refraction (R3), 33b
 reversal and high-amplitude seismic reflections, 372a
 sonobuoy seismic study, 32b-33b
 Prydz Bay: Site 741
 compaction and, 393a
 high-velocity refraction (R5 and R7), 34b, 36b
 scatter, 391a-392a
 sonobuoy seismic study, 33b-34b
 Prydz Bay: Site 742, 146b, 428a, 434a, 436a
 inversion, 400a
 logging data, 8b, 19b
 logging vs. laboratory measurements, 435a
 ray-trace model vs. smoothed log profile, 30b
 reversal, 434a
 small-scale fluctuations, 148b-149b
 sonobuoy seismic study, 29b-30b, 34b-35b
 Prydz Bay: Site 743, 146b, 148b, 469a-470a
 refraction, 46a
 rms and interval, 46a
 sonobuoy, 161a
 sonobuoy stations, 107a, 125a
 Velocity, acoustic, Prydz Bay, unconformity, 11b
 Velocity, interval
 Kerguelen Sediment Ridge: Site 745, 506a
 Prydz Bay, 378a, 400a
 sedimentary section, 398a
 Velocity, refraction
 Kerguelen-Heard Plateau N: Site 736, 125a, 127a
 Kerguelen-Heard Plateau S: Site 738, 233a
 acoustic basement, 232a
 Prydz Bay, 292a, 348a
 sedimentary section, 400a
 Vermiculite, Prydz Bay: Site 739, 86b
 Vesicle filling phases
 Kerguelen-Heard Plateau S: Site 738
 paragenesis A and B, 302b, 307b
 secondary minerals, chemical variation, 308b
 Vestfold Hills
 alkaline mafic igneous rock,
 Cretaceous-Eocene, 7b
 glaciomarine sediments, Pliocene, 886b-887b
 igneous and metamorphic rock, Precambrian age, 64b-66b
 Pliocene marine sediments, interglacial nature, 119b

sedimentary rock, Phanerozoic, 7b
 uplift, 22b
 Vitrinite
 Prydz Bay, 408b
 indigenous nature, 60b
 reflectance values, 410b
 Volcanic basement, Kerguelen-Heard Plateau N: Site 737, refraction velocity, 161a
 Volcanic detritus
 Kerguelen-Heard Plateau N, 172a-173a, 353b
 depositional environment, 130a-131a
 geochemical correlation, 187a-188a
 Volcanic rock, Kerguelen-Heard Plateau S: Site 738, physical properties, 266a
 Volcaniclastic rock
 Kerguelen-Heard Plateau S: Site 738, 241a-243a
 with calcite veins, 244a
 Volcaniclastic sediment, Kerguelen-Heard Plateau S: Site 738, secondary phases, 364b
 Volcanogenic sediment, Kerguelen-Heard Plateau N: Site 737, 217b
 Wacke, arkosic
 Prydz Bay: Site 740, 56b
 petrography, 50b, 51b-52b, 797b
 Water content
 Kerguelen Plateau S: Site 744, 495a, 906b
 with silica variations, 499a
 Kerguelen Sediment Ridge: Site 745, 520a, 522a-524a, 906b
 Kerguelen Sediment Ridge: Site 746, 546a-547a, 906b
 Kerguelen-Heard Plateau N: Site 736, 144a-145a, 148a, 153a, 904b
 Kerguelen-Heard Plateau N: Site 737, 904b-905b
 Kerguelen-Heard Plateau S: Site 738, 262a, 264a, 266a, 905b
 Prydz Bay, plasticity and, 150b
 Prydz Bay: Site 739, 145b, 148b, 319a, 325a, 905b-906b
 silica and, 312a
 Prydz Bay: Site 740, 365a
 diatom ooze, 383b
 Prydz Bay: Site 742, 146b, 428a, 434a
 Prydz Bay: Site 743, 146b, 148b, 469a-470a
 Water/rock ratio
 Kerguelen-Heard Plateau S: Site 738
 for alteration, 314b

basalts, 313b
 Weathering
 Antarctica E, chemical-to-physical transition, 201b
 Kerguelen Plateau S: Site 744, chemical-to-physical transition, 884b
 Kerguelen-Heard Plateau S: Site 738, clay mineralogy effects, 199b
 Prydz Bay
 geochemical effects of, 391b
 stable isotope effects of, 381b
 Weathering, physical
 Kerguelen Plateau S: Site 744, glacial cooling trend, 199b
 Prydz Bay, sedimentary rock source area, 115b
 Weddell Sea
 carbon isotopes, comparison, Prydz Bay and equatorial Pacific Ocean, 704b, 714b
 deep-sea hiatuses, during increased AABW flow, 874b-875b
 glaciation, Pliocene, 181b
 ice rafting, Oligocene, 78b
 ice shelf, expansion, 248b
 ice-raftered debris, 340a
 Oligocene, 873b
 isotopic record, Oligocene inversion, 712b
 organic carbon, Cretaceous, 419b
 organic matter, recycling, 417b
 sedimentary facies, cyclic changes, Pleistocene, 225b
 stable isotopes, 694b
 Weddell Sea Bottom Water, grain size effects, 225b
 Wilkes Land margin, organic matter, recycling, 417b
 Zinc
 Kerguelen-Heard Plateau S: Site 738
 Cretaceous/Tertiary boundary, 724b-725b
 scandium-normalized distribution, 724b
 Zirconium/nickel ratio
 Kerguelen-Heard Plateau S: Site 738, 294b
 in MORB-type basalts, 317b, 318b
 Zirconium/yttrium ratio, Kerguelen-Heard Plateau S: Site 738, comparison, other MORB-type basalts, 318b

SITE INDEX

| | | |
|---|--|--|
| CIROS-1 | Site 237 stable isotopes Paleogene, 695b | glaciation Miocene/Pliocene unconformity, 118b surface-water masses, 872b |
| diamicrites grain-size distribution, 120b-121b | | |
| diatoms, 577b reworking, 744b | | |
| glaciation advances and retreats, 872b | | |
| Oligocene–Miocene record, 79b, 118b | | |
| Oligocene/Miocene unconformity, 872b | | |
| stratigraphic intervals, 119b | | |
| surface-water masses, 872b | | |
| CIROS-2 | Site 251 surface-water masses, 872b | Site 274 <i>Eucyrtidium antiquum</i> , n. sp., 546b ice rafted debris, 248b Oligocene, 873b surface-water masses, 872b |
| glaciation advances and retreats, 79b | | |
| glaciogenic sediments Pliocene, 119b | | |
| MSSTS-1 | Site 252 surface-water masses, 872b | Site 275 surface-water masses, 872b |
| glacial sediments, 872b | | |
| glaciation Miocene/Pliocene unconformity, 118b | | |
| glaciomarine sediments Oligocene, 79b | | |
| stratigraphic intervals, 119b | | |
| surface-water masses, 872b | | |
| Site 11 | Site 253 carbon isotopes <i>Cibicidoides</i> spp., 709b stable isotopes Paleogene, 695b | Site 277 oxygen isotopes Eocene–Holocene latitudinal gradient, 707b, 709b foraminifers, 620b, 869b stable isotopes Eocene/Oligocene boundary hiatuses, 710b- 711b surface-water masses, 872b |
| stratigraphic intervals, 119b | | |
| Site 19 | Site 264 surface-water masses, 872b | Site 278 <i>Botryostrophus kerguelensis</i> , n. sp., 545b <i>Denticulopsis dimorpha</i> Acme Zone, 556b <i>Denticulopsis hustedtii</i> first occurrence, 559b <i>Denticulopsis maccollumii</i> Partial Range Zone, 556b diatoms datum events, stratigraphic constraints, 588b Miocene sequence, 577b surface-water masses, 872b |
| oxygen isotopes Eocene cooling trend, 882b | | |
| Site 20 | Site 265 surface-water masses, 872b | |
| oxygen isotopes Eocene–Holocene latitudinal gradient, 709b | | |
| Site 21 | Site 266 <i>Actinocyclus frysellaee</i> Interval Zone, 555b <i>Denticulopsis hustedtii</i> first occurrence, 559b <i>Denticulopsis maccollumii</i> Partial Range Zone, 556b diatoms datum events, stratigraphic constraints, 588b Miocene sequence, 577b surface-water masses, 872b | |
| oxygen isotopes Eocene–Holocene latitudinal gradient, 709b | | |
| Site 77 | Site 267 surface-water masses, 872b | |
| location, 714b | | |
| oxygen isotopes benthic foraminifers, 715b | | |
| comparison, high-latitude sites, 713b | | |
| Site 95 | Site 268 surface-water masses, 872b | |
| oxygen isotopes Eocene–Holocene latitudinal gradient, 709b | | |
| Site 152 | Site 269 surface-water masses, 872b | |
| oxygen isotopes Eocene–Holocene latitudinal gradient, 709b | | |
| Site 217 | Site 270 glacial history Miocene, 248b glacial sediments, 872b Oligocene, 872b glaciation Miocene/Pliocene unconformity, 118b glaciomarine sediments Oligocene, 79b stratigraphic intervals, 119b surface-water masses, 872b | |
| stable isotopes Paleogene, 695b | | |
| Site 219 | Site 271 glacial sediments, 872b surface-water masses, 872b | Site 279 oxygen isotopes foraminifers, 869b surface-water masses, 872b |
| carbon isotopes <i>Cibicidoides</i> spp., 709b | | |
| Site 220 | Site 272 glacial history Miocene, 248b glacial sediments, 872b glaciation Miocene/Pliocene unconformity, 118b surface-water masses, 872b | Site 280 <i>Eucyrtidium biconicum</i> last common occurrence, 525b <i>Lithomelissa haackeli</i> last common occurrence, 527b surface-water masses, 872b |
| stable isotopes Paleogene, 695b | | |
| Site 273 | Site 273 glacial history Miocene, 248b glacial sediments, 872b | Site 281 <i>Eucyrtidium biconicum</i> last common occurrence, 525b <i>Lithomelissa haackeli</i> last common occurrence, 527b oxygen isotopes foraminifers, 869b |
| | | |

SITE INDEX

- Prunopyle* sp., 521b, 524b
surface-water masses, 872b
- Site 282
surface-water masses, 872b
- Site 283
surface-water masses, 872b
- Site 284
surface-water masses, 872b
- Site 322
surface-water masses, 872b
- Site 323
Prinsium martinii, 502b
surface-water masses, 872b
- Site 324
surface-water masses, 872b
- Site 325
surface-water masses, 872b
- Site 326
surface-water masses, 872b
- Site 327
surface-water masses, 872b
- Site 328
diatoms
zonation, 551b
surface-water masses, 872b
- Site 329
calcareous nannofossils
dominant species fluctuation, 474b
surface-water masses, 872b
- Site 330
location, 417b
surface-water masses, 872b
- Site 331
surface-water masses, 872b
- Site 350
Rhizosolenia oligocenica Partial Range Zone, 557b
- Site 356
oxygen isotopes
Eocene–Holocene latitudinal gradient, 709b
- Site 357
oxygen isotopes
Eocene–Holocene latitudinal gradient, 709b
- Site 358
Eucampia antarctica
distribution, sediment and phytoplankton, 675b–676b
surface-water masses, 872b
- Site 362
carbon isotopes
Cibicidoides spp., 709b
oxygen isotopes
Eocene cooling trend, 882b
- Site 363
oxygen isotopes
Eocene cooling trend, 882b
- Site 364
oxygen isotopes
Eocene–Holocene latitudinal gradient, 709b
- Site 511
calcareous nannofossils, 470b
location, 417b
- Site 513
Amphymenium challengerae, 517b
calcareous nannofossils, 470b
ice rafting, 248b
Rhizosolenia oligocenica Partial Range Zone, 557b
- Site 514
Lithelius nautiloides
first common occurrence, 525b
- Site 516
calcareous nannofossils, 474b
magnetostratigraphy
biostratigraphic correlation, 851b
Reticulofenestra reticulata, 470b
strontium isotopes
carbonate, 733b
Oligocene–Miocene sediments, 732b
- Site 522
location, 714b
oxygen isotopes
benthic foraminifers, 715b
comparison, high-latitude sites, 713b
comparison, low- and mid-latitude sites, 713b
Oligocene event, 736b
strontium isotopes
carbonate, 733b
Oligocene–Miocene sediments, 732b
- Site 525
magnetostratigraphy
biostratigraphic correlation, 851b, 852b
oxygen isotopes
comparison, high-latitude sites, 713b
Eocene–Holocene latitudinal gradient, 709b
- Site 527
location, 714b
magnetostratigraphy
biostratigraphic correlation, 851b, 852b
oxygen isotopes
benthic foraminifers, 715b
comparison, high-latitude sites, 713b
Eocene values, 713b
Eocene–Holocene latitudinal gradient, 709b
- Site 528
oxygen isotopes
Eocene–Holocene latitudinal gradient, 709b
- Site 529
location, 714b
oxygen isotopes
benthic foraminifers, 715b
comparison, high-latitude sites, 713b
Eocene–Holocene latitudinal gradient, 709b
- Site 548
oxygen isotopes
Eocene–Holocene latitudinal gradient, 709b
- Site 550
oxygen isotopes
Eocene–Holocene latitudinal gradient, 709b
- Site 552
oxygen isotopes
Neogene maxima, 885b
- Site 563
strontium isotopes
carbonate, 733b
Oligocene–Miocene sediments, 732b
- Site 574
oxygen isotopes
benthic foraminifers, 715b
location, 714b
oxygen isotopes
comparison, high-latitude sites, 713b
- Site 577
calcareous nannofossils
comparison, Site 738, 504b
carbon isotopes
benthic foraminifers, 714b
Nuttallides truempyi, 704b, 706b, 708b
Coccolithus pelagicus
first occurrence, 497b
magnetic properties
Cretaceous/Tertiary boundary, 758b
magnetostratigraphy
biostratigraphic correlation, 852b
oxygen isotopes
Eocene–Holocene latitudinal gradient, 709b
Nuttallides truempyi, 706b, 708b
- Site 588
calcareous-biosiliceous ooze transition
surface-water temperature correlation, 888b
oxygen isotopes
Neogene, 887b
- Site 590
calcareous nannofossils
datum events, 735b
calcareous-biosiliceous ooze transition
surface-water temperature correlation, 888b
oxygen isotopes
Neogene, 887b
strontium isotopes, 735b
carbonate, 733b
Miocene sediments, 732b
- Site 594
Lithomelissa cheni, n. sp., 544b
- Site 638
Nephrolithus frequens
first appearance datum, 452b
- Site 642
Gilbert Chronozone
normal zones, 753b
- Site 682
redox and carbon flux data, 402b
- Site 683
redox and carbon flux data, 402b
- Site 684
redox and carbon flux data, 402b

- Site 685
redox and carbon flux data, 402b
- Site 686
redox and carbon flux data, 402b
- Site 687
redox and carbon flux data, 402b
- Site 688
redox and carbon flux data, 402b
- Site 689
Antarctissa cylindrica
last appearance, 516b
calcareous nannofossils
Danian, 502b–503b
dominant species fluctuation, 474b
Eocene disconformity, 472b
Ceratocyrtis mashae, 529b
Chiasmolithus altus, 470b
Cycladophora spongothorax
first common occurrence, 518b
Denticulopsis dimorpha Acme Zone, 556b
Denticulopsis maccollumii Partial Range Zone, 556b
Globoturborotalita labiacrassata
last appearance datum, 437b
iridium
Cretaceous/Tertiary boundary, 719b
location, 714b
nannofossil-diatom ooze transition, 202b
oxygen isotopes
benthic foraminifers, 715b
Eocene cooling trend, 883b
Eocene values, 713b
Paleogene, 885b
redox and carbon flux data, 402b
Reticulofenestra reticulata, 470b
stable isotopes
Oligocene inversions, 712b
strontium isotopes
benthic foraminifers, 736b
surface-water masses, 872b
Thalassiosira torokina
first occurrence, 575b
- Site 690
Antarctissa cylindrica
last appearance, 516b
calcareous nannofossils
Danian, 495b, 497b, 502b
dominant species fluctuation, 474b
Eocene disconformity, 472b
carbon isotopes
benthic foraminifers, 714b
Nuttallides truempyi, 704b, 706b, 708b
Ceratocyrtis mashae, 529b
Chiasmolithus altus, 470b
clay mineralogy
Eocene, 874b
Coccolithus pelagicus
first occurrence, 497b
Cycladophora spongothorax
first common occurrence, 518b
Denticulopsis dimorpha Acme Zone, 556b
Denticulopsis maccollumii Partial Range Zone, 556b
Globoturborotalita labiacrassata
last appearance datum, 437b
gray clay layer
Cretaceous/Tertiary boundary, 849b
Hornbrookina
first occurrence, 500b
- iridium
Cretaceous/Tertiary boundary, 719b
location, 495b
magnetobiostratigraphy, 501b
nannofossil-diatom ooze transition, 202b
Nephrolithus frequens
first appearance datum, 452b
oxygen isotopes
Eocene–Holocene latitudinal gradient, 707b, 709b
Nuttallides truempyi, 706b, 708b
redox and carbon flux data, 402b
stable isotopes
Eocene/Oligocene boundary hiatuses, 710b–711b
Oligocene inversions, 712b
surface- to intermediate-water gradient, 704b
surface-water masses, 872b
- Site 692
location, 417b
organic carbon, 419b
surface-water masses, 872b
- Site 693
Antarctissa cylindrica
last appearance, 516b
Cycladophora spongothorax
first common occurrence, 518b
ice-raftered debris
Oligocene, 873b
location, 417b
surface-water masses, 872b
- Site 694
location, 417b
surface-water masses, 872b
- Site 695
Antarctissa cylindrica
last appearance, 516b
surface-water masses, 872b
- Site 696
Antarctissa cylindrica
last appearance, 516b
surface-water masses, 872b
- Site 697
surface-water masses, 872b
- Site 698
redox and carbon flux data, 402b
surface-water masses, 872b
- Site 699
redox and carbon flux data, 402b
surface-water masses, 872b
- Site 700
Nephrolithus frequens
first appearance datum, 452b
redox and carbon flux data, 402b
surface-water masses, 872b
- Site 701
redox and carbon flux data, 402b
surface-water masses, 872b
tephra
geochemistry, 335b
- Site 702
carbon isotopes
Cibicidoides spp., 706b
- surface-water masses, 872b
- Site 703
surface-water masses, 872b
- Site 704
redox and carbon flux data, 402b
surface-water masses, 872b
- Site 705
redox and carbon flux data, 402b
- Site 707
redox and carbon flux data, 402b
- Site 708
redox and carbon flux data, 402b
- Site 709
redox and carbon flux data, 402b
- Site 710
redox and carbon flux data, 402b
- Site 711
redox and carbon flux data, 402b
- Site 713
redox and carbon flux data, 402b
- Site 714
redox and carbon flux data, 402b
- Site 715
redox and carbon flux data, 402b
- Site 717
redox and carbon flux data, 402b
- Site 718
redox and carbon flux data, 402b
- Site 719
redox and carbon flux data, 402b
- Site 720
redox and carbon flux data, 402b
- Site 721
redox and carbon flux data, 402b
- Site 722
iridium
Cretaceous/Tertiary boundary, 720b
redox and carbon flux data, 402b
- Site 723
redox and carbon flux data, 402b
- Site 724
redox and carbon flux data, 402b
- Site 725
redox and carbon flux data, 402b
- Site 727
redox and carbon flux data, 402b
- Site 728
redox and carbon flux data, 402b
- Site 730
redox and carbon flux data, 402b

SITE INDEX

- Site 731
redox and carbon flux data, 402b
- Site 736
ammonia
 pore waters, 395b
 sedimentation rate and, 402b-403b
ash layers
 geochemistry, 326b, 328b-333b
 petrography, 325b
 sources, 329b, 333b
basement description convention, 23a-24a
bathymetry, 6a, 45a, 349b
benthic foraminifers, 133a-134a, 612b,
 615b-616b, 621b
 paleoenvironmental factors, 616b, 619b-620b,
 622b
biogenic sediments, 874b
biogeography, 874b
biology and oceanography, 35a-37a
biostratigraphic datum levels, 837b
biostratigraphy, 24a, 26a-33a, 131a-135a,
 156a, 836b, 838b
calcareous nannofossils, 131a, 469b, 814b, 816b
calcium, pore water, 397b-398b
calcium carbonate, 142a
carbonate, 354b-355b, 357b
Chaetoceros resting spores, 600b, 604b
clay mineralogy, 874b
core handling, 15a-17a
coring summary, 128a
diatoms, 134a, 557b-558b, 560b-561b, 605b
 zonation, 551b-557b, 816b-817b
dikes, 24a, 44a
drilling deformation, 15a, 17a, 19a-20a,
 128a-129a
Eucampia antarctica
 distribution, sediment and phytoplankton,
 675b-676b, 678b
 morphology, summer vs. winter stages, 676b-
 677b, 678b-679b
 size relationships, 677b-678b, 680b
extrusives, 24a, 44a
geochemical logs, 907b-908b, 913b, 915b,
 917b, 919b-920b, 923b, 926b
geographic setting, 5a
geological evolution, 7a
geological setting, 124a, 353b
glaciation
 deep-sea hiatus association, 874b-875b
 isotopic evidence, 869b-870b
 pollen record, 873b-875b
 sediment record, 870b-873b
glaciogenic sediments
 genetic classification, 22a-23a
glass
 major element geochemistry, 337b-343b
 morphology, 344b
grain-size distribution
 tephra layers, 328b
granular sediment classification, 21a-22a, 23a,
 43a-44a
heat flow, 153a, 156a
hydrocarbon gases, 34a, 140a
ice-raftered debris, 873b
inorganic carbon, 142a, 904b
inorganic geochemistry, 35a, 137a-140a, 141a
interstitial-water geochemistry, 140a, 372b
 high-resolution sampling technique, 348b,
 393b-399b
 major cation concentrations, 351b, 929b
 solute distribution, 358b
lithostratigraphy, 156a
 depositional environment, 130a-131a
lithologic Units I-II, 129a-130a
living bacteria, 687b-689b
location, 124a, 300b, 324b, 350b, 376b, 428b,
 468b, 513b, 751b, 814b, 871b
logging, 39a-40a
 synthetic seismogram, 40a
magnesium, pore water, 395b, 397b
magnetic properties, 33a-34a, 45a, 135a-137a,
 138a
 susceptibility, 34a, 137a, 139a
magnetobiochronology, 815b, 825b, 827b
metal contents, 356b
navigation, 45a, 47a, 48a-85a, 87a
objectives, 7a, 124a
operations, 125a-128a
organic carbon, 113b, 354b-355b, 357b, 904b
organic geochemistry, 34a-35a, 140a, 142a,
 156a
oxygen isotopes, 887b
paleceanography, 124a
paleoclimatology, 156a
palyontology, 135a
physical properties, 37a-38a, 144a-153a, 156a
phytoplankton, 140a, 142a-144a
planktonic foraminifers, 131a-133a, 814b, 816b
radiolarians, 134a-135a, 514b-516b
 evolutionary lineages, 529b-531b
 systematics, 531b-537b
 zonation, 44a, 814b, 816b, 817b
Rock-Eval pyrolysis, 35a, 140a, 143a
salinity, 107a
sediment accumulation rate, 139a
sediment classification, 21a
sediment composition, 353b-355b
sedimentation rate, 137a, 401b-404b
seismic stratigraphy, 153a-155a
 reflection profiling, 45a, 86a, 90a, 106a
shear strength, 145a, 148a, 149a
silica, pore water, 398b
silicoflagellate, 933b-934b
sonobuoy seismic recording, 45a-46a, 107a,
 124a-125a, 126a, 154a
stratigraphic column, 327b
sulfate
 pore waters, 398b-399b
 sedimentation rate and, 402b
surface-water masses, 872b
tectonic history, 5a, 7a
temperature, 107a, 150a-153a
tephra, geochronology, 336b
thermal conductivity, 150a
total alkali vs. silica, 333b
velocity, 107a, 126a, 148a, 150a
 measurement errors, 46a
water content, 904b
- Site 737
ammonia
 pore waters, 395b
 sedimentation rate and, 402b-403b
ash layers
 geochemistry, 326b, 328b-333b
 petrography, 325b
 sources, 329b, 333b
authigenic silicate formation, 218b-219b
barite, 217b-218b
basement description convention, 23a-24a
bathymetry, 6a, 45a, 349b
benthic foraminifers, 179a
biogenic sediments, 874b
biogeography, 874b
biology and oceanography, 35a-37a
biostratigraphic datum levels, 840b
biostratigraphy, 24a, 26a-33a, 173a-181a, 218a
Neogene, 839b, 842b
Paleogene, 840b, 843b
burrows, 171a
calcareous nannofossils, 176a-177a,
 470b-471b, 474b, 888b
zonation, 814b, 816b
calcium, pore water, 397b-398b
calcium/magnesium ratio, 193a
carbonate, 354b-355b, 359b
Chaetoceros resting spores, 600b, 604b
clay mineralogy, 874b
core handling, 15a-17a
coring summary, 164a
density-porosity relationship, 275b
diagenesis, 172a
diatom ooze
 initiation of deposition, 195b
diatoms, 179a-180a, 558b-559b, 562b-563b,
 564b
 zonation, 551b-557b, 816b-817b
dikes, 24a, 44a
drilling deformation, 15a, 17a, 19a-20a, 164a
electrical resistivity, 366b
Eucampia antarctica
 distribution, sediment and phytoplankton,
 675b-676b, 678b
 morphology, summer vs. winter stages, 676b-
 677b, 678b-679b
 size relationships, 677b-678b, 680b
extrusives, 24a, 44a
feldspar, 217b
geochemical logs, 908b-909b, 913b-914b,
 915b-916b, 917b-918b, 920b-921b,
 923b-924b, 926b-927b
geographic setting, 5a
geological evolution, 7a
geological setting, 353b
geophysical logs, 265b
glaciation
 deep-sea hiatus association, 874b-875b
 isotopic evidence, 869b-870b
 pollen record, 873b-875b
 sediment record, 870b-873b
glaciogenic sediments
 genetic classification, 22a-23a
glass geochemistry, 337b-343b
grain-size distribution
 tephra layers, 328b
granular sediment classification, 21a-22a, 23a,
 43a-44a
heat flow, 198a-199a
hydrocarbon gases, 34a, 188a
ice-raftered debris, 873b
ichnofabric, 636b-637b, 638b
inorganic carbon, 193a-194a, 196a
inorganic geochemistry, 35a, 185a-188a, 192a,
 219a
interstitial-water geochemistry, 191a,
 354b-355b, 357b, 359b-363b, 372b
chemical shifts, 361b
high-resolution sampling technique, 348b,
 393b-399b, 903b
major cation concentrations, 351b, 362b,
 929b-930b
solute distribution, 360b
laminations, 170a
lithologic column, 266b
lithology
 log-inferred, 272b, 274b-277b, 285b
 summary, 359b
lithostratigraphy, 212b-213b, 216a, 218a, 264b
 depositional environment, 172a-173a
 lithologic Units I-VI, 165a-172a

location, 212b, 264b, 300b, 324b, 350b, 376b, 428b, 468b, 513b, 751b, 814b, 871b
 logging, 39a-49a
 operations, 202a-205a
 quality, 205a
 summary, 221a-227a
 synthetic seismogram, 40a, 212a-214a, 215a
 units 1-6, 207a-210a, 275b-277b
 magnesium, pore water, 395b, 397b
 magnetic properties, 33a-34a, 45a, 181a-183a, 184a, 185a, 186a, 188a, 189a, 754b, 759b
 susceptibility, 34a, 187a
 magnetobiochronology, 815b, 828b-829b, 831b, 833b-836b
 magnetostratigraphy, 183a, 190a, 218a, 754b, 757b, 760b
 Neogene, 839b, 842b
 subchron boundaries, 841b
 microfaults, 171a
 mineralogy, 214b-215b
 lithologic correlation, 276b-277b, 278b-279b
 navigation, 45a, 47a, 48a-85a, 87a
 objectives, 7a, 160a
 opal, 218b
 operations, 162a-164a
 organic carbon, 354b-355b, 359b, 904b-905b
 organic geochemistry, 34a-35a, 188a-191a, 192a-193a, 219a
 oxygen isotopes, 887b
 palynology, 180a
 physical properties, 37a-38a, 194a-203a
 geotechnical units G1-G3, 195a
 phytoplankton, 191a, 193a-194a
 planktonic foraminifers, 178a-179a, 814b, 816b
 prograding sequence, 217a
 quartz, 217b
 radiolarians, 180a, 516b-518b
 evolutionary lineages, 529b-531b
 systematics, 531b-537b
 zonation, 44a, 814b, 816b, 817b
 redox and carbon flux data, 402b
 Rock-Eval pyrolysis, 35a, 189a-191a, 196a, 197a, 198a, 199a
 salinity, 107a
 sediment classification, 21a
 sediment composition, 357b
 sedimentary structures, 172a
 sedimentation rate, 183a, 185a, 191a, 401b-404b
 seismic stratigraphy
 reflection profiling, 45a, 86a, 91a-92a, 106a
 units S1 and S2, 214a-216a
 shear strength, 195a, 197a
 silica, pore water, 398b
 sonobuoy seismic recording, 45a-46a, 107a, 160a-162a, 163a
 stratigraphic column, 327b
 sulfate
 sedimentation rate and, 402b
 pore waters, 398b-399b
 surface-water masses, 872b
 tectonic history, 5a, 7a
 temperature, 107a, 198a-199a, 205a, 206a
 tephra, geochronology, 336b
 thermal conductivity, 197a-198a, 205a
 total alkali vs. silica, 333b
 trace fossils, 636b-637b, 638b
 velocity, 107a, 197a, 204a
 measurement errors, 46a
 water content, 904b-905b
 X-ray diffraction, 221b

Site 738
 alteration, 307b, 313b-315b

ammonia
 pore waters, 395b
 sedimentation rate and, 402b-403b
 antimony
 Cretaceous/Tertiary boundary, 724b-725b
 aqueous silica, 367b
 arsenic
 Cretaceous/Tertiary boundary, 724b-725b
 authigenic silicate formation, 218b-219b, 220b
 barite, 217b-218b
 barium
 Cretaceous/Tertiary boundary, 725b
 basalt
 classification, 302b-303b
 isotopic composition, 294b-295b, 296b-298b
 major and trace element concentration, 294b, 295b, 301b-302b, 312b, 313b
 Mesozoic Southern Hemisphere comparisons, 315b, 317b
 mineralogy, 293b-294b
 petrography, 300b-301b
 rare earth elements, 318b
 secondary minerals, 316b
 source region, 317b-319b
 basement
 alteration, 315b
 geochemistry, 309b-311b
 stratigraphy, 300b
 basement description convention, 23a-24a
 bathymetry, 6a, 45a, 349b
 benthic foraminifers, 250a-251a, 613b-614b, 615b
 comparison, Site 744, 486a
 Cretaceous/Tertiary boundary, 857b-859b
 distribution, 455b
 Maestrichtian-Danian, 457b
 paleoenvironmental factors, 622b
 vertical ranges, 616b-617b
 biogenic sediments, 874b
 biogenic silica, 884b
 biogeography, 874b
 biology and oceanography, 35a-37a
 biostratigraphic datum levels, 833b
 biostratigraphy, 24a, 26a-33a, 244a-252a, 279a, 831b, 832b
 Maestrichtian-Eocene, 835b
 calcareous nannofossils, 244a-246a, 472b-474b, 477b, 481b, 482b, 888b
 Cretaceous/Tertiary boundary, 455b
 Danian, 497b-499b, 508b
 light-microscopic analysis, 856b-857b
 paleobiogeographic patterns, 502b, 504b
 quantitative analyses, 502b, 503b
 zonation, 500b-502b, 814b, 816b
 calcium, pore water, 397b-398b
 calcium carbonate, 261a
 calcium/magnesium ratio, 258a
 carbon isotopes
 Cretaceous/Tertiary boundary, 859b
 Eocene-Oligocene transition, 710b-712b
 foraminifers, 700b-703b
 Oligocene deep-water association, 712b-714b
 Paleocene-Eocene, 698b
 Paleocene-Miocene, 704b-709b
 paleoenvironmental implications, 714b-715b
 surface- to intermediate-water gradients, 697b, 704b, 709b-710b
 carbonate, 187b, 205b, 354b-355b, 364b
 Cretaceous/Tertiary boundary, 854b
 cesium
 Cretaceous/Tertiary boundary, 726b, 728b-729b
Chaetoceros resting spores, 600b-601b, 604b
 chromium
 Cretaceous/Tertiary boundary, 724b
 chronostratigraphy, 438b
 clay layer
 SEM analyses, 857b
 clay mineralogy, 114b, 194b, 874b
 Cretaceous/Tertiary boundary, 854b
 environmental implications, 199b-200b
 Eocene, 882b
 Oligocene, 884b
 source, 198b-199b
 clinopyroxene
 chemical composition, 303b, 304b
 cobalt
 Cretaceous/Tertiary boundary, 724b
 core handling, 15a-17a
 coring summary, 234a
 CTD profile, 109a, 110a
 density-porosity relationship, 281b
 diatoms, 251a-252a, 264a
 in Quaternary sediments, 261a
 zonation, 551b-557b, 816b-817b
 dikes, 24a, 44a
 dinoflagellates
 Cretaceous, 631b-633b
 drilling deformation, 15a, 17a, 19a-20a
 element stratigraphy
 calcite-free data, 728b
 Cretaceous/Tertiary boundary, 721b-729b
 elemental ratios, 727b-728b
Eucampia antarctica
 distribution, sediment and phytoplankton, 675b-676b, 678b
 morphology, summer vs. winter stages, 676b-677b, 678b-679b
 size relationships, 677b-678b, 680b
 extrusives, 24a, 44a
 feldspar, 217b, 302b-303b
 geochemical logs, 909b-910b, 924b, 927b
 geographic setting, 5a
 geological evolution, 7a
 geological setting, 353b
 geophysical logs, 267b
 glacial history, 201b-202b, 880b-888b
 microfossil assemblage changes, 884b
 Miocene-Pliocene transition, 248b-249b
 glaciation
 deep-sea hiatus association, 874b-875b
 isotopic evidence, 869b-870b
 onset, 201b
 pollen record, 873b-875b
 sediment record, 870b-873b
 glaciogenic sediments
 genetic classification, 22a-23a
 grain-size distribution
 nonbiogenic matter, 190b-191b, 192b, 206b-207b
 granular sediment classification, 21a-22a, 23a, 43a-44a
 green clay lamina
 Cretaceous/Tertiary boundary, 866b-867b
 gray clay layer, 850b
 Cretaceous/Tertiary boundary, 862b-865b, 866b
 hafnium
 Cretaceous/Tertiary boundary, 726b, 728b-729b
 hydrocarbon gases, 34a
 ice loading events, 885b-886b
 ice rafting, 195b, 197b-198b
 ice-rafted debris, 120b, 873b, 882b, 884b, 886b
 ichnofabric
 Cretaceous/Tertiary boundary, 637b-639b
 inorganic carbon, 259a-260a, 261a, 905b
 inorganic geochemistry, 35a, 254a-257a

SITE INDEX

interstitial-water geochemistry, 257a, 258a, 372b-373b
 high-resolution sampling technique, 393b
 399b, 348b, 903b
 major cation concentrations, 351b, 366b, 930b
 solute distribution, 365b
iridium
 accumulation rates, 855b
 anomaly, 726b-729b
 in the biosphere, 855b-856b
 Cretaceous/Tertiary boundary, 720b, 723b, 854b-855b
 in the hydrosphere, 855b-856b
iron
 Cretaceous/Tertiary boundary, 723b-724b
lithologic column, 268b
lithology, 187b, 300b, 696b
 log-inferred, 277b-279b, 285b
 summary, 364b
lithostratigraphy, 264b, 279a, 307b
 Cretaceous/Tertiary boundary, 849b
 hiatuses, 243a-244a, 245a, 246a
 lithologic Units I-VIII, 235a-238a
living bacteria, 687b-689b
location, 185b, 186b, 212b, 230a, 264b, 293b, 300b, 324b, 350b, 376b, 428b, 468b, 495b, 513b, 694b, 714b, 751b, 814b, 871b
logging, 39a-40a, 280a
 operations, 269a-270a
 summary, 283a-288a
 synthetic seismogram, 40a, 272a-275a, 277a, 278a
 units 1-4, 270a-272a, 277b-279b
magma evolution, 317b
magnesium, pore water, 395b, 397b
magnetic properties, 33a-34a, 45a, 252a-253a, 254a, 757b-758b, 763b
 Cretaceous/Tertiary boundary, 760b, 764b, 765b, 766b
 susceptibility, 34a, 246a
magnetobiochronology, 454b, 501b, 815b, 822b, 825b, 834b
magnetostratigraphy, 832b
 Cretaceous sequence, 758b-759b
 Cretaceous/Tertiary boundary, 835b, 851b-853b
 subchron boundaries, 833b
mineralogy, lithologic correlation, 280b-281b
nannofossil-diatom ooze transition, 195b, 202b
navigation, 45a, 47a, 48a-85a, 87a
objectives, 7a, 230a, 495b
ocean particulate fluxes, 109a-110a
olivine pseudomorphs, 302b
opal, 190b, 218b
operations, 232a-235a
organic carbon, 354b-355b, 364b, 905b
organic geochemistry, 34a-35a, 257a, 258a, 259a-260a
oxygen isotopes
 comparison, low- and mid-latitude sites, 713b
 Cretaceous/Tertiary boundary, 859b
 deep-water association, 712b-714b
 Eocene-Holocene latitudinal gradient, 707b-708b, 709b
 Eocene-Miocene, 708b-709b, 882b
 Eocene-Oligocene transition, 710b-712b
 foraminifers, 700b-703b
 Oligocene, 712b-714b, 884b
 Paleocene-Eocene, 698b, 704b-708b
 paleoenvironmental implications, 714b-715b
 surface- to intermediate-water gradients, 697b, 704b, 709b-710b
pack ice coverage, 110a, 112a-113a
paleoceanography, 279a

palynology, 252a
 pegmatoid schlieren, 299b, 301b, 322b
petrogenesis, 315b
physical properties, 37a-38a, 261a-269a
 geotechnical units G1-G6, 262a, 279a
phytoplankton, 257a, 260a-261a
planktonic foraminifers, 250a, 428b-430b
 biogeography, 455b
 Cretaceous, 452b-455b, 457b-459b, 469b
 Cretaceous/Tertiary boundary, 455b-459b, 857b-859b
 recycled, 457b-459b, 460b
 zonation, 431b-435b, 814b, 816b
polar ordination values, 502b, 504b
preglacial conditions, 200b-201b
quartz, 217b
radiolarians, 252a, 518b
 evolutionary lineages, 529b-531b
 Paleogene datum events, 527b-529b
 systematics, 531b-537b
 zonation, 44a, 814b, 816b, 817b
rare earth elements, 315b
 Cretaceous/Tertiary boundary, 722b, 726b
 redox and carbon flux data, 402b
Rock-Eval pyrolysis, 35a
salinity, 107a
scandium
 Cretaceous/Tertiary boundary, 722b-723b, 728b-729b
sediment accumulation rate
 Cretaceous/Tertiary boundary, 851b-853b
sediment classification, 21a
sediment composition, 187b-190b, 204b, 363b-364b
 environmental implications, 194b-195b
 nonbiogenic matter, 190b
sedimentation rate, 253a-254a, 256a, 401b-404b
 logging units and, 272a-273a
seismic stratigraphy, 276a-277a, 279a, 280a
 reflection profiling, 45a, 86a, 93a, 106a
 units I-V, 276a
Shannon-Weaver diversity index, 502b, 504b
shear strength, 264a, 267a, 268a
sheet silicates, 301b
 chemical composition, 305b-307b
silica, pore water, 398b
sonobuoy seismic recording, 45a-46a, 107a, 230a-232a
sulfate
 pore waters, 398b-399b
 sedimentation rate and, 402b
surface-water masses, 872b
tantalum
 Cretaceous/Tertiary boundary, 726b, 728b-729b
tectonic history, 5a, 7a, 280a
temperature, 107a
terrigenous clay, 196b, 208b-209b
thorium
 Cretaceous/Tertiary boundary, 725b-726b, 728b-729b
uranium
 Cretaceous/Tertiary boundary, 725b-726b, 728b-729b
velocity, 107a, 264a, 266a-267a, 268a-269a
 measurement errors, 46a
vesicle fillings, 302b, 307b
 chemical composition, 308b
water content, 905b
weather summary, 262a
white ooze lamina
 Cretaceous/Tertiary boundary, 866b
X-ray diffraction, 222b

zinc
 Cretaceous/Tertiary boundary, 724b-725b
Site 739
 acoustic units PS.1-PS.6, 7b-13b, 27b-28b
 areal distribution, 13b-17b
 lithostratigraphic correlation, 99b-101b
 aliphatic hydrocarbons, 411b, 412b, 413b
 alkalinity, 388b-389b
 amino acids, 114a-115a, 116a, 117a
 ammonia
 pore waters, 395b
 sedimentation rate and, 402b-403b
Atterberg limits, 150b-151b
basement description convention, 23a-24a
bathymetry, 11a, 45a, 78b, 291a
bedrock topography, 64b
benthic foraminifers, 304a
biogenic sediments, 112b-113b, 874b
biogenic silica fluxes, 114a
biogeography, 874b
biology and oceanography, 35a-37a
biostratigraphy, 24a, 26a-33a, 304a-305a, 339a, 844b
 summary, 306a-307a
bioturbation, 109b
calcareous nannofossils, 304a, 477b, 478b-480b, 481b, 888b
biostatigraphy, 743b
 zonation, 814b, 816b
calcium, pore water, 397b-398b
carbon isotopes, 742b
carbonate, 378b
 thermodynamic modeling, 389b-391b
Chaetoceros resting spores, 601b, 604b
clasts
 metamorphic, 137b, 138b
 provenance, 138b-139b
 sedimentary, 134b, 135b, 137b
 shapes, 98b
clay mineralogy, 84b, 86b, 87b, 113b-114b, 874b
coal-bearing sediments, 57b-60b
combustible carbon and nitrogen, 114a-115a
 stable isotopic fractionation, 115a, 117a
core handling, 15a-17a
coring summary, 295a
CTD profile, 109a, 110a
data processing and availability, 23b-24b
density-porosity relationship, 283b, 285b
diagenesis, 303a
diamictites, 120b-121b, 147b
diamicts, 88b-89b, 107b
 grain fabrics, 91b-92b
diatoms, 304a-305a, 318a, 576b-577b, 584b-585b, 605b
 analytical methods, 646b-648b
biostratigraphy, 743b-744b
chlorophyll a and density of, 653b, 659b
distribution patterns, 651b-655b, 661b-663b
empty/full ratio, 655b, 656b, 657b, 663b
integrated abundances, 648b-651b, 652b, 664b
phytoplankton vs. sediments, 668b-673b
in Quaternary sediments, 315a
reworking, 744b
spring blooms, 671b-672b
standing stations, 655b-660b, 664b
zonation, 551b-557b, 816b-817b
dikes, 24a, 44a
drifting sediment trap deployments, 111a
drilling deformation, 15a, 17a, 19a-20a, 295a
erosion, ice thickness and, 178b-179b
Eucampia antarctica

distribution, sediment and phytoplankton, 675b-676b, 678b
 morphology, summer vs. winter stages, 676b
 677b, 678b-679b
 size relationships, 677b-678b, 680b
 extractable organic matter
 gas chromatograms, 412b, 413b
 source and maturity, 408b-410b, 411b-412b
 extrusives, 24a, 44a
 flat-lying sequence, 107b, 876b
 gastropod shell, 131b
 geochemical logs, 910b-911b, 914b, 916b,
 918b, 921b-922b, 924b-925b, 927b-928b
 geographic setting, 80b-81b
 geological evolution, 20b-22b
 geological setting, 5b-7b, 8a-9a, 46b, 58b, 79b,
 80b-81b, 133b, 170b, 375b-376b, 377b,
 407b
 geophysical logs, 177b, 272b
 glacial depositional environment, 18b-20b
 glacial history, 78b-80b, 115b-122b, 180b,
 875b-877b, 880b-888b
 glacial sediments
 solute diffusion, 387b-388b
 glacial transport, 71b-73b, 114b-115b
 glaciation
 deep-sea hiatus association, 874b-875b
 isotopic evidence, 869b-870b
 onset, 201b
 pollen record, 873b-875b
 sediment record, 870b-873b
 glaciogenic sediments
 consolidation, 151b-152b, 155b-157b, 174b-
 176b
 depositional environment, 114b-115b
 genetic classification, 22a-23a, 101b, 103b
 glaciomarine deposition, 80b
 grain orientation, 105b
 grain-size distribution, 81b, 149b-150b
 diamictites, 120b-121b
 by facies type, 89b-90b, 94b
 granular sediment classification, 21a-22a, 23a,
 43a-44a
 gravel, grain-size distribution, 83b
 hinterland geology, 64b-68b
 hydrocarbon gases, 34a
 ice loading events, 176b, 178b, 179b-181b,
 885b-886b
 ice rafted debris, 73b-74b, 873b, 886b
 inorganic carbon, 313a-314a, 905b-906b
 inorganic geochemistry, 35a, 310a-313a
 interstitial-water geochemistry, 311a,
 377b-378b, 380b-383b
 high-resolution sampling technique, 376b,
 393b-399b, 903b
 major cation concentrations, 380b, 391b,
 930b-931b
 solute distribution, 382b
 Lambert Glacier-Amery Ice Shelf System
 flow pattern and particle paths, 70b-71b
 ice dynamics, 62b-64b
 structural glaciology, 68b-70b
 lithofacies, 86b, 88b-92b, 103b, 105b, 107b
 lithologic column, 273b
 lithology
 log-inferred, 283b-285b, 286b-287b
 lithostratigraphy, 93b, 95b, 127b-128b, 269b,
 294a-304a, 339a, 877b
 hiatuses, 95b, 97b, 99b
 lithologic Units I-V, 296a-303a
 mixed facies Unit III, 109b
 living bacteria, 687b-689b
 location, 6b, 135b, 264b, 300b, 376b, 428b,
 468b, 513b, 647b, 668b, 751b, 814b, 871b

logging, 39a-40a
 correlation, Site 742, 442a
 lithologic correlation, 328a
 operations, 328a
 quality, 328a
 summary, 342a-344a
 synthetic seismogram, 40a, 331a-332a
 units 1-4, 283b-285b, 329a-331a
 magnesium, pore water, 395b, 397b
 magnetic properties, 33a-34a, 45a, 305a,
 308a-310a
 magnetobiochronology, 815b, 837b
 mineralogy, lithologic correlation, 286b
 mollusk shells, 740b
 isotopic values, 739b-745b
 morphology, 7a-8a
 navigation, 45a, 47a, 48a-85a, 87a, 88a
 objectives, 13a, 290a
 ocean particulate fluxes, 109a-110a, 114a
 opal, 84b
 operations, 292a-294a
 organic carbon, 378b, 905b-906b
 glacial constraints, 388b
 organic geochemistry, 34a-35a, 313a-314a
 oxygen isotopes, 742b, 887b
 pack ice coverage, 110a, 112a-113a
 paleoclimatic background, 9a-10a, 13a
 palynology, 942b, 944b-945b
 permeability, 152b, 154b
 physical properties, 37a-38a, 315a-328a, 339a
 geotechnical units G1-G7, 319a, 325a-326a
 lithologic correlation, 145b
 shipboard analysis, 147b-149b, 160b-163b
 phytoplankton, 314a-315a
 planktonic foraminifers, 304a, 814b, 816b
 plant pigments, 118a-120a
 prograding sequence, 876b, 880b
 radiolarians, 305a, 518b-519b
 evolutionary lineages, 529b-531b
 systematics, 531b-537b
 zonation, 44a, 814b, 816b, 817b
 redox and carbon flux data, 402b
 rhythmites, 108b
 Rock-Eval pyrolysis, 35a, 380b, 409b
 salinity, 107a
 sand/mud ratio, 96b
 sediment classification, 21a
 sediment composition, 81b-84b, 85b,
 379b-380b
 sedimentary succession, 106b
 sedimentation rate, 310a, 401b-404b
 seismic stratigraphy, 9a, 11a, 12a, 333a-338a,
 339a-340a
 correlation, Site 742, 446a-447a
 disrupted sequence, 19b
 reflection profiling, 7b, 10b, 36b-37b, 45a,
 86a, 94a, 106a
 units I-III, 334a-336a
 seismic velocity log, 30b-31b, 35b
 shear strength, 154b-156b
 silica, pore water, 398b
 slumping, 112b
 sonobuoy seismic recording, 45a-46a, 107a,
 290a-293a, 294a
 refraction data, 28b-29b
 stable isotopes, 114a-115a
 stratigraphic intervals, 119b
 stratigraphic log, 102b, 381b, 581b, 878b
 strontium isotopes, 844b
 interstitial waters, 391b
 mollusk shells, 739b-742b
 pore waters, 740b-741b
 structural features, 17b-18b
 sulfate

pore waters, 398b-399b
 sedimentation rate and, 402b
 surface-water masses, 872b
 temperature, 107a
 velocity, 46a, 107a, 327a
 water content, 905b-906b
 weather summary, 315a

Site 740
 acoustic units PS.1-PS.6, 7b-13b, 27b-28b
 areal distribution, 13b-17b
 lithostratigraphic correlation, 99b-101b
 alkalinity, 388b-389b
 amino acids, 114a-115a, 116a, 117a
 ammonia
 sedimentation rate and, 402b-403b
 arkosic wackes, 56b
 Atterberg limits, 150b-151b
 basement description convention, 23a-24a
 bathymetry, 11a, 45a, 78b
 bedrock topography, 64b
 biogenic sediments, 112b-113b, 874b
 biogenic silica fluxes, 114a
 biogeography, 874b
 biology and oceanography, 35a-37a
 biostratigraphy, 24a, 26a-33a, 356a-357a
 calcareous nannofossils, 477b, 814b, 816b
 calcium/magnesium ratio, 362a
 carbonate, 378b
 thermodynamic modeling, 389b-391b
Chaetoceros resting spores, 601b, 604b
 channel deposits, 53b-54b
 clasts, 138b-139b
 clay mineralogy, 84b, 86b, 113b-114b, 874b
 claystone mineralogy, 51b
 coal-bearing sediments, 57b-60b
 combustible carbon and nitrogen, 114a-115a
 core handling, 15a-17a
 CTD profile, 109a, 110a
 data processing and availability, 23b-24b
 depositional environment, preglacial, 18b
 diamictites, 120b-121b
 diamiccts, 88b-89b
 diatoms, 605b
 analytical methods, 646b-648b
 chlorophyll *a* and density of, 653b, 659b
 distribution patterns, 651b-655b, 661b-663b
 empty/full ratio, 655b, 656b, 657b, 663b
 integrated abundances, 648b-651b, 652b,
 664b
 phytoplankton vs. sediments, 668b-673b
 spring blooms, 671b-672b
 standing stations, 655b-660b, 664b
 in surface sediments, 365a
 zonation, 551b-557b, 816b-817b
 dikes, 24a, 44a
 drifting sediment trap deployments, 111a
 drilling deformation, 15a, 17a, 19a-20a
 erosion, ice thickness and, 178b-179b
Eucampia antarctica
 distribution, sediment and phytoplankton,
 675b-676b, 678b
 morphology, summer vs. winter stages, 676b-
 677b, 678b-679b
 size relationships, 677b-678b, 680b
 extrusives, 24a, 44a
 fining-upward sequences, 47b, 48b, 52b, 353a,
 356a
 flat-lying sequence, 107b
 geographic setting, 80b-81b
 geological evolution, 20b-22b
 geological setting, 5b-7b, 8a-9a, 46b, 58b, 79b,
 80b-81b, 133b, 170b, 375b-376b, 377b
 geophysical logs, 177b

SITE INDEX

glacial history, 78b-80b, 115b-122b, 180b, 875b-877b, 880b-888b
 glacial sediments
 solute diffusion, 387b-388b
 glacial transport, 71b-73b, 114b-115b
 glaciation
 deep-sea hiatus association, 874b-875b
 isotopic evidence, 869b-870b
 pollen record, 873b-875b
 sediment record, 870b-873b
 glaciogenic sediments
 consolidation, 151b-152b, 155b-157b, 174b-176b
 depositional environment, 114b-115b
 genetic classification, 22a-23a, 101b, 103b
 glaciomarine deposition, 80b
 grain-size distribution, 81b, 149b-150b
 diamictites, 120b-121b
 granular sediment classification, 21a-22a, 23a, 43a-44a
 hinterland geology, 64b-68b
 hydrocarbon gases, 34a, 363a
 hydrogen isotopes
 interstitial waters, 384b
 ice loading, 368a
 ice loading events, 176b, 178b, 179b-181b, 885b-886b
 ice rafted debris, 73b-74b, 873b, 886b
 inorganic carbon, 363a
 inorganic geochemistry, 35a, 53b, 358a, 360a-363a, 374a
 interstitial-water geochemistry, 360a, 361a, 377b-378b, 383b
 high-resolution sampling technique, 376b
 major cation concentrations, 380b, 391b
 solute distribution, 384b
 Lambert Glacier-Amery Ice Shelf System
 flow pattern and particle paths, 70b-71b
 ice dynamics, 62b-64b
 structural glaciology, 68b-70b
 lithofacies, 86b, 88b-92b, 103b, 105b, 107b
 lithostratigraphy, 125b, 349a-356a
 hiatuses, 95b, 97b, 99b
 lithologic Units I-III, 350a-351a, 372a, 374a
 living bacteria, 687b-689b
 location, 6b, 135b, 300b, 376b, 428b, 468b, 513b, 647b, 668b, 747b, 751b, 796b, 814b, 871b
 logging, 39a-40a
 synthetic seismogram, 40a
 magnetic properties, 33a-34a, 45a, 357a-358a, 359a
 alternating-field demagnetization, 773b
 intensity changes, 804b
 red beds, 798b-804b
 magnetobiochronology, 815b, 836b
 meteoric water incursion, 388b
 mineralogy
 garnet grains, 52b
 morphology, 7a-8a
 navigation, 45a, 47a, 48a-85a, 87a, 88a
 objectives, 13a, 345a-346a, 372a
 ocean particulate fluxes, 109a-110a, 114a
 operations, 349a
 organic carbon, 113b, 378b
 glacial constraints, 388b
 organic geochemistry, 34a-35a, 53b, 362a, 363a, 374a-375a
 oxygen isotopes
 interstitial waters, 384b
 pack ice coverage, 110a, 112a-113a
 paleoclimatic background, 9a-10a, 13a
 paleogeographic reconstruction, 804b, 806b, 807b

palynology, 941b, 942b
 permeability, 152b, 154b
 petrography, 50b-52b
 physical properties, 37a-38a, 365a-369a, 370a, 371a, 375a
 shipboard analysis, 147b-149b, 163b-164b
 phytoplankton, 363a-365a
 planktonic foraminifers, 814b, 816b
 plant pigments, 118a-120a
 productivity, 364a-365a
 prograding sequence, 876b
 radiolarians, 44a, 814b, 816b, 817b
 red beds, 47b, 54b, 354a, 356a, 374a
 Amery Group correlation, 797b, 804b
 distribution, 806b
 magnetic properties, 797b-804b, 805b
 redox and carbon flux data, 402b
 Rock-Eval pyrolysis, 35a
 salinity, 107a
 sandstone mineralogy, 51b
 sediment classification, 21a
 sediment composition, 81b-84b, 383b
 sedimentary succession, 45b-50b
 sedimentation rate, 358a, 401b-434b
 seismic stratigraphy, 9a, 11a, 12a, 372a, 373a
 reflection profiling, 7b, 10b, 36b-37b, 45a, 86a, 95a, 106a
 units I-III, 369a-370a
 seismic velocity log, 32b-33b
 shear strength, 154b-156b
 sonobuoy seismic recording, 28b-29b, 45a-46a, 107a, 346a-349a
 stable isotopes, 114a-115a
 structural features, 17b-18b
 sulfate, sedimentation rate and, 402b
 sulfur content, 113b
 surface-water masses, 872b
 temperature, 107a
 unconsolidated sediments
 Holocene chronology, 747b-750b
 velocity, 46a, 107a

Site 741
 acoustic units PS.1-PS.6, 7b-13b, 27b-28b
 areal distribution, 13b-17b
 lithostratigraphic correlation, 99b-101b
 aliphatic hydrocarbons, 411b, 415b, 418b-423b
 alkalinity, 388b-389b
 amino acids, 114a-115a, 116a, 117a
 ammonia, sedimentation rate and, 402b-403b
 Atterberg limits, 150b-151b
 basement description convention, 23a-24a
 bathymetry, 11a, 45a, 78b
 bedrock topography, 64b
 benthic foraminifers, 382a-383a
 biogenic sediments, 112b-113b, 874b
 biogenic silica fluxes, 114a
 biogeography, 874b
 biology and oceanography, 35a-37a
 biostratigraphy, 24a, 26a-33a, 382a-384a
 calcareous nannofossils, 477b, 814b, 816b
 carbonate, 378b
 Chaetoceros resting spores, 601b, 604b
 clasts, 138b-139b
 clay mineralogy, 84b, 86b, 113b-114b, 874b
 coal-bearing sediments, 57b-60b, 795b
 combustible carbon and nitrogen, 114a-115a
 core handling, 15a-17a
 CTD profile, 109a, 110a
 data processing and availability, 23b-24b
 depositional environment, preglacial, 18b
 diamictites, 120b-121b
 diamicts, 88b-89b
 diatoms, 383a-384a, 605b

distribution patterns, 651b-655b
 analytical methods, 646b-648b
 chlorophyll *a* and density of, 653b, 659b
 distribution patterns, 661b-663b
 empty/full ratio, 655b, 656b, 657b, 663b, 670b-671b
 integrated abundances, 648b-651b, 652b, 664b
 phytoplankton vs. sediments, 668b-673b
 spring blooms, 671b-672b
 standing stations, 655b-660b, 664b
 in surface sediments, 390a
 zonation, 551b-557b, 816b-817b
 dikes, 24a, 44a
 drifting sediment trap deployment, 111a
 drilling deformation, 15a, 17a, 19a-20a
 erosion, ice thickness and, 178b-179b
Eucampia antarctica
 distribution, sediment and phytoplankton, 675b-676b, 678b
 morphology, summer vs. winter stages, 676b, 677b, 678b-679b
 size relationships, 677b-678b, 680b
 extractable organic matter
 source and maturity, 410b-412b, 414b
 extrusives, 24a, 44a
 fecal pellets, 670b-671b
 fining-upward sequences, 382a
 flat-lying sequence, 107b
 fluvio-lacustrine sediments, 59b
 geographic setting, 80b-81b
 geological evolution, 20b-22b
 geological setting, 5b-7b, 8a-9a, 46b, 58b, 79b, 80b-81b, 133b, 170b, 375b-376b, 377b, 407b
 geophysical logs, 177b
 glacial history, 78b-80b, 115b-122b, 180b, 875b-877b, 880b-888b
 glacial sediments, solute diffusion, 387b-388b
 glacial transport, 71b-73b, 114b-115b
 glaciation
 deep-sea hiatus association, 874b-875b
 isotopic evidence, 869b-870b
 pollen record, 873b-875b
 sediment record, 870b-873b
 glaciogenic sediments
 consolidation, 151b-152b, 155b-157b, 174b-176b
 depositional environment, 114b-115b
 genetic classification, 22a-23a, 101b, 103b
 glaciomarine deposition, 80b
 grain-size distribution, 81b, 149b-150b
 diamictites, 120b-121b
 granular sediment classification, 21a-22a, 23a, 43a-44a
 hinterland geology, 64b-68b
 hydrocarbon gases, 34a
 ice loading events, 176b, 178b, 179b-181b, 885b-886b
 ice rafted debris, 73b-74b, 873b, 886b
 inorganic carbon, 386a
 inorganic geochemistry, 35a, 384a, 395a
 interstitial-water geochemistry, 377b-378b, 384b-385b, 386a
 high-resolution sampling technique, 376b
 major cation concentrations, 380b, 391b
 Lambert Glacier-Amery Ice Shelf System
 flow pattern and particle paths, 70b-71b
 ice dynamics, 62b-64b
 structural glaciology, 68b-70b
 lithofacies, 86b, 88b-92b, 103b, 105b, 107b
 lithostratigraphy, 93b, 125b-126b, 394a-395a
 hiatuses, 95b, 97b, 99b
 lithologic Units I-IV, 380a-381a

- location, 6b, 135b, 300b, 376b, 417b, 428b, 468b, 513b, 647b, 668b, 751b, 796b, 814b, 871b
logging, 39a-40a
synthetic seismogram, 40a
magnetic properties, 33a-34a, 45a, 384a-385a
magnetobiochronology, 815b, 836b-837b
morphology, 7a-8a
navigation, 45a, 47a, 48a-85a, 87a, 88a
objectives, 13a, 377a-378a, 394a
ocean particulate fluxes, 109a-110a, 114a
operations, 380a
organic carbon, 113b, 378b
glacial constraints, 388b
organic geochemistry, 34a-35a, 385a-387a, 395a
pack ice coverage, 110a, 112a-113a
paleoclimatic background, 9a-10a, 13a
palyontology, 384a, 941b, 942b-943b
permeability, 152b, 154b
physical properties, 37a-38a, 389a, 390a-392a, 395a
geotechnical units G1-G3, 390a-391a
shipboard analysis, 147b-149b, 164b
phytoplankton, 388a, 390a
planktonic foraminifers, 382a-383a, 814b, 816b
oxygen isotopes, 249b
plant pigments, 118a-120a
prograding sequence, 876b
radiolarians, 44a, 814b, 816b, 817b
redox and carbon flux data, 402b
Rock-Eval pyrolysis, 35a, 386a-387a, 388a, 409b
salinity, 107a
sediment classification, 21a
sediment composition, 81b-84b, 383b-384b
sedimentation rate, 384a, 401b-404b
seismic stratigraphy, 9a, 11a, 12a, 392a-394a
correlation, Site 742, 446a
reflection profiling, 7b, 10b, 25b, 36b-37b, 45a, 86a, 96a, 106a
units I-II, 392a-393a
seismic velocity log, 33b-34b
shear strength, 154b-156b
sonobuoy seismic recording, 28b-29b, 45a-46a, 107a, 378a-380a
stable isotopes, 114a-115a
steroid hydrocarbons, 422b
structural features, 17b-18b
sulfate, sedimentation rate and, 402b
sulfur content, 113b
surface-water masses, 872b
temperature, 107a
triterpenoid hydrocarbons, 421b
velocity, 46a, 107a
wind and weather summary, 388a
- Site 742**
acoustic units PS.1-PS.6, 7b-13b, 27b-28b
areal distribution, 13b-17b
lithostratigraphic correlation, 99b-101b
aliphatic hydrocarbons, 418b-423b
alkalinity, 388b-389b
ammonia
sedimentation rate and, 402b-403b
amphibolite facies, 453a
Atterberg limits, 150b-151b
basement description convention, 23a-24a
bathymetry, 11a, 45a, 78b
bedrock topography, 64b
benthic foraminifers, 413a
biogenic sediments, 112b-113b, 874b
biogenic silica fluxes, 114a
biogeography, 874b
- biology and oceanography, 35a-37a
biostratigraphy, 24a, 26a-33a, 413a-415a, 845b
calcareous nannofossils, 413a, 477b, 814b, 816b
calcium/magnesium ratio, 420a
carbonate, 378b
thermodynamic modeling, 389b-391b
Chaetoceros resting spores, 601b, 604b
clasts
metamorphic, 137b, 138b
provenance, 138b-139b
sedimentary, 135b, 137b
shapes, 97b
clay mineralogy, 84b, 86b, 113b-114b, 874b
coal-bearing sediments, 57b-60b
combustible carbon and nitrogen
stable isotopic fractionation, 115a, 117a
core handling, 15a-17a
coring summary, 402a
CID profile, 109a, 110a
data processing and availability, 23b-24b
density-porosity relationship, 281b
depositional environment, 448a-451a
glacial, 18b-20b
diamictites, 120b-121b
diamicts, 88b-89b, 107b
grain fabrics, 91b-92b
diatoms, 413a-414a, 605b
analytical methods, 646b-648b
chlorophyll *a* and density of, 653b, 659b
distribution patterns, 651b-655b, 661b-663b
empty/full ratio, 655b, 656b, 657b, 663b, 670b-671b
integrated abundances, 648b-651b, 652b, 664b
phytoplankton vs. sediments, 668b-673b
reworking, 744b
spring blooms, 671b-672b
standing stations, 655b-660b, 664b
in surface sediments, 425a
zonation, 551b-557b, 816b-817b
dikes, 24a, 44a
drifting sediment trap deployments, 111a
drilling deformation, 15a, 17a, 19a-20a, 403a
erosion, ice thickness and, 178b-179b
Eucampia antarctica
distribution, sediment and phytoplankton, 675b-676b, 678b
morphology, summer vs. winter stages, 676b-677b, 678b-679b
size relationships, 677b-678b, 680b
extrusives, 24a, 44a
fecal pellets, 670b-671b
flat-lying sequence, 107b, 876b
geographic setting, 80b-81b
geological evolution, 20b-22b
geological setting, 5b-7b, 8a-9a, 46b, 58b, 79b, 80b-81b, 133b, 170b, 375b-376b, 377b
geophysical logs, 177b, 270b
glacial history, 78b-80b, 115b-122b, 180b, 875b-877b, 880b-888b
Miocene-Pliocene transition, 248b-249b
glacial sediments, solute diffusion, 387b-388b
glacial transport, 71b-73b
glaciation
deep-sea hiatus association, 874b-875b
isotopic evidence, 869b-870b
onset, 201b
pollen record, 873b-875b
sediment record, 870b-873b
glaciogenic sediments
consolidation, 151b-152b, 155b-157b, 174b-176b
depositional environment, 114b-115b
genetic classification, 22a-23a, 101b, 103b
- glaciomarine deposition, 80b
grain orientation, 99b, 105b
grain-size distribution, 81b, 149b-150b
diamictites, 120b-121b
by facies type, 94b
gravel, 82b
granular sediment classification, 21a-22a, 23a, 43a-44a
granulite facies, 453a-454a
gravel, 82b
clast petrography, 452a-453a
greenschist to epidote-amphibolite facies, 453a
hinterland geology, 64b-68b
hydrocarbon gases, 34a, 420a-421a, 422a, 423a
hydrogen isotopes, 387b
ice grounding, 452a
ice loading events, 176b, 178b, 179b-181b, 885b-886b
ice rafted debris, 73b-74b, 873b, 886b
inorganic carbon, 423a
inorganic geochemistry, 35a, 415a, 418a-420a
interstitial-water geochemistry, 377b-378b, 385b, 418a, 419a
high-resolution sampling technique, 376b
major cation concentrations, 380b, 391b
solute distribution, 387b
Lambert Glacier-Amery Ice Shelf System
flow pattern and particle paths, 70b-71b
ice dynamics, 62b-64b
structural glaciology, 68b-70b
lithofacies, 86b, 88b-92b, 103b, 105b, 107b, 110b
lithologic column, 271b
lithology
log-inferred, 279b-283b, 285b-286b
lithostratigraphy, 93b, 126b-127b, 269b, 402a-413a, 448a-451a, 877b
hiatuses, 95b, 97b, 99b
lithologic Units I-VI, 402a-412a
mixed facies Unit III, 108b
mudstone Subunit VIA, 111b
living bacteria, 687b-689b
location, 6b, 135b, 264b, 300b, 376b, 397a, 417b, 428b, 468b, 513b, 647b, 668b, 751b, 814b, 871b
logging, 39a-40a, 451a
correlation, Site 739, 442a
operations, 435a-436a
summary, 455a-458a
synthetic seismogram, 40a
units 1-4, 281b-283b, 436a-437a
magnetic properties, 33a-34a, 45a, 415a, 416a-417a, 451a, 769b, 770b
lithologic correlation, 759b-760b, 768b
magnetobiochronology, 815b, 837b-838b, 843b
magnetostratigraphy, 845b
mineralogy, lithologic correlation, 282b-283b, 284b
morphology, 7a-8a
navigation, 45a, 47a, 48a-85a, 87a, 88a
objectives, 13a, 397a-398a
ocean particulate fluxes, 109a-110a, 114a
opal, 84b
operations, 400a, 402a
organic carbon, 113b, 378b
glacial constraints, 388b
organic geochemistry, 34a-35a, 420a-422a, 423a, 424a
oxygen isotopes, 387b
pack ice coverage, 110a, 112a-113a
paleoclimatic background, 9a-10a, 13a
palyontology, 414a-415a, 941b-942b, 943b-944b
permeability, 152b, 154b
physical properties, 37a-38a, 425a-435a, 451a

SITE INDEX

geotechnical units G1–G6, 426a, 428a
 lithologic correlation, 146b
 shipboard analysis, 147b–149b, 164b–166b
 phytoplankton, 423a, 425a
 planktonic foraminifers, 413a, 814b, 816b
 plant pigments, 118a–120a
 porosity, 274b
 prograding sequence, 876b, 880b
 radiolarians, 44a, 414a, 814b, 816b, 817b
 redox and carbon flux data, 402b
 Rock-Eval pyrolysis, 35a, 380b, 422a, 424a, 426a
 salinity, 107a
 sand/mud ratio, 95b
 sediment classification, 21a
 sediment composition, 81b–84b, 385b
 sedimentary succession, 105b
 sedimentation rate, 401b–404b, 415a
 seismic stratigraphy, 9a, 11a, 12a
 correlation, Sites 739 and 741, 446a–447a
 reflection profiling, 7b, 10b, 36b–37b, 45a, 86a, 97a–99a, 106a
 units I–VI, 443a–446a, 451a–452a
 seismic structures, 446a
 seismic velocity log, 29b–30b, 34b–35b
 shear strength, 154b–156b
 slumping, 112b
 sonobuoy seismic recording, 45a–46a, 107a, 398a–410a, 401a, 447a
 refraction data, 28b–29b
 steroid hydrocarbons, 422b
 stratigraphic intervals, 119b
 stratigraphic summary, 878b
 structural features, 17b–18b
 sulfate, sedimentation rate and, 402b
 sulfur content, 113b
 summary stratigraphic log, 101b, 386b
 surface-water masses, 872b
 synthetic seismogram, 437a–443a
 temperature, 107a
 thermal conductivity, 428a, 431a
 triterpenoid hydrocarbons, 421b
 velocity, 46a, 107a
 wind and weather summary, 426a

Site 743

acoustic units PS.1–PS.6, 7b–13b, 27b–28b
 areal distribution, 13b–17b
 lithostratigraphic correlation, 99b–101b
 alkalinity, 388b–389b
 ammonia, sedimentation rate and, 402b–403b
 Atterberg limits, 150b–151b
 basement description convention, 23a–24a
 bathymetry, 11a, 45a, 78b
 bedrock topography, 64b
 benthic foraminifers, 465a
 biogenic sediments, 112b–113b, 874b
 biogeography, 874b
 biology and oceanography, 35a–37a
 biostratigraphy, 24a, 26a–33a, 465a
 calcareous nannofossils, 477b, 814b, 816b
 carbon isotopes, 740b, 742b
 carbonate, 378b
 thermodynamic modeling, 389b–391b
Chaetoceros resting spores, 601b
 clasts
 provenance, 138b–139b
 sedimentary, 135b, 137b
 clay mineralogy, 84b, 86b, 113b–114b, 874b
 coal-heaving sediments, 57b–60b
 core handling, 15a–17a
 CTD profile, 109a, 110a
 data processing and availability, 23b–24b
 depositional environment, 475a

diamictites, 120b–121b
 diamicts, 88b–89b
 diatoms, 465a
 analytical methods, 646b–648b
 chlorophyll *a* and density of, 653b, 659b
 distribution patterns, 651b–655b, 661b–663b
 empty/full ratio, 655b, 656b, 657b, 663b
 integrated abundances, 648b–651b, 652b, 664b
 phytoplankton vs. sediments, 668b–673b
 spring blooms, 671b–672b
 standing stations, 655b–660b, 664b
 zonation, 551b–557b, 816b–817b
 dikes, 24a, 44a
 drilling deformation, 15a, 17a, 19a–20a, 461a–462a
 erosion, ice thickness and, 178b–179b
Eucampia antarctica
 distribution, sediment and phytoplankton, 675b–676b, 678b
 morphology, summer vs. winter stages, 676b–677b, 678b–679b
 size relationships, 677b–678b, 680b
 extrusives, 24a, 44a
 flat-lying sequence, 107b
 geographic setting, 80b–81b
 geological evolution, 20b–22b
 geological setting, 5b–7b, 8a–9a, 46b, 58b, 79b, 80b–81b, 133b, 170b, 375b–376b, 377b
 geophysical logs, 177b
 glacial history, 78b–80b, 115b–122b, 180b, 875b–877b, 880b–888b
 glacial sediments, solute diffusion, 387b–388b
 glacial transport, 71b–73b
 glaciation
 deep-sea hiatus association, 874b–875b
 isotopic evidence, 869b–870b
 pollen record, 873b–875b
 sediment record, 870b–873b
 glaciogenic sediments
 consolidation, 151b–152b, 155b–157b, 174b–176b
 depositional environment, 114b–115b
 genetic classification, 22a–23a, 101b, 103b
 glaciomarine deposition, 80b
 grain-size distribution, 81b, 149b–150b
 diamictites, 120b–121b
 by facies type, 94b
 granular sediment classification, 21a–22a, 23a, 43a–44a
 hinterland geology, 64b–68b
 hydrocarbon gases, 34a, 466a
 ice loading events, 176b, 178b, 179b–181b, 885b–886b
 ice-raftered debris, 73b–74b, 873b, 886b
 inorganic carbon, 466a
 inorganic geochemistry, 35a, 465a–466a
 interstitial-water geochemistry, 377b–378b, 385b–387b, 466a
 high-resolution sampling technique, 376b
 major cation concentrations, 380b, 391b
 Lambert Glacier–Amery Ice Shelf System
 flow pattern and particle paths, 70b–71b
 ice dynamics, 62b–64b
 structural glaciology, 68b–70b
 lithofacies, 86b, 88b–92b, 103b, 105b, 107b
 lithostratigraphy, 95b, 128b, 460a–465a, 485a
 hiatuses, 95b, 97b, 99b
 lithologic Units I–III, 462a
 location, 6b, 135b, 300b, 376b, 428b, 459a, 468b, 513b, 647b, 668b, 751b, 814b, 871b
 logging, 39a–40a
 synthetic seismogram, 40a
 magnetic properties, 33a–34a, 45a, 465a
 magnetobiochronology, 815b, 846b
 mollusk shells, 740b
 morphology, 7a–8a
 navigation, 45a, 47a, 48a–85a, 87a, 89a
 objectives, 13a, 459a–460a
 operations, 460a, 474a–475a
 organic carbon, 113b, 378b
 glacial constraints, 388b
 organic geochemistry, 34a–35a, 466a, 467a, 475a
 oxygen isotopes, 740b, 742b
 pack ice coverage, 110a, 112a–113a
 paleoclimatic background, 9a–10a, 13a
 permeability, 152b, 154b
 physical properties, 37a–38a, 468a–473a, 475a
 geotechnical units G1–G3, 469a–470a
 lithologic correlation, 146b
 shipboard analysis, 147b–149b, 166b–167b
 phytoplankton, 466a–468a
 planktonic foraminifers, 465a, 814b, 816b
 prograding sequence, 459a, 474a, 876b
 radiolarians, 44a, 814b, 816b, 817b
 redox and carbon flux data, 402b
 Rock-Eval pyrolysis, 35a, 380b, 422a, 424a, 426a
 sediment classification, 21a
 sediment composition, 81b–84b, 385b
 sedimentation rate, 401b–404b
 seismic stratigraphy, 9a, 11a, 12a, 475a
 reflection profiling, 7b, 10b, 36b–37b, 45a, 86a, 100a–101a, 106a
 units I–II, 473a
 shear strength, 154b–156b
 sonobuoy seismic recording, 28b–29b, 45a–46a, 460a
 structural features, 17b–18b
 sulfate, sedimentation rate and, 402b
 sulfur content, 113b
 surface-water masses, 872b
 thermal conductivity, 470a, 473a
 velocity, 46a
 wind and weather summary, 469a

Site 744

ammonia
 pore waters, 395b
 sedimentation rate and, 402b–403b
 authigenic silicate formation, 218b–219b
 barite, 217b–218b
 basement description convention, 23a–24a
 bathymetry, 6a, 45a, 349b
 bathythermograph, 494a
 benthic foraminifers, 486a–487a, 614b–615b, 618b
 comparison, Site 738, 486a
 paleoenvironmental factors, 622b
 vertical ranges, 619b
 biogenic sediments, 874b
 biogenic silica, 884b
 biogeography, 874b
 biology and oceanography, 35a–37a
 biostratigraphic datum levels, 826b–828b
 Paleogene, 830b
 biostratigraphy, 24a, 26a–33a, 484a–488a, 824b, 825b
 calcareous nannofossils, 474b–476b, 482b, 484a, 503a, 888b
 magnetostratigraphic correlation, 475b
 zonation, 814b, 816b
 calcium, pore water, 397b–398b
 calcium carbonate, 898b–899b
 carbon isotopes
 Eocene–Miocene, 709b, 936b–939b
 Eocene–Oligocene transition, 710b–712b
 foraminifers, 696b, 698b–699b

- Oligocene deep-water association, 712b-714b
 Paleocene–Eocene, 698b, 704b-708b
 paleoenvironmental implications, 714b-715b
 surface- to intermediate-water gradients,
 697b, 704b, 709b-710b
 carbonate, 187b, 189b-190b, 367b, 935b
 carbonate carbon, 493a-494a
Chaetoceros resting spores, 601b, 604b
 chronostratigraphy, 437b-438b
 clay mineralogy, 114b, 194b, 874b
 environmental implications, 199b-200b
 Eocene, 882b
 Oligocene, 884b
 source, 198b-199b
 core handling, 15a-17a
 coring summary, 480a
 diatoms, 487a-488a, 559b, 564b-573b, 605b
 magnetostratigraphic correlation, 586b-587b
 stratigraphic recovery, 573b, 574b
 in surface sediments, 494a-495a
 zonation, 551b-557b, 816b-817b
 dikes, 24a, 44a
 drifting sediment trap deployments, 111a
 drilling deformation, 15a, 17a, 19a-20a
Eucampia antarctica
 distribution, sediment and phytoplankton,
 675b-676b, 678b
 morphology, summer vs. winter stages, 676b-
 677b, 678b-679b
 size relationships, 677b-678b, 680b
 extrusives, 24a, 44a
 feldspar, 217b
 geochemical logs, 911b, 925b, 928b
 geographic setting, 5a
 geological evolution, 7a
 geological setting, 353b
 glacial history, 201b-202b, 248b
 microfossil assemblage changes, 884b
 glaciation
 deep-sea hiatus association, 874b-875b
 isotopic evidence, 869b-870b
 onset, 201b
 pollen record, 873b-875b
 sediment record, 870b-873b
 glaciogenic sediments
 genetic classification, 22a-23a
 grain-size distribution
 nonbiogenic matter, 191b, 193b, 207b-208b
 granular sediment classification, 21a-22a, 23a,
 43a-44a
 heat flow, 499a
 hydrocarbon gases, 34a, 491a
 ice loading events, 885b-886b
 ice rafting, 195b, 197b-198b
 ice-raftered debris, 120b, 873b, 882b, 884b, 886b
 inorganic carbon, 493a-494a, 906b
 inorganic geochemistry, 35a, 490a-491a
 interstitial-water geochemistry, 367b, 369b,
 372b-373b, 491a, 492a
 high-resolution sampling technique, 348b,
 393b-399b, 903b
 major cation concentrations, 351b, 369b, 931b
 solute distribution, 368b
 lithology, 187b, 696b
 lithostratigraphy, 213b-214b, 479a-484a,
 502a-503a, 771b-772b, 935b
 lithologic Units I-II, 480a, 482a-484a
 Miocene hiatuses, 936b
 living bacteria, 687b-689b
 location, 185b, 186b, 212b, 300b, 324b, 350b,
 376b, 428b, 468b, 513b, 694b, 714b, 731b,
 751b, 814b
 logging, 39a-40a
 synthetic seismogram, 40a
- magnesium, pore water, 395b, 397b
 magnetic properties, 33a-34a, 45a, 488a-490a,
 503a, 783b-794b
 alternating-field demagnetization, 772b, 773b-
 780b
 susceptibility, 34a
 magnetic susceptibility logs, 895b, 896b-897b,
 898b
 magnetobiochronology, 815b, 819b-822b
 magnetostratigraphy, 824b, 825b
 biostratigraphic correlation, 781b-782b
 Paleogene, 830b
 polarity reversals, 772b, 781b
 subchron boundaries, 826b-828b
 mineralogy, 215b-217b
 nannofossil-diatom ooze transition, 195b, 202b,
 503a
 navigation, 45a, 47a, 48a-85a, 87a, 89a
 objectives, 478a
 ocean particulate fluxes, 109a-110a, 114a
 opal, 190b, 218b, 882b
 operations, 478a-479a
 organic carbon, 113b, 190b, 367b, 906b
 organic geochemistry, 34a-35a, 491a, 493a,
 503a
 oxygen isotopes, 826b-828b
 comparison, low- and mid-latitude sites, 713b
 deep-water association, 712b-714b
 Eocene, 882b
 Eocene–Holocene latitudinal gradient, 707b
 708b
 Eocene–Oligocene transition, 708b-709b,
 710b-712b
 foraminifers, 696b, 698b-699b
 Neogene, 887b, 936b-939b
 Oligocene, 712b-714b, 884b
 Paleocene, 698b, 704b-708b
 paleoenvironmental implications, 714b-715b
 surface- to intermediate-water gradients,
 697b, 704b, 709b-710b
 pack ice coverage, 110a, 112a-113a
 physical properties, 37a-38a, 495a-500a, 501a,
 503a
 geophysical units G1–G2, 495a
 phytoplankton, 493a-494a
 planktonic foraminifers, 431b, 435b-437b,
 485a-486a, 814b, 816b
 plant pigments, 118a-120a
 preglacial conditions, 200b-201b
 quartz, 217b
 radiolarians, 488a, 519b
 evolutionary lineages, 529b-531b
 Neogene, 522b
 Oligocene, 523b
 Oligocene/Miocene boundary, 529b
 Paleogene datum events, 527b
 paleomagnetic correlation, 530b, 531b
 systematics, 531b-537b
 zonation, 44a, 814b, 816b, 817b
 redox and carbon flux data, 402b
 Rock-Eval pyrolysis, 35a
 sediment classification, 21a
 sediment composition, 187b-190b, 204b-205b,
 367b
 environmental implications, 194b-195b
 nonbiogenic matter, 190b
 sedimentation rate, 401b-404b, 490a
 seismic stratigraphy
 reflection profiling, 45a, 86a, 102a, 106a
 units I-IV, 500a-501a
 silica, pore water, 398b
 silicoflagellates, 933b-934b
 sonobuoy seismic recording, 45a-46a, 478a,
 479a
- strontium isotopes, 826b-828b
 foraminifer preservation, 732b-733b
 Oligocene–Miocene, 733b-736b
 Paleogene, 830b
 sulfate
 pore waters, 398b-399b
 sedimentation rate and, 402b
 surface-water masses, 872b
 tectonic history, 5a, 7a
 temperature, 499a, 500a
 terrigenous clay, 197b, 209b-210b
 thermal conductivity, 500a
 velocity, 46a
 water content, 906b
 wind and weather summary, 494a
 X-ray diffraction, 223b
- Site 745
 ammonia
 pore waters, 395b
 sedimentation rate and, 402b-403b
 ash layers
 geochemistry, 326b, 328b-333b
 petrography, 325b
 sources, 329b, 333b
 basement description convention, 23a-24a
 bathymetry, 6a, 45a, 349b
 bathythermograph, 525a
 benthic foraminifers, 514a
 biogenic sediments, 874b
 biogeography, 874b
 biology and oceanography, 35a-37a
 biostratigraphic datum levels, 819b
 biostratigraphy, 24a, 26a-33a, 514a-516a, 530a
 bottom current influence, 530a
 calcareous nannofossils, 514a, 814b, 816b
 calcium, pore water, 397b-398b
 calcium/magnesium ratio, 522a
 carbonate, 369b
 carbonate critical depth, 230b
Chaetoceros resting spores, 601b, 604b
 clasts, 257b, 530a
 clay mineralogy, 237b, 242b, 244b, 874b
 source, 228b-229b
 core handling, 15a-17a
 coring summary, 510a
 deep-water circulation, 232b
 diatomaceous clay, 252b
 diatoms, 514a-515a, 526a, 558b, 573b-575b,
 576b, 605b
 magnetostratigraphic correlation, 586b-587b
 in surface sediments, 520a
 zonation, 551b-557b, 816b-817b
 dikes, 24a, 44a
 downslope transport, 230b-231b
 drilling deformation, 15a, 17a, 19a-20a,
 508a-509a
 environmental setting, 244b, 246b-247b
Eucampia antarctica
 distribution, sediment and phytoplankton,
 675b-676b, 678b
 morphology, summer vs. winter stages, 676b-
 677b, 678b-679b
 size relationships, 677b-678b, 680b
 extrusives, 24a, 44a
 geochemical logs, 912b, 914b, 916b, 922b,
 925b, 928b, 945b
 geographic setting, 5a
 geological evolution, 7a
 geological setting, 353b
 glacial history, 247b-249b
 glaciation
 deep-sea hiatus association, 874b-875b
 isotopic evidence, 869b-870b

SITE INDEX

pollen record, 873b-875b
 sediment record, 870b-873b
 glaciogenic sediments
 genetic classification, 22a-23a
 glass geochemistry, 337b-343b
 grain-size distribution, 227b-228b, 229b
 tephra layers, 328b
 terrigenous component, 236b
 granular sediment classification, 21a-22a, 23a,
 43a-44a
 gravel, 243b
 hydrocarbon gases, 34a, 519a
 ice rafting, 195b, 231b-232b, 530a
 ice-rafterd debris, 120b, 247b, 873b
 inorganic carbon, 522a, 906b
 inorganic geochemistry, 35a, 516a-518a
 interstitial-water geochemistry, 369b-371b,
 372b-373b, 520a, 521a
 high-resolution sampling technique, 348b,
 393b-399b, 903b
 major cation concentrations, 351b, 371b, 931b
 solute distribution, 370b
 lithostratigraphy, 508a-514a, 530a
 lithologic Unit I, 509a-513a
 location, 226b, 239b, 300b, 324b, 350b, 376b,
 428b, 468b, 505a, 513b, 751b, 814b
 logging, 39a-40a
 synthetic seismogram, 40a
 magnesium, pore water, 395b, 397b
 magnetic properties, 33a-34a, 45a, 516a, 517a,
 518a, 752b-753b, 754b, 757b, 761b
 susceptibility, 34a
 magnetic susceptibility logs, 895b, 900b-901b
 magnetobiochronology, 815b, 817b-818b, 820b
 magnetostratigraphy, 752b-753b, 756b, 760b
 subchron boundaries, 819b
 navigation, 45a, 47a, 48a-85a, 89a
 objectives, 7a, 505a-506a
 operations, 507a, 538a
 organic carbon, 369b, 906b
 organic geochemistry, 34a-35a, 518a-519a,
 522a, 523a, 530a
 oxidation-reduction potential, 371b
 oxygen isotopes, 887b
 physical properties, 37a-38a, 520a, 522a-525a,
 527a-530a, 532a
 phytoplankton, 519a-520a
 planktonic foraminifers, 514a, 814b, 816b
 productivity, 229b-230b
 quartz, 237b
 radiolarians, 515a-516a, 516b, 519b-521b,
 524b-526b
 evolutionary lineages, 529b-531b
 Miocene-Holocene datum events, 525b
 systematics, 531b-537b
 zonation, 44a, 814b, 816b, 817b
 redox and carbon flux data, 402b
 Rock-Eval pyrolysis, 35a, 519a, 523a, 524a
 salinity, 107a
 sediment classification, 21a
 sediment composition, 227b, 235b, 254b-256b,
 369b
 glacial/interglacial cycles, 240b, 241b
 sedimentary cycles, 226b-229b
 origin, 229b-232b
 sedimentation rate, 401b-404b, 516a
 seismic stratigraphy, 525a-526a, 530a
 reflection profiling, 45a, 86a, 103a-104a,
 106a
 silica, pore water, 398b
 sonobuoy seismic recording, 45a-46a, 107a,
 506a, 507a, 508a, 509a, 534a, 537a-538a
 stratigraphic column, 327b
 sulfate

pore waters, 398b-399b
 sedimentation rate and, 402b
 surface-water masses, 872b
 tectonic history, 5a, 7a
 temperature, 107a, 524a, 533a
 tephra, geochronology, 336b
 terrigenous clay, 237b, 245b, 259b-260b
 terrigenous sand fraction, 229b
 terrigenous sediment
 grain-size distribution, 240b, 242b, 243b,
 258a
 thermal conductivity, 524a
 total alkali vs. silica, 333b, 334b
 velocity, 46a, 107a
 water content, 906b
 wind and weather summary, 525a

Site 746

ammonia, pore water, 395b
 ash layers
 petrography, 325b
 geochemistry, 326b, 328b-333b
 sources, 329b, 333b
 basement description convention, 23a-24a
 bathymetry, 6a, 45a, 349b
 biogenic sediments, 874b
 biogeography, 874b
 biology and oceanography, 35a-37a
 biostratigraphic datum levels, 822b
 biostratigraphy, 24a, 26a-33a, 542a-544a, 821b
 calcareous nannofossils, 476b, 542a, 888b
 zonation, 814b, 816b
 calcium, pore water, 397b-398b
 carbonate, 369b
 carbonate critical depth, 230b
 clasts, 257b
 clay mineralogy, 242b, 244b, 874b
 source, 228b-229b
 core handling, 15a-17a
 deep-water circulation, 232b
 diatomaceous clay, 252b-253b
 diatoms, 542a-543a, 575b, 578b-579b
 magnetostratigraphic correlation, 586b-587b
 stratigraphic recovery, 580b
 zonation, 551b-557b, 816b-817b
 dikes, 24a, 44a
 downslope transport, 230b-231b
 drilling deformation, 15a, 17a, 19a-20a
 environmental setting, 244b, 246b-247b
Eucampia antarctica
 distribution in plankton, 675b-676b
 distribution, sediment and phytoplankton,
 678b
 morphology, summer vs. winter stages, 676b-
 677b, 678b-679b
 sexual cycle, 679b-680b
 size relationships, 677b-678b
 extrusives, 24a, 44a
 feldspar, 237b
 geochemical logs, 912b, 914b, 916b, 922b,
 925b, 928b, 946b
 geographic setting, 5a
 geological evolution, 7a
 geological setting, 353b
 glacial history, 247b-249b
 glaciation
 deep-sea hiatus association, 874b-875b
 isotopic evidence, 869b-870b
 pollen record, 873b-875b
 sediment record, 870b-873b
 glaciogenic sediments
 genetic classification, 22a-23a
 glass geochemistry, 337b-343b
 grain-size distribution, 227b-228b

granular sediment classification, 21a-22a, 23a,
 43a-44a
 gravel, 244b
 hydrocarbon gases, 34a, 545a
 ice rafting, 195b, 231b-232b, 541a-542a
 ice-rafterd debris, 120b, 247b, 873b
 inorganic carbon, 548a, 906b
 inorganic geochemistry, 35a, 544a-545a
 interstitial-water geochemistry, 369b-371b,
 372b-373b, 547a
 high-resolution sampling technique, 348b,
 393b-399b, 903b
 major cation concentrations, 351b, 371b, 931b
 solute distribution, 370b
 lithostratigraphy, 550a
 location, 226b, 239b, 300b, 324b, 350b, 376b,
 428b, 468b, 513b, 751b, 814b
 logging, 39a-40a
 synthetic seismogram, 40a
 magnesium, pore water, 395b, 397b
 magnetic properties, 33a-34a, 45a, 544a, 545a,
 546a, 547a, 753b, 756b, 757b
 magnetobiochronology, 815b, 818b-819b,
 823b, 843b
 magnetostratigraphy, 550a-551a, 754b, 755b,
 760b, 821b
 subchron boundaries, 822b
 navigation, 45a, 47a, 48a-85a, 89a
 objectives, 7a, 537a
 organic carbon, 369b, 906b
 organic geochemistry, 34a-35a, 545a, 548a,
 551a
 oxidation-reduction potential, 371b
 oxygen isotopes, 887b
 physical properties, 37a-38a, 545a-548a, 549a,
 550a
 planktonic foraminifers, 814b, 816b
 productivity, 229b-230b
 radiolarians, 526b-527b, 543a-544a
 evolutionary lineages, 529b-531b
 Miocene-Holocene datum events, 525b
 systematics, 531b-537b
 zonation, 44a, 814b, 816b, 817b
 Rock-Eval pyrolysis, 35a
 sea-surface temperature, 550a-553a
 sediment classification, 21a
 sediment composition, 227b, 256b, 369b, 551a
 glacial/interglacial cycles, 240b, 242b
 sedimentary cycles, 226b-229b
 origin, 229b-232b
 sedimentation rate, 544a
 seismic stratigraphy, 539a, 548a
 reflection profiling, 45a, 86a
 silica, pore water, 398b
 sonobuoy seismic recording, 45a-46a
 stratigraphic column, 327b
 sulfate

Site 747

Globigerina eugapertura
 last appearance datum, 437b
Globorotalia zealandica
 last appearance datum, 436b
Globoturborotalita labiacrassata

last appearance datum, 437b
inorganic carbon, 904b-905b
lava, Cretaceous, 293b
location, 293b
objectives, 7a
surface-water masses, 872b

Site 748

Globigerina euapertura
last appearance datum, 437b
Globoturborotalita labiacrassata
last appearance datum, 437b
redox and carbon flux data, 402b
Subbotina linaperta
surface-water masses, 872b

Site 749

location, 293b
lava, Cretaceous, 293b

redox and carbon flux data, 402b
surface-water masses, 872b

Site 750

lava, Cretaceous, 293b
location, 293b
surface-water masses, 872b

Site 751

carbonate concentration minima, 195b
redox and carbon flux data, 402b
surface-water masses, 872b

Site 752

gray clay layer
Cretaceous/Tertiary boundary, 849b
redox and carbon flux data, 402b

Site 753

redox and carbon flux data, 402b

Site 754

redox and carbon flux data, 402b

Site 756

redox and carbon flux data, 402b

Site 757

redox and carbon flux data, 402b

Site 758

redox and carbon flux data, 402b

PALEONTOLOGICAL INDEX

Abathomphalus mayaroensis Total Range Zone, Kerguelen-Heard Plateau S: Site 738, 452b, 459b
Abathomphalus mayaroensis Zone, Kerguelen-Heard Plateau S: Site 738, 250a
abisectus, *Cyclicargolithus*
Kerguelen-Heard Plateau N: Site 737, 489b
first appearance datum, 177a
acarinata, *Acarinina*, Kerguelen-Heard Plateau S: Site 738, 443b
Acarinina bullbrookii Interval Zone, Kerguelen-Heard Plateau S: Site 738, 434b-435b
Acarinina collectae Partial Range Zone, Kerguelen-Heard Plateau S: Site 738, 435b
Acarinina matthewsae Interval Zone, Kerguelen-Heard Plateau S: Site 738, 435b
Acarinina mckannai Interval Zone, Kerguelen-Heard Plateau S: Site 738, 433b
Acarinina pentacamerata Interval Subzone, Kerguelen-Heard Plateau S: Site 738, 434b
Acarinina praepentacamerata Interval Zone, Kerguelen-Heard Plateau S: Site 738, 433b-434b
Acarinina primitiva Interval Zone, Kerguelen-Heard Plateau S: Site 738, 434b
Acarinina sp., Kerguelen-Heard Plateau S: Site 738, 443b
Acarinina spp., Kerguelen Plateau S, stable isotopes, 69b, 707b
Acarinina wilcoxensis Interval Zone, Kerguelen-Heard Plateau S: Site 738, 434b
Actinocyclus frysella Interval Zone
Kerguelen Plateau, 553b, 555b-556b
Kerguelen Plateau S: Site 744, 565b
Kerguelen Sediment Ridge: Site 746, 575b
Kerguelen-Heard Plateau N: Site 737, 559b
Actinocyclus ingens Partial Range Zone
Kerguelen Plateau, 551b
Kerguelen Plateau S: Site 744, 565b
Kerguelen Sediment Ridge: Site 745, 574b
Kerguelen-Heard Plateau N: Site 736, 558b
Actinocyclus ingens-Denticulopsis maccollumii
Partial Range Zone
Kerguelen Plateau, 556b
Kerguelen Plateau S: Site 744, 565b
Kerguelen-Heard Plateau N: Site 737, 556b
aculeata, *Bulimina*
Kerguelen Plateau, 622b

Kerguelen-Heard Plateau N: Site 736, 134a, 615b, 616b, 628b
alabamensis, *Bifarina*, Kerguelen-Heard Plateau S: Site 738, 464b
alata, *Rhizosolenia*
Prydz Bay: Site 739
chains, 315a
large diameter, 315a
Prydz Bay: Site 742, 423a
alazanensis, *Bulimina*, Kerguelen Plateau S: Site 744, 627b
alsatica, *Turrilina*, Kerguelen Plateau S: Site 744, 614b, 715b
altus, *Chiasmolithus*
Kerguelen-Heard Plateau N: Site 737, 470b, 489b, 490b
last occurrence, 833b
Kerguelen-Heard Plateau S: Site 738, last occurrence, 474b
Amauropsis, Prydz Bay: Site 739, 131b
amekiensis, *Cassigerinelloita*
Kerguelen Plateau S, 439b
Kerguelen-Heard Plateau S: Site 738, 449b
Amphyipyndax tylotus Zone, Kerguelen-Heard Plateau S: Site 738, 252a, 518b
amphitrite, *Lithomelissa*, Kerguelen-Heard Plateau S: Site 738, last common occurrence, 527b
amphitrite, *Lychnocanoma*, Kerguelen-Heard Plateau S: Site 738, first common occurrence, 527b
ampliapertura, *Globigerina*, Kerguelen Plateau S: Site 744, last appearance datum, 486a
angiporoidea, *Subbotina*
Kerguelen Plateau S: Site 744, 437b
Kerguelen-Heard Plateau S: Site 738, 447b
angulata, *Nitzschia*, Kerguelen-Heard Plateau N: Site 737, 180a
angulosa, *Trifarina*, Kerguelen-Heard Plateau N: Site 736, 615b
angustumbilicata, *Tenuitella*, Kerguelen Plateau S: Site 744, 449b
antarctica antarctica, *Eucampia*
Kerguelen-Heard Plateau N: Site 736, 682b
summer vegetative stage, 684b, 685b
winter growth stage, 683b
Prydz Bay
morphology, 676b

summer vegetative stage, 676b-677b, 679b
antarctica, *Eucampia*
Kerguelen Plateau S: Site 744, northern form, 493a-494a
Kerguelen Sediment Ridge, glacial/interglacial cycles, 230b, 232b
Kerguelen-Heard Plateau N: Site 737, living Pleistocene-Holocene, 194a
Kerguelen-Heard Plateau S: Site 738, 260a
Prydz Bay
integrated abundance, 678b
morphology, 679b
plankton vs. sediment concentration, 675b-676b, 678b
sexual cycle, 679b-680b
subspecies size relationships, 677b-678b, 680b
vs. north Kerguelen Plateau, 679b
winter growth stage, 678b-679b
Prydz Bay: Site 739, southern form, 315a
Prydz Bay: Site 740, winter forms, 364a, 365a, 369a
Prydz Bay: Site 741, winter forms, 388a
antarctica, *Martinotilla*, Kerguelen-Heard Plateau N: Site 736, 134a
antarctica, *Prunopyle*, Kerguelen Sediment Ridge: Site 745, first occurrence, 521b
antarctica recta, *Eucampia*
Kerguelen-Heard Plateau N: Site 736
summer vegetative stage, 685b
winter growth stage, 683b
Prydz Bay
morphology, 676b
summer vegetative stage, 677b, 679b
antarctica, *Saccospirys*, Kerguelen-Heard Plateau N: She 736, first common occurrence, 516b

PALEONTOLOGICAL INDEX

Kerguelen-Heard Plateau N: Site 736, 135a, 516b
 Prydz Bay; Site 742, 414a

Antarctiella denticulata Zone, Kerguelen Sediment Ridge: Site 745, 515a–516a, 519b
Antarctiella ewingi interval, Prydz Bay: Site 742, 414a

anthoporus, *Reinhardtites*, Kerguelen-Heard Plateau S: Site 738, first occurrence, 474b
antillea, *Orthomorphina*, Kerguelen-Heard Plateau S: Site 738, 627b
antiquum, *Eucyrtidium*, n. sp.
 Kerguelen Plateau, 535b–536b
 Kerguelen Plateau S: Site 744, 546b
 Kerguelen-Heard Plateau S: Site 738, first common occurrence, 529b
apanthesma, *Acarinina*, Kerguelen-Heard Plateau S: Site 738, 446b
aperitus, *Markalius*, Kerguelen-Heard Plateau S: Site 738, 509b
Appendicisporites- type spores, Prydz Bay: Site 741, 943b
appressus, *Planorotalites*, Kerguelen-Heard Plateau S: Site 738, 461b, 465b
araneafera, *Mitrocalpis*, Kerguelen-Heard Plateau N: Site 736, first common occurrence, 516b
archipilium, *Dictyophimus*
 Kerguelen Plateau S: Site 744, first common occurrence, 529b
 Kerguelen-Heard Plateau S: Site 738, first common occurrence, 527b
Archipilium sp., Kerguelen-Heard Plateau S: Site 738, 544b
arenacea, *Miliammina*, Prydz Bay: Site 740, 357a
asperum, *Calcyclus*
 Kerguelen Plateau S: Site 744, 546b
 Kerguelen-Heard Plateau S: Site 738, last common occurrence, 527b–528b
atlanticum bulbosum, *Chaetoceros*, Prydz Bay: Site 740, 601b
aurelia, *Clathrocyclus*
 Kerguelen Plateau, 544b
 Kerguelen-Heard Plateau S: Site 738, last common occurrence, 527b
australiformis, *Planorotalites*
 Kerguelen-Heard Plateau S: Site 738, 448b
 first appearance datum, 437b
australis, *Archaeoglobigerina*, Kerguelen-Heard Plateau S: Site 738, 461b, 463b
babylonis, *Lynchnocanoma*, Kerguelen-Heard Plateau S: Site 738, last common occurrence, 527b
barbadiensis, *Discoaster*, Kerguelen-Heard Plateau N: Site 737, 177a, 493b
barbata, *Centosphaera*, Kerguelen-Heard Plateau S: Site 738, 867b
barboi, *Rhizosolenia*
 Kerguelen Sediment Ridge: Site 745, 574b
 Kerguelen-Heard Plateau N: Site 736, last occurrence, 134a, 558b, 828b
barleanus, *Melonis*, Kerguelen-Heard Plateau N: Site 736, 628b
barri, *Globigerinatethka*, Kerguelen-Heard Plateau S: Site 738, 449b
barronii, *Nitzschia*
 Kerguelen Sediment Ridge: Site 745, first occurrence, 818b
 Kerguelen-Heard Plateau N: Site 737, 598b
beccariiformis, *Gavelinella*
 Kerguelen Plateau S, stable isotopes, 69b
 Kerguelen-Heard Plateau S: Site 738, 455b, 858b

last appearance datum, 437b
 Benthic foraminifers
 Antarctica, oxygen isotopes, 78b
 Kerguelen Plateau
 faunal change, Eocene/Oligocene boundary, 619b–620b
 long-ranging species, 622b
 Miocene productivity changes, 620b
 paleoenvironmental factors, 616b, 619b–622b
 sediment/water interface, 635b
 Kerguelen Plateau S
 stable isotopes, 694b, 704b–710b
 stable isotopes, Eocene/Oligocene transition, 710b–712b
 Kerguelen Plateau S: Site 744
 abundance, 613b, 618b
 analytical methods, 32a
 Antarctic Circumpolar Current influence, 620b, 622b
 assemblages 1–3, 486a–487a
 assemblages 1–4, 614b–615b
 bathyal-abyssal depositional environment, 615b
 biostratigraphic summary, 485a
 comparison, Kerguelen-Heard Plateau S: Site 738, 487a, 615b, 620b
 datum events, 31a
 Eocene/Oligocene boundary, 614b, 615b
 long stratigraphic ranges, 487a
 Oligocene–Miocene hiatus, 487a
 Oligocene–Miocene unconformity, 620b
 radiolarian-nannofossil correlation, 530b
 stable isotopes, 698b–699b, 936b–938b
 strontium isotopes, 732b–736b
 zonation, 26a–28a
 Kerguelen Sediment Ridge: Site 745
 analytical methods, 32a
 biostratigraphy, 514a
 datum events, 31a
 zonation, 26a–28a
 Kerguelen Sediment Ridge: Site 746
 analytical methods, 32a
 datum events, 31a
 zonation, 26a–28a
 Kerguelen-Heard Plateau N: Site 736
 abundance, 612b–613b, 621b
 analytical methods, 32a
 biostratigraphy, 133a–134a, 615b
 carbonate dissolution and organic supply, 616b, 620b
 datum events, 31a
 mid-bathyal water depth, 615b
 planktonic/benthic ratio, 131a
 upwelling and surface-water productivity, 156a
 Weddell Sea–Heard Island comparison, 615b–616b
 zonation, 26a–28a
 Kerguelen-Heard Plateau N: Site 737
 analytical methods, 32a
 assemblages 1–4, 179a
 biostratigraphic summary, 174a–176a
 datum events, 31a
 zonation, 26a–28a
 Kerguelen-Heard Plateau S: Site 738
 abundance, 614b
 analytical methods, 32a
 Antarctic Circumpolar Current influence, 620b, 622b
 assemblages 1–5, 250a–251a, 613b–614b
 bathyal-abyssal depositional environment, 615b
 biostratigraphic summary, 248a–249a

comparison, Kerguelen Plateau S: Site 744, 487a, 615b, 620b
 Cretaceous/Tertiary boundary, 455b, 857b–858b
 Danian, 457b
 datum events, 31a
 Eocene/Oligocene boundary, 251a, 613b, 615b
 Oligocene/Miocene unconformity, 613b, 620b
 stable isotopes, 700b–703b
 vertical ranges, 616b–617b
 zonation, 26a–28a
 Zones P18–P20, 251a
 Prydz Bay
 analytical methods, 32a
 datum events, 31a
 oxygen isotopes, 120b
 zonation, 26a–28a
 Prydz Bay: Site 739
 biostratigraphy, 304a
 sediment/water interface, 304a
 Prydz Bay: Site 740, biostratigraphy, 356a
 Prydz Bay: Site 741, biostratigraphy, 382a–383a
 Prydz Bay: Site 742, biostratigraphy, 413a
 Prydz Bay: Site 743, biostratigraphy, 465a
Bianholithus sparsus Zone, Kerguelen-Heard Plateau S: Site 738, 500b–501b
biconicum, *Eucyrtidium*, nov. comb.
 Kerguelen Plateau, 538b
 Kerguelen Sediment Ridge: Site 745, last common occurrence, 525b
bicornis, *Clathrocyclas*, last occurrence, 135a
bicornis, *Cycladophora*
 Kerguelen Sediment Ridge: Site 745, first common occurrence, 521b
 Kerguelen-Heard Plateau N: Site 736, first common occurrence, 516b
bidens, *Chiasmolithus*
 Kerguelen-Heard Plateau S: Site 738, 512b
 first occurrence, 497b
bifax, *Discoaster*, Kerguelen-Heard Plateau N: Site 737, 178a, 470b, 492b
bijugatus, *Zygrhablithus*, Kerguelen-Heard Plateau S: Site 738, 491b
bisecta filewiczii, *Reticulofenestra*
 Kerguelen Plateau S: Site 744, 491b
 Kerguelen-Heard Plateau N: Site 737, 490b
bisecta, *Reticulofenestra*
 Kerguelen Plateau S: Site 744, 491b
 last occurrence, 474b
Thalassiosira spinosa correlation, 821b
 Kerguelen-Heard Plateau N: Site 737, 490b
 Kerguelen-Heard Plateau S: Site 738, first occurrence, 472b
bisulcus, *Prinsius*
 Kerguelen-Heard Plateau S: Site 738, 505b, 511b
 first occurrence, 497b
 paleobiogeographic pattern, 502b
Blackites spinosus Zone, Kerguelen-Heard Plateau S: Site 738, 472b, 822b
Bogorovia veniamini-Rocella gelida Zones, Kerguelen-Heard Plateau N: Site 737, 180a
brasili, *Nothofagus*
 Prydz Bay: Site 741, 943b
 Prydz Bay: Site 742, 944b
brazieri, *Globoturborotalita*
 Kerguelen Plateau S, 440b
 Kerguelen Plateau S: Site 744, 437b, 447b
brevis, *Subbotina*
 Kerguelen Plateau S, 440b
 Kerguelen-Heard Plateau S: Site 738, 447b
broedermanni, *Acarinina*, Kerguelen-Heard Plateau S: Site 738, 444b

bullbrookii, *Acarinina*, Kerguelen-Heard Plateau S: Site 738, 444b
first appearance datum, 250a
bulloides, *Globigerina* Kerguelen-Heard Plateau N: Site 736, 132a
Kerguelen-Heard Plateau N: Site 737, 178a
bulloides, *Sphaeroidina*, Kerguelen-Heard Plateau N: Site 736, 628b
burckiana, *Thalassiosira* Kerguelen Sediment Ridge: Site 746, 575b
Kerguelen-Heard Plateau N: Site 737, 180a, 597b
first appearance datum, 185a
Prydz Bay: Site 739, 579b

Calcareous nannofossils Kerguelen Plateau, zonation, 468b, 814b, 816b
Kerguelen Plateau S: Site 744 analytical methods, 31a–32a
biostratigraphic summary, 485a
biostratigraphy, 474b–476b, 484b, 824b
combined CP17–CP19 zones, 484a
datum events, 30a–31a, 482b
Eocene/Oligocene boundary, 484a
magnetostratigraphic correlation, 475b
Oligocene/Miocene boundary, 474b, 484a
radiolarian-foraminifer correlation, 530b
Subzone CP15b, 484a
zonation, 26a–28a
Zone CP16, 484a
Zones CN1–CN5a, 476b
Kerguelen Sediment Ridge: Site 745 analytical methods, 31a–32a
biostratigraphy, 514a
datum events, 30a–31a
zonation, 26a–28a
Kerguelen Sediment Ridge: Site 746, 542a analytical methods, 31a–32a
biostratigraphy, 476b–477b
datum events, 30a–31a
zonation, 26a–28a
Kerguelen-Heard Plateau N: Site 736 analytical methods, 31a–32a
biostratigraphy, 131a, 469b
datum events, 30a–31a
upwelling and productivity changes, 156a
zonation, 26a–28a
Kerguelen-Heard Plateau N: Site 737 analytical methods, 31a–32a
biostratigraphic summary, 174a–176a
biostratigraphy, 177a–178a, 470b–471b
datum events, 30a–31a, 482b
disconformity, diatom-nannofossil ooze transition, 470b
Eocene/Oligocene boundary, 177a–178a, 218a, 470b
Oligocene–Miocene hiatus, 177a
zonation, 26a–28a
Kerguelen-Heard Plateau S: Site 738 analytical methods, 31a–32a
biostratigraphic summary, 248a–249a
biostratigraphy, 244a–247a
Campanian–Maestrichtian, 474b
combined CP2–CP3 zones, 473b
combined CP6–CP7 zones, 473b
combined CP10–CP11 zones, 473b
Cretaceous/Tertiary boundary, 245a–247a, 455b, 473b, 497b, 500b, 502b, 851b, 856b–857b
Danian, 501b, 508b
datum events, 30a–31a, 482b
Eocene hiatus, 244a
Eocene/Oligocene boundary, 245a, 472b, 474b
Maestrichtian high-latitude assemblage, 474b

magnetostratigraphic correlation, 474b, 501b
paleobiogeographic patterns, 502b, 503b
Paleogene/Neogene boundary disconformity, 472b
Pleistocene surface water influence, 472b
polar ordination analysis, 502b, 504b
relative abundance, 503b
reworked Cretaceous species, 502b
reworked species, 244a–245a
Santonian, 474b
Shannon-Weaver diversity values, 502b, 504b
stratigraphic distribution patterns, 497b–500b
Subzone CP1b, 473b
water temperature influence, 472b, 481b
zonation, 26a–28a, 500b–502b
Zone CP1b, 246a
Zone CP8, 473b
Zone CP16, 474b
Zone CP4, 473b
Zone NC23, 497b
Zones CP2–CP11, 245a
Zones NC12–NC17, 247a
Zones NC19b–NC23, 247a
Prydz Bay analytical methods, 31a–32a
biostratigraphy, 26a
datum events, 30a–31a
zonation, 26a–28a, 814b, 816b
Prydz Bay: Site 739
biostratigraphic summary, 481b
biostratigraphy, 304a, 477b, 743b
comparison, Kerguelen Plateau, 477b
distribution, 478b–480b
Paleocene, comparison, Weddell Sea, 481b
reworked, 304a
reworked Eocene, 477b
Prydz Bay; Site 742
biostratigraphy, 413a, 477b
depositional environment, 113b
californica, *Rouxia*, Kerguelen-Heard Plateau N: Site 737, 596b
callopismia, *Anthocyrtella* Kerguelen Sediment Ridge: Site 745 last appearance datum, 519b
last common occurrence, 516a
callousus, *Dictyophimus*, Kerguelen-Heard Plateau S: Site 738, first common occurrence, 527b
callousus, *Toweius*, Kerguelen-Heard Plateau S: Site 738, 493b
Calocyclus polyporus Zone, Kerguelen Plateau S: Site 744, 488a
Calocyclus semipolita evolutionary lineage, Kerguelen Plateau, 529b–530b, 532b
campanula, *Cycladophora*, Kerguelen Plateau S: Site 744, first common occurrence, 529b
campilacantha, *Actinomma*, n. sp. Kerguelen Plateau, 533b
Kerguelen Plateau S: Site 74, 543b
carnarvonensis, *Samlandia*, Kerguelen-Heard Plateau S: Site 738, 631b–632b
castracanei, *Chaetoceros*, Kerguelen Sediment Ridge: Site 745, long chains, 520a
Chaetoceros Kerguelen Sediment Ridge: Site 745, 520a
Prydz Bay, 672b
Austral spring blooms, 661b
resting spores, 679b
Chaetoceros sp. Kerguelen-Heard Plateau N: Site 736, 600b
resting spores, 607b, 608b, 609b
Prydz Bay, resting spores, 673b
Prydz Bay: Site 740, 601b
Prydz Bay: Site 743, 601b

challengerae, *Amphymenium*, Kerguelen-Heard Plateau N: Site 737, 517b–518b
chapmani, *Planorotalites*, Kerguelen-Heard Plateau S: Site 738, 448b
characteristicus, *Hemiaulus*, Prydz Bay: Site 739, 743b–744b
chascanova, *Acarinina*, Kerguelen-Heard Plateau S: Site 738, 444b
cheni, *Lithomelissa*, n. sp. Kerguelen Plateau, 533b, 544b
Kerguelen Plateau S: Site 744, last common occurrence, 529b
Chiasmoliths, Kerguelen Plateau, size-temperature relationships, 481b
Chiasmolithus bidens Zone, Kerguelen-Heard Plateau S: Site 738, 501b–502b
Chiasmolithus danicus Zone, Kerguelen-Heard Plateau S: Site 738, 501b
Chiasmolithus oamaruensis Zone Kerguelen-Heard Plateau N: Site 737, 470b
Kerguelen-Heard Plateau S: Site 738, 472b
Chiloguembelina, Kerguelen-Heard Plateau S: Site 738, 858b
Chiloguembelina sp. Kerguelen-Heard Plateau S: Site 738, 461b
biserial heterohelicids, 455b, 457b, 458b, 460b
Chiloguembelinids, Kerguelen-Heard Plateau S: Site 738, 457b
Chondrites Kerguelen-Heard Plateau N: Site 737, 636b, 639b, 641b
claystone burrows, 171a
Kerguelen-Heard Plateau S: Site 738, 636b, 637b
Cibicidoides, Kerguelen Plateau S, stable isotopes, 69b
Cibicidoides spp. Kerguelen Plateau S, stable isotopes, 706b, 708b, 709b, 884b
Kerguelen Plateau S: Site 744
stable isotopes, 936b–937b, 938b
strontium isotopes, 736b
ciperoensis, *Sphenolithus* Kerguelen-Heard Plateau N: Site 737, 470b
first appearance datum, 177a
circularis, *Saturnalis*, Kerguelen Sediment Ridge: Site 745, last common occurrence, 516a
circummodifer, *Rugotruncana*, Kerguelen-Heard Plateau S: Site 738, 461b, 463b
circumtabulata, *Aliscysta*, Kerguelen-Heard Plateau S: Site 738, 632b
circumtexta, *Streblacantha*, Kerguelen Sediment Ridge: Site 745, 520b
citae, *Globotruncanella*, Kerguelen-Heard Plateau S: Site 738, 452b, 461b, 463b
clausa, *Thrysocystis*, Kerguelen Plateau S: Site 744, 488a
closterium, *Nitzschia* Prydz Bay, 648b, 649b, 672b
austral spring bloom, 660b–661b, 671b
chlorophyll *a* and density of, 653b
distribution, 651b, 661b
seasonal variation, 669b
vs. *Nitzschia cylindrus*, 662b–663b
coalingensis, *Acarinina* Kerguelen Plateau S, 439b
Kerguelen-Heard Plateau S: Site 738, 444b
collectaea, *Acarinina* Kerguelen Plateau S, 439b
Kerguelen-Heard Plateau S: Site 738, 444b
columbiana, *Guembelitria*, Kerguelen-Heard Plateau S: Site 738, 448b

PALEONTOLOGICAL INDEX

complanata, *Stainforthia*, Kerguelen-Heard Plateau N: Site 736, 628b
compressus, *Planorotalites* Kerguelen-Heard Plateau S: Site 738, 465b
 first appearance datum, 457b
concinnus, *Neochiastostygus*, Kerguelen-Heard Plateau S: Site 738, 509b
congruenta, *Orthopithonella*, Kerguelen-Heard Plateau S: Site 738, 862b, 863b, 864b, 867b
conica, *Cycladophora*, Kerguelen-Heard Plateau S: Site 738, first common occurrence, 529b
conica, *Lynchnocanoma*, Kerguelen Plateau S: Site 744, last common occurrence, 529b
conithorax, *Saccospiris*, Kerguelen Sediment Ridge: Site 745, first common occurrence, 525b
connecta, *Globigerina*, Kerguelen-Heard Plateau N: Site 737, 178a
Conosphaeridium striatoconus Zone, Kerguelen-Heard Plateau S: Site 738, 631b
constans, *Biscutum*, Kerguelen-Heard Plateau S: Site 738, 509b, 510b
contiguum, *Pteropilum*, Kerguelen Plateau S: Site 744, 544b
Coptospora paradoxa Zone, Prydz Bay: Site 741, 943b
corethron, Prydz Bay, Austral spring blooms, 661b
coronata, *Triceraspis* Kerguelen Sediment Ridge: Site 745, last common occurrence, 526b
 Kerguelen-Heard Plateau N: Site 737, last common occurrence, 516b
coronum, *Biscutum*, Kerguelen-Heard Plateau S: Site 738, 474b
Coscinodiscus elliptopora–Actinocyclus ingens Zone Kerguelen Plateau S: Site 744, 487a
 Kerguelen Sediment Ridge: Site 745, 514a
 Kerguelen-Heard Plateau N: Site 736, 134a
 Kerguelen-Heard Plateau S: Site 738, 244a, 251a, 261a
Coscinodiscus kolbei–Rhizosolenia barboi Zone, Kerguelen Sediment Ridge: Site 745, 514a
Coscinodiscus lewisiatus Zone, Kerguelen-Heard Plateau N: Site 737, 180a
Coscinodiscus superbus Zone, Kerguelen Plateau S: Site 744, 488a
Coscinodiscus vulnificus Partial Range Zone Kerguelen Plateau, 552b
 Kerguelen Plateau S: Site 744, 565b
 Kerguelen Sediment Ridge: Site 745, 574b
 Kerguelen-Heard Plateau N: Site 736, 552b
Coscinodiscus vulnificus Zone Kerguelen Sediment Ridge: Site 745, 514a
 Kerguelen-Heard Plateau N: Site 736, 134a
Cosmoldiscus insignis Zone Kerguelen Plateau S: Site 744, 487a
 Kerguelen-Heard Plateau N: Site 736, 134a
costata, *Odontochitina*, Kerguelen-Heard Plateau S: Site 738, 631b
costatum, *Skeletonema*, Prydz Bay, winter stage morphology, 679b
crassaformis, *Globorotalia*, Kerguelen-Heard Plateau N: Site 736, 133a
creta, *Alabamina*, Kerguelen-Heard Plateau S: Site 738, 858b
cretacea, *Guembelitria*, Kerguelen-Heard Plateau S: Site 738, 463b
cretacea, *Prediscosphaera*, Kerguelen-Heard Plateau S: Site 738, 862b, 864b
cribrinata, *Chiloguembelina*, Kerguelen-Heard Plateau S: Site 738, 448b, 455b, 457b, 464b, 858b
criophilum, *Corethron*

Prydz Bay, 650b
 abundance, 653b, 654b
 auxospores, 670b
 distribution, 655b, 661b, 669b–670b
 inner vs. outer bay, 662b
 life stages, 672b
 seasonal variation, 669b
 Prydz Bay: Site 739, 315a
 Prydz Bay: Site 740, 364a
 Prydz Bay: Site 741, auxospore formation, 388a
 Prydz Bay: Site 742, auxospores, 423a
cristata, *Nannotetra*, Kerguelen-Heard Plateau S: Site 738, 472b
Crucicentricula nicobarica Partial Range Zone Kerguelen Plateau, 556b
 Kerguelen Plateau S: Site 744, 565b
Cruciplacolithus, Kerguelen-Heard Plateau S: Site 738, *Prinsius dimorphous* relationship, 502b
Cruciplacolithus tenuis Zone, Kerguelen-Heard Plateau S: Site 738, 501b
cubensis, *Chiloguembelina* Kerguelen Plateau S, stable isotopes, 69b, 708b
 Kerguelen Plateau S: Site 744, last appearance datum, 614b
 Kerguelen-Heard Plateau S: Site 738, 628b
cubensis, *Bolivinopsis* Kerguelen Plateau S: Site 744, last appearance datum, 614b
 Kerguelen-Heard Plateau S: Site 738, 628b
cubensis, *Karreriella*, Kerguelen-Heard Plateau S: Site 738, 627b
cucullaris, *Ceratocyrtis*, Kerguelen Plateau S: Site 744, 529b
cuneicamerata, *Acarinina*, Kerguelen-Heard Plateau S: Site 738, 444b
curta, *Nitzschia* Prydz Bay, 649b, 672b
 abundance, 653b
 austral spring bloom, 671b
 seasonal variation, 669b
 Prydz Bay: Site 741, 601b
curvata, *Rhizosolenia*, Kerguelen-Heard Plateau N: Site 737, living Pleistocene–Holocene, 194a
Cycladophora spongotorax Zone, Kerguelen-Heard Plateau N: Site 737, absence of, 831b
Cyclampterium milowii group, Kerguelen Plateau S: Site 744, last occurrence, 488a
cylindrica, *Antarctissa* Kerguelen Sediment Ridge: Site 745, last appearance datum, 519b–520b
 Kerguelen-Heard Plateau N: Site 736, last appearance datum, 516b
cylindrica, *Nitzschia* Kerguelen Sediment Ridge: Site 746, first occurrence, 575b
 Kerguelen-Heard Plateau N: Site 737, 598b
 Prydz Bay: Site 739, 579b
cylindrurus, *Nitzschia* Kerguelen Sediment Ridge: Site 745, 520a
 Prydz Bay, 648b–649b, 672b
 austral spring bloom, 660b–661b, 671b
 distribution, 651b, 661b
 empty/full ratio, 652b–653b
 seasonal variation, 669b
 vs. *Nitzschia closterium*, 662b–663b
 Prydz Bay: Site 741, 601b
Cymatodon, n. gen., Kerguelen Plateau, 536b
Cyrtocapsella tetrapera Zone, Kerguelen Plateau S: Site 744, 488a, 519b

daniae, *Cribrosphaerella*, Kerguelen-Heard Plateau S: Site 738, 512b, 864b
danicus, *Chiasmolithus* Kerguelen-Heard Plateau S: Site 738, 245a, 502b, 512b
 first occurrence, 497b, 852b
daubjergensis, *Globoconusa* Kerguelen-Heard Plateau S: Site 738, 464b, 858b
 first appearance datum, 457b, 459b
 last appearance datum, 250a
davissii, *Reticulofenestra* Kerguelen-Heard Plateau N: Site 737, 491b
 Kerguelen-Heard Plateau S: Site 738, 472b
 Prydz Bay: Site 739, 304a, 477b, 743b
 Prydz Bay: Site 742, 413a, 477b, 838b, 881b
davisiiana, *Cycladophora* Kerguelen Sediment Ridge: Site 745, first occurrence, 525b
 Kerguelen-Heard Plateau N: Site 736, first occurrence, 137a
davisiiana, *Theocalyptra* Kerguelen Sediment Ridge, glacial/interglacial cycles, 230b, 234b
 Kerguelen Sediment Ridge: Site 745, 516a
 Kerguelen-Heard Plateau N: Site 736, first occurrence, 135a
decora, *Periphera*, Kerguelen-Heard Plateau S: Site 738, first common occurrence, 527b
delicatulum, *Actinomma*, nov. comb., Kerguelen Plateau, 537b
denticulata, *Antarctissa*, Kerguelen Sediment Ridge: Site 745, first common occurrence, 525b
denticuloides, *Nitzschia* Kerguelen Plateau S: Site 744, first occurrence, 565b
 Kerguelen-Heard Plateau N: Site 737, 598b
 first occurrence, 559b
 last occurrence, 180a, 559b
Denticulopsis dimorpha Acme Zone Kerguelen Plateau, 556b
 Kerguelen Plateau S: Site 744, 565b
 Kerguelen-Heard Plateau N: Site 737, 559b
Denticulopsis hustedtii Zone Kerguelen Sediment Ridge: Site 745, 515a
 Kerguelen Sediment Ridge: Site 746, 543a
 Kerguelen-Heard Plateau N: Site 737, 180a
 Prydz Bay: Site 739, 305a
Denticulopsis hustedtii–Nitzschia grossepunctata Partial Range Zone Kerguelen Plateau, 556b
 Kerguelen Plateau S: Site 744, 565b
Denticulopsis hustedtii–Denticulopsis lauta Zone Kerguelen Plateau S: Site 744, 487a
 Kerguelen Sediment Ridge: Site 746, 543a
 Kerguelen-Heard Plateau N: Site 737, 180a
 Kerguelen-Heard Plateau S: Site 738, 251a
Denticulopsis hustedtii/Nitzschia reinholdii zonal boundary, Kerguelen-Heard Plateau N: Site 737, 180a
Denticulopsis hyalina Zone, Kerguelen-Heard Plateau N: Site 737, 180a
Denticulopsis maccollumii Partial Range Zone Kerguelen Plateau, 556b
 Kerguelen Plateau S: Site 744, 565b
Denticulopsis praedimorpha Partial Range Zone Kerguelen Plateau, 556b
 Kerguelen Plateau S: Site 744, 565b
 Kerguelen-Heard Plateau N: Site 737, 556b
depressa, *Spirolocolina*, Kerguelen-Heard Plateau N: Site 736, 628b
Desmospyris spongiosa Zone Kerguelen Plateau S: Site 744, 488a, 519b

PALEONTOLOGICAL INDEX

Kerguelen Sediment Ridge: Site 745, 516a, 526b
 Kerguelen Sediment Ridge: Site 746, 514b, 526b
 Kerguelen-Heard Plateau N: Site 737, 181a
 Kerguelen-Heard Plateau S: Site 738, 252a, 518b
diablocensis, *Theocyrtis*
 Kerguelen Plateau S: Site 744, 546b
 Kerguelen-Heard Plateau S: Site 738, first common occurrence, 529b
diasystus, *Discoaster*, Kerguelen-Heard Plateau S: Site 738, first occurrence, 473b
 Diatoms
 Kerguelen Plateau
 high productivity correlation, 603b
 sedimentological and environmental changes influencing, 601b
 spore percentage, 601b
 zonation, 551b-557b, 816b-817b
 Kerguelen Plateau S: Site 744, 566b-569b, 570b-572b, 601b, 606b
 abundance, 604b
 analytical methods, 33a
 biostratigraphic summary, 485a, 554b
 biostratigraphy, 487a-488a, 559b, 565b, 573b, 824b
 magnetostratigraphic correlation, 559b, 565b, 573b, 588b-589b
 Miocene hiatuses, 487a, 565b
 stratigraphic recovery, 573b, 574b
 in surface sediments, 494a-495a
 zonation, 26a-28a, 31a
 Kerguelen Sediment Ridge: Site 745, 526a, 576b, 601b, 606b
 abundance, 604b
 analytical methods, 33a
 biostratigraphic summary, 554b
 biostratigraphy, 514a-515a, 573b-575b, 578b
 magnetostratigraphic correlation, 573b-575b
 Miocene/Pliocene boundary, 515a
 zonation, 26a-28a, 31a
 Kerguelen Sediment Ridge: Site 746, 542a-543a, 580b-581b
 analytical methods, 33a
 biostratigraphic summary, 554b
 biostratigraphy, 575b
 magnetostratigraphic correlation, 575b
 Miocene/Pliocene boundary, 542a-543a
 stratigraphic recovery, 582b
 zonation, 26a-28a, 31a
 Kerguelen-Heard Plateau N: Site 736, 560b-561b, 600b, 605b
 abundance, 604b
 analytical methods, 33a
 biostratigraphic summary, 554b
 biostratigraphy, 134a, 557b-558b
 low diversity assemblages, 156a
 zonation, 26a-28a, 31a
 Kerguelen-Heard Plateau N: Site 737, 562b-563b, 564b, 600b
 abundance, 604b
 analytical methods, 33a
 biostratigraphic summary, 174a-176a, 554b
 biostratigraphy, 179a-180a, 558b-559b
 burrowed vs. nonburrowed intervals, 180a
 Miocene hiatuses, 559b
 Miocene/Pliocene boundary, 559b
 Pliocene-Quaternary hiatus, 179a, 558b
 zonation, 26a-28a, 31a
 Kerguelen-Heard Plateau S: Site 738, 600b-601b
 abundance, 604b
 analytical methods, 33a
 biostratigraphic summary, 248a-249a
 biostratigraphy, 251a-252a

Neogene hiatuses, 252a
 in Quaternary sediments, 261a
 zonation, 31a
 Prydz Bay, 112b
 abundance, absolute vs. relative, 651b, 652b
 analytical methods, 33a, 646b-648b
 austral spring bloom, 660b-661b, 671b
 chlorophyll *a* and density of, 656b-657b, 659b-660b, 661b, 663b, 672b
 in clayey sediments, 672b
 distribution, 651b-655b, 661b-663b
 empty/full ratio, 650b-652b, 655b, 663b, 670b-671b
 high productivity correlation, 603b
 inner vs. outer bay, 662b, 671b
 integrated abundances, 648b-651b
 large- vs. small-celled, 662b, 663b, 669b
 pack ice effect, 663b
 physical-biological structure relationship, 651b, 663b
 plankton vs. sediment concentration, 669b, 673b, 675b-676b
 principal component analysis, 660b, 663b, 665b
 sedimentological and environmental changes influencing, 601b
 sequential changes during growing season, 662b-663b
 spore percentage, 601b
 standing stock, 655b-660b
 station clustering, 661b-662b, 665b
 summer surface water layer, 653b, 655b
 in surface sediments, 668b-673b
 terrigenous influx reflection, 672b-673b
 vertical stratification, 662b
 zonation, 26a-28a, 31a, 816b-817b
 Prydz Bay: Site 739, 139b, 586b-587b, 601b, 605b
 abundance, 604b
 biostratigraphy, 304a-305a, 577b-579b, 743b-744b
 Oligocene succession, 743b-744b
 Oligocene/Miocene disconformity, 744b
 Oligocene-Miocene hiatus, 305a
 in Quaternary sediments, 315a
 reworked Oligocene species, 579b
 reworking, 744b
 stratigraphic log, 583b
 strontium isotopes, 579b
 Prydz Bay: Site 740, 601b, 605b
 abundance, 604b
 biostratigraphy, 356a
 in surface sediments, 365a
 Prydz Bay: Site 741, 601b, 605b-606b
 abundance, 604b
 biostratigraphy, 383a-384a
 in surface sediments, 390a
 Prydz Bay: Site 742, 601b, 606b
 abundance, 604b
 biostratigraphy, 413a-414a
 in surface sediments, 425a
 Prydz Bay: Site 743, 601b
 biostratigraphy, 465a
 living Holocene-Pleistocene, 191a-194a
 in surface sediments, 468a
 Southern Ocean
 datum events, 552b-553b
 zonation, 548b-551b, 555b
 zonation
 Ciesielski, 550b
 Gersonde and Burkle, 550b-551b
 Kerguelen Plateau vs. other Southern Ocean schemes, 549b-550b
 McCollum, 548b-549b

Schrader, 549b
 Weaver and Gombos, 549b-550b
dichaeta, *Chaetoceros*
 Prydz Bay, 650b
 abundance, 653b, 654b
 distribution, 655b-656b, 661b
 inner vs. outer bay, 662b
 seasonal variation, 669b
 Prydz Bay: Site 740, 601b
dictyocha, Kerguelen-Heard Plateau S: Site 738, 251a
Dictyomitra sp., Kerguelen-Heard Plateau S: Site 738, 252a
dimorpha areolata, *Denticulopsis*, Kerguelen Sediment Ridge: Site 746, 575b
dimorpha, *Denticulopsis*
 Kerguelen Plateau S: Site 744
 first occurrence, 565b
 last occurrence, 565b
 Kerguelen Sediment Ridge: Site 746, 543a
 last common occurrence, 575b
 Kerguelen-Heard Plateau N: Site 737, 180a, 559b, 598b
dimorphosus, *Prinsius*
 Kerguelen-Heard Plateau S: Site 738, 497b, 505b, 509b
Cruciplacolithus relationship, 502b
 Dinoflagellates
 Kerguelen Plateau S: Site 744
 analytical methods, 33a
 zonation, 31a
 Kerguelen Sediment Ridge: Site 745
 analytical methods, 33a
 zonation, 31a
 Kerguelen Sediment Ridge: Site 746
 analytical methods, 33a
 zonation, 31a
 Kerguelen-Heard Plateau N: Site 736
 analytical methods, 33a
 zonation, 31a
 Kerguelen-Heard Plateau N: Site 737
 analytical methods, 33a
 zonation, 31a
 Kerguelen-Heard Plateau S: Site 738
 analytical methods, 33a
 Cretaceous sediments, 633b
 zonation, 31a
 Prydz Bay
 analytical methods, 33a
 zonation, 31a
Discoaster bifax Subzone, Kerguelen-Heard Plateau S: Site 738, 245a
Discoaster saipanensis Zone, Kerguelen-Heard Plateau S: Site 738, disconformity, 472b
Discoaster sp.
 Kerguelen-Heard Plateau N: Site 737, 492b, 493b
 Kerguelen-Heard Plateau S: Site 738, 491b
Discoaster spp., Kerguelen Plateau, Eocene, 481b
Discoaster sublodoensis Zone, Kerguelen-Heard Plateau S: Site 738, 245a, 472b
dissimilis, *Catapsydrax*
 Kerguelen Plateau S: Site 744, 447b
 last appearance datum, 437b
Distephanus speculum halionoma Subzone, Kerguelen Plateau S: Site 744, 821b
distichus, *Ellipsolithus*, Kerguelen-Heard Plateau S: Site 738, 493b
donahuensis, *Nitzschia*, Kerguelen-Heard Plateau N: Site 737, 598b
draco, *Bolinoides*, Kerguelen-Heard Plateau S: Site 738, 858b
druggii, *Manumiella*, Kerguelen-Heard Plateau S: Site 738, 631b-632b

PALEONTOLOGICAL INDEX

dubius, *Neococcolithes*, Kerguelen-Heard Plateau S: Site 738, 492b
duplicifysa, *Lithomelissa*, n. sp.
 Kerguelen Plateau, 534b
 Kerguelen Plateau S: Site 744, 544b
 Kerguelen-Heard Plateau S: Site 738, first common occurrence, 529b

earlandi, *Angulogerina*, Kerguelen-Heard Plateau N: Site 736, 134a, 628b
 Ebridians, Prydz Bay: Site 739, 139b
echinatus, *Catapsydrax*, Kerguelen Plateau S: Site 744, 447b
edwardsii, *Cruciplacolithus*
 Kerguelen-Heard Plateau S: Site 738, 497b, 511b
 first occurrence, 852b
ehrenbergi, *Lithomelissa*
 Kerguelen Plateau, 544b
 Kerguelen-Heard Plateau S: Site 738, last common occurrence, 527b
ehrenbergii, *Cribrosphaerella*, Kerguelen-Heard Plateau S: Site 738, 862b
Ehrenbergina sp., Kerguelen Plateau S: Site 744, 629b
elegans, *Pseudotextularia*, Kerguelen-Heard Plateau S: Site 738, 461b, 463b
elliptopora, *Coscinodiscus*
 Kerguelen Sediment Ridge: Site 745
 first occurrence, 818b
 last occurrence, 574b
 Kerguelen-Heard Plateau N: Site 736, 134a
 first occurrence, 558b
 last occurrence, 558b
elongata, *Buliminella*, Kerguelen-Heard Plateau N: Site 736, 134a
elongata, *Lychnocanoma*, Kerguelen Plateau S: Site 744, first common occurrence, 529b
ebulliooides, *Eoglobigerina*
 Kerguelen-Heard Plateau S: Site 738, 464b
 first appearance datum, 457b
eocaena, *Subbotina*, Kerguelen-Heard Plateau S: Site 738, 447b
Eoglobigerina fringa Partial Range Zone, Kerguelen-Heard Plateau S: Site 738, 250a
Eoglobigerina fringa Zone, Kerguelen-Heard Plateau S: Site 738, 457b
Eoglobigerina? sp., Kerguelen-Heard Plateau S: Site 738, 461b
esnaensis, *Acarinina*
 Kerguelen-Heard Plateau S: Site 738, 443b
 Danian/Selandian boundary, 457b
euapertura, *Globigerina*
 Kerguelen Plateau S: Site 744, 438b, 447b
 first occurrence, 437b
Eucyrtidium calvertense Zone
 Kerguelen Sediment Ridge: Site 745, 524b-525b
eugubina, *Parvulorugoglobigerina*, Kerguelen-Heard Plateau S: Site 738, 858b
Eurystomoskevos, n. gen., Kerguelen Plateau, 536b
exigua, *Epistominella*
 Kerguelen Plateau S: Site 744, 486a, 620b, 622b, 630b
 Kerguelen-Heard Plateau S: Site 738, 250a
expansus, *Chiasmolithus*, Kerguelen-Heard Plateau S: Site 738, 492b

falconensis, *Globigerina*, Kerguelen-Heard Plateau N: Site 737, 178a
 Fecal pellets, Prydz Bay, 670b-671b
fenestratus, *Clausicoccus*
 Kerguelen Plateau, 482b-483b
 Kerguelen Plateau S: Site 744, 491b

Kerguelen-Heard Plateau N: Site 737, 470b
 last occurrence, 470b
floralis, *Lithostrinus*, Kerguelen-Heard Plateau S: Site 738, last occurrence, 474b
floridanus, *Cyclicargolithus*
 Kerguelen-Heard Plateau N: Site 737, 489b
 Kerguelen-Heard Plateau S: Site 738, 476b
 Foraminifers. See Benthic foraminifers; Planktonic foraminifers
foremanae, *Lithielius*, Kerguelen-Heard Plateau S: Site 738, last common occurrence, 527b-528b
formosus, *Coccilithus*
 Kerguelen-Heard Plateau N: Site 737, 491b
 last occurrence, 470b
 Kerguelen-Heard Plateau S: Site 738, 472b
fraga, *Thalassiosira*, Kerguelen Plateau S: Site 744, first occurrence, 565b
frakesi, *Prunopyle*, Kerguelen-Heard Plateau S: Site 738, last common occurrence, 527b-528b
frequens, *Nephrolithus*
 Kerguelen-Heard Plateau S: Site 738, 497b, 862b
 first appearance datum, 452b
 first occurrence, 474b, 851b
fringa, *Eoglobigerina*, Kerguelen-Heard Plateau S: Site 738, 457b, 461b, 464b
Frondicularia spp., Kerguelen-Heard Plateau S: Site 738, 858b
fryxellae, *Actinocyclus*, n. sp.
 Kerguelen Plateau, 585b
 Kerguelen Plateau S: Site 744, 592b
 Kerguelen Sediment Ridge: Site 746, 592b
 Kerguelen-Heard Plateau N: Site 737, first occurrence, 559b
fulgens, *Nannotetrina*, Kerguelen-Heard Plateau S: Site 738, 472b

galeatus, *Pseudodictyophimus*, n. sp.
 Kerguelen Plateau, 534b
 Kerguelen Plateau S: Site 744, 544b
 Kerguelen-Heard Plateau S: Site 738, first common occurrence, 527b
gelida, *Reticulofenestra*
 Kerguelen Plateau, 483b
 Kerguelen Plateau S: Site 744, 474b
 Kerguelen-Heard Plateau N: Site 737, 177a, 488b
gelida, *Rocella*
 Kerguelen Plateau S: Site 744
 first occurrence, 565b
 last occurrence, 565b
gemma, *Tenuitella*, Kerguelen Plateau S: Site 744, 449b
gersondei, *Thalassiosira*, n. sp.
 Kerguelen Plateau, 585b
 Kerguelen Sediment Ridge: Site 746, 575b, 593b, 594b, 596b
 first occurrence, 575b
glabrum, *Isabelidinium*, Kerguelen-Heard Plateau S: Site 738, 631b
Globigerina brevis Zone, Kerguelen Plateau S: Site 744, 486a
Globigerinatheka index Partial Range Zone, Kerguelen-Heard Plateau S: Site 738, 435b
Globococonusa daubjergensis Interval Subzone, Kerguelen-Heard Plateau S: Site 738, 432b
Globococonusa daubjergensis Partial Range Subzone, Kerguelen-Heard Plateau S: Site 738, 459b
Globorotaloides suteri Partial Range Zone
 Kerguelen Plateau S: Site 744, 435b-436b
 Kerguelen-Heard Plateau S: Site 738, 435b
Globotruncanella sp., Kerguelen-Heard Plateau S: Site 738, 461b, 463b

Globoturborotalita labiacrassata Interval Zone, Kerguelen Plateau S: Site 744, 436b-437b
globulosa, *Heterohelix*
 Kerguelen-Heard Plateau S: Site 738, 463b
 recycled, 457b-458b
globulosus, *Gyroidinoides*, Kerguelen-Heard Plateau S: Site 738, 455b
globulosus, *Heterohelix*, Kerguelen-Heard Plateau S: Site 738, 858b
glomerata, *Adereotryma*, Prydz Bay: Site 740, 357a
golowniki, *Actinomma*
 Kerguelen Plateau S: Site 744, last common occurrence, 519b
 Kerguelen-Heard Plateau N: Site 737, 518b
gortanii, *Subbotina*, Kerguelen-Heard Plateau S: Site 738, 447b
gracilis, *Morozovella*
 Kerguelen-Heard Plateau S: Site 738, 446b
 high-latitude incursion, 439b
grande, *Lychnocanium*
 Kerguelen Sediment Ridge: Site 745, last common occurrence, 526b
 Kerguelen-Heard Plateau N: Site 737, last common occurrence, 516b
grandis, *Chiasmolithus*, Kerguelen-Heard Plateau S: Site 738, 492b
gravida, *Thalassiosira*, Prydz Bay: Site 739, 305a, 315a
grosspunctata, *Nitzschia*, Kerguelen Plateau S: Site 744, last occurrence, 565b
Guembelitrioides sp.
 Kerguelen Plateau S, 440b
 Kerguelen-Heard Plateau S: Site 738, 446b, 465b

haeckeli, *Lithomelissa*, Kerguelen-Heard Plateau S: Site 738, last common occurrence, 527b
hanckenina, Kerguelen-Heard Plateau S: Site 738, first appearance datum, 438b
hardingae, *Subbotina*, Kerguelen-Heard Plateau S: Site 738, 447b
harstenii, *Hemidiscus*, Prydz Bay: Site 739, 305a
havanensis, *Orthomorphina*, Kerguelen-Heard Plateau S: Site 738, 627b
hayesi, *Prunopyle*, Kerguelen-Heard Plateau S: Site 738, last common occurrence, 527b-528b
hebetata hiemalis, *Rhizosolenia*, Kerguelen-Heard Plateau N: Site 737, 598b
hebetata semispina, *Rhizosolenia*
 Kerguelen Sediment Ridge: Site 745, 520a
 Kerguelen-Heard Plateau S: Site 738, 260a
hedbergella sp., Kerguelen-Heard Plateau S: Site 738, 461b, 463b
Helicosphaera sp., Kerguelen-Heard Plateau S: Site 738, 493b
Heliotystols sp., Kerguelen-Heard Plateau S: Site 738, first common occurrence, 527b
Helotholus praevena-Pseudocubus vema evolutionary transition
 Kerguelen Sediment Ridge: Site 745, 525b-526b
 Kerguelen-Heard Plateau N: Site 737, 516b
Helotholus vema Zone
 Kerguelen Plateau S: Site 744, 488a, 519b
 Kerguelen Sediment Ridge: Site 746, 526b, 544a-545a
 Kerguelen-Heard Plateau N: Site 736, 135a
 Kerguelen-Heard Plateau N: Site 737, 181a
Helotholus vema/Desmospyris spongiosa zonal boundary, Kerguelen Plateau S: Site 744, 519b
hendeyi, *Chaetoceros*, Kerguelen Plateau S: Site 744, 494a
Heronallenia sp., Kerguelen Plateau S: Site 744, 629b

hesslandii, *Reticulofenestra*
Kerguelen Plateau S: Site 744, last appearance
datum, 484a
Kerguelen-Heard Plateau N: Site 737, 177a,
489b

Heterohelix, Kerguelen-Heard Plateau S: Site
738, 858b

heteropolaria, *Rouxia*
Kerguelen Sediment Ridge: Site 745, first
occurrence, 818b
Kerguelen-Heard Plateau N: Site 737, last
occurrence, 558b

hister, *Gondwanaria*, Kerguelen Plateau S: Site
744, first common occurrence, 529b

hornibrooki, *Subbotina*, Kerguelen Plateau S,
440b

Hornibrookina
Kerguelen-Heard Plateau S: Site 738
Cretaceous/Tertiary boundary, 504b
first appearance datum, 497b
first occurrence, 500b-501b

Hornibrookina teuriensis Zone, Kerguelen-Heard
Plateau S: Site 738, 501b

hughesi, *Diatrurus*, Kerguelen-Heard Plateau N:
Site 737, reworked, 181a

hughesii, *Cicatricosporites*, Prydz Bay: Site
741, 943b

hustedtii, *Denticulopsis*
Kerguelen Plateau S: Site 744
first occurrence, 565b
reworked, 487a
Kerguelen Sediment Ridge: Site 745, 514a
Kerguelen Sediment Ridge: Site 746, last
common occurrence, 575b
Kerguelen-Heard Plateau N: Site 737, 598b
first common occurrence, 559b
last common occurrence, 180a, 559b
Prydz Bay: Site 739, 305a

hyalina, *Denticulopsis*, Kerguelen-Heard Plateau
N: Site 737, first occurrence, 180a

hyalinus, *Asteromphalus*, Kerguelen Sediment
Ridge: Site 745, 519a-520a

imbricata, *Siphocampe*, Kerguelen Plateau, 545b

imitatus, *Planorotalites*, Kerguelen-Heard Pla-
teau S: Site 738, 448b, 457b

impensus, *Globigerinoides*
Kerguelen-Heard Plateau S: Site 738, 452b,
455b, 461b, 463b
recycled, 458b

imperatrix, *Coccineis*
Prydz Bay, 673b
Prydz Bay: Site 741, 390a

inca, *Astrophacus*, nov. comb.
Kerguelen Plateau, 537b

Kerguelen-Heard Plateau S: Site 738, last
common occurrence, 527b

incognita, *Paragloborotalia*, Kerguelen Plateau
S: Site 744, first occurrence, 437b

inconstans, *Morozovella*
Kerguelen-Heard Plateau S: Site 738, 465b
first appearance datum, 457b

index, *Globigerinatheka*
Kerguelen Plateau S: Site 744, last appearance
datum, 438b

Kerguelen-Heard Plateau S: Site 738, 449b
last appearance datum, 250a

inerme, *Corethron*
Kerguelen-Heard Plateau S: Site 738, 260a

Prydz Bay: Site 739, 315a

Prydz Bay: Site 740, 364a

Prydz Bay: Site 741, auxospore formation, 388a

Prydz Bay: Site 742, auxospores, 423a

inxpectata, *Lamprocyclas*, n. sp.

Kerguelen Plateau, 534b
Kerguelen Plateau S: Site 744, 529b, 545b
inflatum, *Eucyrtidium*, Kerguelen Sediment
Ridge: Site 745, last common occurrence,
526b

ingens, *Actinocyclus*
Kerguelen Plateau S: Site 744
first occurrence, 565b
reworked, 487a
Kerguelen-Heard Plateau N: Site 736, last
occurrence, 134a, 137a, 558b, 828b
Kerguelen-Heard Plateau N: Site 737, 596b
ingens ovalis, *Actinocyclus*, Kerguelen Sediment
Ridge: Site 746, 575b, 596b

interfrigidaria, *Nitzschia*, Kerguelen-Heard Pla-
teau N: Site 737, first occurrence, 180a

Inoceramus, Kerguelen-Heard Plateau S: Site
738, 637b

Inoceramus prisms, Kerguelen-Heard Plateau S:
Site 738, 251a

insignis, *Cosmiodiscus*
Kerguelen Sediment Ridge: Site 745, last
occurrence, 574b
Kerguelen-Heard Plateau N: Site 736, 558b,
828b
last common occurrence, 134a

insolita, *Praetenuitella*
Kerguelen Plateau S: Site 744, last appearance
datum, 436b
Kerguelen-Heard Plateau S: Site 738, 449b

interfrigidaria, *Nitzschia*, Kerguelen-Heard Pla-
teau N: Site 737, first appearance datum, 185a

intermedia, *Acarinina*, Kerguelen-Heard Plateau
S: Site 738, 444b

intermedius, *Abathomphalus*, Kerguelen-Heard
Plateau S: Site 738, 452b, 463b

interposita, *Acarinina*, Kerguelen-Heard Plateau
S: Site 738, 444b

intersectus, *Cosmiodiscus*, Kerguelen-Heard Pla-
teau N: Site 737, 597b

inura, *Thalassiosira*
Kerguelen Sediment Ridge: Site 745, 575b
Kerguelen-Heard Plateau N: Site 737, 597b

inversus, *Markalius*
Kerguelen Plateau S: Site 744, 491b
Kerguelen-Heard Plateau S: Site 738, 473b,
509b, 512b, 866b
Cretaceous/Tertiary boundary, 856b-857b

iridea, *Nonionella*, Kerguelen-Heard Plateau N:
Site 736, 615b, 629b

Isthmolithus recurvus Zone, Kerguelen-Heard Pla-
teau S: Site 738, 472b

jacksonii, *Thalassiosira*
Kerguelen Sediment Ridge: Site 746, first
occurrence, 575b
Kerguelen-Heard Plateau N: Site 737, 597b

januaria, *Nitzschia*, Kerguelen Sediment Ridge:
Site 746, 596b

japonica, *Cyrtocapsella*, Kerguelen-Heard Pla-
teau N: Site 737, 181a

jouseana, *Synedra*, Kerguelen Plateau S: Site
744, first occurrence, 573b

juvenile, *Globigerina*, Kerguelen-Heard Plateau
N: Site 737, 178a-179a

juvenilis, *Globigerinita*, Kerguelen Plateau S: Site
744, 439b

kennettii, *Asteromphalus*
Kerguelen Sediment Ridge: Site 746, 575b
last occurrence, 819b

kerguelensis, *Actinomma*, n. sp.
Kerguelen Plateau, 531b, 533b
Kerguelen Plateau S: Site 74, 543b

kerguelensis, *Botryostrobus*, n. sp.
Kerguelen Plateau, 535b
Kerguelen Plateau S: Site 744, 545b
Kerguelen-Heard Plateau S: Site 738, first
common occurrence, 527b

kerguelensis, *Nitzschia*
Kerguelen Plateau S: Site 744, 601b
in surface sediments, 494a
Kerguelen Sediment Ridge: Site 745, 526a, 601b
first occurrence, 818b

Kerguelen-Heard Plateau N: Site 737, living
Pleistocene-Holocene, 193a, 194a

Kerguelen-Heard Plateau S: Site 738, 601b

Prydz Bay: Site 739, 305a, 578b

Prydz Bay: Site 741, 601b

Prydz Bay: Site 743, 465a, 468a, 601b, 846b

kleinpelli, *Helolithus*, Kerguelen-Heard Plateau
S: Site 738, 493b

kolbei, *Coscinodiscus*, Kerguelen-Heard Plateau
N: Site 736, last occurrence, 134a

kolbei, *Thalassiosira*
Kerguelen Sediment Ridge: Site 745, last
occurrence, 574b

Kerguelen-Heard Plateau N: Site 736, last
occurrence, 828b

Kozloviella minor Zone, Kerguelen Plateau S:
Site 744, 487a

kuepperi, *Discoaster*, Kerguelen-Heard Plateau S:
Site 738, 492b

labiacrassata, *Globoturborotalita*
Kerguelen Plateau S, 440b

Kerguelen Plateau S: Site 744, 437b, 447b

lacunosa, *Pseudoemiliania*, Kerguelen Plateau S:
Site 744, 474b

laevigata, *Bolivinoides*, Kerguelen-Heard Plateau
S: Site 738, 858b

lamarckiana, *Gyroidina*, Kerguelen Plateau S:
Site 744, 629b

Lapideacassis sp., Kerguelen-Heard Plateau S:
Site 738, 510b

laticonus, *Diatrurus*, Kerguelen-Heard Plateau N:
Site 737, reworked, 181a

lauta, *Denticulopsis*, Kerguelen-Heard Plateau N:
Site 737, 180a, 598b

lentiginosa, *Thalassiosira*
Kerguelen Sediment Ridge: Site 745, 520a

Kerguelen-Heard Plateau N: Site 737, living
Pleistocene-Holocene, 194a

lepidula, *Stilosomella*, Kerguelen-Heard Plateau
S: Site 738, 627b

leptoporus, *Calcidiscus*
Kerguelen Plateau S: Site 744, 490b
first occurrence, 484a

Kerguelen-Heard Plateau N: Site 737, 177a

leroyi, *Globigerina*, Kerguelen-Heard Plateau S:
Site 738, 447b

linaperta, *Globigerina*, Kerguelen-Heard Plateau
N: Site 737, 178a

linaperta, *Subbotina*
Kerguelen Plateau S, 440b-441b

Kerguelen-Heard Plateau S: Site 738, 447b
last appearance datum, 438b

lingulatus, *Sceptroneis*, Prydz Bay: Site 739, 881b

Lithomitrella sp., Kerguelen-Heard Plateau N:
Site 737, reworked, 181a

lodoensis, *Discoaster*, Kerguelen-Heard Plateau
S: Site 738, 493b

longissima, *Thalassiothrix*, Prydz Bay: Site 742,
414a

longithorax, *Cyrtocapsella*, Kerguelen Plateau S:
Site 744; last common occurrence, 529b

Lychnocanoma sp., Kerguelen Plateau, 545b

PALEONTOLOGICAL INDEX

- maccollumii*, *Denticulopsis*, Kerguelen Plateau S: Site 744, first occurrence, 565b
macytrei, *Calcidiscus*, Kerguelen Plateau S: Site 744, 490b
magnicrassus, *Toweius*, Kerguelen-Heard Plateau S: Site 738, 493b
magnicrus, *Kamptnerius*, Kerguelen-Heard Plateau S: Site 738, first occurrence, 474b
mahooodii, *Thalassiosira*, n. sp.
 Kerguelen Plateau, 585b
 Kerguelen Sediment Ridge: Site 746, 593b, 594b, 595b, 596b
maleinterpretaria, *Nitzschia*
 Kerguelen Plateau S: Site 744, 487a
 first occurrence, 565b
Manumiella druggii Interval Zone, Kerguelen-Heard Plateau S: Site 738, 632b
Manumiella sp., Kerguelen-Heard Plateau S: Site 738, 631b-632b
Manumulla druggii Interval Zone, Kerguelen-Heard Plateau S: Site 738, 252a
margarita, *Alisocystis*, Kerguelen-Heard Plateau S: Site 738, 632b
mariae, *Eucyrtidium*, n. sp., Kerguelen Plateau, 536b, 546b
marina, *Nitzschia*, Kerguelen-Heard Plateau N: Site 737, 598b
martini, *Prinsius*
 Kerguelen-Heard Plateau S: Site 738, 502b, 505b, 511b
 first occurrence, 497b
 paleobiogeographic pattern, 502b
mashae, *Ceratocyrtis*, Kerguelen-Heard Plateau S: Site 738, first common occurrence, 529b
mateola, *Archaeoglobigerina*, Kerguelen-Heard Plateau S: Site 738, 461b, 463b
matthewsae, *Acarinina*, Kerguelen-Heard Plateau S: Site 738, 444b
mawsonii, *Phyllocladidites*, Prydz Bay: Site 742, 944b
mayaroensis, *Abathomphalus*
 Kerguelen-Heard Plateau S: Site 738, 250a, 455b, 463b, 858b
 first appearance datum, 759b
 first occurrence, 851b
mckannai, *Acarinina*
 Kerguelen Plateau S, 439b
 Kerguelen-Heard Plateau S: Site 738, 443b
medierraneus, *Leptocylindrus*
 Kerguelen-Heard Plateau S: Site 738, 260a
 Prydz Bay: Site 739, 315a
 Prydz Bay: Site 743, 468a
Melosira architecturalis Zone, Kerguelen Plateau S: Site 744, 488a
micra, *Pseudohastigerina*
 Kerguelen Plateau S: Site 744, 436b
 Kerguelen-Heard Plateau S: Site 738, 438b, 448b
Microachrydites-Hoegisporis Superzones, Prydz Bay: Site 741, 384a
microtrias, *Stellarina*
 Prydz Bay: Site 741, 390a
 Prydz Bay: Site 743, 468a
minor, *Cyathidites*, Prydz Bay: Site 742, 415a
mioceneica, *Nitzschia*
 Kerguelen Sediment Ridge: Site 746, 575b
 Kerguelen-Heard Plateau N: Site 737, 180a, 598b
mioceneica, *Thalassiosira*
 Kerguelen Plateau S: Site 744, 487a
 Kerguelen Sediment Ridge: Site 745, 515a
 Kerguelen Sediment Ridge: Site 746, 575b
 last occurrence, 542a
 Kerguelen-Heard Plateau N: Site 737, 597b
- last occurrence, 559b
mohleri, *Discoaster*, Kerguelen-Heard Plateau S: Site 738, 245a
 Mollusks, Prydz Bay: Site 739, strontium isotopes, 739b-742b
monikae, *Prunopyle*, nov. comb.
 Kerguelen Plateau, 539b
 Kerguelen-Heard Plateau S: Site 738, first common occurrence, 529b
monmouthisis, *Hedbergella*, Kerguelen-Heard Plateau S: Site 738, 464b
Monosulcites sp., Prydz Bay: Site 742, 415a
montiparum, *Eucyrtidium*
 Kerguelen Plateau S: Site 744, 546b
 Kerguelen-Heard Plateau S: Site 738, 546b
moriformis, *Sphenolithus*
 Kerguelen-Heard Plateau N: Site 737, 470b
 Kerguelen-Heard Plateau S: Site 738, 492b
Morozovella angulata Zone, Kerguelen-Heard Plateau S: Site 738, 250a
Morozovella inconstans Interval Subzone, Kerguelen-Heard Plateau S: Site 738, 432b, 457b
Morozovella uncinata Zone, Kerguelen-Heard Plateau S: Site 738, 250a
Morozovella uncinata-Igorina spiralis Partial Range Zone, Kerguelen-Heard Plateau S: Site 738, 457b
morsei, *Chiloguembelina*, Kerguelen-Heard Plateau S: Site 738, 448b
multipora, *Pontosphaera*, Kerguelen-Heard Plateau N: Site 737, 489b
multiradiatus, *Discoaster*, Kerguelen-Heard Plateau S: Site 738, 245a, 473b, 493b
multispinata, *Shackolina*, Kerguelen-Heard Plateau S: Site 738, recycled, 458b-459b
multispinatus, *Globigerinelloides*, Kerguelen-Heard Plateau S: Site 738, 858b
multispinus, *Globigerinelloides*
 Kerguelen-Heard Plateau S: Site 738, 463b
 recycled, 457b, 459b
munda, *Tenuitella*, Kerguelen-Heard Plateau S: Site 738, 449b
- nana*, *Paraturborotalia*, Kerguelen-Heard Plateau S: Site 738, 449b
 Nannofossils. See Calcareous nannofossils
Nannotetraena fulgens Zone, Kerguelen-Heard Plateau S: Site 738, 472b
nanum, *Gartnerago*, Kerguelen-Heard Plateau S: Site 738, last occurrence, 474b
nativa, *Thalassiosira*, Kerguelen-Heard Plateau N: Site 737, 597b
nautioides, *Lithelius*
 Kerguelen Sediment Ridge: Site 745, first common occurrence, 525b
 Kerguelen-Heard Plateau N: Site 736
 first common occurrence, 516b
 first occurrence, 135a
Naviculopsis biapiculata Zone, Kerguelen Plateau S: Site 744, 821b
neglectus, *Chaetoceros*
 Kerguelen-Heard Plateau N: Site 736, 600b
 resting spores, 609b
 Prydz Bay, 650b, 670b
 abundance, 653b, 654b
 distribution, 655b, 661b
 seasonal variation, 669b
 Prydz Bay: Site 740, 364a, 601b
 resting spores, 364a
 Prydz Bay: Site 741, resting spores, 388a, 390a
 Prydz Bay: Site 742, resting spores, 425a
Neoraistrickia sp., Prydz Bay: Site 742, 415a
nicobarica, *Crucidenticula*
- Kerguelen Plateau S: Site 744, first occurrence, 565b
 Kerguelen-Heard Plateau N: Site 737, 598b
 last occurrence, 559b
nitida, *Acarinina*, Kerguelen-Heard Plateau S: Site 738, 444b
Nitzschia, Prydz Bay, Austral spring blooms, 661b
Nitzschia angulata Zone
 Kerguelen Plateau S: Site 744, 487a
 Kerguelen Sediment Ridge: Site 745, 514a
 Kerguelen-Heard Plateau S: Site 738, 251a
Nitzschia barronii Partial Range Zone
 Kerguelen Plateau, 552b-553b
 Kerguelen Plateau S: Site 744, 565b
 Kerguelen Sediment Ridge: Site 745, 574b-575b
 Kerguelen-Heard Plateau N: Site 737, 558b
Nitzschia denticuloides Partial Range Zone
 Kerguelen Plateau, 556b
 Kerguelen Plateau S: Site 744, 565b
 Kerguelen-Heard Plateau N: Site 737, 556b
Nitzschia denticuloides Zone
 Kerguelen Plateau S: Site 744, 487a
 Kerguelen-Heard Plateau N: Site 737, 180a
Nitzschia grossepunctata Zone
 Kerguelen Plateau S: Site 744, 487a
 Kerguelen-Heard Plateau N: Site 737, 180a
Nitzschia interfrigidaria Zone
 Kerguelen Plateau S: Site 744, 487a
 Kerguelen Sediment Ridge: Site 745, 514a
 Kerguelen-Heard Plateau N: Site 737, 179a, 180a
Nitzschia interfrigidaria-Coscinodiscus insignis Partial Range Zone
 Kerguelen Plateau, 552b
 Kerguelen Plateau S: Site 744, 565b
 Kerguelen Sediment Ridge: Site 745, 574b
 Kerguelen-Heard Plateau N: Site 736, 552b
 Kerguelen-Heard Plateau N: Site 737, 558b
Nitzschia interfrigidaria-Coscinodiscus vulnificus Zone
 Kerguelen Plateau S: Site 744, 487a
 Kerguelen-Heard Plateau N: Site 736, 134a
Nitzschia interpretaria-Coscinodiscus rhombicus Zone, Kerguelen Plateau S: Site 744, 487a
Nitzschia miocenica zone, Kerguelen-Heard Plateau N: Site 737, 180a
Nitzschia porteri Zone, Kerguelen-Heard Plateau N: Site 737, 180a
Nitzschia praefrigidaria Zone, Kerguelen Sediment Ridge: Site 745, 514a
Nitzschia reinholdii Zone
 Kerguelen Plateau S: Site 744, 487a
 Kerguelen Sediment Ridge: Site 745, 514a
 Kerguelen Sediment Ridge: Site 746, 542a
 Kerguelen-Heard Plateau N: Site 737, 180a
 Kerguelen-Heard Plateau S: Site 738, 251a
Nitzschia spp., Prydz Bay: Site 739, ribbon-colony-forming, 315a, 365a
Nitzschia weaveri zone, Kerguelen-Heard Plateau N: Site 736, 134a
nitzschiooides, *Thalassionema*, Kerguelen-Heard Plateau N: Site 737, living Pleistocene-Holocene, 193a
nodulifer, *Azpeitia*, Kerguelen Sediment Ridge: Site 746, 542a
nodulosa, *Bigeneria*, Kerguelen Plateau S: Site 744, 627b
nova, *Clathrocyclas*, Kerguelen-Heard Plateau S: Site 738, last common occurrence, 527b
oamaruensis, *Chiasmolithus*
 Kerguelen Plateau S: Site 744, 484a, 492b
 Kerguelen-Heard Plateau N: Site 737, 178a
 first occurrence, 835b

- Kerguelen-Heard Plateau S: Site 738, first occurrence, 472b
 Prydz Bay: Site 739, 743b
oamaruensis, *Reticulofenestra*
 Kerguelen Plateau S: Site 744, 491b
 Kerguelen-Heard Plateau N: Site 737, 470b
 Kerguelen-Heard Plateau S: Site 738, 472b, 474b
oblonga, *Cassidulina*, Kerguelen-Heard Plateau N: Site 736, 628b
obtusa, *Pleurostomella*, Kerguelen Plateau S: Site 744, 627b
oceania, *Gephyrocapsa*, Kerguelen-Heard Plateau N: Site 736, 131a, 469b
ochotensis, *Actinocyclus*, Kerguelen-Heard Plateau N: Site 737, 596b
Odontella, Prydz Bay, resting spores, 679b
oestrupii, *Thalassiosira*, Kerguelen Sediment Ridge: Site 745, 575b
oligocenicus, *Asteromphalus*, Kerguelen Plateau S: Site 744, first occurrence, 573b
onusta, *Reticulofenestra*, Kerguelen-Heard Plateau S: Site 738, 490b
 Ophiroid, Kerguelen-Heard Plateau N: Site 736, 135a
Orthopithonella sp., Kerguelen-Heard Plateau S: Site 738, 863b
orthostylus, *Tribachiatius*, Kerguelen-Heard Plateau S: Site 738, 472b-473b, 493b
osculosus, *Spongodiscus*, Kerguelen Plateau S: Site 744, 529b
osloensis, *Pullenia*, Kerguelen Plateau S: Site 744, 629b
pachyderma, *Neogloboquadrina*
 Kerguelen-Heard Plateau N: Site 736 sinistrally coiled, 132a, 528b
 Kerguelen-Heard Plateau N: Site 737, 178a
 Kerguelen-Heard Plateau S: Site 738, 250a
 Prydz Bay: Site 741, 382a
 Prydz Bay: Site 743, 465a, 846b
pachyderma, *Siphocampe*
 Kerguelen Plateau, 545b
 Kerguelen Plateau S: Site 744, 529b
pacifica, *Globobulimina*, Kerguelen-Heard Plateau N: Site 736, 134a
Palaeohystrichophora infusoroides Zone, Kerguelen-Heard Plateau S: Site 738, 631b
 Palynology
 Antarctica, 873b-874b
 Kerguelen-Heard Plateau N: Site 736, biostratigraphy, 135a
 Kerguelen-Heard Plateau S: Site 738, 252a biostratigraphy, 631b-632b
 Prydz Bay, 941b-945b
 Prydz Bay: Site 739, 139b
 Prydz Bay: Site 740, 54b
 Prydz Bay: Site 741, 384a
 Prydz Bay: Site 742, 139b, 414a-415a reworked Cretaceous species, 415a
 Prydz Bay: Site 743, 139b
pannosus, *Phimopollenites*, Prydz Bay: Site 742, 944b
paradoxa, *Coptospora*, Prydz Bay: Site 741, 943b
parca, *Brownsonia*, Kerguelen-Heard Plateau S: Site 738, 474b
parvulus, *Asteromphalus*, Kerguelen Sediment Ridge: Site 745, 520a
patagonica, *Subbotina*, Kerguelen-Heard Plateau S: Site 738, 446b
pauperata, *Laticarinina*, Kerguelen Plateau S: Site 744, 628b
pelagicus, *Coccolithus*
- Kerguelen-Heard Plateau N: Site 736, 131a, 469b
 Kerguelen-Heard Plateau S: Site 738, 474b, 502b, 512b assemblage dominance fluctuation, 481b first occurrence, 497b paleobiogeographic pattern, 502b
pellitum, *Tectacodinium*, Kerguelen-Heard Plateau N: Site 736, 135a
pentacamerata, *Acarinina*
 Kerguelen-Heard Plateau S: Site 738, 444b first appearance datum, 250a
pentagonum, *Ceratium*, Kerguelen-Heard Plateau N: Site 737, living Pleistocene-Holocene, 194a
peragalli, *Rouxia*, Kerguelen-Heard Plateau N: Site 737, 596b
peregrina, *Stichocorys*
 Kerguelen Sediment Ridge: Site 746, 544a
 Kerguelen-Heard Plateau N: Site 737, 181a last common occurrence, 517b
perlóngus, *Rhabdosphaera*, Kerguelen-Heard Plateau S: Site 738, 492b
perplexa, *Reticulofenestra*
 Kerguelen Plateau, 483b
 Kerguelen-Heard Plateau N: Site 736, 131a
 Kerguelen-Heard Plateau N: Site 737, 177a, 470b
 Kerguelen-Heard Plateau S: Site 738, 474b, 488b
pertusus, *Toweius*
 Kerguelen-Heard Plateau S: Site 738, 505b, 511b, 512b first occurrence, 497b
petaloidea, *Globotruncanella*, Kerguelen-Heard Plateau S: Site 738, 452b, 463b, 858b
petrushevskaae, *Eurystomoskevos*, n. gen., n. sp., Kerguelen Plateau S: Site 744, 536b, 545b
Phaeocystis
 Prydz Bay, Austral spring blooms, 661b
 Prydz Bay: Site 740, 365a
 Prydz Bay: Site 742, 425a
physothorax, *Dictyoprora*, n. sp.
 Kerguelen Plateau, 535b
 Kerguelen Plateau S: Site 744, 545b first common occurrence, 529b
pierinae, *Axoprunum*, Kerguelen-Heard Plateau S: Site 738, last common occurrence, 527b
pierinae, *Axoprunum* group, Kerguelen Plateau, 537b
pitomorphus, *Phormostichoartus*
 Kerguelen Sediment Ridge: Site 745 last appearance datum, 519b last common occurrence, 516a
 Planktonic foraminifers
 Antarctica, oxygen isotopes, 78b
 Kerguelen Plateau, zonation, 814b, 816b
 Kerguelen Plateau S stable isotopes, 69b, 707b stable isotopes, Eocene/Oligocene transition, 710b-712b
 Kerguelen Plateau S: Site 744 analytical methods, 32a Antarctic surface water influence, 438b-439b biostratigraphic summary, 485a biostratigraphy, 435b, 485a-486a, 824b datum events, 31a, 436b Eocene-Holocene ranges, 429b Eocene/Oligocene boundary, 438b, 486a high-latitude cooling trend, 439b Oligocene hiatus, 486a Oligocene-Neogene, 437b Oligocene/Miocene boundary, 438b, 486a Paleocene/Eocene boundary, 438b radiolarian-nannofossil correlation, 530b
- stable isotopes, 698b-699b, 936b-938b strontium isotopes, 732b-736b zonation, 26a-28a
 Kerguelen Sediment Ridge: Site 745 analytical methods, 32a biostratigraphy, 514a datum events, 31a zonation, 26a-28a
 Kerguelen Sediment Ridge: Site 746 analytical methods, 32a datum events, 31a zonation, 26a-28a
 Kerguelen-Heard Plateau N: Site 736 analytical methods, 32a biostratigraphy, 133a datum events, 31a planktonic/benthic ratio, 131a upwelling and surface-water productivity, 156a zonation, 26a-28a
 Kerguelen-Heard Plateau N: Site 737 analytical methods, 32a biostratigraphic summary, 174a-176a biostratigraphy, 178a-179a datum events, 31a, 178a zonation, 26a-28a
 Kerguelen-Heard Plateau S: Site 738 analytical methods, 32a Antarctic surface water influence, 438b-439b Austral Realm correlation, 455b, 459b-460b biogeography, 455b biostratigraphic summary, 248a-249a biostratigraphy, 247a, 250a, 428b, 431b Cretaceous, 452b, 455b Cretaceous ranges, 453b Cretaceous/Tertiary boundary, 250a, 455b-460b, 857b-858b Danian ranges, 453b datum events, 31a, 455b Eocene/Oligocene boundary, 250a high-latitude cooling trend, 439b magnetostratigraphic correlation, 454b Oligocene/Miocene unconformity, 250a, 428b Paleocene-Eocene ranges, 429b Paleocene/Eocene boundary, 250a preservation, 452b recycled Cretaceous species, 457b-459b reworked Cretaceous, 428b stable isotopes, 700b-703b zonation, 26a-28a zonation, Cretaceous, 459b Zone P9, 250a Zones P3b-P6, 250a Zones P10-P12, 250a
 Prydz Bay analytical methods, 32a datum events, 31a oxygen isotopes, 120b zonation, 26a-28a, 814b, 816b
 Prydz Bay: Site 739, biostratigraphy, 304a
 Prydz Bay: Site 740, 368a biostratigraphy, 356a
 Prydz Bay: Site 741, biostratigraphy, 382a-383a
 Prydz Bay: Site 742, biostratigraphy, 413a
 Prydz Bay: Site 743, biostratigraphy, 465a
planoconicus, *Planorotalites*
 Kerguelen Plateau S, 440b
 Kerguelen-Heard Plateau S: Site 738, 448b
Planolites
 Kerguelen-Heard Plateau N: Site 737, 639b claystone burrows, 171a
 Kerguelen-Heard Plateau S: Site 738, 637b
Planorotalites australiformis Interval Zone, Kerguelen-Heard Plateau S: Site 738, 434b

PALEONTOLOGICAL INDEX

- Planorotalites imitalus* Interval Zone, Kerguelen-Heard Plateau S: Site 738, 432b-133b, 457b
- Planorotalites pseudomenardii* Zone, Kerguelen-Heard Plateau S: Site 738, 250a
- Planorotalites* sp., Kerguelen-Heard Plateau S: Site 738, 448b, 461b, 464b
- pliocenica*, *Anthocystidium*, Kerguelen-Heard Plateau N: Site 737, 181a
- pliocenica*, *Cycladophora*
Kerguelen Plateau S: Site 744, last common occurrence, 519b
- Kerguelen Sediment Ridge: Site 745
last appearance datum, 520b-521b
- Prunopyle tetrapila* co-occurrence, 521b, 524b
- Kerguelen-Heard Plateau N: Site 736, last common occurrence, 516b
- Podocyrtis chalara*/*Podocyrtis goetheana* Zones, Kerguelen-Heard Plateau S: Site 738, 252a
- polyacantha*, *Prunopyle*, Kerguelen-Heard Plateau S: Site 738, last common occurrence, 527b-528b
- polyacantha*, *Prunopyle*, nov. comb., Kerguelen Plateau, 539b
- polycamera*, *Eoglobigerina*, Kerguelen-Heard Plateau S: Site 738, 465b
- popofskii*, *Actinomma*, nov. comb., Kerguelen Plateau, 537b
- Porosira*, Prydz Bay, resting spores, 679b
- praeangulata*, *Acarinina*, Kerguelen-Heard Plateau S: Site 738, 443b
- praebifax*, *Discoaster*, Kerguelen-Heard Plateau S: Site 738, 245a, 472b
- praebulloides*, *Globigerina*, Kerguelen-Heard Plateau N: Site 737, 178a
- praeconvexa*, *Thalassiosira*, Kerguelen Sediment Ridge: Site 746, 575b
- praecurta*, *Nitzschia*, Kerguelen Sediment Ridge: Site 746, 598b
- praecylindrus*, *Nitzschia*, Kerguelen Sediment Ridge: Site 746, 596b
- praedimorpha*, *Denticulopsis*
Kerguelen Plateau S: Site 744, first occurrence, 565b
Kerguelen-Heard Plateau N: Site 737, 598b
last occurrence, 559b
- praeinterfrigidaria*, *Nitzschia*
Kerguelen-Heard Plateau N: Site 737, 598b
first occurrence, 558b
- Prydz Bay: Site 739, 305a, 578b
- praeoestrupii*, *Thalassiosira*
Kerguelen-Heard Plateau N: Site 737, 180a
first occurrence, 559b
last occurrence, 559b
- praepentacamerata*, *Acarinina*, Kerguelen-Heard Plateau S: Site 738, 444b
- praereticulata*, *Neoflabellina*, Kerguelen-Heard Plateau S: Site 738, 858b
- praescitula*, *Globorotalia*, Kerguelen-Heard Plateau N: Site 737, 178a
- Praetenuitella* sp., Kerguelen-Heard Plateau S: Site 738, 448b, 449b
- praetextum*, *Perichlamidium*, Kerguelen-Heard Plateau S: Site 738, first common occurrence, 529b
- praevema*, *Helotholus*
Kerguelen Plateau S: Site 744, 519b
Kerguelen Sediment Ridge: Site 746, 526b, 544a
Kerguelen-Heard Plateau N: Site 737, 181a
- primitiva*, *Acarinina*, Kerguelen-Heard Plateau S: Site 738, 444b
- primus*, *Cruciplacolithus*
Kerguelen-Heard Plateau S: Site 738, 473b
- first occurrence, 473b, 501b
- primus*, *Sphenolithus*, Kerguelen-Heard Plateau S: Site 738, 473b
- Prinsius martinii* Zone, Kerguelen-Heard Plateau S: Site 738, 501b
- prionotocodon*, *Lamprocyclas*, n. sp.
Kerguelen Plateau, 534b-535b
Kerguelen Plateau S: Site 744, 545b
- producta*, *Reticulofenestra*
Kerguelen Plateau, Neogene, 481b-482b
Kerguelen-Heard Plateau N: Site 736, 469b
Kerguelen-Heard Plateau N: Site 737, 177a, 488b
Prydz Bay: Site 739, 304a, 477b
- productus*, *Ellipsoplacolithus*, Kerguelen Plateau, 483b
- Prunopyle trypopyrena* lineage, Kerguelen Plateau, 530b-531b
- pseudobulloides*, *Subbotina*
Kerguelen-Heard Plateau S: Site 738, 464b, 858b
first appearance datum, 457b
first occurrence, 851b, 852b
- pseudocontinuosa*, *Globigerinatheka*, Kerguelen-Heard Plateau S: Site 738, 449b
- pseudocontinuosa*, *Paragloborotalia*, Kerguelen Plateau S, 440b
- pseudoeocaena*, *Subbotina*, Kerguelen-Heard Plateau S: Site 738, 447b
- Pseudohastigerina* sp.
Kerguelen Plateau S, 440b
Kerguelen-Heard Plateau S: Site 738, 448b
- pseudonana*, *Nitzschia*, Kerguelen Plateau S: Site 744, 494a
- pseudopunctata*, *Bolivina*, Kerguelen-Heard Plateau N: Site 736, 627b
- pseudotopilensis*, *Acarinina*, Kerguelen-Heard Plateau S: Site 738, 443b
- pseudoumbilica*, *Reticulofenestra*, Kerguelen-Heard Plateau N: Site 737, 177a
- pseudoveduezeliana*, *Subbotina*, Kerguelen Plateau S: Site 744, 447b
- pusilla*, *Igorina*, Kerguelen-Heard Plateau S: Site 738, 444b
- pusilla*, *Morozovella*, Kerguelen-Heard Plateau S: Site 738, first appearance datum, 250a
- pusillum*, *Ophthalmidium*, Kerguelen Plateau S: Site 744, 628b
- Pyxilla* group, Prydz Bay: Site 739, 744b
- Pyxilla* sp., Prydz Bay: Site 742, 414a, 881b
- quadrilobatus*, *Globigerinoides*, Kerguelen-Heard Plateau N: Site 736, 133a
- quadrilocula*, *Eoglobigerina*, Kerguelen-Heard Plateau S: Site 738, 465b
- quatra*, *Morozovella*, Kerguelen-Heard Plateau S: Site 738, 446b
- radicula*, *Stichocorys*, Kerguelen Plateau, 539b
- Radiolarians
Antarctica, zonation, 514b
Kerguelen Plateau
evolutionary lineages, 529b-531b
zonation, 814b, 816b, 817b
- Kerguelen Plateau S: Site 744
analytical methods, 32a-33a
biostratigraphic summary, 485a
biostratigraphy, 488a, 519b, 824b
datum events, 31a
nannofossil-foraminifer correlation, 530b
Neogene, 522b
Oligocene, 523b
Oligocene/Miocene boundary, 519b
paleomagnetic correlation, 531b
- zonation, 26a-28a, 29a-31a
- Zone NR3/NR4 interval, 519b
- Kerguelen Sediment Ridge: Site 745
analytical methods, 32a-33a
biostratigraphy, 515a-516a, 519b-521b, 524b
526b
- datum events, 31a, 525b
- Pliocene/Pleistocene boundary, 520b
- zonation, 26a-28a, 29a-31a
- Zone NR3/NR4 interval, 520b, 521b
- Kerguelen Sediment Ridge: Site 746
analytical methods, 32a-33a
biostratigraphy, 526b-527b, 543a-544a
datum events, 31a, 525b
- zonation, 26a-28a, 29a-31a
- Kerguelen-Heard Plateau N: Site 736
analytical methods, 32a-33a
biostratigraphy, 134a-135a, 514b-516b
datum events, 31a
- low diversity assemblages, 156a
- zonation, 26a-28a, 29a-31a
- Kerguelen-Heard Plateau N: Site 737
analytical methods, 32a-33a
biostratigraphic summary, 174a-176a
biostratigraphy, 180a-181a, 516b-518b
datum events, 31a
- Oligocene hiatus, 831b, 833b
- zonation, 26a-28a, 29a-31a
- Kerguelen-Heard Plateau S: Site 738
analytical methods, 32a-33a
biostratigraphic summary, 248a-249a
biostratigraphy, 252a, 518b
datum events, 31a
- datum events, Eocene, 527b
- datum events, Eocene/Oligocene boundary, 527b-528b
- datum events, Oligocene/Miocene boundary, 528b-529b
- Eocene-Oligocene, 520b-521b
- Eocene/Oligocene boundary, 518b
- Miocene hiatus, 252a, 518b
- Miocene-Quaternary, 518b
- nannofossil-foraminifer correlation, 528b
- NR3/NR4 Zone, 252a, 516a
- opal correlation, 190b
- Pliocene-Pleistocene hiatus, 244a, 252a, 518b
- winnowing effects, 252a
- zonation, 26a-28a, 29a-31a
- Zone NR3/NR4 interval, 518b
- Prydz Bay
analytical methods, 32a-33a
datum events, 30a
- zonation, 26a-28a, 29a-31a, 814b, 816b, 817b
- Prydz Bay: Site 739, biostratigraphy, 305a, 518b-519b
- Prydz Bay: Site 742, biostratigraphy, 414a
- radondoensis*, *Thecocrys*, Kerguelen-Heard Plateau N: Site 736, last occurrence, 135a
- recurvus*, *Isthmolithus*
Kerguelen Plateau S: Site 744, 484a, 491b
Kerguelen-Heard Plateau N: Site 737, 178a, 470b, 491b
first occurrence, 834b-835b
- Kerguelen-Heard Plateau S: Site 738
first occurrence, 472b
reworked, 474b
- Prydz Bay: Site 739, 477b, 743b, 837b, 880b
- redhosus*, *Botryostrobus*, n. sp.
Kerguelen Plateau, 535b
- Kerguelen Plateau S: Site 744, 545b
first common occurrence, 529b
- reinholdii*, *Synedra*, Kerguelen Plateau S: Site 744, 493a

reissi, *Planorotalites*, Kerguelen-Heard Plateau S: Site 738, 446b
reticulata, *Reticulofenestra* Kerguelen Plateau S: Site 744, 490b, 492b
 Kerguelen-Heard Plateau N: Site 737, 178a, 470b
 Kerguelen-Heard Plateau S: Site 738, 472b, 490b
Isthmolithus recurvus overlap, 474b
 last appearance datum, 245a
 Prydz Bay: Site 739, 477b
 Prydz Bay: Site 742, 477b
Reticulofenestra Kerguelen-Heard Plateau S: Site 738 assemblage dominance fluctuations, 474b, 481b
Reticulofenestra oamaruensis Zone, Kerguelen-Heard Plateau S: Site 738, 472b
Rhephax, Prydz Bay: Site 739, high species diversity, 304a
Rhizosolenia, Prydz Bay, Austral spring blooms, 661b
Rhizosolenia barboi Partial Range Zone Kerguelen Plateau, 551b
 Kerguelen Plateau S: Site 744, 565b
 Kerguelen Sediment Ridge: Site 745, 574b
 Kerguelen-Heard Plateau N: Site 736, 558b
Rhizosolenia barboi–Nitzschia kerguelensis Zone Kerguelen Sediment Ridge: Site 745, 514a
 Kerguelen-Heard Plateau N: Site 736, 134a
Rhizosolenia gravida Zone, Prydz Bay: Site 739, 579b, 743b
Rhizosolenia oligocenica Partial Range Zone Kerguelen Plateau, 557b
 Kerguelen Plateau S: Site 744, 573b
rhodospyroides, *Desmospyris*, Kerguelen Sediment Ridge: Site 745, 526b
ritscheri, *Nitzschia*, Prydz Bay: Site 741, 601b
ritscheri, *Thalassiosira* Prydz Bay: Site 740, silicified, 365a
 Prydz Bay: Site 741, 390a
 Prydz Bay: Site 743, 468a
robusta, *Antarctissa*, Kerguelen-Heard Plateau N: Site 737, first common occurrence, 518b
robusta, *Gublerina*, Kerguelen-Heard Plateau S: Site 738, 858b
robusta, *Lithomelissa*, Kerguelen-Heard Plateau S: Site 738, first common occurrence, 529b
robusta, *Theocyrtis*, Kerguelen Plateau, 546b
robusta, *Theocyrtis*, nov. comb., Kerguelen Plateau, 539b
Rocella gelida Partial Range Zone Kerguelen Plateau, 557b
 Kerguelen Plateau S: Site 744, 565b
Rocella gelida Zone, Kerguelen Plateau S: Site 744, 487a
Rocella vigilans Partial Range Zone Kerguelen Plateau, 557b
 Kerguelen Plateau S: Site 744, 565b, 573b
 Prydz Bay: Site 739, 579b
rugosa, *Rugoglobigerina*, Kerguelen-Heard Plateau S: Site 738, 461b, 463b

saipanensis, *Discoaster* Kerguelen Plateau S: Site 744, 484a, 492b
 Kerguelen-Heard Plateau N: Site 737 last appearance datum, 177a–178a
 last occurrence, 470b
 Kerguelen-Heard Plateau S: Site 738, 474b
samudrovia, *Reticulofenestra* Kerguelen-Heard Plateau S: Site 738, 474b
 Prydz Bay: Site 739, 743b
Saturnalis circularis Zone, Kerguelen Plateau S: Site 744, 488a

schraderi, *Rocella* Kerguelen Plateau S: Site 744 first occurrence, 565b
 last occurrence, 565b
scitula gigantea, *Globorotalia*, Kerguelen-Heard Plateau N: Site 737, 178a
scitula, *Globorotalia*, Kerguelen-Heard Plateau N: Site 737, 178a
seelandica, *Manumiella*, Kerguelen-Heard Plateau S: Site 738, 632b
semicosata, *Bulimina* Kerguelen Plateau S: Site 744, 822b
 Kerguelen-Heard Plateau S: Site 738, 627b
semipolita, *Calocyclus* Kerguelen Plateau, 546b
 Kerguelen-Heard Plateau S: Site 738 first common occurrence, 527b
 last appearance datum, 529b
senni, *Muricoglobigerina*, Kerguelen-Heard Plateau S: Site 738, 449b
setigera, *Solenocola* Kerguelen-Heard Plateau S: Site 738, 260a
 Prydz Bay: Site 739, 315a
sigmoidea, *Zygodiscus*, Kerguelen-Heard Plateau S: Site 738, 510b
Silicoflagellates Kerguelen Plateau, Oligocene–Quaternary, 933b–934b
 Prydz Bay: Site 739, 139b
simplicissima, *Eoglobigerina* Kerguelen-Heard Plateau S: Site 738, 464b first appearance datum, 457b
sinolampas, *Cymatodon*, n. gen., n. sp. Kerguelen Plateau, 536b–537b
 Kerguelen Plateau S: Site 744, 546b
 Kerguelen-Heard Plateau S: Site 738, first common occurrence, 529b
Siphocampe arachnea group, Kerguelen Plateau, 539b
Skolithos Kerguelen-Heard Plateau N: Site 737, 641b
 Kerguelen-Heard Plateau S: Site 738, 637b
sliteri, *Hedbergella*, Kerguelen-Heard Plateau S: Site 738, 461b, 463b
socialis, *Parvicorbicula*, Kerguelen-Heard Plateau S: Site 738, 260a
soldadoensis, *Acarinina* Kerguelen-Heard Plateau S: Site 738, 444b last appearance datum, 250a
solutus, *Chiasmolithus* Kerguelen-Heard Plateau N: Site 737, 470b last appearance datum, 178a
 Kerguelen-Heard Plateau S: Site 738, 492b last appearance datum, 245a
 last occurrence, 472b
sparsus, *Biantholithus* Kerguelen-Heard Plateau S: Site 738, 497b, 500b
 first occurrence, 473b
speculum, *Distephanus* Kerguelen-Heard Plateau S: Site 738, 260a
 Prydz Bay: Site 743, 468a
sphaerocephalus, *Lithomelissa*, Kerguelen-Heard Plateau S: Site 738, last common occurrence, 527b–528b
spinosa, *Thalassiosira*, Kerguelen Plateau S: Site 744, *Reticulofenestra bisecta* correlation, 821b
spinuloinflata, *Acarinina*, Kerguelen-Heard Plateau S: Site 738, 444b
spinulosa, *Morozovella*, Kerguelen-Heard Plateau S: Site 738, 446b
spinulosa, *Stylospira*, Kerguelen-Heard Plateau S: Site 738, last common occurrence, 527b

spiralis, *Igorina* Kerguelen Plateau S, 440b
 Kerguelen-Heard Plateau S: Site 738, 444b, 465b
 first occurrence, 457b
spongiosa, *Desmospyris* Kerguelen Sediment Ridge: Site 745, 516a last common occurrence, 525b
 Kerguelen-Heard Plateau N: Site 736 last common occurrence, 516b
 last occurrence, 135a
 Kerguelen-Heard Plateau N: Site 737, first common occurrence, 516b
Spongomelissa dilli Zone, Kerguelen Plateau S: Site 744, 488a
spongothorax, *Cycladophora* Kerguelen Plateau S: Site 744, first common occurrence, 519b
 Kerguelen-Heard Plateau N: Site 737, 518b
spumellaroides, *Thalassiosira*, Kerguelen Plateau S: Site 744, last occurrence, 565b
staurophora, *Micula*, Kerguelen-Heard Plateau S: Site 738, first occurrence, 474b
Stichocorys peregrina Zone Kerguelen Plateau S: Site 744, 488a, 519b
 Kerguelen Sediment Ridge: Site 746, 514a, 526b
 Kerguelen-Heard Plateau N: Site 737, 181a, 517b–518b
 Kerguelen-Heard Plateau S: Site 738, 252a, 518b
stigi, *Ceratocyrtis*, nov. comb., Kerguelen Plateau, 537b
Stilostomella spp., Kerguelen-Heard Plateau S: Site 738, 251a
stoveri, *Prediscosphaera*, Kerguelen-Heard Plateau S: Site 738, 862b
strabocella, *Acarinina*, Kerguelen-Heard Plateau S: Site 738, 444b
Stylatractus universis Zone, Kerguelen Sediment Ridge: Site 745, 516a, 519b
Subbotina angiporoidea Interval Zone Kerguelen Plateau S: Site 744, 436b
 Kerguelen-Heard Plateau S: Site 738, 435b
Subbotina inaequispira Partial Range Subzone, Kerguelen-Heard Plateau S: Site 738, 434b
Subbotina pseudobulloides Interval Range Zone, Kerguelen-Heard Plateau S: Site 738, 459b, 460b
Subbotina pseudobulloides Partial Range Zone, Kerguelen-Heard Plateau S: Site 738, 431b–432b
Subbotina spp., Kerguelen Plateau S, stable isotopes, 69b, 707b
Subbotina trinidadensis Zone, Kerguelen-Heard Plateau S: Site 738, 250a
subbotinae, *Morozovella* Kerguelen-Heard Plateau S: Site 738, 446b high-latitude incursion, 439b
subcarinata, *Pullenia*, Kerguelen Plateau S: Site 744, 629b
subcarinatus, *Globigerinelloides*, Kerguelen-Heard Plateau S: Site 738, 463b, 858b
subcurvata, *Nitzschia* Prydz Bay: Site 739, 314a, 315a
 Prydz Bay: Site 743, 467a
subhumerus, *Cycladophora*, Kerguelen-Heard Plateau S: Site 738, first common occurrence, 529b
sublodoenensis, *Discoaster*, Kerguelen-Heard Plateau S: Site 738, 472b
subpertusus, *Coccilithus*, Kerguelen-Heard Plateau S: Site 738, paleobiogeographic pattern, 502b

PALEONTOLOGICAL INDEX

subpertusus, *Coccolithus*, n. comb., Kerguelen-Heard Plateau S: Site 738, 504b
subspinosa, *Stilostomella*, Kerguelen-Heard Plateau S: Site 738, 627b
superba, *Euphausia*, Prydz Bay: Site 740, 364a
superba, *Stephanopyxis*, Prydz Bay: Site 739, 744b
suteri, *Globorotaloides*
 Kerguelen Plateau S, stable isotopes, 708b
 Kerguelen Plateau S: Site 744, strontium isotopes, 732b, 738b
 Kerguelen-Heard Plateau S: Site 738, 449b

tabularis, *Azpeitia*, Kerguelen-Heard Plateau N: Site 737, living Pleistocene–Holocene, 193a
tainemplektia, *Stylocictya*, n. sp.
 Kerguelen Plateau 533b
 Kerguelen-Heard Plateau S: Site 738, 543b
tanyacantha, *Actinomma*, Kerguelen-Heard Plateau N: Site 737, reworked, 181a
tenuijunctus, *Dactyliosolen*, Kerguelen Sediment Ridge: Site 745, 520a
tenuis, *Cruciplacolithus*
 Kerguelen-Heard Plateau S: Site 738, 246a, 473b, 497b
 first occurrence, 501b, 851b–852b
 in green clay layer, 857b
testarugosa, *Globorotaloides*
 Kerguelen Plateau S, 439b
 Kerguelen-Heard Plateau S: Site 738, 449b
tetrapila, *Prunopyle*, Kerguelen Sediment Ridge: Site 745, *Cycladophora pliocenica* co-occurrence, 521b, 524b
teuria, *Zeauvigerina*, Kerguelen-Heard Plateau S: Site 738, 452b, 461b, 464b
teuriensis, *Hornbrookina*
 Kerguelen-Heard Plateau S: Site 738, 477b, 481b, 504b–505b, 510b
 last occurrence, 497b, 500b
teuscheri, *Eucyrtidium*, Kerguelen-Heard Plateau N: Site 737, 181a
Thalassinoides
 Kerguelen-Heard Plateau N, depositional environment, 639b
 Kerguelen-Heard Plateau N: Site 737, 639b, 641b
 claystone burrows, 171a, 173a
 geometry, 636b
 Kerguelen-Heard Plateau S: Site 738, 637b
 geometry, 636b
Thalassiosira
 Prydz Bay
 Austral spring blooms, 661b
 resting spores, 679b
Thalassiosira convexa Zone, Kerguelen-Heard Plateau N: Site 737, 180a
Thalassiosira fraga Partial Range Zone
 Kerguelen Plateau, 557b
 Kerguelen Plateau S: Site 744, 565b
Thalassiosira inura Partial Range Zone
 Kerguelen Plateau, 553b
 Kerguelen Sediment Ridge: Site 745, 575b
 Kerguelen-Heard Plateau N: Site 737, 558b
Thalassiosira kolbei Partial Range Zone
 Kerguelen Plateau, 551b–552b
 Kerguelen-Heard Plateau N: Site 736, 558b–559b
Thalassiosira lentiginosa Partial Range Zone
 Kerguelen Plateau, 551b
 Kerguelen Plateau S: Site 744, 559b
 Kerguelen Sediment Ridge: Site 745, 574b
 Kerguelen-Heard Plateau N: Site 736, 558b
 Kerguelen-Heard Plateau N: Site 737, 558b
Thalassiosira lentiginosa Zone

Kerguelen Plateau S: Site 744, 487a
 Kerguelen Sediment Ridge: Site 745, 514a
 Kerguelen-Heard Plateau N: Site 736, 134a
 Prydz Bay: Site 739, 305a
 Prydz Bay: Site 740, 358a
 Prydz Bay: Site 741, 384a
 Prydz Bay: Site 742, 413a, 414a
Thalassiosira oestrupii Partial Range Zone
 Kerguelen Plateau, 553b
 Kerguelen Sediment Ridge: Site 745, 575b
 Kerguelen-Heard Plateau N: Site 737, 558b
Thalassiosira sp.
 Kerguelen Plateau S: Site 744, 592b
 Prydz Bay: Site 741, resting spores, 390a
Thalassiosira spinosa Partial Range Zone
 Kerguelen Plateau, 557b
 Kerguelen Plateau S: Site 744, 565b
Thalassiosira spumellaroides Partial Range Zone
 Kerguelen Plateau, 557b
 Kerguelen Plateau S: Site 744, 565b
Thalassiosira torokina Partial Range Zone
 Kerguelen Plateau, 553b
 Kerguelen Sediment Ridge: Site 745, 575b
 Kerguelen Sediment Ridge: Site 746, 575b
 Kerguelen-Heard Plateau N: Site 737, 559b
 Prydz Bay: Site 739, 579b
thalmanni, *Bolivina*, Kerguelen Plateau S: Site 744, 627b
thaumasia, *Lophophphaena*, n. sp.
 Kerguelen Plateau, 534b
 Kerguelen Plateau S: Site 744, 544b
 last common occurrence, 529b
Theocalyptra spongoothorax Zone
 Kerguelen Plateau S: Site 744, 488a
 Kerguelen Sediment Ridge: Site 746, 514a, 527b
 Kerguelen-Heard Plateau N: Site 737, 181a
 Kerguelen-Heard Plateau S: Site 738, 252a
Theocotyle cryptocephala Zone, Kerguelen-Heard Plateau S: Site 738, 252a
Thoracosphaera, Kerguelen-Heard Plateau S: Site 738, paleobiogeographic pattern, 502b
Thoracosphaera sp.
 Kerguelen-Heard Plateau S: Site 738, 510b
 Cretaceous/Tertiary boundary, 856b–857b
Thoracosphaera spp., Kerguelen-Heard Plateau S: Site 738, 473b
titan, *Prunopyle*
 Kerguelen Plateau S: Site 744, 519b
 Kerguelen Sediment Ridge: Site 745, last occurrence, 524b–525b
 Kerguelen-Heard Plateau N: Site 736, last common occurrence, 516b
 Kerguelen-Heard Plateau N: Site 737, first common occurrence, 517b
torokina, *Thalassiosira*
 Kerguelen Sediment Ridge: Site 746, first occurrence, 575b
 Kerguelen-Heard Plateau N: Site 737, 597b
Toweius sp.
 Kerguelen-Heard Plateau S: Site 738, 473b
 in green clay layer, 857b
 Trace fossils
 Kerguelen-Heard Plateau N, to interpret water depth, 639b
 Kerguelen-Heard Plateau N: Site 737, 636b–637b, 638b, 639b
 Kerguelen-Heard Plateau S: Site 738, 636b–637b, 638b–639b
 Cretaceous/Tertiary boundary, 637b, 639b
Transversopontia sp., Kerguelen-Heard Plateau N: Site 737, 491b
triangularis, *Subbotina*, Kerguelen-Heard Plateau S: Site 738, 447b, 465b
Triceraspyris coronata Zone

Kerguelen Sediment Ridge: Site 745, 526b
 Kerguelen-Heard Plateau N: Site 737, 181a, 517b
 Kerguelen-Heard Plateau S: Site 738, 252a, 518b
trifulta, *Thalassiosira*, Kerguelen Plateau S: Site 744, 494a
trilobum, *Pterocanium*, Kerguelen-Heard Plateau N: Site 736, last appearance datum, 516b
triloculinoides, *Subbotina*
 Kerguelen-Heard Plateau S: Site 738, 464b
 first appearance datum, 457b
triplex, *Acarinina*, Kerguelen-Heard Plateau S: Site 738, 444b
trivalvis, *Subbotina*, Kerguelen-Heard Plateau S: Site 738, 465b
Trochammina, Prydz Bay: Site 740, 357a
Tropidoneis, Prydz Bay, Austral spring blooms, 661b
truempyi, *Nuttallides*
 Kerguelen Plateau S, stable isotopes, 69b, 704b, 706b, 707b, 708b
 Kerguelen Plateau S: Site 744, 615b, 616b
 last appearance datum, 614b
 Kerguelen-Heard Plateau S: Site 738, 455b, 613b, 615b, 616b, 630b, 858b
 last appearance datum, 251a
tryopyrena, *Prunopyle*, n. sp.
 Kerguelen Plateau, 533b
 Kerguelen Plateau S: Site 744, 543b
 Kerguelen-Heard Plateau S: Site 738, first common occurrence, 529b
tsugaruensis, *Actinocyclus*, Kerguelen Sediment Ridge: Site 746, 592b
tumida, *Thalassiosira*
 Kerguelen Plateau S: Site 744, 493a
 cold-water form, 494a
 Kerguelen-Heard Plateau N: Site 737, living Pleistocene–Holocene, 194a
 Kerguelen-Heard Plateau S: Site 738, 260a
 Prydz Bay: Site 739, silicified valves, 315a
 Prydz Bay: Site 740, 364a
 silicified, 365a
 Prydz Bay: Site 743, 468a
typaniformis, *Fasciculithus*, Kerguelen-Heard Plateau S: Site 738, first occurrence, 497b, 852b
umbilica, *Reticulofenestra*
 Kerguelen-Heard Plateau N: Site 737, 470b
 last appearance datum, 177a
 last occurrence, 470b
 Kerguelen-Heard Plateau S: Site 738, 474b, 492b
 first appearance datum, 245a
 first occurrence, 472b
 Prydz Bay: Site 739, 477b, 743b
umbilicatum, *Astrononion*, Kerguelen Plateau S: Site 744, 630b
umbonatus, *Oridorsalis*, Kerguelen Plateau S: Site 744, 630b
umbonifera, *Nuttallides*
 Kerguelen Plateau S: Site 744, 486a, 620b, 622b, 630b, 822b
 Eocene/Oligocene boundary, 616b
 Kerguelen-Heard Plateau S: Site 738
 Antarctic Bottom Water initiation effect, 251a
 Eocene/Oligocene boundary, 616b
ungerianus, *Cibicidoides*, Kerguelen Plateau S: Site 744, 629b
unicavus, *Catapsydrax*, Kerguelen-Heard Plateau S: Site 738, 447b
universus, *Stylatractus*
 Kerguelen Sediment Ridge: Site 745, 516a

last appearance datum, 519b
utilisindex, *Subbotina*
 Kerguelen Plateau S, 440b-441b
 Kerguelen-Heard Plateau S: Site 738, 447b

vanheurckii, *Tropidoneis*
 Prydz Bay
 inner vs. outer bay, 662b
 seasonal variation, 669b

varianta, *Subbotina*, Kerguelen-Heard Plateau S:
 Site 738, 446b

velascoensis, *Subbotina*, Kerguelen-Heard Plateau S: Site 738, 446b

vema, *Pseudocubus*
 Kerguelen Sediment Ridge: Site 745, last common occurrence, 525b
 Kerguelen-Heard Plateau N: Site 736
 last common occurrence, 516b
 last occurrence, 135a

veniamini, *Bogorovia*
 Kerguelen-Heard Plateau N: Site 737, 180a
 last occurrence, 833b-834b

venusta, *Quinqueloculina*, Kerguelen Plateau S: Site 744, 629b

vigilans, *Rocella*, Kerguelen Plateau S: Site 744,
 last occurrence, 565b
vitrea, *Epistominella*, Kerguelen Plateau S: Site 744, 630b
vulnificus, *Coscinodiscus*
 Kerguelen Sediment Ridge: Site 745, last occurrence, 574b
 Kerguelen-Heard Plateau N: Site 736, last occurrence, 134a, 558b

waiparensis, *Chiloguembelina*, Kerguelen-Heard Plateau S: Site 738, 464b
weaveri, *Nitzschia*
 Kerguelen-Heard Plateau N: Site 736
 first occurrence, 134a
 last common occurrence, 134a
 Kerguelen-Heard Plateau N: Site 737, 598b
 first occurrence, 180a, 558b
weissflogii, *Odontella*
 Kerguelen Plateau S: Site 744, 494a
 Prydz Bay: Site 739, 315a
wighamii, *Chaetoceros*
 Prydz Bay, inner vs. outer bay, 662b
 Prydz Bay: Site 740, 364a
 Prydz Bay: Site 742, resting spores, 425a

wilcoxensis, *Acarinina*
 Kerguelen Plateau S, 439b
 Kerguelen-Heard Plateau S: Site 738, 444b
 first appearance datum, 437b
wilcoxensis, *Chiloguembelina*, Kerguelen-Heard Plateau S: Site 738, 448b
wilcoxensis, *Pseudohastigerina*
 Kerguelen Plateau S, 440b
 Kerguelen-Heard Plateau S: Site 738, 448b
 first appearance datum, 437b
woodi connecta, *Globigerina*, Kerguelen-Heard Plateau N: Site 737, 178a
wuellerstorfi, *Cibicidoides*, Kerguelen Plateau S: Site 744, 629b

zealandica, *Globorotalia*, Kerguelen Plateau S: Site 744, 437b
Zoophycos
 Kerguelen-Heard Plateau N, distribution, 639b
 Kerguelen-Heard Plateau N: Site 737, 636b, 639b
 claystone burrows, 171a
 Kerguelen-Heard Plateau S: Site 738, 636b, 637b, 641b