

INDEX TO VOLUME 121

This index provides coverage for both the *Initial Reports* and *Scientific Results* portions of Volume 121 of the *Proceedings of the Ocean Drilling Program*. Index entries to page numbers in the *Initial Reports* are preceded by A with a colon (A:), and those to the *Scientific Results* (this book), by B with a colon (B:).

Previous indexes to ODP Vols. 101–119 and 124 were presented mostly in three parts: (1) a subject index, (2) a site index, and (3) a paleontological index. Beginning with this volume, only two parts are presented. Entries for the site index have been incorporated with the Subject Index, and the paleontological index is now called the Taxonomic Index. For information on electronic formats available, please contact the Data Librarian, Ocean Drilling Program, 1000 Discovery Drive, College Station, Texas 77845-9547, U.S.A.

The index was prepared by Wm. J. Richardson Associates, Inc., under subcontract to the Ocean Drilling Program. The index is based on a hierarchy of entries: (1) a main entry, defined as a key word or concept followed by a reference to the page on which that word or concept appears; (2) a subentry, defined as a further elaboration on the main entry followed by a page reference; and (3) a sub-subentry, defined as an even further elaboration on the main entry or subentry followed by a page reference.

The Subject Index follows a standard format. Geographic and individual names are referenced in the index only if they are subjects of discussion. As mentioned above, the Subject Index now includes entries for the sites discussed in the volume. This index also includes broad fossil groups, such as foraminifers and radiolarians.

The Taxonomic Index is an index relating to significant findings and/or substantive discussions, not of species names *per se*. This index covers three varieties of information: (1) individual genera and species that have been erected or emended formally, (2) biostratigraphic zones, and (3) fossils depicted in illustrations. Taxonomic entries consisting of both genera and species are listed alphabetically *by species*; biostratigraphic zones, however, are listed alphabetically *by genus*.

The indexes cover text figures and tables but not core description forms (“barrel sheets”) or core photographs. Also excluded are bibliographic references, names of individuals, and routine front and back matter.

For further information, contact the Chief Production Editor, Ocean Drilling Program, at the address given in the second paragraph above.

SUBJECT INDEX

- Acoustic impedance
 Broken Ridge, A: 151, 163, 185, 254
 ash layers, A: 152
 correlation, angular unconformity, A:255
 Eocene angular unconformity, A:255
 lithologic correlation, A:79, B:744
 physical properties correlation, A:218–219
 seismic correlation, A: 186, B:669, 679
 units 1–3, B:744
 Ninetyeast Ridge, A:292-294, 340, 404, 416, 431
 seismic correlation, A:347–348, 357, 448
 Unit A vs. Unit B, A:289
 Site 752, A: 163
 seismic correlation, B:679
 Site 757, A:347-348
 seismic correlation, A:357
 Site 758, A:416
 seismic correlation, A:448
- Acoustic stratigraphy, Broken Ridge, units A–C, A:75
- Adzhidere, virtual geomagnetic pole, B:385, 387
- Africa E, rifting, active, A:8
- Africa S, Oligocene aridity, and dust flux reduction, Broken Ridge, B:220
- African Plate, absolute motion, Fleitout's model, B:772
- Alkalinity
 Broken Ridge, A:136
 calcite-gypsum supersaturation and, B:448
 Ninetyeast Ridge, A:398
 calcite-gypsum supersaturation and, B:448
 calcium concentration and, A:286, 399
 gradient, effect of volcanic alteration on, A:334
- Alteration
 Broken Ridge, pre- vs. post-rift, B:522
 Ninetyeast Ridge
 ash-to-clay, A:374
 basalts, A:321–322, 324, 391, 393
 basement and sediment cover effects, A:532–533
 degree of, intersite variation, A:333
 eruption level and state of, B:543
 ignition losses from, B:563
 nonoxidative, Ninetyeast Ridge, A:329
 open-system, Ninetyeast Ridge, B:563
 oxidative, Ninetyeast Ridge, B:563
 oxidative/nonoxidative zones, Ninetyeast Ridge, A:277–279, 324, B:625
 oxides, Ninetyeast Ridge, A:333
 oxygen isotope changes with, Site 752, B:452–453
 post-magmatic, Ninetyeast Ridge, A:329, B:563–565, 614–615, 625
 pre- vs. post-rift, Site 752, B:522
 secondary mineral associations, Ninetyeast Ridge, A:330
 temperature of, carbonate veins, Ninetyeast Ridge, B:453
- Alteration, deutenc, Ninetyeast Ridge, basalts, B:534–535
- Alteration, hydrothermal
 Ninetyeast Ridge, oxidative vs. non-oxidative temperatures, B:565
 post-magmatic, Ninetyeast Ridge, B:629
 temperature of, Ninetyeast Ridge, B:535
- Alteration, low-temperature
 basalt, Ninetyeast Ridge, B:509
 lavas, Ninetyeast Ridge, B:722
- magmatic processes, Ninetyeast Ridge, B :625
 magnetic processes, Ninetyeast Ridge, B:539, 541
 Ninetyeast Ridge, B:534
 seawater, Ninetyeast Ridge, B:564-565
- Aluminum oxide
 basalts, Ninetyeast Ridge, A:331, B:567, 570
 incompatible element comparison, Ninetyeast Ridge, B:635, 637
 low-pressure melt segregation, Ninetyeast Ridge, B:641
 parental magma, Ninetyeast Ridge, B:629
 tephra layers, Ninetyeast Ridge, B:277
 vs. calcium oxide, Ninetyeast Ridge, B:277, 285
 vs. iron oxide, Ninetyeast Ridge, B:277, 285, 574
 vs. magnesium oxide, Ninetyeast Ridge, A:329
 vs. modal plagioclase, basalts, Ninetyeast Ridge, B:580
 vs. zirconium, basalts, Ninetyeast Ridge, B:569, 577
- Aluminum oxide/calcium oxide ratio
 basalt, Ninetyeast Ridge, A:331
 low-pressure melt segregation, Ninetyeast Ridge, B:641
 vs. magnesium oxide, Ninetyeast Ridge, A:329
- Amphibole
 fractionation, Site 758, B:287
 tephra layers, Site 758, B:284
- Amsterdam Island
 lava, isotopic ratios, A:526
 paleolatitude, B:802
 tholeiitic basalt, A:526
- Amsterdam/St. Paul islands
 basalts, correlation, Ninetyeast Ridge; B:562, 579
 ridge-building plumes, B:559
- Amsterdm–St. Paul hot spot, paleolatitude, B:801
- Andesite, oceanic, Ninetyeast Ridge, B:572, 577
 incompatible-element-enriched, B :592
- Anhyseretic remanent magnetization (ARM)
 with magnetite dominant, Site 758, B:379
 vs. susceptibility, Site 758, B:379, 382, 384
- Anisotropy
 of anhysteretic remanence, Site 752, B:374
 chalks and limestones, Maestrichtian, Broken Ridge, A:498
 Site 752, B:376
- Anisotropy of ARM, Broken Ridge, B:361, 372–374, 376
- Ankerite, Site 758, A:374
- Antarctic Bottom Water, flow through Vema Gap, A:470
- Antarctic Circumpolar Current
 Broken Ridge, erosion and influence of, A:84, 90
 Ninetyeast Ridge, Oligocene, A:18
- Antarctic Circumpolar Ocean, silica productivity, Kerguelen Plateau record, B:938
- Antarctic Ice Sheet, oxygen isotope enrichment and, Ninetyeast Ridge, B:247
- Antarctica, ice volume growth, global paleoclimatic changes. B:943
- Apatite
 Site 755, A:238
 tephra layers, Site 758, B:284
- Apparent polar wander path (APWP), Ninetyeast Ridge, B:782
- Argon dating, Indian Ocean, basalt, B:509-516, 799
- Arid conditions, India, wind intensity and, A:20
- Ascension Island, tholeiitic basalt, A:526
- Ash
 Broken Ridge, A:112, 120, 191–192, 194-195, 199, 237–238, 460, 463-464, 468, 471, 499-500, 525, B:939
 accumulation rate, A:471, B:940
 acoustic impedance, A:153
 calcareous chalk contact, A:156
 Campanian/Maestrichtian boundary, A:491
 cementation, A:498
 consolidation rate, B:260
 Cretaceous/Tertiary boundary, A:149, 512, B:398-399, 428, 941
 deposition environment, B:939
 flux rates, A:465, 468, B:428, 939-940
 geochemistry, A:135–136, 472-477
 graded bedding characteristics, B:937
 magnetic properties, A:206, 247, 475, 482, 488, 491
 physical property effects, A:498, B:268
 seismic reflection profiling, B:744
 source, A:465, B:477, 939
 temporal variation, A:475
 thermal conductivity and, A:147
 transport, redistribution, and deposition, A:465
 Turonian-Eocene flux rate, A:90
 types of, A:471
 vs. calcium carbonate, B:521
 vs. Kerguelen Plateau, A:463
- Kerguelen/Ninetyeast Ridge, sequence of hot-spot-produced, A:262
- Nirtetyeast Ridge, A:531, B:229
 alteration, basalt contacts, A:523
 correlation, Toba ash, A:520
 density, A:288
 intercalated, A:523
 magnetic properties, B:779
 paleolatitude, B:801
 physical properties, B:257
 scoured basal contacts, A:521–522
 source, B :477, 633
 vs. Hawaiian Islands, B:633
- oxygen isotopes with alteration of, Indian Ocean, B:452-453
- Site 752, B:398–399
 Cretaceous/tertiary boundary, A:507, 513, B:424
 geochemistry, A:472-475, 480-482
 iridium/cesium ratio, B:417
 magnetic susceptibility, A:482-483
 origin, B:731
 planktonic productivity and, B:428-429
 temporal variations, A:474-475
 thickness of, A:507
- Site 754, A:199
- Site 757
 comparison, Broken Ridge, A:306
 deposition environment, A:311, 350
 geochemistry, A:327
 lapilli, A:311
 magnetic properties, B:784
- Site 758, A:368, 372, 374, 415, B:286-287
 carbonate content, A:401
 depositional environment, vs. lavas, A:360
 host sediment contacts, B:276

- incompatible element ratios, A:525
 layers, B:285–286
 magnetic properties, A:423, B:276, 307–309, 796, 798, 801
 magnetic susceptibility, B:307–309
 massive layers vs. concentrated beds, A:415
 maximum age range, B:275
 microscopic description, B:277
 mineralogy, B:277
 paleolatitude, B:802
 Sunda arc origin, B:822
 thickness of, B:276
 Toba ash content, B:489
 trace elements, A:393
 X-ray fluorescence, A:403
- Ash, glauconite-bearing, Site 757, magnetic susceptibility, A:331
 Ash-chert-chalk sequence, Site 752, Cretaceous/Tertiary boundary, B:424
 Asian accretionary wedge, emplacement onto Greater India, A:16
 Asifabad, Intertrappean Beds, paleoryctic mammals, B:818
 Atlantic Ocean, carbonate veins, isotopic analysis, B:447
 Atlantic Ocean, equatorial, geomagnetic polarity transition, Jaramillo Subchronozone, B:377–388
 Atlantic Ocean N, deep-water temperature, vs. equatorial Indian Ocean surface temperature, B:304
 Atlantic Ocean SE, carbonate compensation depth, B:212
 Aulis Volcanics/Taltung Formation, paleolatitude, B:802
 Australasian microtektites
 Cretaceous/Paleogene boundary
 flow structures and bubble cavities, B:490
 North American goyazite/smectite spherules and, B:490–491, 493
 planktonic foraminiferal influence, B:489–492
 geochemistry, B:494
 Site 758, B:503
 Australia, paleomagnetic pole positions, B:879–880
- Barium
 Site 752, B:439
 Cretaceous/Tertiary boundary, B:424
 vs. zirconium, basalts, Ninetyeast Ridge, B:565
 Barium/lanthanum ratio, Ninetyeast Ridge, B:640
 Barium/niobium ratio, Ninetyeast Ridge, B:568
 Dupal signature basalts, B:638–639
 vs. lanthanum content, B:640
- Basalt
 Broken Ridge
 age determination, B:510–512, 514, 516
 iridium/cesium ratio, B:417
 petrography, B:507–509
 flows vs. sills, B:547, 556
 Ninetyeast Ridge, B:612–614
 age determination, B:510, 513–516
 alteration, B:534–535, 563–565
 chemical type II, B:576–577
 comparison, Kerguelen Archipelago and Plateau, B:592
 depositional environment, B:509
 eruptive environment, A:13
 geochemistry, B:561–562
 groundmass, B:658
 intersite differences, A:330
 magnetic properties, B:526–537, 542–543, 561, 779
 microprobe analysis, B:535–540
 multiple mantle sources, A:525
 optical examination, B:534–535
 petrography, A:330–331, B:507–509
 sampling techniques, B:779–780
 sheet flows, B:526
 subaerial eruption, B:229
 tholeiitic vs. alkaline, B:622
 trapped melt, B:658
 Site 756
 alteration, A:274, 277–278
 color and alteration mineral association, A:274
 derivation, A:277
 geochemistry, A:275, 277, 279–281, B:567–569
 iron-enriched, B:567, 571
 logging data, A:274
 magnetic properties, A:281–282, 287–288
 major oxides, B:567
 mineralogy, A:275, 282–283
 modality, A:276–277
 petrography, A:275, B:621–624
 physical properties, A:297
 seismic reflection profiling, A:295–296
 Site 757 compared, A:335
 subaerial eruption, A:260, 265, 267, 274–275, 524
 velocity, A:289, 293
 Site 757
 alteration, A:321–322, 324
 chemical types I and II, B:569–571
 depositional environment, A:350
 geochemistry, A:306, B:569–571
 incompatible elements, A:332
 intersite differences, A:324, 326
 magnetic mineralogy, A:333
 magnetic properties, B:792
 major oxide content, A:331
 modality, A:323–325
 petrography, A:322–326, B:624–629
 post-magmatic alteration, B:625
 X-ray fluorescence, A:326–331
 Site 758, A:360, 374
 Cretaceous/Tertiary boundary, A:360
 deep water eruption, A:375
 geochemistry, A:400, 402–403, B:571–572
 horizontal and vertical stress, B:704–706
 magnetic properties, A:394–396, 410
 major oxides, A:401
 mineralogy, B:567
 modality, A:394–395
 petrography, A:392–393, B:629–633
 physical properties, A:404
 quench structures, A:391
 recovery, A:365–366
 seismic reflection profiling, A:421–422
 sheet flows, A:391–393
 X-ray diffraction, A:396
 Basalt, massive
 Ninetyeast Ridge, petrology, B:526
 Site 758
 chill zones, A:389–390
 mineralogy, A:392–393
 petrography, A:387–389
 thickness of zones, A:390
 vesicle zones, A:389
 Basalt, olivine-rich, Ninetyeast Ridge, alteration, B:564
 Basalt, pillow, Ninetyeast Ridge, A:391–392, B:526
 Basalt, plagioclase-phyric
 Site 757, A:311, 321, B:624–625
 Site 758, B:567
 Basalt, tholeiitic, A:526
 Ninetyeast Ridge, A:277
 Dupal isotopic signature, B:640
 formation, A:526
 incompatible-element-enriched, B:592
 incompatible elements, A:525–526, 534
 spreading-ridge axis sites and, A:279
 Basalt sheet flows, mineralogy, Site 758, A:392–393
 Basaltic flow units
 Ninetyeast Ridge, A:531
 thickness, B:632
 ophiolite studies, B:556
 Site 756, orientation, A:275
 Site 757, A:321–322
 alteration, A:321
 paleolatitudes, B:801
 units F1–F19, A:321–322
 units F8 and F13–F14, B:628
 units F18–F19, B:627–629
 Site 758, A:387–391, B:629
 altered sediments surrounding, B:549
 contacts, A:389–390, B:548–549
 grain size variation, B:549
 paleolatitudes, B:802
 quench zones, B:549, 630
 seismic reflection profiling, A:421–422
 spatial distribution, B:547–549
 thickness of, A:390
 units F1–F4, B:629–631
 vertical thickness, B:548
- Basement, basaltic
 Broken Ridge, A:5
 Kerguelen-Heard Plateau, A:5
 Ninetyeast Ridge, A:5
 acid-leaching procedures, B:592–593, 596
 age, B:526, 561, 811
 geochemistry, A:393, 404, 526, 528, B:594–595
 incompatible element ratios, A:527
 intersite variations, A:526, 528, B:562
 magnetic properties, B:792, 796, 798, 801
 major oxide content, A:327, 329, 335
 neodymium isotopes, B:596
 paleolatitudes, B:811
 physical properties, A:337, 339, 344
 radiogenic isotopes, A:13–14
 relation to Amsterdam–St. Paul hot spot, B:811
 source, B:811
 strontium isotopes, B:593, 596
 trace elements, A:13, 330, 334
 Site 756
 flow units, A:274
 macroscopic description, A:273–275
 Basement, magnetic, Broken Ridge
 depth estimates, Werner deconvolution, B:683–584, 687–689, 692
 sediment thickness, B:683–684
- Bengal Fan
 formation, A:17, B:310
 erogenic uplift, sediment deposition and, B:822
 petrology, B:822
 terrestrial deposition, Ninetyeast Ridge, A:401, 424
- Benthic foraminifers. *See also under particular sites and by taxa in Taxonomic Index*
 Broken Ridge
 Antarctic ice sheet expansion and abundance of, A:485
 bathyal-to-abyssal Eocene faunas, A:486

- Cretaceous/Tertiary boundary, B:5–13, 16, 19, 731
depth migration, B:19, 21
diversity, paleoenvironmental differences, B:9
faunal analysis, B:9–10, 12–13, 16
Maestrichtian–Eocene, B:5, 7, 10–11
Midway-type fauna, B:21
Oligocene–Pleistocene, B:32–43, 50–52
oxygen isotopes, B:4, 19
Paleocene/Eocene boundary, A:487, B:5
Paleocene extinction event, B:3, 19, 21
Paleogene, A:486–487
planktonic/benthic ratios, A:477–478
stratigraphic range, B:6, 8–9, 17
varimax assemblages, stratigraphic distribution, B:17
- Indian Ocean
carbon isotope correlation, B:4
water paleodepth estimates, A:43, 45
relation to known water masses, B:43, 50
Site 709, isotope stratigraphy, B:242–244
Site 752, A:43–45, 129, B:722
Antarctic-source water influence, B:19, 21
ash layer, B:13
biogenic productivity and increase in, B:16
bottom-water temperature and extinction of, B:19
carbon isotopes, B:16, 19, 21, 888
carbon vs. oxygen isotope records, B:20
Eocene/Paleocene boundary, A:129
faunal turnover, Zone CP8, B:6
oxygen isotopes, B:214, 886–888
planktonic/benthic ratio, A:129, B:16, 81
Velasco-type fauna, B:9, 19, 21
warm, saline bottom water model, B:21
water paleodepth estimates, A:121, 123
- Site 753, A:178–179
Indian Deep Water association, A:178
planktonic/benthic ratio, B:81
- Site 754, A:204–205
agglutinated forms, A:204
bathyal-to-abyssal deposition, A:205
bottom water cooling and Neogene changes in, B:51–52
carbonate content and productivity of, B:51
comparison, Site 756, B:38
diversity, paleoenvironmental differences, B:9
Eocene–Pleistocene, B:32
faunal analysis, B:9–10, 12–13, 16, 38–39, 59–60, 69–70
faunal associations, A:204
Indian Deep Water association, A:204
Maestrichtian–Eocene, B:6–7
Oligocene–Pleistocene, B:34–39
oxygen isotopes, B:214, 218
Paleocene/Eocene boundary, B:5
planktonic/benthic ratio, B:81
stratigraphic range, B:39
three-stage species diversity, B:32
- Sites 754/756 compared
comparison, Site 754, B:38
corrosive water effects, B:51
deep water associations, B:43, 50
Neogene changes, paleoceanographic association, B:50
Neogene stratigraphic distribution, B:50
North Atlantic Deep Water association, B:52
oxygen isotopes, B:50–52
saline water event, B:52
- Site 755, A:246
planktonic/benthic ratio, B:81–82
- Site 756, A:272–273
bottom water cooling and Neogene changes in, B:51–52
carbonate content and productivity of, B:51
faunal analysis, B:39, 43, 61, 69
Oligocene–Pleistocene, B:32–33, 38, 44–52
planktonic/benthic ratio, B:38
species diversity, B:58
stratigraphic ranges, B:54
- Site 757, A:320
Indian Deep Water association, A:320
neritic environment, A:314–315
planktonic/benthic ratio, B:83
temporal variation, A:315
water paleodepth estimates, A:319–320
- Site 758, A:385–386
Indian Bottom Water influence, A:385
isotope stratigraphy, B:245
mid-latitude to tropical-subtropical transition, A:376
planktonic/benthic ratio, B:84–85
preservation, A:375
- Bermuda Seamount
suboptical exsolution, B:535
transition and upper dike complex, magnetic mineralogy, B:539
- Biostratigraphy
Site 752, A:121, 123–130, B:364, 722
Site 753, A:173–179, B:368, 731–732
Site 754, A:195–205, B:369
Site 755, A:242–246, B:370
Site 756, A:267–273
Site 757, A:311–320
Site 758, A:375–387
paleoenvironmental implications, A:375–376
- Biotite, Site 758
fractionation, B:287
tephra layers, B:284
- Bioturbation, Broken Ridge, Turonian tuffs, A:464
- Bituminite, Site 755, A:496
- Bottaccione Gorge, clay, Cretaceous/Tertiary boundary, B:415
- Bottom-water circulation, Indian Ocean, cold water influx and, A:18
- Bouguer anomaly, Broken Ridge, B:684
- Bouvet hot spot, predicted path, B:516
- Bouvet Island, tholeiitic basalt, A:526
- Bowlingite, Ninetyeast Ridge, pseudomorphs in basalt, B:563
- Brahmaputra River, sediment output, B:298
- Breccia, volcanic clay, Site 758, A:373–374
- Broken Ridge
alteration, pre- vs. post-rift, B:522
ash layers, trace elements, A:474, 476–479
asymmetry, A:151
basalt, B:507–509
age determination, B:510–516
trace elements, B:594–595
bathymetry, B:664
biostratigraphy, A:43–45
benthic foraminifers, A:485–487
calcareous nannofossils, A:478–483
diatoms, A:487–488
planktonic foraminifers, A:483–485, B:921
carbonate accumulation rate, B:470, 473, 475, 477
carbonate veins, alteration temperatures, B:453–455
chalk-limestone transition, physical properties, A:497–498
coarse fraction, B:937
compaction history, B:253–260
- Cretaceous–Paleocene reconstruction, A:525
Cretaceous/Tertiary boundary, A:460, 463
depositional environment, A:467–469, B:941
crust/mantle boundary, uplift and tilt, A:9
currents, Neogene, A:469–470
deformation, pre- vs. post-rift, B:522
depositional environment, A:10
depositional model, B:747, 753, 760
diagenesis, Kerguelen Plateau compared, A:463–464
- dipping and truncated sequence
correlation across ridge, A:72
deposition, B:744
magnetic properties, A:488–491
magnetostratigraphy, A:488–489, B:359, 363
paleolatitudes during formation, B:371
physical properties, A:497–498
seismic reflection profiling, B:743–744, 756–757
seismic stratigraphy, A:72–75, 153, 156, 226, 255
- downlapping sequence, B:744
- Eocene angular unconformity, A:8, 71, 86, 112, 153, 171, 191, 457, 471, B:447
correlation, circum-Antarctic seaway opening, A:500
limestone pebble layer, A:469
magnetic susceptibility, A:179
physical properties, A:495
seismic reflection profiling, A:5, 186, 254, 255, B:668–669
seismic stratigraphy, A:10
shallow-water deposits, A:460
uplift and formation of, B:438, 943
variation in depth to, A:84
- eolian dust, B:219–221
- fining-upward sequence, A:116, 194
interface, post-rift sequence, A:10
- flexural strength, pre-rift, B:691
- formation, A:86, 457, 465, 470, B:516, 681, 747, 753, 760
flexural rebound model, A:86, 90
rifting/drifting process, B:359
tectonic and stratigraphic model, A:89–90
- geochemistry, inorganic, A:57, 491–492
geochemistry, organic, A:56–57, 492–495
geological setting, A:71–72, B:467, 721
geophysical logging, A:59–60
geothermal gradient, B:522
- gravity anomalies
magnetic response, B:682
topographic effects, B:684–692
topographic relationship, A:86–87
- hot-spot origin, B:507, 514
paleolatitudes evidence, B:367, 371
- lipid geochemistry, B:473
- lithologic column, B:935
- lithologic summary, B:254–255, 729, 732, 734, 736
- lithospheric extension models, A:5–12
- lithostratigraphy, B:359, 398, 437–438, 447
Kerguelen Plateau compared, B:938
lithologic Unit I, A:488
lithologic Unit II, A:488–489, 491
magnetostratigraphic correlation, A:488–491
- Oligocene disconformity, A:457
seismic correlation, B:668–669, 745
summary, A:461
- low-molecular weight hydrocarbons, B:457–465
- magnetic basement depth of, B:683–684

- magnetic properties, A:50-56, 63
 demagnetization, B:361-363
 inclination, B:367, 371-374
- Oligocene disconformity, A:469, 471, B:211, 229
 seismic reflection profiling, B:668-669
- onlapping sequence
 depositional environment, B:747
 seismic reflection profiling, B:746-747, 758-759
- organic carbon, B:457-465
 accumulation rate, B:470, 477, 481
- organic matter, type of, B:472-473
 origin, flexural vs. thermal, B:690
- paleoceanography, A:465-471, B:936, 939-943
 Maestrichtian-Eocene, B:34
- paleodepth, A:81, B:724, 746
 paleodepths, across Eocene unconformity, B:747, 753
- paleoenvironment, A:469, B:936
 paleogeographic reconstruction, A:8
 paleolatitudes, B:367-368, 469, 724, 941
 pre-rift sequence, B:359
- pebble layer, A:460
- pelagic cap, A:185-186, 457-458, 460, B:936-937
 depositional environment, B:753
 magnetic properties, A:488
 physical properties, A:495-497, B:254, 264
 seismic reflection profiling, A:84-86, 153, 185-186, 225-226, 233, 254-255, B:747
 subsidence and accumulation of, A:90
 thickness of, B:936
- physical properties, A:57-59, 495-496
 consolidation effects, B:253-260
 silica mineralogy and, B:521-522
- physiography, B:468
- porosity, magnetic properties and, A:484, B:373
- productivity, B:420-421, 472
 following Cretaceous/Tertiary boundary event, B:427
 nutrient source, B:940
 Santonian-Maestrichtian, A:500
 sea level and, B:753
 subsidence and increase in, A:90
- prograding downlapping wedge, A:75, 79, 81-82
- ridge evolution models, A:89-90
- rift basin
 depth of, vs. new ocean/old ocean boundary, B:691
 formation, B:692
- rifting
 Kerguelen Plateau separation, A:5, B:437-438
 sediment composition and mechanisms for, B:444
- Santonian-Eocene hiatus, A:71
- sediment thickness, eastern vs. western ridge, A:75, 77
- sedimentary sequence, A:457-463
 Kerguelen Plateau compared, A:463-454
- sedimentation rate, B:735-740
- seismic stratigraphy, B:398, 663-670
- silica mineralogy, B:521-522
 lithification effects, B:261-269
- sonobuoy seismic recording, A:80, B:682
- stable isotopes
 mass balance calculations, B:452-453
 pore waters, B:448-452
- subsidence history, A:87-89
- tectonic history, A:5-12, 457, B:933-934
- tephra
 geochemistry, A:472-475
 lithology, A:471-472
 magnetic susceptibility, A:475
- topography
 lithospheric strength in determining, A:9, 11, 86-87
 modeled vs. observed, B:691-692
 uplift, passive mechanism, B:437
 velocity, B:682-683
 water depth, pre-rift, A:10
- Broken Ridge/Kerguelen-Heard Plateau formation, A:71, B:747
 rifting, B:747
- Bromine, Site 752, Cretaceous/Tertiary boundary, B:424
- Brunhes/Matuyama boundary, Site 758, oxygen isotope stratigraphy, B:306
- Brunhes/Matuyama chronozone boundary, microtektites, B:489-490
- Bunbury continental flood basalts
 hot spot origin, A:71
 paleolatitudes, B:371, 802
- Bytownite crystals, Site 758, source, B:287
- Calcareous nannofossils. *See also under particular sites and by taxa in Taxonomic Index*
- Broken Ridge
 Antarctic Circumpolar Current influence, A:479
 Cretaceous, A:482-483
 dating of uplift and subaerial erosion from, A:476-477
 dipping units, A:480-481
 low Miocene diversity, A:479
 Maestrichtian/Danian boundary, A:481
 Neogene assemblage, A:478
 preservation, A:476
 shallow-water origin, A:480
 Southern Hemisphere high-latitude affinity, A:482
- Cretaceous/Tertiary boundary, Site 752, B:396-398, 401-407
- Site 752, A:123-127, B:722
 ash layer, B:396, 398-399
 austral affinity, B:144
 carbonate and productivity of, B:398-399
 Cretaceous, A:511-512, B:143-144, 400-401, 406
 Cretaceous/Tertiary boundary, A:127, 481-482, 507, 509, 511-512, B:396-398, 401-408, 730-731
 Danian-Eocene unconformity, A:126
 datum levels, B:725-726
 foraminifers age correlation, A:121
 globotruncanid extinction, B:418-419
 high-latitude affinity, A:512
 high-latitude assemblage, A:127
 Maestrichtian, B:146-147
 magnetostratigraphic correlation, B:725-731
 Maud Rise compared, A:512
 Miocene/Pliocene boundary, A:123
 Oligocene/Miocene boundary, A:123, 479
 Paleocene/Eocene boundary, A:127, B:727
 Pliocene/Pleistocene boundary, A:123
 preservation, B:143
 survivor forms, A:512, B:400, 406-408
 Tertiary species, A:512, B:400, 407
 zonation, A:123-125, B:143-144
- Site 753, A:175-176
 datum levels, A:177, B:727
- Eocene-Miocene unconformity, A:176
 magnetostratigraphic correlation, B:732
 Miocene/Pliocene boundary, A:175
 reworked Eocene-Oligocene species, A:175-176
- Site 754, A:198-200
 biostratigraphic summary, B:162
 Campanian, A:482
 Cretaceous, B:144-145
 datum levels, B:727
 dipping units, A:480
 Eocene unconformity, A:195
 Maestrichtian, B:148, 150-153
 magnetostratigraphic correlation, B:733, 735
 Miocene-Pliocene diversity increase, A:198
 Oligocene disconformity, A:200
 Oligocene/Miocene boundary, A:479, B:733
 open-ocean pelagic deposition, A:205
 Pliocene/Pleistocene boundary, A:199
 zonation, B:144-145
- Site 755, A:242, 244-246
 ash layers, B:145
 Coniacian/Santonian boundary, B:145
 Cretaceous, B:145
 datum levels, B:727
 magnetostratigraphic correlation, B:735
 Santonian/Coniacian boundary, B:735
 Turonian-Campanian, B:154-156
 Turonian/Coniacian boundary, A:244
 zonation, B:143, 145
- Site 756, A:268, 270
 Eocene/Oligocene boundary, A:270
 Oligocene/Miocene boundary, A:270
 Pliocene/Pleistocene boundary, A:268
 reworked species, A:270
 strontium isotopes, B:927, 929
- Site 757, A:315-318
 datum levels, A:316
 dissolution, A:315
 Eocene/Oligocene boundary, A:316
 Oligocene hiatus, A:315
 Oligocene/Miocene boundary, A:316
 strontium isotopes, B:927, 930
 zonation, A:318
- Site 758, A:376-382
 Campanian-Maestrichtian, B:159-161
 Campanian/Maestrichtian boundary, A:376
 Cretaceous, A:380-382, B:145, 149
 Cretaceous/Tertiary disconformity, A:380
 datum levels, A:382
 Eocene-Oligocene disconformity, A:376
 lithostratigraphic correlation, B:149
 Miocene/Pliocene boundary, A:376
 Oligocene/Miocene boundary, A:376
 Paleocene-Oligocene hiatus, A:376
 paleoenvironmental implications, A:375-376
 preservation, A:375
 zonation, A:376, B:145, 149
- Southern Hemisphere, zonation, Campanian-Maestrichtian, B:142
- strontium isotopes
 Site 744, B:922
 Site 748, B:927
 zonation, A:43, 46-47, B:143
- Calcareous sediments
 Broken Ridge
 compression index, B:258
 physical properties, B:254
- Ninetyeast Ridge
 compaction rate, B:258, 260

- compression index, B:258
- Calcite
 accumulation rate, B:470, 475
 Broken Ridge, A:138, 463, 494, B:272
 physical properties, A:498, B:268
 recrystallization, A:237, B:268
 Ninetyeast Ridge, A:324
 in basalts, A:391
 formation, B:563
 secondary, B:625
 supersaturation, A:335, B:448
 Calcite, sparry, Broken Ridge, A:243
 Calcite/dolomite ratio, Broken Ridge, A:494
 Calcium
 Broken Ridge
 ash content and gradient changes, A:136, 491-492
 intersite correlation, A:180
 pore water concentration, A:213, B:448, 450
 with slow water convection, B:452
 vertical gradient, A:494
 volcanic alteration effects, B:448
 Ninetyeast Ridge, A:260, 290
 ash abundance and, A:283-284
 effect of alteration on, A:334, 399, 531-532, B:448
 faulting and, A:286
 lithologic and topographic correlations, A:533
 pore water concentration, B:448, 450-451
 vertical gradient, A:334, 398, 534
 Site 756, A:284, 290
 Calcium carbonate
 Broken Ridge, A:141-142, 183, 215-216, 251, 497-498, B:79, 81-82, 426-428
 accumulation rate, A:465, B:940
 ash layers, A:473
 Cretaceous/Tertiary boundary, A:471, B:398-399, 941, 943
 flux rate, A:90, 468-469, B:939, 941, 943
 magnetic susceptibility and, A:483
 mass accumulation rate, A:507
 vs. ash content, B:521
 Ninetyeast Ridge, A:290-291, 343-344, 372, 420-424, B:28, 82-85, 299, 316-328, 490-491, 495
 across Eocene unconformity, A:368
 Brunhes dissolution cycle, B:312
 coarse fraction and dissolution of, B:307
 comparison, Brown University and ODP statistics, B:315
 composite depth record, B:306-308, 310-313, 340-355
 dilution, Pleistocene record, B:311
 glacial/interglacial cycles, B:311
 mass accumulation rate, B:299, 303, 307, 309
 mid-Pleistocene shift, B:313
 Miocene, B:246
 Northern Hemispheric glaciation effects, B:313
 sedimentation rate correlation, B:227
 stratigraphy, B:307
 Site 758, A:422-423, B:307
 standard sediment sample, B:892
 Calcium gradient, ash, Site 757, A:336
 Calcium/magnesium ratio
 Broken Ridge, with volcanic alteration, B:449
 Ninetyeast Ridge, B:452
 oxygen isotope values and, B:448
 Calcium oxide
 Broken Ridge, ash layers, A:473
 Ninetyeast Ridge
 basement basalts, A:335
 vs. iron oxide, B:285
 vs. iron oxide, tephra layers, B:277
 Calcium oxide/aluminum oxide ratio, Ninetyeast Ridge, B:285
 tephra layers, B:277
 vs. magnesium oxide and strontium content, B:639
 Calcium/potassium release, Ninetyeast Ridge, basalt, B:515
 Cambay graben, paleoposition, B:818
 Cape Verde Rise
 basalt
 intrusive vs. extrusive, B:550
 petrography, B:549-550
 carbonaceous shale, contact metamorphic effects, B:550
 methane/ethane ratio, with distance from basaltic sill, B:550
 Carbon, inorganic
 Broken Ridge, A:183, 215-216, 251
 Ninetyeast Ridge, A:290, 343
 Carbon, organic
 accumulation rate, B:470, 475
 Broken Ridge, A:137-138, 141-142, 212, 215-216, 250-251, 493-494, B:457, 460-461, 474
 Cretaceous/Tertiary boundary, A:138
 quality of, A:212-215
 changes with paleolatitude and paleotectonic movement, B:467
 Ninetyeast Ridge, A:287, 290-291, 337, 343-344, 400-401, 420-422, 424, B:457, 461, 474
 productivity rate, B:472
 open ocean deposition, B:477, 481
 sedimentation rate and, B:472, 481
 Site 752, A:137-138
 Site 754, A:218-219
 Site 758, A:422, 424
 Miocene-Pliocene calcium carbonate flux rate, B:309
 sediment accumulation rate and, B:822
 source of, B:481
 terrigenous component, B:472
 Carbon, total
 Broken Ridge, A:141-142, 215-216, 218, 251
 Ninetyeast Ridge, A:290, 343, 420-421
 Carbon dioxide, Ninetyeast Ridge
 lavas, B:614
 outgassing, B:455
 Carbon isotopes
 Broken Ridge, B:231-233
 Maestrichtian-Eocene, B:885-886, 888
 pelagic oozes, B:237-238
 inter-reservoir transfer and enrichment of, B:234
 Ninetyeast Ridge, B:233-234, 236
 carbonate veins, B:453-455
 composite depth record, B:306-308, 310-311, 340-355
 depletion, Messinian desiccation event, B:250-251
 glacial/interglacial cycles, B:304
 pelagic oozes, B:239-240
 planktonic foraminifers, B:250, 329-339
 pelagic oozes
 Broken Ridge, B:232-234
 Ninetyeast Ridge, B:232-234
 Site 752, Maestrichtian-Eocene, B:885-889
 Site 758, stratigraphy, B:245, 248-251, 304-307
 standard sediment sample, B:892
 Carbon preference index, B:483
 Carbonate
 Broken Ridge, A:42, 138, 141-142, 181, 215, 237, 250, 494-495, B:433
 accumulation rate, B:472-473, 747
 Cretaceous-Paleogene, B:937
 Cretaceous/Tertiary boundary, A:460, 470-471, 495, B:475
 density and, B:269
 negative organic carbon correlation, A:212
 oceanic circulation and flux rate changes, A:466, B:941
 open-ocean sedimentation, B:475
 Santonian deposition, B:747
 sea level and productivity of, B:753
 vs. volcanogenic sediment content, B:433
 Cretaceous-to-Quaternary increase, B:470
 Ninetyeast Ridge, A:42, 337, B:471, 493-494
 accumulation rate, B:472-473
 across Toba ash layer, B:489
 depositional environment, B:475
 magnetic properties and, A:360
 Miocene cooling trend, B:475, 477
 vs. Broken Ridge sediments, A:401-402
 porosity and, B:256
 Site 752, A:138
 Site 753, A:181
 Site 754, A:215-216
 Site 756, A:287
 Site 758, A:372, 401-402, 424
 volcanic influence, B:475, 477
 Carbonate carbon, changes with paleolatitudes and paleotectonic movement, B:467
 Carbonate compensation depth
 Atlantic Ocean SE, B:212
 Indian Ocean, B:309
 Carbonate-free residue, Broken Ridge
 biogenic compounds, B:271
 elemental composition, B:262
 Carbonate platform
 Broken Ridge
 Maestrichtian, A:465-467, B:940
 response to sea-level change, B:743, 753, 761
 Vail model, B:753, 761
 Carnegie Ridge, carbonate pelagic sediments, winnowing, B:212
 Central Indian Basin
 magnetostratigraphy, B:765
 tectonic chart, B:765-766
 Central Water Zone, Indian Ocean, productivity, A:18
 Cerium, Site 752, Cretaceous/Tertiary boundary, B:426
 Cesium, Site 752
 as clay source indicator, B:417-418
 in Cretaceous/Tertiary boundary clays, B:421
 Cesium/lead ratio, Ninetyeast Ridge
 intersite differences, B:637
 vs. lanthanum content, B:640
 Cesium/yttrium ratio, Broken Ridge, ash layers, A:474
 Chagos Bank
 basement, age, B:770
 hot spot origin, relative motion models, B:770
 Chagos-Laccadive Ridge, formation, Réunion hot spot role, B:770
 Chagos-Maldives-Laccadive ridge system, Réunion hot spot, predicted path, B:514
 Chalcedony, Site 754, A:195
 Chalcopyrite, Site 758, B:629
 Chalk, Broken Ridge
 accumulation rate, A:464, B:938-939
 Cretaceous/Tertiary boundary, B:731
 high density/low porosity spikes, A:150

- laminae, B:937
- Chalk, calcareous
- Broken Ridge, A:194
 - carbonate content, B:475
 - compression index, B:260
 - dissolution, B:753
 - magnetic susceptibility, A:206
 - physical properties, A:497
 - porosity, B:258
 - volcanic ash contact, A:156
 - Site 758, A:360, 368
 - magnetic susceptibility, A:394
- Chalk, foraminiferal, Site 758, A:368
- Chalk, nannofossil
- Site 752, A:460
 - Maestrichtian–Eocene, A:75
 - Site 758, A:360, 368, 373
 - logging data, A:410
 - magnetic susceptibility, A:394, 424
- Chalk-chert-ash layer, Broken Ridge, Cretaceous/Tertiary boundary, A:471
- Chalk-limestone-chert sequence, Broken Ridge, accumulation rate, A:464-465, B:939
- Chalk-limestone transition, Site 754
- logging data, A:221
 - seismic reflection profiling, A:226
- Chemical sediments, classification, A:42
- Chert
- Broken Ridge, A:194-195, 462-463
 - Cretaceous–Paleogene, A:460, B:937
 - Cretaceous/Tertiary boundary, A:507
 - logging data, A:149, 221
 - magnetic susceptibility, A:206
 - physical property effects, B:268
 - seismic reflection profiling, A:226, B:683
 - Ninetyeast Ridge
 - carbonate content, B:471
 - chemical gradient effect, A:399, 410
 - pebbles, A:368
 - remanence effects of, A:395
- China Sea S, rifting, A:17
- Chloride
- Broken Ridge
 - pore water concentration, A:213
 - with volcanic alteration, B:449
 - Ninetyeast Ridge, B:452
 - pore water concentration, B:448
- Chlorinity
- pore waters, B:448, 450-451
 - Site 757, A:335
 - Site 758, A:398
 - calcium concentration and, A:399
- Chlorite, Broken Ridge, B:521
- Chromium
- Broken Ridge, ash layers, A:474
 - Ninetyeast Ridge, A:280
 - basalts, A:333, 399
 - vs. iron oxide/magnesium oxide ratio, B:567-568, 571
 - vs. zirconium, B:574
- Circulation, atmospheric
- Pacific Ocean S, sea-surface productivity and, A:20
 - polar-to-equator temperature gradient, 211a
- Circulation, oceanic
- Broken Ridge
 - biogenic productivity and, A:466-467
 - grain size with changes in, A:470, B:943
 - ice volume increase and, A:500
 - Indian Ocean, driving mechanisms, A:18
 - Ninetyeast Ridge, sedimentation rate effects, A:517
 - Clay
 - Site 752, Cretaceous/Tertiary boundary, origin, B:419-420
 - Site 758, geochemical logging effect, A:410
 - Clay, calcareous, Site 758, A:360
 - magnetic susceptibility, A:394, 424
 - Clay, terrigenous, Ninetyeast Ridge, A:534
 - Clay, volcanic, Site 758, drilling biscuits, A:373-374
 - Clay mineralogy, Broken Ridge, B:521-522
 - mixed-layer smectite-chlorites, B:521
 - Claystone, kaolinitic, Site 758, iridium content, B:490-491
 - Clinofolds, prograding, Broken Ridge, B:747, 753
 - Clinoptilolite, calcium-rich, Broken Ridge, B:521
 - Clinopyroxene
 - Ninetyeast Ridge, B:570-571, 623, 656
 - phenocryst phases, A:275, 323
 - segregation, B:624
 - Site 757, phenocryst phases, B:625
 - Coarse fraction, Site 758
 - composite depth section, B:306-310, 316-328, 340-355
 - glacial/interglacial cycles, B:311
 - winnowing effect, B:309
 - Cobalt, Site 758, in sulfides, B:629
 - Cobalt/iron ratio, Site 752, Cretaceous/Tertiary boundary, B:420-421
 - Cocos Ridge, carbonate sediment winnowing, B:212
 - Coiba Ridge, carbonate sediment winnowing, B:212
 - Compaction
 - Broken Ridge
 - intersite comparison, B:258
 - physical properties and, B:253-254, 256-258
 - rate of, B:254, 256
 - shallow inclinations from, B:372-374
 - lithologic correlation, B:253
 - Ninetyeast Ridge, physical properties and, A:343, B:253-254, 257-258
 - Compression index
 - Broken Ridge, B:260
 - calcareous sediments vs. pelagic ooze, B:258
 - Consolidation
 - Broken Ridge
 - diagenetic effects on, B:258
 - void ratio vs. applied stress, B:259
 - tests, B:258
 - Convection systems, Ninetyeast Ridge, geochemical effects of, A:533
 - Coral Sea, spreading end, B:816
 - Costa Rica Rift
 - geochemistry, vertical gradient uniformity, A:283
 - stable isotopes, pore waters, B:447
 - stress, direction of, B:697
 - Cretaceous/Tertiary boundary. *See also under minerals; particular fossil species; sediment categories; sites*
 - Broken Ridge
 - magnetic properties, B:941
 - sedimentation rate, B:739-740
 - fossil transition, Site 752, B:939
 - Site 752, A:122-123, 463, 500, 507-516, B:492, 730, 740, 937, 941
 - biostratigraphy, A:507-512, B:366
 - clay, origin of, B:419-420
 - depositional environment, A:507
 - geochemistry, B:417-418, 424-429
 - lithologic expression, A:112, 120-123
 - lithology, B:366, 415-416, 423-424
 - lithostratigraphy, A:507
 - logging, A:513-516
 - magnetic properties, A:510, 512-513
 - magnetostratigraphy, A:133, B:363, 366, 730-731
 - physical properties, A:148
 - productivity, B:420-421
 - sedimentation rate, A:507
 - Site 754, magnetostratigraphy, B:735
 - Site 758
 - lithostratigraphy, A:372
 - magnetic properties, B:795
 - Crozet Basin, oceanic crust, B:765
 - Crust, oceanic
 - Broken Ridge
 - gravity anomalies and thickness of, A:86
 - northward drift, Eocene, B:220
 - oxygen isotopes with alteration of, B:452
 - pre-rift, vs. post-rift Southeast Indian Ridge, B:690
 - Ninetyeast Ridge
 - age, B:774
 - oxygen isotopes with alteration of, B:452
 - Cryptoturbidites, Ninetyeast Ridge, B:242
 - Daito Basin
 - basalt
 - geochemistry, B:550-551
 - induration of adjacent sediments, B:551
 - magnetic properties, B:550-551
 - basalt sill, age, B:550
 - interbedded claystone-basaltic units, B:550
 - Deccan Traps
 - diastrophism, India-Asia convergence and, B:819
 - extrusion
 - correlation, Chronozone 31N, B:818
 - paleolatitude, B:778, 805
 - volcanic source migration and, B:819
 - hot spot origin, relative motion models, B:770
 - paleoposition, B:816-818
 - radiometric age, B:770
- Deformation, Broken Ridge, pre- vs. post-rift, B:522
- Deformation rate analysis, Site 758
- for *in-situ* stress estimation, B:699-701, 706
 - Kaiser effect of acoustic emissions, B:698
 - strain difference function, B:698, 700-701, 706-707, 714-715
- Density
- Broken Ridge, A:140, 144, 163, 182-183, 217, 250-251, 500, 505, B:472
 - carbonate content and, B:269
 - dipping and truncated sequence, A:497
 - glauconite and, A:252
 - laboratory vs. logging data, A:149-150, 160
 - opal-A to -CT transition, B:256-257
 - pelagic cap, A:495, 497, B:254, 264
 - silica diagenesis and, A:498
 - vs. velocity, B:521, 684
 - Ninetyeast Ridge, A:287-288, 292, 337, 402-404, B:316-328, 472
 - basalt, A:297, 344
 - calcareous chalks, B:256
 - GRAPE vs. discrete, B:299
 - lithologic Unit I, A:343
 - Site 752, laboratory vs. logging measurements, A:160
- Depositional environment
- open-ocean transition, Site 757, A:350
 - Site 753, A:171
 - Site 756, A:260
 - Site 757, A:311

- Site 758, Paleocene, A:401
 Diatoms. *See also under particular sites and by taxa in Taxonomic Index*
- Broken Ridge
 Eocene, A:476
 zonation, A:45
- Ninetyeast Ridge, zonation, A:45
 Site 752, A:129-130, B:722
 biostratigraphic events, B:173-175
 datum levels, B:179
 Eocene-Paleocene, A:129
 Eocene/Paleocene boundary, A:130
 magnetostratigraphic correlation, B:172-173
 Neogene, A:487-488
 Paleocene-Eocene, B:171-178
 Paleogene, A:488
 preservation, B:263-264, 268-269
 recrystallized, B:271
 zonation, A:129-130, B:175
- Site 753
 Eocene, A:179
 Neogene, A:487-488
 Paleogene, A:488
- Site 754, A:205
 Cretaceous, A:488
- Site 755, A:246
- Site 757, A:320
- Site 758, A:386
 Campanian, B:199
 datum levels, B:191
 Miocene-Oligocene, B:212-213
 Miocene/Oligocene boundary, B:199
 Miocene/Oligocene hiatus, B:199
 Neogene, B:192
 Oligocene-Miocene, B:195-197
 paleoenvironmental implications, A:375-376
 Pliocene-Quaternary, B:190-195, 209-211
 preservation, A:375
 stratigraphic range, B:198, B:193
 zonation, A:376, B:191, 194-195, 198-199
- Dipping and truncated sequence. *See* Broken Ridge, dipping and truncated sequence
- Dogie Creek (Wyoming)
 goyazite spherules, Cretaceous/Paleogene boundary, B:497, 503
 kaolinitic spherules
 altered to ferruginous smectite, B:491
 Cretaceous/Paleogene boundary, B:490
- Dolomite, Broken Ridge, A:462-463, 494, B:470
- Downwelling, Ninetyeast Ridge, oxygen isotopes and rate of, B:452
- Dupal anomaly, Ninetyeast Ridge, B:601, 603, 609, 638-639
- Earthquakes, Ninetyeast Ridge, focal mechanisms, A:13
- East Pacific Rise
 basalt sheet flow, B:556
 hydrothermal deposition, tectonic reorganizations and, B:444
- Eastman-Whipstock Multishot core orientation
 Site 757, A:331
 Site 758, magnetic properties, A:398, 418
- Emperor Seamount, primary oxide composition, B:539
- Eocene angular unconformity. *See* Broken Ridge, Eocene angular unconformity
- Eolian dust
 Broken Ridge, B:219-221
 Site 756, A:262, B:219-221
 Site 757, A:309
 Site 758, accumulation rate, A:363
- Equatorial Water Zone, Indian Ocean, productivity, A:18
- Erosion, Broken Ridge
 prograding vs. limestone-chert sequences, A:84
 sediment volume, B:746-747
 subaerial, A:469
- Ethane, Site 758, A:419
- Europium anomaly
 Ninetyeast Ridge, B:635
 basalts, B:570-571, 627
 intrasite variation, B:575-576
 Site 757, B:628
- Eustatic sea level
 Broken Ridge
 coastal onlap shift with changes in, A:81
 downlapping, B:747
 grain size with changes in, B:943
 isotopic correlation, B:943
 Oligocene disconformity and fall of, A:90, B:438
 stratigraphic response, B:753
 Site 758, sedimentary record correlation, B:823
 stratigraphic response to, carbonate vs. terrigenous margins, B:743, 753
- Evaporates, Leg 21, A:42
- Fault, normal, Broken Ridge, instantaneous slip, A:11
- Fault, transform, Ninetyeast Ridge, A:260-261
- Faulting, Site 756, calcium gradient and, A:286
- Feldspar, Ninetyeast Ridge, B:568-569
 inclusions, B:571
 tephra layers, B:284
- Feldspar, plagioclase, Ninetyeast Ridge, B:567
- Feldspar, potassium, Ninetyeast Ridge, B:567, 625, 656-657
- Ferrobasalt, Ninetyeast Ridge, B:526
 isotopic variations, A:526
- Ferrotholeiite, Ninetyeast Ridge, incompatible-element-enriched, B:592
- Fish teeth, Site 756, B:83
- Flexural coupling, Broken Ridge, new ocean/old ocean boundary, B:692
- Florida escarpment, prograding clinofolds, B:753
- Foch Island, tholeiites
 eruption, B:599
 isotopic ratios, B:598-599
 Ninetyeast Ridge compared, B:598-599
- Foraminifers. *See* Benthic foraminifers; Planktonic foraminifers. *See also individual taxa in Taxonomic Index*
- Formation factor
 Broken Ridge, A:143, 148-149, 183, 185, 219-221, 227, 252
 vs. porosity, A:150
 Ninetyeast Ridge, A:288-289, 294-295, 342, 350, 352, 404, 407, 435-436
 lithologic Unit I, A:343
 Site 756, A:294-295
 Site 757, A:350, 352
 Site 758, A:435-436
- Fossil zones. *See in Taxonomic Index*
- Galapagos Islands, tholeiites and ferrobasalts, Ninetyeast Ridge correlated, A:526, B:579
- Galapagos Rift, basalt sheet flows, B:556
- Ganges-Brahmaputra river system
 sediment output, Himalayan uplift rate and increase in, B:310-311
 seismic reflection profiling, correlation, Middle Siwalkiks deposition, B:825
- Ganges River, sediment output, B:298
- Geochemistry, igneous, Ninetyeast Ridge, B:561
 Amsterdam-St. Paul hot spot signature, B:802
 fractionation effects, B:572
 incompatible elements, A:332, 393, 403
 trace elements, A:393
 vs. Kerguelen lavas, B:639-642, 644-645
- Geochemistry, inorganic
 Broken Ridge, A:56-57, 135-136, 138, 181, 183, 212-213, 247, 250, 491-492
 ash layers, A:472-475
 compositional end members, B:444
 lateral gradient variations, A:491-492
 logging data, B:895-905
 major elements, A:473-474
 trace elements, A:474, 476-477
 compositional end members, Site 752, B:439-444
 Ninetyeast Ridge, A:56-57, 283-284, 333-336, 398-400, B:586-588
 alteration effects, A:531
 basalts, A:275, 277, 279, 326-331, B:567-572
 crystal-melt segregation, A:529
 gradient uniformity, A:283
 incompatible elements, A:327, 329, 331, 525
 intersite differences, A:329, 525-526, B:633, 635, 637-638
 lavas, B:594-595, 616-620
 logging data, B:906-908, 910-912
 long-term trends, A:525-529
 reference standards, B:585, 614, 891-893
 short-term trends, A:529-530
 southern ridge sites compared, A:279
 tephra layers, B:277-284
 tungsten carbide vs. agate crushed aliquots, B:615
- Site 752
 compositional end members, B:424-430, 439-444
 factor analysis, B:424-425, 428-429, 432, 439-440
 hydrothermal end members, B:442-443
 mineralogical and physical properties correlation, B:265
 rifting mechanisms related to, B:444
 Site 758, A:398-400
- Geochemistry, lipid, B:743
- Geochemistry, organic
 Broken Ridge, A:57, 492-495
 bulk composition, A:493-494
 Ninetyeast Ridge, A:57, 286-287, 336-337, 400-402
 Site 752, A:136-138, 141-143
 Site 753, A:181
 Site 754, A:212-216
 Site 755, A:250-251
 Site 756, A:286-287, 290-291
 Site 757, A:336-337, 343-344
 Site 758, A:400-402, 420-421
- Geomagnetic polarity transition
 behavioral models, B:377
 Site 758, B:388
 directional variation, B:382-383, 386, 391
 Jaramillo Subchronozone, B:377
 looping behavior, B:385
- Geothermal gradient, Broken Ridge, calculated from diagenetic minerals, B:522
- Glacial/interglacial cycles
 Ninetyeast Ridge, stable isotope stratigraphy, B:452
 Site 758
 calcium carbonate MAR, B:307, 311

- planktonic foraminifers, B:130
sedimentation rate, A:363
stable isotope stratigraphy, B:304
- Glaciation**
Ninetyeast Ridge, Milankovitch cycles, B:823
Northern Hemisphere, initiation, B:312-313
- Glass, basaltic, Site 758, alteration to clay, A:521**
- Glass, volcanic**
Broken Ridge
alteration, A:472, B:521
discoloration, A:471
geochemistry, A:471-472
Ninetyeast Ridge
basaltic flow unit boundaries, B:549
recrystallized, A:324
- Glass shards**
Broken Ridge, A:472-473
Ninetyeast Ridge, A:520
geochemistry, B:277, 284
major oxides, B:277-283
morphological categories, B:277
- Glaucinites**
Broken Ridge, A:73, 240, 463, 465
accumulation rate, B:939
density and, A:252
depositional environment, A:89
formation environment, A:465, B:939-940
in Turonian tuffs, A:464
Ninetyeast Ridge, A:311
- Gold, Site 752, Cretaceous/Tertiary boundary, B:425, 429**
- Gondwana dike and sills, paleomagnetic pole positions, correlation, Rajmahal Trap poles, B:820**
- Gough islands, lava, Dupal isotopic signature, B:599**
- Goyazite spherules, Site 758, Cretaceous/Paleogene boundary, B:491-492**
- Grain size**
Broken Ridge, A:42, B:217
density and, A:495
dissolution effects, B:214
lithostratigraphic correlation, B:214
oceanic circulation effects, A:470
pelagic ooze, A:84, 86, 458, 460, 462, B:211-215, 937
with sea-level changes, B:943
seismic correlation, A:254
southern hemispheric ice buildup and, B:943
terrigenous organic particles, B:472, 476, 481
transport distance and, B:481
winnowing effects, B:212, 936
Ninetyeast Ridge, A:42
seismic correlation, A:294, 303
terrigenous organic particles, B:476, 481
terrigenous vitrinite and inertinite, B:472
transport distance and, terrigenous particles, B:481
- Site 752**
seismic stratigraphy and, A:153
unconsolidated sediment, A:116, 120
- Site 753, A:176**
seismic stratigraphy and, A:185
- Site 754, A:194, 198**
pelagic oozes, B:213-215
seismic reflection and, A:225
winnowing and, B:212
- Site 755, A:242**
- Site 756, A:267**
sedimentation rate and, A:265
- Site 757, A:348**
lithologic Unit I, A:311
- seismic correlation, A:357
seismic reflection profile correlation, A:348, 357
- Site 758, A:373-374**
ARM ratios and, B:379, 382
ash layers, B:277
basaltic flow units, B:549
lithologic Unit I, A:373-374
mineralogical correlation, B:379, 382
unconsolidated sediment, A:368
- Grainstone, shelly foraminifers, Site 755, A:238**
- Granular sediment, classification, A:40-42**
- GRAPE density**
Broken Ridge, A:57
pelagic cap, A:495, 499
Ninetyeast Ridge, A:57, 347, 402, 430-431
- Gravel, Broken Ridge, A:5, 71**
logging data, A:149
- Gravity anomaly, Broken Ridge, A:84-86, B:682**
magnetic response, B:682, 685
new ocean/old ocean boundary, B:691
topographic effects, B:684, 690-692
- Great Bahama Bank, prograding clinofolds, B:753**
- Green Tuff region (Japan), volcanoclastic sediment, burial diagenetic zoning, B:519**
- Guaymas Basin**
basaltic units
alteration temperature, B:553-554
porosity of surrounding sediments, B:553
geochemistry, organic, basalt layers, B:554
heat flow, B:552-553
hydrocarbon distribution, B:554
hydrothermal alteration, B:554
interstitial-water chemistry, basalt-seawater interaction, B:554
mineral assemblages, temperature of formation, B:553-554
oxygen isotopes, B:554-555
porosity, vertical variation, B:553
regional relationships, B:552
thermal metamorphism, sill contacts, B:555
thermogenic component, B:554
- Gubbio (Italy), iridium, Cretaceous/Tertiary boundary, B:415**
- Guinea, Gulf of, magnetic anomaly inversion, B:543**
- Gypsum**
Broken Ridge, supersaturation, B:448
Ninetyeast Ridge, supersaturation, A:335, 399, B:448
- Hafnium/lanthanum ratio, Ninetyeast Ridge vs. samarium/lanthanum ratio, B:644 vs. samarium/lanthanum trend, B:642**
- Hafnium/samarium ratio, Ninetyeast Ridge intersite differences, B:637 vs. lanthanum content, B:640**
- Hawaiian-Emperor seamounts, basalt, correlation, Ninetyeast Ridge, B:526**
- Hawaiian Islands**
lava geochemistry, with distance from spreading center, A:528-529
volcanism
magma mixing, temporal variation, B:642, 644
tholeiitic-to-alkalic transition, A:526, B:612, 633
- Heard Island**
basalt
lead isotopes, B:569
thorium/tantalum ratio, B:644
magmatic activity, B:609
- Heat flow**
Broken Ridge, A:143-144, 171, 498-499, 506
silica diagenesis and, B:266
Site 752, A:143-144
- Hell Creek (Montana), kaolinitic spherules, Cretaceous/Paleogene boundary, B:490**
- Hematite, Ninetyeast Ridge, secondary, B:542**
- Hess Rise**
basement, age, A:465
carbonate
accumulation rate, A:466
flux values, A:112
volcanism, B:933
- Himalayan Arc, oroclinal bending, extensional tectonism and, B:827, 829**
- Himalayan orogeny**
initial collision, A:16-17
Neogene susceptibility record, Site 758, A:424, 426-427
- Himalayan Syntaxis, Western**
Jammu-Kashmir region, Siwalik sedimentation, B:827, 829
uplift and thrusting, Ninetyeast Ridge susceptibility correlation, B:829
- Himalayan-Tibetan convergence zone**
initial contact, B:777
Paleogene, B:778
- Himalayas, uplift**
after Indian-Asian collision, A:363
sedimentation rate impact, A:20
- Himalayas, Higher**
tectonic activity, Ninetyeast Ridge susceptibility correlation, B:826
uplift, Siwaliks sedimentation following, B:824
- Himalayas, NW, Siwalik sedimentation, dating of, B:825**
- Hopanoids, B:481**
- Horst-graben features, Kerguelen-Heard Plateau, A:89**
- Hot spot**
Atlantic Ocean S, plate rotation over, B:517
Broken Ridge
as ash layer source, A:474
subsidence rate with proximity to, A:89
Indian Ocean, plate rotation over, B:517
Kerguelen-Heard Plateau, Ninetyeast Ridge formation and, A:517
Kerguelen/Ninetyeast Ridge
ridge formation and, A:13
volcanism, A:262
motion models, B:770-773
Ninetyeast Ridge
age, A:5
Cretaceous-Oligocene locations, B:769
distance from spreading axis, A:528-530, 534, B:591, 599, 642
episodic behavior, A:528
incompatible element association, A:330
in ridge formation, A:523-524
- Hunza region, uplift, dating, B:826**
- Hydrocarbon gases**
Broken Ridge, A:137, 212, 250
Ninetyeast Ridge, A:286, 336, 400
Site 752, A:137, 140-141
Site 758, A:419
- Hydrocarbons, low-molecular weight**
Broken Ridge, B:460-462
age variation in, B:461, 465
hydrogen stripping/thermo vaporization technique, B:457, 459-460
Ninetyeast Ridge, B:462-465
intersite variation, B:463, 465
vs. Broken Ridge sites, B:463

- Hydrocarbons, saturated
Broken Ridge
biological markers, B:483
gas chromatograms, B:478-480, 484-486
Ninetyeast Ridge
biological markers, B:483
gas chromatograms, B:478-480, 484-486
thermal stress characteristics, B:481
Hydrocarbons, steroid, B:473
Hydrogen index
Broken Ridge, A:218, 493, B:472, 476
vs. oxygen index, A:495
Ninetyeast Ridge, B:472, 476
Site 754, vs. oxygen index, A:219
Hydrothermal deposition, Site 752, active rifting
and increase in, B:444
- Iceland, tholeiites and ferrobasalts, Ninetyeast
Ridge correlated, B:579
Iceland-Faeroes volcanoes, tholeiitic basalt, A:526
Iddingsite, Ninetyeast Ridge, pseudomorphs in basalt, B:563
Igneous petrology
Broken Ridge, A:45-50
Ninetyeast Ridge, A:45-50, 273-279,
320-331, 387-393, 523-531, B:567
southern ridge sites compared, A:277, 279
Site 756, A:273-283
Site 757, A:320-335
Site 758, A:386-394, 399-404
Ilmenite, Ninetyeast Ridge, A:323, B:658
trellis lamellae, A:333, B:542
Ilmenite-magnetite grains, Site 756, B:545
Impedance, Broken Ridge, A:252
lithologic correlation, A:142
Incompatible elements, Ninetyeast Ridge
aluminum oxide comparison, B:637
basalts, B:564, 571, 574-575, 579-581, 621,
625, 631, 633, 645
lavas, B:627-628
oceanic island basalt characteristics, B:639
potassium oxide correlation, B:629
rubidium correlation, B:629
vs. lanthanum content, B:630-631, 640-641
India-Asia convergence
completion of suturing, paleolatitude
constraints, B:812, 816
Cretaceous/Tertiary boundary
faunal evidence, B:818-819
initial collision, B:820-821
paleomagnetic record, B:816-818
evolution
Neogene, B:778-779
stages of, A:453, B:876
geological record, B:819
initial collision, B:777-778, 811-812
Deccan Traps diastrophism and, B:819-
820
magnetic susceptibility record, Site 758,
A:423-427
paleolatitude, B:778, 811-812
secondary components, collision-related,
B:877
sedimentary record, Ninetyeast Ridge,
B:821-829
Indian/Antarctic plate boundary, migration,
through Kerguelen Fracture Zone, A:537
Indian-Australian plates, relative motion models,
B:770, 881
Indian Bottom Water, Site 758, carbonate dissolution
and corrosiveness of, B:307
Indian Deep Water, Site 758, carbonate dissolution
and corrosiveness of, B:307
Indian-Eurasian convergence, A:20
- Indian Ocean
carbon isotope stratigraphy
Chron 6 shift, B:251
levels A-H, B:249-250
carbonate compensation depth, corrosiveness
with deepening of, B:309
geochemistry, standard sediment sample,
B:891-893
heat flow, vs. age, A:506
igneous geochemistry, A:14
lead isotopes, B:606
magnetic anomalies, A:6, 518
paleoceanic circulation, Cretaceous/Tertiary
boundary, B:820-821
paleoceanographic zones, A:17-18
Paleocene/Eocene climatic change, B:816
Indian Ocean, equatorial
plate boundary formation, correlation,
Ninetyeast Ridge NRM intensity,
B:826
sea-surface temperature, vs. North Atlantic
abyssal temperature, B:304
Indian Ocean, protomagnetic anomaly 34, A:30
paleocurrent, Cretaceous, B:487
physiography, A:31
Indian Ocean E
magnetic anomaly 13, A:25, 27
magnetic anomaly 21, A:28
magnetic anomaly 24, A:29
Indian Ocean E, subtropical, A:24
Indian Ocean NE
coarse fraction, long-term changes, B:309
Eocene/Paleocene boundary hiatus, B:816
tephra, B:273-274
Indian Ocean S, oceanic island basalts, isotopic ratios, B:592
Indian Ocean SE, geological setting, A:71
Indian plate
Asian collision, northward motion after,
A:306-307, 362-363
equatorial crossing, B:811
northward motion, B:516, 777
Cenozoic, A:19
Cretaceous/Tertiary boundary, B:818
Morgan/Patriat pole positions, B:875
paleolatitudinal control, B:811
relation to Australian motion, A:13
slowdown, B:778, 812, 816
paleomagnetic pole positions, B:874, 879-880
Indonesian Arc, ash
Ninetyeast Ridge sites, A:374
tephrochronology, A:363-364
Indus-Tsangpo suture zone, secondary magnetization
component, B:811-812
Inertinite, A:496, B:472, 476
Inertinite/vitrinite ratio, B:472, 481
Interstitial-water chemistry
Broken Ridge, A:138, 182, 212, 215, 250
interhole variation, A:491
oxygen isotopes, B:448-452
volcanic alteration effects, A:135, B:448
Ninetyeast Ridge, A:289, 334, 341-342,
418-419, 531-532
oxygen isotopes, B:448-452
volcanic alteration effects, B:448
Site 756
diffusion barrier, 60 mbsf, A:286
isotopic values, B:449, 452
lateral variation, A:283
vertical advection, A:284, 286
Site 757, interhole variation, A:335-336
Site 758, vertical gradient, A:398
Intertrappean Beds, microvertebrate fauna, Laurasian
affinities, B:818
- Iridium
Broken Ridge, magnetic susceptibility, B:492
Cretaceous/Tertiary boundary, B:415, 417,
731,913-916
origin, B:420, 423
Ninetyeast Ridge
kaolinic claystone, B:490-491
magnetic susceptibility, B:492
Site 752
deposition of ash and, B:420
measurement procedures, B:416-417
nanofossil extinction and, B:418-419
non-Cretaceous/Tertiary boundary peaks,
B:421-422
Iridium/cesium ratio, Site 752, B:420
altered ash and basalt, B:417
Iron, Site 752, source of, B:440
Iron/cesium ratio, Site 752
Cretaceous/Tertiary boundary, B:417-420
vs. iridium, B:421
Iron oxide, Ninetyeast Ridge
basalts, B:567, 571, 574
tephra layers, B:277
vs. aluminum oxide, B:277, 285, 574
vs. calcium oxide, B:277, 285
vs. iron oxide/magnesium oxide ratio, B:574
Iron oxide/magnesium oxide ratio
Ninetyeast Ridge
basalts, B:526, 574
vs. nickel and chromium content, B:567-
568, 571
vs. silicon dioxide, A:279
Site 758, vs. silicon dioxide, A:400
Iron pyrite, Site 758, basalts, A:391
Iron-stained sediments, Site 752, hydrothermal
source, B:444
Iron-titanium oxides, Ninetyeast Ridge, B:534
Isothermal remanence acquisition
Broken Ridge, B:361, 363
Site 758, B:379
- Japan arc, hydroclastic tuff, A:522
Japan Sea, opal-A to -CT conversion, B:522
Jaramillo Subchronozone, Site 758
geomagnetic polarity transition, B:377
looping behavior, B:384-385
magnetic properties, B:379
oxygen isotope stratigraphy, B:306
- Kalahari region (Africa), Quaternary rainfall, Quaternary
increase, B:221
Kali Gandaki Graben, Himalayan tectonism and,
B:426, 827
Kaolinite spherules, North America, Cretaceous/Paleogene
boundary, B:490
Karakorum Batholith, tectonic activity
dating, B:826
Ninetyeast Ridge susceptibility breaks and,
B:826
Karakorum-Jiali Fault Zone, deformation, Himalayan
Arc bending and, B:827
Kerguelen Archipelago
basalt, isotopic ratios, Ninetyeast Ridge
compared, B:598-599
evolution, spreading ridge migration and,
B:599
lavas
barium/niobium ratio, B:638-639
Dupal isotopic signature, B:599
volcanism, tholeiitic-to-alkaline transition,
B:640-642, 644-645
Kerguelen-Broken Ridge Plateau
constructional volcanism, B:933-934
formation, constructional volcanism in, B:943

- oceanic circulation, biogenic productivity and, B:940
 spreading center, northward migration, B:768–769
 tectonic history, Eocene split, B:933–934
- Kerguelen Fracture Zone, B:765
- Kerguelen-Heard Plateau
 constructional volcanism, A:499, B:559
 hot spot, B:507
 in Ninetyeast Ridge formation, B:639
 paleogeographic reconstruction, A:8, B:516
 subsidence, productivity and, B:747
 tectonic history, A:5
 debris flow record, B:747
- Kerguelen-Heard Plume, migration, A:534–535
- Kerguelen hot spot
 location, Cretaceous, B:371
 origin, B:514, 516
 predicted path, B:516
 in ridge formation, B:516
- Kerguelen Island
 basalt, thorium/tantalum ratio, B:644
 lavas, Dupal isotopic signature, B:645
 magmatic activity, B:603
- Kerguelen/Ninetyeast hot spot
 kinematic models, B:770–773
 migration, B:769–773, 776
 present-day location, B:770–773
- Kerguelen Plateau
 Broken Ridge, transport, redistribution, and deposition, A:465
 chalk, Santonian–Turonian, B:938
 deformation, continental lithospheric component, B:644
 diatom ooze, B:938
 formation, A:465, 470
 glauconite, environment of formation, A:465
 glauconite sandstone, B:938
 hot spot origin, B:514
 lava, lead isotopes, B:599
 limestone, B:938
 lithostratigraphy, Broken Ridge compared, A:463–464, B:938
 nannofossil ooze, B:938
 neodymium/lead isotope ratio, Ninetyeast Ridge compared, B:599
 paleoeposition, B:941
 planktonic foraminifers, B:921
 separation from Broken Ridge, A:5
 strontium isotopes, B:921–924
 nannofossil correlation, B:924, 927
 tholeiitic basalt, A:526
 uplift, Maestrichtian, A:467
- Kohistan, Northern, suture zone, secondary magnetization component, B:811–812
- La Palma Island, primary oxide composition, B:539
- Laccadive Ridge
 basement, age, B:770
 hot spot origin, relative motion models, B:770
 hot spot track, B:770
- Lamellae, Broken Ridge, ash layers, A:460, 462
- Lamellae, trellis, Ninetyeast Ridge, B:545
 ilmenite, B:545
 titanomagnetite, B:535
- Lanthanum, Ninetyeast Ridge
 compositional uniformity, B:629
 vs. incompatible elements, B:630–631, 640
 vs. thorium and tantalum, basalts, B:578
 vs. ytterbium, basalts, B:568
- Lanthanum/cesium ratio, Ninetyeast Ridge, vs. lanthanum content, B:642
- Lanthanum/samarium ratio, Ninetyeast Ridge
 vs. lanthanum content, B:642
 vs. strontium isotopes, B:599, 608
- Lanthanum/tantalum ratio, Ninetyeast Ridge
 basalts, B:568, 571
 basalts, lithosphere-derived, B:581
 intersite comparison, B:510, 570
 oceanic island basalt characteristics, B:644
 vs. tantalum, B:582
 vs. thorium/tantalum ratio, B:581, 584, 641, 644
- Lanthanum/ytterbium ratio, Ninetyeast Ridge
 basalts, B:571
 basalts, type I vs. type II, B:570
 vs. lanthanum content, B:642
- Lapilli, Site 757
 in ash layers, A:311
 basalt contacts, A:321
 physical properties, A:337, 344
- Lapilli, accretionary, Site 757, A:522–523
- Lava
 Broken Ridge
 evolved, A:13–15
 oceanic andesite overlay, A:13
- Ninetyeast Ridge
 alteration, brown vs. blue-green, B:624
 asthenospheric and plume-related components, B:641
 Dupal isotopic signature, B:645
 enriched mantle component, B:580
 evolved, A:524
 formation, intersite variation, B:600–601
 geochemistry, A:525, B:633
 Indian Ocean basalts compared, B:598–601
 isotopic ratios, B:603
 mineral abundance and composition of, B:633
 petrology, B:616–620
 post-magmatic alteration, B:614–615
 two-component magma mixing, B:599, 633, 641
 vs. St. Paul/Amsterdam islands, A:526
 pillowed vs. non-pillowed, B:547
 Site 756, subaerial flows, A:274–275
 Site 757
 heterogeneous flow, B:597
 mineral abundance and composition of, B:625
 petrography, B:567
 plagioclase-rich, A:529
 subaerial flows, A:322
 two-component magma mixing, B:628–629
- Site 758
 depositional environment, vs. ash, A:360
 secondary mineralogy, B:577
- Lead, Ninetyeast Ridge, B:600–602
- Lead isotopes
 Indian Ocean, mid-ocean ridge basalts, A:15
 Ninetyeast Ridge, B:580, 605
 alteration effects, B:596–598
 basalts, B:569–573, 575, 579
 intersite variation, B:598
 Kerguelen lava ratios compared, B:599, 604, 640, 642
 lavas, A:529, B:600–603
 phonolites vs. basalts, B:599
 source components, B:633
 vs. neodymium isotopes, B:578, 581
 vs. strontium isotopes, A:529, B:607
- Ninetyeast Ridge vs. Kerguelen Archipelago, A:526
- Lead/neodymium ratio, Ninetyeast Ridge, vs. lanthanum content, B:640
- Lepispheeres
 Broken Ridge
 calcareous sediments, B:267
 opal-CT, B:272
 Ninetyeast Ridge, calcareous sediments, B:257
- Lhagoi-Kangri belt, formation factors, B:824
- Limestone
 Broken Ridge
 ash-bearing vs. ash-barren, A:81
 carbonate content, B:475
 depositional environment, A:469
 induration, A:81, 462
- Site 754
 ash-barren vs. ash-bearing, A:226
 mottling, A:194
 seismic reflection profiling, A:226
- Site 755, A:465
 Santonian, B:937
- Site 756, seismic reflection profiling, A:293
- Limestone, ashy
 Broken Ridge, A:239
 magnetic susceptibility, A:206
- Site 755
 Santonian, B:440
 Site 752 compared, B:440, 442
- Limestone, foraminifers, Site 756, A:264, B:229
- Limestone-chert sequence, Broken Ridge, A:5
 contact, A:199
 Santonian, A:112
 seismic stratigraphy, A:72–75
 uplift, A:9
- Limestone-ooze transition, Broken Ridge, seismic reflection profiling, B:682
- Limestone pebbles
 Site 752, A:116, 119, B:937
 Oligocene disconformity, B:936
- Site 753, A:176
- Lingzizong Formation, magnetic properties, Paleocene, B:816
- Lipids, geochemistry, B:473
- Liptinite, B:472–473
- Lithification, Site 752, causes of, B:267–268
- Lithologic column, Broken Ridge, B:723
- Lithosphere, strength of, Broken Ridge, A:8–9
- Lithosphere extension, Broken Ridge, B:663, 721
 isostatic response, A:8
 kinematic model, A:8–9, 11
 simple normal-faulting model, A:9–12
 sublithospheric mantle and initiation of, A:7–8
- Lithostratigraphy
 Broken Ridge, B:219–220
 Eocene hiatus, Site 758, A:375
 Eocene–Miocene hiatus, Site 753, A:86
 Eocene–Oligocene hiatus, Site 752, A:86
 interhole correlation, Site 758, A:375
- lithologic Unit I
 Site 752, A:111, 116, B:732
 Site 753, A:171, 173
 Site 754, A:191, 194, B:733
 Site 755, A:237–239, B:735
 Site 756, A:260, 265
 Site 757, A:306, 311
 Site 758, A:360, 367–368
- lithologic Unit I/Unit II contact, Site 755, A:242
- lithologic Unit II
 Site 752, A:111–112, 116, 119–121, B:732
 Site 753, A:171, 173–179
 Site 754, A:191, 194–195, B:733
 Site 755, A:237, 239–240, 242, B:735
 Site 756, A:260, 265–266
 Site 757, A:306, 311

- Site 758, A:360, 368, 372-373
 lithologic Unit III
 Site 757, A:306, 311
 Site 758, A:360, 373-374
 lithologic Unit IV, Site 758, A:360, 374
 lithologic Unit V, Site 758, A:360, 374-375
 Maestrichtian-Eocene hiatus, Site 754, A:86
 magnetic properties and, Site 758, B:882
 NRM intensity variations, Site 758, A:395
 physical properties correlation
 Site 754, A:219
 Site 757, A:343-344
 Santonian-Miocene hiatus, Site 755, A:86
 seismic correlation
 Broken Ridge, A:165, 188, 235, 257
 Site 752, A:151, 165, 257
 Site 753, A:165, 185, 188, 257
 Site 754, A:224-225, 235, 257
 Site 755, A:254-255, 257
 Site 756, A:293-295
 Site 757, A:345, 347-350, 357
 Site 758, A:344, 414, 418-419
 Site 752, A:116-119
 Site 753, A:172-173
 Site 754, A:196-198, B:219-220
 Site 755, A:238-242
 Site 756, A:265-267, B:219-220
 Site 757, A:311
 Site 758, A:367-371
 stratigraphic inversions, Site 758, B:242
 volcanogenic components, Broken Ridge,
 B:519-520
- Logging
 gamma ray, Site 754, A:220
 geochemical
 Site 752, A:148-149, 154-156, 158-160,
 B:895-901
 Site 754, A:219, 221, 224, 230-232,
 B:902-905
 Site 758, A:366-367, 409-410, 438-442,
 B:906-908, 910-912
 geochemical, vs. core-derived lithology, B:908
 geophysical
 Site 752, A:148-151
 Site 754, A:219, 221
 Site 758, A:408-410, 414
 lithoporosity, Site 752, A:148-149, 154-156,
 158-159
 objectives, Site 758, A:363-364
 resistivity, Site 754, A:230
 seismic stratigraphic
 Site 752, A:148
 Site 758, A:366-367, 409-410, 414, 438-
 440
 Site 752, A:167-169
 synthetic seismogram
 Site 752, A:150, 161-162
 Site 754, A:221, 232-233
 Louisville Ridge, lavas, geochemistry, A:528-529
- Madingley Rise
 carbon isotope stratigraphy, B:243, 246, 250
 Oligocene/Miocene boundary, B:248-249
 magnetostratigraphy, B:242
 oxygen isotope stratigraphy, B:243, 246
 oxygen isotopes, Eocene/Oligocene boundary,
 B:247
 Maestrichtian/Eocene unconformity, physical
 properties, Site 754, A:217
 Magellan Rise, carbonate accumulation rate,
 A:466
 Cretaceous, B:940
 Magma
 Broken Ridge, source, A:474
 melt generation
 influence of lithospheric extension on,
 B:579
 mantle temperature influence, B:579
 Ninetyeast Ridge, source, B:603, 633, 642
 Magma, parental, Ninetyeast Ridge, A:277, 329-
 330, 393, B:628
 tephra layers, B:287
 Magma chamber, Ninetyeast Ridge, A:14-15
 Magma mixing, Ninetyeast Ridge
 MORB- vs. OIB, A:526, 528, 530-531
 ocean island and mid-ocean ridge basalts,
 B:592, 637
 two-component, B:599, 628-629, 633, 642
 Magnesium
 Broken Ridge
 ash content and gradient changes, A:136,
 492
 intersite variation, A:180
 pore water concentration, A:213, B:448,
 450
 Ninetyeast Ridge
 ash abundance and, A:283-284, 334
 between-hole variation, A:284
 calcium and variations in, A:286
 effect of alteration on, A:399
 pore water concentration, B:448, 450-451
 vertical gradient, A:398
 Magnesium oxide, Ninetyeast Ridge, A:281
 basalts, A:393, B:571
 basement basalts, A:335
 fractionation effects, B:622, 624
 lavas, B:626-628
 residue-melt segregation, B:633
 in tholeiitic basalts, A:277
 vs. aluminum oxide, A:329
 vs. calcium oxide/aluminum oxide ratio, B:639
 vs. nickel content, basalts, A:402
 vs. nickel content, lavas, B:621, 629
 vs. sodium oxide, basement basalts, A:401
 Magnetic anomalies
 Broken Ridge, A:5
 Ninetyeast Ridge, B:542-543
 Magnetic anomaly 13, A:27
 Magnetic anomaly 21, A:28
 Magnetic anomaly 24, A:29
 Magnetic anomaly 34, A:30
 Magnetic mineralogy, basalts
 Site 756, A:282-283
 Site 757, A:333
 Magnetic properties
 alteration effects, Ninetyeast Ridge,
 B:539-542
 basalts
 Ninetyeast Ridge, B:526-534
 Site 756, B:530
 Site 757, B:531
 Site 758, B:532
 Broken Ridge, A:63
 alternating-field demagnetization, A:51, 55
 ash layers, A:482
 compaction-related errors, B:371-372, 374
 declination bias, B:360
 demagnetization, type I and II behaviors,
 B:361-362, 367
 gravity anomaly, B:682
 inclination, B:367, 371, 373
 lithologic correlation, A:475
 natural remanent magnetization, A:50,
 492-493
 postdepositional remanence, B:372-373
 rifting overprint, A:491
 shallow inclinations, origin, B:371-374
 susceptibility, A:50, 468, 475, 482-483,
 491-493
 Broken Ridge/Kerguelen-Heard Plateau,
 seafloor-spreading anomaly, A:71
 directional disturbance, A:51
 Ninetyeast Ridge
 alternating-field demagnetization, A:51,
 55, B:780-783
 direction of 180 degree component,
 B:783-784, 798
 Eulerian rotation poles, B:782
 grain size and, B:822
 gravity anomaly, B:561
 hysteresis loop measurements, B:533-534,
 536-537
 inclination, B:803, 805
 inclination, basalts, B:533
 linear regression analysis, B:805, 809-810
 multiple overprint components, B:779,
 781, 783, 795, 802
 natural remanent magnetization, A:50,
 B:526-533, 855
 primary magnetization components,
 B:783-792
 shallow inclinations, effects of secondary
 mineralogy on, B:542
 susceptibility, A:50, B:533, 799, 823-827,
 829, 855
 viscous remanent magnetization, B:783
 Site 752, A:130-137
 anhysteretic remanent magnetization,
 A:133, 135
 correlation, volcanic ash content, A:130
 lithologic Unit I vs. Unit II, A:130
 logging data, A:513-514
 susceptibility, A:130
 Site 753, A:179-182
 susceptibility, A:178-179
 Site 754, A:192, 205-212
 secondary magnetizations, B:367
 Site 757 correlation, A:332
 susceptibility, A:206
 Site 755, A:238, 246-249
 susceptibility, A:247
 Site 756, A:101, 279-283, 285-288, B:784
 alternating-field demagnetization, B:784,
 793-796, 798-799, 802-803
 inclination, B:785-786
 mean inclinations, B:800-801
 natural remanent magnetization, A:281-
 282, 285, 287-288, B:797
 seismic correlation, A:294
 susceptibility, A:279-284
 thermal demagnetization, B:784
 thermal stability, B:797, 803
 Site 757, A:52-53, 101, 105, 331-333,
 338-340, B:784, 792
 alternating-field demagnetization, B:806-
 809, 812-815, 817, 820-821, 824
 825
 inclination, B:781, 786-792, 801
 natural remanent magnetization, B:805,
 818
 polarity, A:339
 remanence, A:331-333, 338-340
 susceptibility, A:331, 336-337
 thermal demagnetization, B:784, 792, 804
 thermal stability, B:816
 viscous remanent magnetization, A:332
 Site 758, A:394-398, 410-411, B:792-797
 alternating-field demagnetization, B:701,
 827-829, 840, 842-844, 846-849,
 851-854
 APC-cored sediments, B:792-794, 801

- ash layer, B:796, 801
 basement flows, B:796-797, 801
 between-hole correlation, A:396-398
 carbonate content and, A:360
 declinations, B:691, 705, 710-712
 demagnetization, B:379, 381, 385
 inclination, B:382-385, 793-794
 inclinations, B:801-803
 interhole correlation, A:396-398
 Jaramillo polarity transition, B:379
 lithostratigraphic correlation, A:395, 397, B:882
 low-stability components, B:379
 median destructive field, A:395
 natural remanent magnetization, A:55, B:379, 782, 826, 828
 NRM/ARM ratio, B:379, 382
 NRM intensity, B:383
 Olduvai/Jaramillo transition zone, B:388
 Oligocene-Pliocene hiatus, B:795
 oxygen isotope stratigraphy and, tephra layers, B:285
 paleolatitude, A:395-396
 primary magnetization components, B:793, 796, 841
 remanence, A:394-396, 410
 susceptibility, A:394, 397-398, 405-407, 416-417, 423-424, 426-427, 449-450, B:276, 299-230, 300, 306-313, 340-355, 379, 384, 805, 825-826, 829
 susceptibility breaks, A:424-427, 452
 susceptibility units K1-K9, boundary breaks, A:426-427, 451-452, B:881
 thermal demagnetization, B:379
 thermal stability, B:796, 822, 839, 845-846, 850, 855
 viscous remanent magnetization, B:701, 709
 XCB-cored sediments, B:795-796, 801
 time scale, A:50-51
- Magnetite**
 Cretaceous/Tertiary boundary, Site 752, A:512
 Site 757, A:323
- Magnetostratigraphy. See also Cretaceous/Tertiary boundary**
 Broken Ridge, B:729
 chronozone boundaries, B:727-728
 dipping and truncated sequence, A:488-491, B:359
 GRTS correlation, B:363
 pelagic cap, A:488
 reversal boundaries, B:365
- Ninetyeast Ridge**
 anomalies, B:767
 Brunhes Chron to Chron 6, B:797
 chronozones, B:798-799
 normal polarity interval, B:797
 Site 752, A:131-134, 136, B:363-365, 725-731
 biostratigraphic correlation, A:132-133, B:730
 chron boundaries, B:363, 365, 727-728, 730-731
 Eocene/Paleocene boundary, B:363
 extra polarity events, A:489, 491
 low-stability overprints, B:363
 normal interval in reversed zone, A:133
 polarity reversals, A:112
 Site 754 correlations, A:206, 210, 212
 Site 753, A:171, 179-180, B:365, 367-368, 732
 Chronozone C20N/C20R transition, B:732
- seismic correlation, B:365, 367
 Site 754, A:206, 210, 212, B:369, 733-735
 biostratigraphic correlation, A:206, 210, 212
 Campanian/Maestrichtian boundary, B:367
 chron boundaries, B:367, 733, 735
 Site 755, A:247, B:367, 370, 735-736
 Site 756, A:282
 Site 757, A:332
 Site 758, A:395, 407-409, 413, B:779
 analytical methods, B:377-379
 between-hole correlation, A:397
 Brunhes Chron to Chron 6, B:245
 declination drift, A:413
 normal/reversed polarity difference, B:794, 801
 polarity reversals, B:302
 reversal boundaries, B:855
- Mammoth Subchronozone, Site 758, sedimentation rate, B:302**
- Manganese, Site 752, B:439**
 Cretaceous/Tertiary boundary, B:424
- Manganese/iron ratio, Site 752, B:440**
 of end-members, B:425
 hydrothermal sediments, B:443
- Manihiki Rise**
 basement, age, A:465
 carbonate accumulation rate, A:466
 Cretaceous, B:940
 constructional volcanism, B:933
- Mantle geochemistry, Ninetyeast Ridge**
 isotopic and incompatible element ratios, A:526
 multiple sources, A:525
- Mantle plumes, Ninetyeast Ridge, role in origin of, B:578-581**
- Maquiang Volcanics, formation factors, B:824**
- MAR. See Mass accumulation rate**
- Marion hot spot, predicted path, B:516**
- Marpo Limestone, erosion, Ninetyeast Ridge Eocene unconformity compared, B:819**
- Mascarene Plateau**
 formation, Réunion hot spot role, B:770
 Réunion hot spot, predicted path, B:514
- Mass accumulation rate**
 Cretaceous/Tertiary boundary, Site 752, B:426-427
 Site 752, B:426-428
 geochemical end members, B:439, 442
- Matuyama/Brunhes transition, Site 758, virtual geomagnetic pole, B:377**
- Matuyama/Gauss transition, Site 758, oxygen isotope stratigraphy, B:306**
- Mauna Kea Volcano**
 lanthanum content, B:643
 volcanic edifice cross section, A:17
- Mauritius Island, hot spot origin, relative motion models, B:770**
- McDonald Island, volcanic activity, hot spot role, B:771-772**
- Mesostasis, Ninetyeast Ridge, B:526**
 basalts, A:389
- Messinian desiccation event, Ninetyeast Ridge, carbon isotope depletion, B:250-251**
- Metalliferous sediment**
 Broken Ridge, A:42
 Ninetyeast Ridge, A:42
- Metals, transition, Site 756, B:621**
- Methane**
 Broken Ridge, A:140-141, 212, 215, 492
 Site 758, A:419
- Micrite**
 Site 755, A:73, 239
 Site 757, A:311
- Microtektites**
 Site 752, B:489-492
 Site 758, B:489-493, 495-496, 502
 carbonate content, B:490
- Mid-ocean ridge basalt (MORB)**
 Indian Ocean, radiogenic isotopes, A:15
 Ninetyeast Ridge, Indian plate migration over source of, B:765
- Middle Siwalkiks Sedimentation, initiation, Site 758 susceptibility correlation, A:427**
- Mineralogy**
 Broken Ridge, silica, B:261-262
 Ninetyeast Ridge
 alteration, B:659
 geochemical correlation, basalts, B:633
 oxidative assemblages, basement basalts, B:56
 secondary, A:275, 278, 324, 393, B:542, 563
 shock-metamorphosed, Cretaceous/Paleogene boundary, B:489
 Site 752, chemical and physical properties correlation, B:265
 Site 756
 basalt, A:278
 primary, A:275, 278
 Site 757
 basalt, A:323-324
 geochemical correlation, basalts, B:625
 interflow-unit differences, A:324, 326
 vesicle fillings, A:324-325
 Site 758
 ash layers, B:277
 basalt, A:392
 grain size and, B:379, 382
 mafic mineral fractionation, tephra layers, B:287
 secondary, A:393
- Mineralogy, igneous, Ninetyeast Ridge, B:567**
 crystal settling, A:323
 groundmass phases, A:275, 323-324, 392
 primary, A:392-393
 secondary associations, A:330
- Mineralogy, magnetic, Ninetyeast Ridge**
 basalts, A:282-283, 333, B:534, 539
 magnetic property effects, B:385, 822
 petrological correlation, B:526
- Mineralogy, opaque, Site 756, A:275**
- Moho topography, Broken Ridge, B:692**
 gravity effect, B:684, 690-691
- Monsoon**
 Indian Ocean, climatic evolution, A:359, 363
 Ninetyeast Ridge, climatic fluctuations, A:21
 Monterey carbon excursion, Indian Ocean, B:250
 Monterey Formation, opal-CT, depth of, B:266
 MORB. See Mid-ocean ridge basalt
 Mordenite, Broken Ridge, calcium-rich, B:521
- Nagri-Dhok Pathan Formation transition, formation break, magnetostratigraphic evidence, B:826**
- Nanga Parbat-Haramosh region, uplift, dating, B:826**
- Nannofossils. See Calcareous nannofossils. See also individual taxa in Taxonomic Index**
- Narmada-Son lineament**
 extrusion, B:818
 magmatic activity, B:819-820
- Narsarpur Trap, basalt flow, correlation, Deccan Traps, B:818**
- Naturaliste Plateau**
 formation, hot spot origin, A:71
 glauconite
 in ash layers, B:939

- environment of formation, A:465
hot spot origin, B:514
- Nauru Basin
basaltic sills, B:551-552
geochemistry, phenocryst assemblages, B:552
magnetic properties, basalts, B:551
- Nazareth Bank, hot spot origin, relative motion models, B:770
- Neodymium, Ninetyeast Ridge
basement basalts, A:527
ferrobasalts, A:526
lavas, B:600-602
- Neodymium isotopes
Kerguelen Plateau, A:14
lavas, Ninetyeast Ridge vs. Kerguelen Archipelago, A:526
Ninetyeast Ridge, A:14, B:580, 639
basalts, B:568-569, 571-573, 581
lavas, B:593, 596, 600-602
vs. lead isotopes, B:578, 581
vs. strontium isotopes, B:578, 603
- Neodymium/lead isotope ratio, Ninetyeast Ridge, Kerguelen Plateau compared, B:599
- Neogene/Cretaceous unconformity, Site 755, A:237
- Neritic sediment
Broken Ridge, classification, A:41
Ninetyeast Ridge, classification, A:41
- New England seamounts, lavas, geochemistry, A:528-529
- Nickel
Broken Ridge, ash layers, A:474
Ninetyeast Ridge
in sulfides, B:629
vs. iron oxide/magnesium oxide ratio, B:567-568, 571
vs. magnesium oxide, A:402, B:621, 629
vs. zirconium, basalts, B:574
- Nickel/magnesium oxide ratio, Site 757, lavas, B:628
- Nickel/zirconium ratio, Broken Ridge, inverse correlation, A:474
- Ninetyeast Fracture, orientation relative to adjacent zones, B:765
- Ninetyeast Ridge
alteration, water-dominated, B:565
basal sediments
age, B:768
depositional environment, B:560-561
basalt, trace elements, B:572, 608
basalts, B:507-509
age determination, B:510-516
Dupal isotopic signature, B:638-639
modern ocean islands correlated, B:579, 581
basement, age dating, B:447
biostratigraphy, A:43-45
carbonate accumulation rate, B:470, 473, 475, 477
carbonate veins, alteration temperatures, B:453-455
Cretaceous-Paleocene reconstruction, A:525
Eocene angular unconformity
physical properties across, B:256
Zanskar unconformity correlated, B:819
formation, A:260-261
Dupal anomaly and, B:601, 603, 609
geochemical record, B:639-645
hot spot origin, A:71
Indian plate migration and, B:591
models, B:559-560, 578-581, 768-773
paleomagnetic record, B:766-768
ridge jump, B:779
geochemistry, A:56-57, 531-534
- basalts, B:567-575
intersite differences, B:633-638
topographic relationship, B:447
geological setting, A:13, B:467, 526
hot spot origin, B:507, 514, 516, 811
models, A:517, 523-524, B:561-562, 578-581, 591, 763, 765, 768-770, 774
igneous petrology, A:523-531
Indian plate attachment, paleomagnetic evidence, B:766, 768
- lava
comparison, Indian Ocean basalts, B:598-601
Dupal isotopic signature, B:599, 601, 603, 609, 638-639
post-magmatic alteration, B:614, 621
- lead isotopes, B:596-598
lipid geochemistry, B:473
lithology, B:447
lithostratigraphy, A:517-520
Miocene flux increase, A:517
logging, geophysical, A:59-60
low-molecular weight hydrocarbons, B:457-465
magnetic properties, A:50-56
mirror-image ridge, Antarctic plate, B:770, 774
morphology, A:13, B:560
Wharton Basin structure and, B:765-766
neodymium isotopes, B:596
northward motion, seafloor spreading rate correlation, B:810
northward movement, breakpoints, maximum likelihood determination, B:805, 809-810, 870
oceanic andesites, B:572, 577
organic carbon, B:457-465
accumulation rate, B:470, 472, 481
organic matter, type of, B:472-473
origin
compositional variation, A:529-531
geochemical evidence, B:639-642, 644-645
Indian plate movement association, A:523
models, B:559-560, 811
petrologic and geochemical constraints, A:525-529
volcanologic constraints, A:524-525
paleoceanography, A:17-20
paleoclimatology, A:20
paleodepth, Eocene/Oligocene boundary, B:232
paleolatitude, A:20, B:469, 856-867, 941
ash layers, B:561
breaks in movement rate and, B:799-811
Cretaceous/Tertiary boundary, B:802
mean inclinations and, B:799-803, 871
observed vs. expected, B:818, 878
petrography, B:648-655
petrology, B:575-578
physical properties, A:57-59
productivity, B:472
high- vs. low-productivity zones, A:18
proposed future research, A:531
residual magnetic anomaly, B:768-769
seismic reflection profiling, A:93-94, 98-100, 103-104, 106-107
spreading center
northward migration, B:768-769
tectonic and paleomagnetic effects, B:801
- stable isotopes
mass balance calculations, B:452-453
pore waters, B:448-452
topographic relationship, B:447
- strontium isotopes, B:592, 596
surrounding basins, tectonic chart, B:765-766
tectonic chart, B:765-766
tectonic history, A:5, 12, 534-535, B:773
tephra, A:520-523
volcanic facies, A:532
- Ninetyeast Transform Fault, history, A:13, 260-261
- Nintoku Seamount, primary oxide composition, B:539
- Niobium, Ninetyeast Ridge, A:281, 327
basement basalts, A:330, 400, 402, 526
vs. vanadium, A:402
vs. zirconium, A:400
- Niobium/zirconium ratio, Ninetyeast Ridge, basaltic ashes and fragments, A:333
- Noble metals, Site 752, Cretaceous/Tertiary boundary, B:425
- Norwegian Sea, carbonate veins, isotopic values, B:453
- Obsidian pebbles, Site 757, A:311
- Ocean island basalts (OIB), Ninetyeast Ridge, A:531, B:579, 581
magma mixing with MORB, B:592, 637
- Olduvai Subchronozone, Site 758
oxygen isotope stratigraphy, B:306
sedimentation rate, B:302-303
- Olefin, Broken Ridge, toluene content correlation, B:461-462
- Olivine, Ninetyeast Ridge
alteration, A:323, B:563-564
phenocrysts, A:275
replacement, B:577
- Ontong-Java Plateau
basement, age, A:465
calcium carbonate, B:246
carbonate accumulation rate, A:466, B:940
constructional volcanism, B:933
oxygen isotope stratigraphy, B:246
- Ooze, calcareous
Broken Ridge
carbonate content, B:475
diagenesis and compaction of, B:258
physical properties, B:254
Ninetyeast Ridge, A:311
physical properties, A:337
- Ooze, carbonate
Broken Ridge
oligotrophic flux rate, B:943-944
winnowing, A:471, B:229
Ninetyeast Ridge, A:311
- Ooze, clayey nannofossil, Ninetyeast Ridge, A:367
- Ooze, foraminifer-nannofossil, Broken Ridge, A:84, 116, 147, 194
- Ooze, foraminifer-nannofossil
Broken Ridge
compression index, B:258
physical properties, A:185
Ninetyeast Ridge, compression index, B:258
- Ooze, foraminifer, Broken Ridge, A:469, B:211-212
- Ooze, nannofossil
Broken Ridge, A:238-239, 457-458, 469
abyssal reworking, Miocene, A:470
accumulation rate, A:464, B:938
physical properties, A:495
Ninetyeast Ridge, A:265, 360, B:229
dark-light bed contacts, A:367-368
magnetic susceptibility, A:394
sedimentation rate, A:517
seismic reflection profiling, A:293, 295-296

- Pacific Ocean, SE, abyssal reworking, Miocene, A:470
- Ooze, pelagic, Broken Ridge depositional environment, B:212 oxygen isotopes, coarse material and, B:217-218
- Ooze, sandy, Broken Ridge, magnetic susceptibility, A:206
- Ooze-chalk-limestone transition, Ninetyeast Ridge, physical properties, B:256
- Ooze-chalk transition Broken Ridge, B:264 physical properties, B:256 Ninetyeast Ridge, seismic reflection profiling, A:414
- Opal Broken Ridge, A:471, B:262, 265-267, 271 A-to-CT transition, A:497, B:522 accumulation rate, A:465, B:939 first occurrence, B:262 flux rate, Cretaceous/Tertiary boundary, A:468, B:941 lepispheres, B:272 physical properties, B:256 quartz transition, A:498, B:266, 268, 521-522 source, B:268 Ninetyeast Ridge, quartz transition, B:257
- Osborne Knoll bathymetry, B:560 ridge jump, B:765 width, A:13
- Oxidation exsolution, Ninetyeast Ridge cooling rate, B:541 magnetic effects, B:535 temperature of, A:323, B:535, 539 titanomagnetite, B:535
- Oxides Broken Ridge basalts, B:621 logging data, B:900-901, 904-905 Ninetyeast Ridge glass shards, B:277-283 high- vs. low-temperature alteration, A:333 morphology, cooling rate and, B:542 pigment, B:542 tephra layers, B:277, 286, 288-291
- Oxides, secondary, Ninetyeast Ridge, low-temperature alteration, B:541-542
- Oxygen index, Broken Ridge, vs. hydrogen index, A:219, 495
- Oxygen isotope stratigraphy, Site 758, B:245, 247-248, 304-307 planktonic vs. benthic foraminifers, B:285 Site 607 correlated, B:305-307 Site 677 correlated, B:304
- Oxygen isotopes Broken Ridge, B:231, 449 eolian dust flux and, B:221 grain size and, B:214-215 Maestrichtian-Eocene, B:885-886, 888 mass balance calculations, B:452-453 Miocene cooling trend, A:211, B:214, 232, 943 pelagic oozes, B:217, 237-238 pore waters, B:448-452 reaction zone thickness, B:453 sea-level changes and, B:943 sedimentation rate correlation, B:226 Ninetyeast Ridge, B:231-234, 236, 329-339, 449 benthic vs. planktonic records, B:247-248 carbonate veins, B:453-455 comparison, SPECMAP global record, B:303-305 composite depth record, B:306-308, 310-311, 340-355 Eocene/Oligocene boundary, B:232 glacial/interglacial cycles, B:304, 452 mass balance calculations, B:452-453 mid-Pleistocene shift, B:313 Miocene cooling trend, B:232, 247 Neogene benthic foraminifer values, B:247 pelagic oozes, B:231-232, 239-240 planktonic foraminifers, B:246, 250 pore waters, B:448-451, 453 pore waters, sinusoidal pattern, B:452 reaction zone thickness, B:453 sedimentation rate correlation, B:227 tephra layers, B:285 polar cooling and, B:211 Site 752 Maestrichtian-Eocene, B:885-889 pelagic oozes, B:231-232 Site 754, pelagic oozes, B:231-232 standard sediment sample, B:892
- P*- wave velocity Broken Ridge, A:57-58, 151, 217, 252 Ninetyeast Ridge, A:57-58, 289, 340, 347, 404, 430-431
- Pacific mud, replicate analyses, B:425
- Pacific Ocean, geomagnetic polarity transition, Jaramillo Subchronozone, B:385
- Pacific Ocean, equatorial, oxygen isotope stratigraphy, B:247
- Pacific Ocean S oceanic island basalts, isotopic ratios, B:592 sea-surface productivity, atmospheric circulation and, A:20 winnowing, pelagic oozes, B:212
- Pacific Ocean SE, eolian dust, Cenozoic accumulation rate, B:220
- Pacific Ocean W, ocean plateaus, tholeiitic, A:526
- Panama Basin Project, carbonate pelagic sediments, winnowing, B:212
- Pebble layer, Broken Ridge, A:462, B:937 eustatic sea level and deposition of, A:469 physical properties, A:495
- Pelagic cap. *See* Broken Ridge, pelagic cap
- Peru-Chile Trench, stress, direction of, B:697
- Phillipsite Ninetyeast Ridge, B:564 ambient alteration, B:565 Site 752, B:271-272
- Phosphorus/neodymium ratio, Ninetyeast Ridge, intersite differences, B:637
- Phosphorus/titanium ratio, Ninetyeast Ridge, basalts, B:569
- Phosphorus/zirconium ratio Ninetyeast Ridge, basalts, B:569-570, 577 Site 756, basalts, B:568
- Physical properties Broken Ridge, A:57-59 compaction and changes in, B:253-254, 256-258 dipping and truncated sequence, A:497-498 grain size correlation, B:256 interhole correlation, A:505 opal-CT-to-quartz transition, B:268 pelagic cap, A:495-497, B:254, 264 with silica mineralogy changes, B:521-522 Ninetyeast Ridge, A:57-59 compaction and changes in, B:253-254, 256-258 laboratory vs. log measurements, B:257 opal-CT-to quartz transition, B:257 units A-F, B:256 seismic correlation, Site 756, A:303 Site 752, A:138-150 across angular unconformity, B:264 lithification effects, B:267-268 lithologic correlation, A:147 lithostratigraphic correlation, A:139 mineralogical and chemical correlation, B:265 units A-E, A: 139, 141, 147-148, B:264-266 Site 753, A:181-185 Site 754, A:191, 216-219, 222, 227 logging correlation, A:221 units A-C, A:216, 219 Site 755, A:250-253 Site 756, A:287-292 seismic correlation, A:294-295, 303 units A-C, A:287-288, 291 Unit B, A:291 Site 757, A:337-347 units A-C, A:337, 339, 343-344 Site 758, A:402-408, 425-426, 429-431, B:704 seismic correlation, A:419 units A-D, A:402-404 units A-E, Site 752, B:264-266
- Phytoliths, Site 758, B:192
- Pillow basalts, Site 758, A:391 mineralogy, A:392-393
- Plagioclase Ninetyeast Ridge, A:520, B:656-657 accumulation, B:575-576, 630-631 phenocrysts, A:275 replacement, B:563 vs. modal plagioclase and aluminum oxide, B:580 Site 757 groundmass phases, A:323 major oxide content and, A:327, 329 megacrysts, B:625 mobility, B:625-627 phenocrysts, A:321-322, B:567, 569 segregation, B:628 vs. strontium content, B:627 Site 758, A:393 accumulation, B:577 ash layers, B:277 fractionation, B:287
- Planktonic foraminifers. *See also under particular sites and in Taxonomic Index*
- Broken Ridge biostratigraphy, B:85-89 Cretaceous/Tertiary boundary, A:484 Eocene-Oligocene disconformity, A:476-477 hardground assemblage, A:477 mid-latitude temperature species, A:484 planktonic/benthic ratios, A:477-478, 484 quantitative analysis, B:79-82 Southern Hemisphere high-latitude affinity, A:483 Southern Ocean high-latitude affinity, A:476 Turonian-Santonian, A:485 zonation, A:43, 46-47 Indian Ocean, foraminifer lysocline, B:127 Indo-Pacific species, spatial distribution, B:138 Ninetyeast Ridge biostratigraphy, B:89-907 quantitative analysis, B:82-83 zonation, A:43, 46-47

- Site 709, isotope stratigraphy, B:242–244
- Site 752, A:128–129, B:93, 722
ash layers, A:510
carbonate content and productivity of, A:510, B:79, 81–82
Cretaceous distribution, B:92
Cretaceous/Tertiary boundary, A:128, B:13
Eocene–Oligocene distribution, B:89
low-productivity interval, A:509
mid-latitude temperate faunas, A:128
Miocene/Pliocene boundary, A:128
nannofossil age correlation, A:121
Paleocene/Eocene boundary, B:727
Paleocene–Eocene distribution, B:90–91
planktonic/benthic ratio, A:128, B:16
sedimentation rate correlation, A:509
zonation, A:124–125
- Site 753, A:177–178, B:94
carbonate content and productivity of, B:81, 83
datum levels, A:177
Eocene–Oligocene pelagic environment, A:178
mid-latitude temperate faunas, A:177
Miocene/Pliocene boundary, A:178
planktonic/benthic ratio, B:81
- Site 754, A:201–204, B:97
Campanian/Maestrichtian boundary, B:89
carbonate content and productivity of, B:81, 84
Cretaceous distribution, B:96
depositional environment, A:205
Miocene/Oligocene boundary, A:203
Miocene/Pliocene boundary, A:202
Paleogene distribution, B:95
planktonic/benthic ratio, B:81
Pliocene/Pleistocene boundary, A:201
preservation, A:204
Southern Hemisphere high latitude affinity, A:201
Southern Ocean high latitude affinity, A:195
- Site 755, A:246, B:98
carbonate content and productivity of, B:81–82, 85
planktonic/benthic ratio, A:246, B:81–82
Turonian–Coniacian, B:89
- Site 756, A:271–272, B:99–100
Broken Ridge fauna compared, A:270
carbonate content and productivity of, B:82–83, 86
Eocene/Oligocene boundary, A:271
Oligocene/Miocene boundary, A:271
planktonic/benthic ratio, B:38
preservation, A:271
- Site 757, A:318–320, B:102
carbonate content and productivity of, B:83, 87
datum levels, A:316
Eocene–Oligocene distribution, B:100–101
mid-latitude temperate faunas, A:318
Miocene/Pliocene boundary, A:319
planktonic/benthic ratio, B:83
zonation, A:318
- Site 758, A:382–384, B:108–109, 128–131
abundance, B:126–127
ash layer, B:489
biological productivity influence, B:134, 136
Brunhes dissolution cycle, B:133
carbonate content and productivity of, B:84–85, 88, 134, 491
coarse fraction association, B:130
Cretaceous distribution, B:106
Cretaceous/Tertiary boundary, A:376, 384
datum levels, A:382
dissolution, B:130–133
ecological factors influencing, B:133–137
Eocene–Oligocene disconformity, A:376
glacial/interglacial cycles, B:130
glauconite-filled, A:383
isotope stratigraphy, B:245
Maestrichtian–Campanian, A:384, B:97
mid-latitude to tropical-subtropical transition, A:376, 384
Miocene/Pliocene boundary, A:382
Oligocene distribution, B:103
Oligocene/Miocene boundary, A:382
Paleocene–Eocene distribution, B:104–105
paleoenvironmental implications, A:375–376
planktonic/benthic ratio, A:383–384, B:84–85
preservation, A:375
relative abundance, B:126–127
resistance species ratio, B:127, 130
salinity gradient effects, B:133–134
Santonian age interval, A:384
sea-surface temperature control, B:133, 137
shallow- vs. deep-dwelling species, B:246
strontium isotopes, B:199
temporal variation, B:127–130
Toba Lake eruption, B:489, 492
winnowing effects, B:84
zonation, A:376
site comparisons, A:483–484
- Porcellanite
Broken Ridge, A:239–240, 460, B:267
Cretaceous/Tertiary boundary, A:507
seismic reflection profiling, A:79, 156
Ninetyeast Ridge, chemical gradient effect, A:399, 410
- Porosity
Broken Ridge, A:182–183, 217, 250–252
carbonate content and, B:256
compaction effects, B:258
magnetic properties and, A:484, B:371–372, 383
vs. applied vertical load, B:258
Ninetyeast Ridge, A:287–288, 292, 337, 402–404
basalt, A:297
carbonate content and, B:256
vs. applied vertical load, B:258
Site 752, A:139, 144
calcareous chalks, B:257
formation factor and, A:143
pelagic cap, B:254, 258
vs. formation factor, A:150
Site 758, stress effect, B:702
- Potassium
Ninetyeast Ridge
alteration effects, B:563
in non-oxidized basalts, B:565
Site 756, basalt, A:277
- Potassium oxide
Ninetyeast Ridge, mobility during alteration, B:615
Site 758, incompatible element correlation, B:629
- Potassium oxide/phosphate ratio, Ninetyeast Ridge, lavas, B:624
- Potassium/rubidium ratio, Ninetyeast Ridge, B:624–625, 629
- Potwar Basin, Siwaliks sedimentation, source uplift and rate of, B:823, 825
- Potwar Basin/Salt Range region, uplift, dating, B:826
- Pristane/phytane ratio, B:473, 483
- Productivity, biogenic. *See* Broken Ridge, productivity
- Pumice blebs, Site 758, A:368
- Pumice shards, Broken Ridge, A:472–473
- Pyrite, Site 758, B:629
- Pyrite, framboidal, Site 755, A:496
- Pyroxene, Ninetyeast Ridge, B:567
basalt phenocrysts, A:323
fractionation, B:576–577
- Quartz
Broken Ridge, B:521
Site 752, Cretaceous/Tertiary boundary, B:916
Site 758
ash layers, B:277
fractionation, B:287
- Quartz, authigenic, Broken Ridge
chemical environment for, B:267–268
source, B:268
- Quartz/opal-CT ratio, Broken Ridge, B:267–268
- Quench structures, Ninetyeast Ridge, A:523
basalts, A:392
pillow basalts, A:391
- Radiolarians. *See also under individual taxa in Taxonomic Index*
Site 752, B:722
Site 758, A:382, 387
- Rajmahal continental flood basalts, hot spot origin, A:71
- Rajmahal Traps
hot spot origin, B:507, 514, 516
paleolatitude, B:802
hot spot location and, B:371
- Rare earth elements
Broken Ridge, ash layers, A:474
Ninetyeast Ridge
basalts, B:572, 574–575
chondrite-normalized, B:576
depleted source component, B:640
fractional crystallization effects, B:624
intersite differences, B:637
Kerguelen Archipelago lavas compared, B:639, 641
- Site 752
Cretaceous/Tertiary boundary, B:425–426, 430, 501
Santonian ash, B:441, 443
- Site 756
basalts, B:568
chondrite-normalized, B:588, 621–622, 634–638
- Site 757
basalts, B:570
chondrite-normalized, B:589
- Site 758
basalts, B:571, 629–631
chondrite-normalized, B:590
Kerguelen Archipelago lavas compared, B:642
- Raton Basin (Colorado), kaolinitic spherules, Cretaceous/Paleogene boundary, B:490
- Rawalpindi Group, age, A:16
- Resistivity
Broken Ridge
chalk-limestone transition, A:221
chert layer, A:149
Site 754, A:230
- Réunion hot spot
predicted path, B:514, 516
vs. present location, B:812, 818–819

- relative motion model, B:770
- Réunion Island
hot spot origin, relative motion models, B:770
volcanoes, tholeiitic basalt, A:526
- Réunion Subchronozone, Ninetyeast Ridge, B:797-798
- Rifting
Broken Ridge, A:5, B:437-438
active, A:7-8
duration, A:10, 172, B:732
geochemical evidence, B:444
history, A:7, 12
magnetic properties and, A:491
paleodepths, B:733, 747
passive, A:8
water depths, A:86
Broken Ridge/Kerguelen Heard Plateau, flexural modeling, A:90
Broken Ridge-Kerguelen Plateau, duration, A:500
China Sea S, A:17
- Rock-Eval pyrolysis
Broken Ridge, A:143, 212-213, 217, 251, B:472-473
Ninetyeast Ridge, A:344, 400, 424, B:472-473
- Rubidium, Ninetyeast Ridge
alteration effects, B:563
basalt, A:277
incompatible element correlation, B:624, 629
lavas, B:600-602
mobility during alteration, B:615
in non-oxidized basalts, B:565
vs. rubidium/zirconium ratio, basalts, B:564
vs. zirconium, basalts, B:564
- Rubidium/zirconium ratio, Ninetyeast Ridge, vs. rubidium, basalts, B:564
- Salinity
Ninetyeast Ridge, A:335
isotopic effects of, B:304
Site 757, A:335
- Samarium, Ninetyeast Ridge, lavas, B:600-602
- Samarium/lanthanum ratio, Ninetyeast Ridge, vs. hafnium/lanthanum ratio, B:644
- Samarium/neodymium ratio
Ninetyeast Ridge, B:639
Site 758, vs. Kerguelen lava ratios, B:642
- Sand
Broken Ridge, Eocene, A:71
Site 754, limestone-chert pebbles in, B:936-937
- Sand, detrital, Broken Ridge, A:5
- Sand, glauconitic, Broken Ridge, B:747
Cretaceous, A:89
- Sandstone, foraminifer, Site 755, B:937
- Sandstone, volcanic, Site 758, A:374
- Saponite, Ninetyeast Ridge, formation environment, B:624
- Scandium, Site 758, basalts, B:571
- Scandium/zirconium ratio, Site 756, basalts, B:568
- Sea-surface circulation, Broken Ridge, ice buildup and intensification of, 9:215
- Sea-surface temperature, Ninetyeast Ridge, warming trends, isotopic effects of, B:247
- Seafloor spreading
Broken Ridge, B:934
Ninetyeast Ridge, magnetic anomalies and rate of, B:810
- Seamounts, basalt, magnetic properties, B:525-526
- Sediment/basalt contact
Site 756, physical properties, A:287-288
Site 758, seismic reflection profiling, A:414
- Sedimentation rate
Broken Ridge, A:464, 466-467, B:737, 934, 936, 938
Cretaceous, B:735, 738
Paleocene-Eocene, B:740
- Ninetyeast Ridge
flux rates, A:519-520
intersite comparison, A:517
Site 752, A:112, 134-135, 138, B:223, 738, 938
magnetostratigraphic correlation, A:138
Paleocene-Eocene, B:740
Site 753, B:739, 938
Site 754, B:224, 938
magnetostratigraphic correlation, A:210, 212
nannofossil vs. foraminiferal disparity, B:738
Site 755, B:938
Site 756, B:224
grain size and, A:265
Site 757, A:311
Site 758, B:242, 302-304
comparison, calcium carbonate MAR, B:303
correlation, susceptibility breaks, B:822-823
impact of Himalayan-Tibetan uplift on, A:363
intervals 1-4, B:302
Jaramillo Subchronozone, B:378
low-rate intervals, B:302
magnetic susceptibility and, A:394
at reversal boundaries, B:302-303
uplift impact, A:20
vs. Site 607, B:306
- Seismic reflection profiling
basalt
Site 756, A:295-296
Site 758, A:421-422
basaltic flow units, Site 758, A:421-422
Broken Ridge, A:9, 33, 63-68, 459, B:262, 469, 673-677, 682-683, 723, 748-752, 755-759, 934
ash, B:744
dipping and truncated sequence, Broken Ridge, A:153, 156, 226, 254-255, B:743-744
Eocene angular unconformity, Broken Ridge, A:186, 254-255
Ganges-Brahmaputra river system, B:825
limestone
Site 754, A:226
Site 756, A:293
limestone-ooze transition, Broken Ridge, B:682
Miocene hiatus, Site 753, A:186
Ninetyeast Ridge, A:33, 93-94, 98-100, 103-104, 106-107
ooze, nannofossil, Ninetyeast Ridge, A:293, 295-296
ooze-chalk transition, Ninetyeast Ridge, A:414
pelagic cap, Broken Ridge, A:153, 185-186, 225-226, 233, B:747
porcellanite, Broken Ridge, A:79, 156
sediment/basalt contact, Site 758, A:414
Site 752, A:114
Site 753, A:185-186, 189
Site 754, A:194
Site 755, A:240, 256-257
Site 756, A:300-302
Site 757, A:354-356, 358
Site 758, A:444-447
- Seismic stratigraphy
Broken Ridge
composite signal, B:665-668
composite signal, stacked, B:672
composite signal, synthetic, B:671
deconvolution, B:667-669
distal variation, A:78
downlapping sequence, B:751
Eocene angular unconformity, A:10
intersite correlation, A:86
lithostratigraphic correlation, A:165, 188, 235, 257, B:668-669, 745
magnetic anomaly 18, A:7-8
onlapping sequence I, A:81, 83, 90
onlapping sequence II, A:84
porcellanite-chert-chalk-limestone transition, B:268
pre-rift/post-rift boundary, A:10-11
primary signal, B:665, 667-670
prograding downlapping wedge, A:75, 79, 81-82
sequence I, B:743-744, 758-759
sequence I/II boundary, B:747, 752
sequence II, B:744, 746-747, 758-759
sequence III, B:747
source-receiver geometry, B:663-665
stratigraphic sequences, A:76
units 1-4, A:5
weakly laminated unit, A:75, 81
deconvolving water-gun data, B:665-668
dipping and truncated sequence, A:72-75, 153, 156, 226, 255, B:751
lithostratigraphic correlation
Site 752, A:151, 165, 188, 235, 257, B:398, 668-669, 745
Site 753, A:165, 185, 188, 235, 257, B:398, 668-669, 745
Site 754, A:165, 188, 224-226, 235, 257, B:398, 668-669, 745
Site 755, A:165, 188, 235, 254-255, 257, B:398, 668-669, 745
Site 756, A:293-294
Site 757, A:344, 347-350, 357
Site 758, A:414, 418-419
pelagic cap, A:84-86, 152, 185-186, 225-226, 254, B:751
physical properties and, Site 756, A:303
reflectors 1-10, Site 758, A:417-419
Site 255, correlation, Site 755, A:252, 254
Site 752, A:151-156, 166
comparison, synthetic seismogram, A:162
with increase in siliceous component, A:153
lithologic correlation, A:79
synthetic seismogram, A:75, 79, 151, 153
Site 753, A:185-187
Site 754, A:224-226, 236
deconvolution, B:678
low-velocity zone, A:221, 225
synthetic seismogram, A:79, 81, 221, 225, 233
Site 755, A:252, 254-255
correlation, Site 255, A:252, 254
reflection profiling, A:240, 256
Site 756, A:291-296, 299-302
physical properties correlation, A:303
units 1A-ID, A:295-296
Site 757, A:344-345, 347-350
reflectors 1-10, A:345, 348-349
site survey, A:344-345
Site 758, A:414, 416-423
impedance and lithologic correlation, A:448
logging data, A:366-367

- physical properties correlation, A:418-119
reflection profiling, A:93-94, 106-107, 443-447
reflectors 1-10, A:417
synthetic seismogram
Site 752, A:151, 153, 161-162
Site 754, A:225, 232-233
- Selenium, Site 752, Cretaceous/Tertiary boundary, B:420
- Shatsky Rise
carbonate accumulation rate, A:466, B:940
following Cretaceous/Tertiary boundary event, B:427
- Shear strength
Broken Ridge, A:58, 143, 148-149, 183, 219-221, 227, 252, 495
Ninetyeast Ridge, A:58, 288, 293-294, 340, 342, 344, 350, 404, 435
- Shell hash, Ninetyeast Ridge, A:521, 523
- Shikoku Basin, basalt, B:550
- Silica
Broken Ridge
biogenic vs. nonbiogenic, B:264, 267
carbonate content and production of, B:475
deposition, Campanian-Maestrichtian, A:467
diagenesis, A:498, 500, B:265-266, 521
mineralogy, B:261-262, 521-522
physical properties, B:521-522
in pore water, B:266
productivity, A:464, B:941
source of, B:262
X-ray diffraction, B:264
X-ray mineralogy, B:261-262, 264
Ninetyeast Ridge
lavas, B:615, 629
vs. iron oxide/magnesium oxide ratio, A:279
Site 752
physical properties effect, A:139
in pore water, B:270
Site 758
Cretaceous, vs. Broken Ridge sediments, A:402
eruption explosivity and content of, B:287
major oxide correlation, tephra layers, B:277
vs. iron oxide/magnesium oxide ratio, A:400
- Silica, authigenic, Broken Ridge, B:264, 267
pre-rift vs. post-rift burial depth, B:261, 266
- Silica, biogenic
Broken Ridge
porosity and consolidation of, B:264-265
preservation and degree of, B:267
source, B:261
Site 758, Cretaceous vs. Paleogene sections, B:477
- Silica diagenesis
Broken Ridge
chemical environment for, B:267
lithification effects, B:261-269
timing, B:266
Site 752, B:266
- Silicates
Broken Ridge, A:42
Ninetyeast Ridge, A:42
- Siliceous microfossils, Broken Ridge
dissolution, B:264
preservation, B:262-264
- Silicoflagellates, Site 758, B:192-193
- Sill, basaltic, Site 758, B:549
- Silt, glauconitic, Broken Ridge, B:747
Cretaceous, A:89
- Site 31, basaltic microlapilli, Miocene-Oligocene, B:491
- Site 77
benthic foraminifers, oxygen isotopes, Oligocene boundary event, B:51
carbon isotope stratigraphy
enriched Miocene interval, B:250
Oligocene/Miocene boundary, B:249
- Site 94, diatoms, Eocene, A:129, B:171, 173
- Site 208, diatoms
Eocene, A:129
Paleocene, B:171, 173, 175
- Site 211
magnetic anomalies, A:24
oceanic island basalts, incompatible element ratios, B:639
- Site 212
magnetic anomalies, A:24
paleolatitude, B:469
- Site 213
brown clay, B:467
hot spots, models, B:771
magnetic anomalies, A:24
paleolatitude, B:469
planktonic foraminifers, Paleocene-Eocene, B:77
ridge formation, paleomagnetic record, B:766-768
siliceous ooze, carbonate-poor, B:467
- Site 214
alteration, post-magmatic, B:563-564
basal sediments, depositional environment, B:560-561
basalts
geochemistry, A:525-530, B:561-562, 565, 572, 574-582, 592
iron oxide/magnesium oxide ratio, B:526
isotopic values, B:572-573, 578-579
subaerial eruption, A:524
basement, A:13
age, B:561
lithology, B:562
bathymetry, B:764
diatoms
Eocene, A:129
Paleocene, B:171, 173
ferrobasalts, A:13
geochemistry, eruption age and, B:633
hot spots
Cretaceous-Oligocene locations, B:769
migration, B:776
models, B:771-773
incompatible elements, basalts, B:577, 580
lavas
evolved, A:14
incompatible element-enriched, A:530
Site 756 comparison, A:277
magnetic anomalies, A:24
magnetic properties, inclination, B:869
neodymium isotopes, vs. strontium isotopes, A:14
oceanic andesite, B:572
titanomagnetite fractionation, B:577
oxygen isotope stratigraphy, B:247
paleolatitude, B:469
petrography, basalts, A:275
planktonic foraminifers, Paleocene-Eocene, B:77
rare earth elements, basalts, B:576
ridge formation, paleomagnetic record, B:766-768
volcanic ash, B:633
- Site 215
bathymetry, B:764
- hot spots, Cretaceous-Oligocene locations, B:769
magnetic anomalies, A:24
magnetic properties, inclination, B:869
oceanic island basalts, incompatible element ratios, B:639
planktonic foraminifers, Paleocene-Eocene, B:77
ridge formation, paleomagnetic record, B:766-768
ridge jump, B:765
- Site 216
alteration, post-magmatic, B:563-564
basalts
geochemistry, A:525-530, B:561-562, 565, 572, 574-582, 592
iron oxide/magnesium oxide ratio, B:526
isotopic values, B:572-574, 578-579
petrography, A:275
basement, A:13
age, B:561
lithology, B:562
bathymetry, B:764
diatoms, Cretaceous, B:189
ferrobasalts, A:13
hot spots
Cretaceous-Oligocene locations, B:769
migration, B:776
models, B:771-773
incompatible elements, basalts, B:577, 580
lavas
comparison, Site 756, A:277
evolved, A:14
incompatible element-enriched, A:530
lead isotopes, B:603
magnetic anomalies, A:24
magnetic properties, inclination, B:869
neodymium isotopes, vs. strontium isotopes, A:14
oxygen isotope stratigraphy, B:247
paleolatitude, B:469, 818
paleomotion, A:18
planktonic foraminifers, Paleocene-Eocene, B:77
Planulina wuellerstorfi assemblage, oxygen isotopic correlation, B:51
rare earth elements, basalts, B:576
ridge activity, earliest record of, B:591
ridge formation, paleomagnetic record, B:766-768
strontium isotopes, B:603
- Site 217
bathymetry, B:764
hot spots
Cretaceous-Oligocene locations, B:769
migration, B:776
models, B:771-773
magnetic anomalies, A:24
magnetic properties, inclination, B:869
organic carbon, B:467
paleolatitude, B:469
planktonic foraminifers, Paleocene-Eocene, B:77
ridge formation, paleomagnetic record, B:766-768
- Site 219, magnetic properties, inclination, B:869
Site 220, magnetic properties, inclination, B:869
Site 221, magnetic properties, inclination, B:869
Site 222, magnetic properties, inclination, B:869
Site 237, oxygen isotope stratigraphy, B:247
Site 238
magnetic properties, inclination, B:869
oxygen isotope stratigraphy, B:247

Site 253

- alteration, post-magmatic, B:564
 basal sediments, ages, B:768
 basalts
 geochemistry, A:525–530, B:561–562, 565, 575–582
 isotopic values, B:572–573, 578–579
 lead isotopes, B:570–571
 basement, A:13
 age, B:561
 lithology, B:562
 bathymetry, B:764
 calcium, Site 757 correlated, B:448
 hot spots
 Cretaceous–Oligocene locations, B:769
 migration, B:776
 models, B:771–773
 incompatible elements, basalts, B:577, 580
 lavas, incompatible element-enriched, A:530
 magnesium, correlation, Site 757, B:448
 magnetic anomalies, A:24
 magnetic properties, B:810
 inclination, B:869
 oxygen isotopes, correlation, Site 757, B:448
 paleolatitude, B:469
 rare earth elements, basalts, B:576
 seismic stratigraphy, A:291
 volcanic ash, B:633
 volcanogenic sediment, inter-element ratios, B:425
- Site 254
 alteration, post-magmatic, B:564
 basal sediments, ages, B:768
 basaltic flow units, submarine eruption, A:524
 basalts
 geochemistry, A:525–530, B:561–562, 565, 574–582, 592
 isotopic values, B:572–573, 575, 578–579
 basement, A:13
 age, B:561
 lithology, B:562
 bathymetry, B:764
 geochemistry, basalts, B:575
 hot spots
 Cretaceous–Oligocene locations, B:769
 migration, B:776
 models, B:771–773
 igneous petrology, geochemistry, A:279
 incompatible elements, basalts, B:577, 580
 lavas
 comparison, Site 756, A:277
 incompatible element-enriched, A:530
 magnetic anomalies, A:24
 magnetic properties, inclination, B:869
 rare earth elements, basalts, B:576
 seismic stratigraphy, A:291
 volcanoes, age of, B:467
- Site 255
 angular unconformity, A:71
 basal sediments, B:507
 bathymetry, A:72, 173, 193, 239, B:438, 744, 764, 934
 biomicritic limestones, correlation, Broken Ridge, A:72
 extension models, A:12
 limestone-chert sequence, A:5, 112
 lithostratigraphy, A:75
 magnetic anomalies, A:24
 pelagic cap, A:5
 pelagic ooze, winnowing, B:212
 planktonic foraminifers, B:78
 Santonian–Eocene hiatus, A:71
 seismic stratigraphy
 correlation, Site 755, A:252, 254

- reflection profiling, A:74
 Sequence I, B:744
 volcanism, Cretaceous, A:465, B:939
- Site 256
 lavas, incompatible element-enriched, A:530
 magnetic anomalies, A:24
 oceanic island basalts, incompatible element ratios, B:639
 paleolatitude, B:469
 volcanism, Cretaceous, A:465, B:939
- Site 257
 alteration mineral paragenesis, A:275
 magnetic anomalies, A:24
 volcanism, Cretaceous, A:465, B:939
- Site 258
 magnetic anomalies, A:24
 volcanism, Cretaceous, A:465, B:939
- Site 259, magnetic anomalies, A:24
- Site 260, magnetic anomalies, A:24
- Site 261, lead isotopes, B:604
- Site 263, magnetic anomalies, A:24
- Site 275, diatoms, Cretaceous, B:189
- Site 289, oxygen isotope stratigraphy, B:246–247
- Site 327, diatoms
 Eocene, A:129
 Paleocene–Eocene, B:173
 zonation, Paleocene, B:175
- Site 339, diatoms, Eocene, A:129, B:171
- Site 343, diatoms, Eocene, A:129, B:171
- Site 348, carbonate veins, isotope values, B:453
- Site 356, *Micula decussata*, Cretaceous/Tertiary boundary, B:406
- Site 366
 benthic foraminifers, oxygen isotopes, Oligocene boundary event, B:50–51
 carbon isotope stratigraphy
 enriched Miocene interval, B:250
 Oligocene/Miocene boundary, B:249
- Site 367, basaltic flow units, age of, B:549–550
- Site 368
 basaltic flow units, age of, B:549–550
 calcareous shales, contact metamorphic effects, B:550
 methane/ethane ratio, with distance from sill, B:550
- Site 384, diatoms
 Eocene, A:129
 Paleocene, B:171, 173
- Site 390, diatoms, Eocene, B:171, 173
- Site 395, carbonate veins, isotope values, B:453–454
- Site 398, diatoms
 Eocene, A:129
 Paleocene, B:171, 173
- Site 417, carbonate veins, isotope values, B:453–454
- Site 442, basaltic flow units, B:550
- Site 443, basaltic flow units, B:550
- Site 444, basaltic flow units, B:550
- Site 446
 basaltic flow units
 magnetic properties, B:550–551
 petrography, B:550–551
 basaltic sill, age of, B:550
- Site 462, basaltic sill, B:551–552
- Site 477
 alteration, low-temperature, B:553–554
 basaltic flow units
 alteration, B:553–554
 regional relationships, B:552–553
 basaltic sill, thermal metamorphism, B:555
 basalts, geochemistry, B:554
 interstitial-water chemistry, B:554
 mineralogy, formation temperature, B:553

- organic geochemistry, B:554
 oxygen isotopes, B:554–555
 porosity, vertical variation, B:553
- Site 478
 alteration, low-temperature, B:553–554
 basaltic flow units
 alteration, B:553–554
 regional relationships, B:552–553
 basaltic sill, thermal metamorphism, B:555
 basalts, geochemistry, B:554
 interstitial-water chemistry, B:554
 mineralogy, formation temperature, B:553
 organic geochemistry, B:554
 oxygen isotopes, B:554–555
 porosity, vertical variation, B:553
- Site 479
 basalts, geochemistry, B:554
 organic geochemistry, B:554
- Site 481
 alteration, low-temperature, B:553–554
 basaltic flow units
 alteration, B:553–554
 regional relationships, B:552–553
 basaltic sill, thermal metamorphism, B:555
 basalts, geochemistry, B:554
 interstitial-water chemistry, B:554
 mineralogy, formation temperature, B:553
 organic geochemistry, B:554
 oxygen isotopes, B:554–555
 porosity, vertical variation, B:553
- Site 501
 calcium, sediment/water interface, A:533
 stable isotopes, pore waters, B:447
- Site 504
 calcium, sediment/water interface, A:533
 magnetic mineralogy, transition zone and upper dike complex, B:539
 ocean-floor basalt, mineralogy, B:56
 stable isotopes, pore waters, B:447
 stress, borehole breakouts, B:697
 titanomagnetites, Curie temperature, B:535
- Site 516, *Rocella princeps*, last occurrence, B:199
- Site 522
 carbon isotope stratigraphy,
 Oligocene/Miocene boundary, B:249
 strontium isotopes, B:921
- Site 524
 calcareous nannofossils, Subzone CP1a, B:408
 diatoms
 Eocene, A:129
 Paleocene, B:171, 173
 Thoracosphaera, Cretaceous/Tertiary boundary, B:406
- Site 558
 benthic foraminifers
 Oligocene faunal change, B:51
 oxygen isotopes, Oligocene boundary event, B:50–51
 carbon isotope stratigraphy, enriched Miocene interval, B:250
- Site 563
 benthic foraminifers, Oligocene faunal change, B:51
 carbon isotope stratigraphy
 enriched Miocene interval, B:250
 Oligocene/Miocene boundary, B:249
Planulina wuellerstorfi assemblage, oxygen isotopic correlation, B:51
- Site 566, carbon isotope stratigraphy, Oligocene/Miocene boundary, B:249
- Site 574
 benthic foraminifers, oxygen isotopes,
 Oligocene boundary event, B:51
 calcium carbonate, B:246

- carbon isotope stratigraphy,
Oligocene/Miocene boundary,
B:248-249
- oxygen isotope stratigraphy, B:246-247
- Site 575
benthic foraminifers, Oligocene faunal
change, B:51
- calcium carbonate, B:246
- carbon isotope stratigraphy,
Oligocene/Miocene boundary, B:249
- oxygen isotope stratigraphy, B:246-247
- Site 577
benthic foraminifers, isotopic records, B:19
- calcareous nannofossils, Subzone CP1a, B:408
- Stensioina beccariiformis* assemblage,
Paleocene decrease, B:19
- Site 586, light-hydrocarbon yields, B:461
- Site 597, ocean-floor basalt, mineralogy, B:56
- Site 603, ferruginous smectite spherules, B:491
- Site 607
oxygen isotope stratigraphy, B:303
- comparison, Site 758, B:304
- magnetostratigraphic correlation, B:306-307
- Spectral Mapping, B:285
- sedimentation rate, comparison, Site 758,
B:306
- Site 608
benthic foraminifers, Oligocene faunal
change, B:51
- Planulina wuellerstorfi* assemblage, oxygen
isotopic correlation, B:51
- strontium isotopes, B:921
- Site 609, magnetostratigraphy, Jaramillo polarity
transition, B:385-388
- Site 610,
benthic foraminifers, Oligocene faunal
change, B:51
- Planulina wuellerstorfi* assemblage, oxygen
isotopic correlation, B:51
- Site 665
magnetic properties, B:378
- demagnetization, B:379-380, 382, 386
- directional changes, B:382-383
- transition field, B:383-385
- magnetostratigraphy, B:377-378
- Jaramillo polarity transition, B:390
- mineralogy, grain size and, B:379
- paleolatitude, magnetic inclination variation
with, B:385
- Site 666, carbon isotope stratigraphy, enriched
Miocene interval, B:250
- Site 677, oxygen isotope stratigraphy, B:285, 303
- Site 758 compared, B:304
- Site 689
Cretaceous/Paleogene boundary, B:492
- epifaunal species, Cretaceous/Tertiary
boundary, B:19
- Epistominella umbonifera*, B:51
- Site 690
Biantholithus sparsus, B:407
- calcareous nannofossils
- Subzone CP1a, B:408
- survivor forms, B:407
- clay, Cretaceous/Tertiary boundary, B:415
- Cretaceous/Paleogene boundary, B:492
- Cruciplacolithus* lineage, B:407
- Discoaster mohleri*, B:728
- epifaunal species, Cretaceous/Tertiary
boundary, B:19
- Epistominella umbonifera*, B:51
- Hornibrookina*, Cretaceous bloom, B:407
- nannofossil chalk, Cretaceous/Tertiary
boundary, B:395
- Prediscosphaera stoveri*, Cretaceous/Tertiary
boundary, B:406
- Thoracosphaera*, Cretaceous/Tertiary
boundary, B:406
- Zygodiscus sigmoides*, acme, B:406
- Site 698, diatoms, Paleocene, B:171, 173
- Site 699, diatoms, Paleocene, B:171, 173
- Site 700, diatoms
- Eocene, A:129
- Paleocene, B:171, 173
- Site 702, diatoms, Paleocene, B:171, 173
- Site 706
basement, ages, B:770
- hot spots
- migration, B:776
- models, B:771-773
- Site 707, winnowing, B:309
- Site 709
calcareous nannofossils, datum events, B:242
- carbon isotope stratigraphy,
Oligocene/Miocene boundary,
B:248-251
- Cibicidoides*, carbon isotopes, B:246
- Globigerinoides sacculifer*, oxygen isotopes,
B:246
- Globoquadrina venezuelana*, oxygen isotopes,
B:246
- oxygen isotope stratigraphy, B:247-249
- planktonic foraminifers
- carbon isotope stratigraphy, B:242, 244
- oxygen isotope stratigraphy, B:242-243
- sedimentation rate, calcareous ooze, B:242
- Site 710, magnetostratigraphy, B:245
- Site 713
basement, ages, B:770
- hot spots
- migration, B:776
- models, B:771-773
- magnetic properties, inclination, B:869
- paleolatitude, B:810
- Site 715
basement, ages, B:770
- hot spots
- migration, B:776
- models, B:772-773
- magnetic properties, inclination, B:869
- paleolatitude, B:810
- Site 716, Brunhes dissolution cycle, B:312
- Site 717
magnetic properties, susceptibility unit breaks,
B:823
- stress folds, A:22
- Site 718, stress folds, A:22
- Site 719
magnetic properties, susceptibility unit breaks,
B:823
- stress folds, A:22
- Site 722
calcium carbonate, mass accumulation rate,
B:309
- organic carbon, mass accumulation rate, B:309
- Site 738
calcareous nannofossils, Subzone CP1a, B:408
- Hornibrookina*, Cretaceous bloom, B:395, 407
- Prediscosphaera stoveri*, Cretaceous/Tertiary
boundary, B:406
- Site 744, strontium isotopes, planktonic for-
aminifers, B:921, 923
- Site 747
debris flow, B:747
- Maestrichtian unconformity, uplift-related,
A:89
- strontium isotopes, planktonic foraminifers,
B:921, 923-924
- uplift, A:12, B:747
- Site 748
calcareous nannofossils, strontium isotopes,
B:927
- carbonate, bioclastic, B:747
- glauconitic sands, A:89
- hot spots, formation, B:514
- Maestrichtian unconformity, uplift-related,
A:89
- strontium isotopes
- calcareous nannofossils, B:927
- planktonic foraminifers, B:921, 924
- tuff, terrigenous plant material, B:747
- uplift, B:747
- volcanic rocks, incompatible
element-enriched, A:474
- Site 750
Cretaceous/Paleogene boundary, B:492
- Futyania petalosa*, B:407
- Hornibrookina*, Cretaceous bloom, B:395, 407
- hot spots, formation, B:514
- Maestrichtian unconformity, uplift-related,
A:89
- tuff, terrigenous plant material, B:747
- Site 751, physical properties, vs. calcium carbon-
ate, B:254
- Site 752, A:111-169
ash layers, magnetic susceptibility, A:475
- bathymetry, A:63, 72, 113, 460, B:262, 360,
438, 458, 664, 682, 744, 764, 914, 934
- benthic foraminifers, A:485-487
- Cretaceous/Tertiary boundary, B:13, 19
- diversity, B:9-10
- faunal analysis, B:9-16
- Paleocene extinction event, B:19-21
- relative abundance, B:5-7
- stratigraphic range, B:6-9, 17
- biomagnetostratigraphy, B:725-731
- biostratigraphy, A:476-488, B:364, 722
- Cretaceous/Tertiary boundary, A:507-512, B:366
- calcareous nannofossils, A:43, 478-483
- ash layer, B:396
- Cretaceous assemblage, B:143-144, 401,
406
- Cretaceous/Tertiary boundary, A:511-512, B:396, 398-408
- Maestrichtian, B:146-147
- Subzone CP1a, B:408
- survivor forms, B:406-407
- Tertiary species, B:407
- zonation, Cretaceous, B:143
- calcareous sediments, lithification process,
B:260
- calcium, pore water concentration, A:494,
B:448-453
- calcium carbonate, A:497-498, B:79, 82,
398-399
- carbon isotopes, B:231, 888
- pelagic oozes, B:237
- carbonate, A:494-495, B:269, 433, 472
- accumulation rate, B:470
- Cretaceous-Paleogene, A:460, 462-463,
B:937
- gravel layers, B:471
- sedimentation, B:473, 475, 477
- chert, B:937
- chloride, pore water concentration, B:448-453
- clay
- Cretaceous/Tertiary boundary, source of,
B:419-420
- mineralogy, B:521
- compaction, physical properties and,
B:253-260

- consolidation tests, B:258–259
 Cretaceous/Paleogene boundary, B:492
 geochemistry, B:499–501
 Cretaceous/Tertiary boundary, A:463,
 507–516, B:730
 biomagnetostratigraphy, B:730–731
 depositional environment, A:467–469,
 507, B:941–943
 geochemistry, B:420, 424
 magnetic properties, A:510, 512–513
 productivity, B:420–421
 sedimentation rate, B:739–740
 trace elements, B:499–500, 913–918
 density, A:500, 504, B:269, 472
 pelagic oozes, A:499
 diatoms, A:45, 487–488
 biostratigraphic events, B:173–175
 datum levels, B:179
 Paleocene–Eocene, B:172–173, 175–178
 preservation, B:268–269
 dipping and truncated sequence, physical
 properties, A:497–498
 Eocene angular unconformity, A:460, 469
 eolian sediment flux, B:220–221
 geochemistry, A:56–57, 492–495
 carbonate-free residues, B:262, 266
 compositional end-members, B:424–426,
 439–440, 442–444
 end-member distribution, B:426–427
 factor analysis, B:424–425, 432
 rifting mechanisms and, B:444
 geological setting, A:71, B:423–424,
 437–438, 460, 721, 913
 geothermal gradient, B:522
 grain size
 pelagic oozes, A:462, B:213–215
 winnowing and, B:212
 heat flow, A:498, 506
 hydrocarbon gases, A:492–493
 igneous petrology, A:45–50
 interstitial-water chemistry, B:448–453
 iridium, B:421–422
 Cretaceous/Tertiary boundary, B:417, 915–
 916
 nannofossil extinction and, B:418–419
 Iridium Coincidence Spectrometer, B:416–417
 latitude and water depth, B:468
 limestone pebbles, B:936
 lipid geochemistry, B:473
 lithologic column, B:723, 935
 lithologic summary, A:461, B:729
 lithology, B:254, 364
 Cretaceous/Tertiary boundary, B:366, 415–
 416
 lithostratigraphy, B:219–220, 519–520, 722
 comparison, Kerguelen Plateau, A:463–
 464, B:938
 Cretaceous/Tertiary boundary, A:507
 seismic correlation, A:188, 235, B:668–
 669, 745
 location, A:6–7, 26, 458, 518, B:4, 79, 172,
 220, 230, 254, 453, 508, 521, 527,
 560, 592, 612, 722
 logging
 Cretaceous/Tertiary boundary, A:513–516
 geochemical, B:895–899
 geochemical, vs. core-derived lithology,
 B:908
 geophysical, A:59–60
 physical properties and, A:495–499
 low-molecular-weight hydrocarbons
 analytical methods, B:459–460
 toluene contents, B:461
 yields, B:460–465
 Maestrichtian carbonate platform, A:465–467,
 B:940–941
 magnesium, pore water concentration,
 B:448–453
 magnetic basement, depth estimates,
 B:683–684
 magnetic properties, A:50–56, 490
 Cretaceous/Tertiary boundary, A:510,
 512–513
 demagnetization, B:361–363
 inclination, B:367, 371–374
 remanence disturbance, A:51
 susceptibility, A:491
 magnetostratigraphy
 chronozone boundaries, B:727
 Cretaceous–Paleogene, B:363–365
 Cretaceous/Tertiary boundary, B:366
 lithologic correlation, A:488–489, 491
 reversal boundaries, B:363
 mass accumulation rates, A:466–467, B:223
 Cretaceous–Paleogene fluxes, A:468
 navigation, A:63, 65, 69–70
 Neogene currents, A:469–470
 objectives, A:12–13, 457, B:721, 936
 opal-CT-to-quartz transition, B:257
 organic carbon, A:493–494, B:474
 accumulation rate, B:470, 472, 475
 analytical methods, B:457
 source, B:477, 481
 organic matter, type of, B:472–473
 oxides, logging data, B:900–901
 oxygen isotopes, B:226, 231–232, 886–888
 alteration effects, B:452–453
 pelagic oozes, B:213–215, 217, 237
 pore water concentration, B:448–453
 paleoceanography, A:465–471, B:936
 Maestrichtian–Eocene, B:3–4
 paleodepths, B:724, 731, 746
 paleoenvironment, Paleogene, A:469
 paleolatitude, B:469, 724
 physical properties, A:57–59
 compaction effects, B:253–260
 laboratory vs. log measurements, B:257
 lithification effects, B:267–268
 logging and, A:495–499
 silica mineralogy and, B:521–522
 units A–E, A:505, B:264–266
 physiography, B:468
 planktonic foraminifers, A:43, 483–484
 Cretaceous, B:92
 Cretaceous–Paleogene, B:81, 85–87, 93
 Eocene–Oligocene distribution, B:89
 Paleocene–Eocene, B:90–91
 porosity, B:256
 productivity, following Cretaceous/Tertiary
 boundary event, B:427
 rare earth elements, B:441
 Cretaceous/Tertiary boundary, B:430, 501
 sedimentation rate, A:464, B:226, 472, 735,
 737–740, 934, 936, 938–939
 seismic stratigraphy, B:80, 439
 onlapping sequences, A:81–84, B:746–747
 prograding downlapping wedge, A:75–81
 Sequence I, B:743–744
 Sequence II, B:744
 seismic velocity, B:682–683
 silica, B:270
 siliceous microfossils, preservation,
 B:262–264
 tephra, lithology, A:471–472
 trace elements, Cretaceous/Paleogene
 boundary, B:499–500
 uplift, A:87–89
 velocity, laboratory vs. logged measurements,
 A:503
 volcanogenic sediment, inter-element ratios,
 B:425, 430
 winnowing, pelagic oozes, A:469–470
 X-ray fluorescence, B:520
- Site 753
 alteration, pre- vs. post-rift, B:522
 anisotropy, B:376
 of anhysteretic remanence, B:374
 ash layers
 geochemistry, A:472–475, 480–482
 magnetic susceptibility, A:475, 482–483
 temporal variations, A:474–475
 basalts
 geochemistry, B:574–575
 isotopic values, B:575
 bathymetry, A:63, 72, 460, B:262, 360, 438,
 458, 664, 682, 744, 764, 934
 benthic foraminifers, A:43–45, 485–487
 Cretaceous/Tertiary boundary, B:13, 19
 diversity, B:9
 faunal analysis, B:9–13, 16
 Paleocene extinction event, B:19–21
 relative abundance, B:5–7
 stratigraphic range, B:6–9, 17
 biomagnetostratigraphy, B:732–733
 biostratigraphy, A:476–488, B:368, 731–732
 calcareous nannofossils, A:43, 478–483
 zonation, Cretaceous, B:143
 calcium, pore water concentration, A:494,
 B:448–453
 calcium carbonate, A:497–498, B:81, 83
 carbonate, A:494–495, B:472
 accumulation rate, B:470
 Cretaceous–Paleogene, A:460, 462–463,
 B:937
 sedimentation, B:473, 475, 477
 chert, B:937
 chloride, pore water concentration, B:448–453
 clay mineralogy, B:521
 compaction, physical properties and,
 B:253–260
 Cretaceous/Tertiary boundary, depositional
 environment, B:941–943
 density, A:500, 504, B:472
 pelagic oozes, A:499
 diatoms, A:45, 487–488
 Eocene angular unconformity, A:460, 469
 geochemistry, A:56–57, 491–495
 carbonate-free residues, B:262
 geological setting, A:71, B:721
 geothermal gradient, B:522
 grain size, pelagic oozes, A:462
 gravity anomaly, A:84–86, B:682
 topography and, B:684, 690–692
 heat flow, A:498, 506
 hydrocarbon gases, A:492–493
 igneous petrology, A:45–50
 interstitial-water chemistry, B:448–453
 latitude and water depth, B:468
 lipid geochemistry, B:473
 lithologic column, B:723, 935
 lithologic summary, A:461
 lithology, B:368
 lithostratigraphy, B:519
 comparison, Kerguelen Plateau, A:463–
 464, B:938
 seismic correlation, A:235, B:668–669,
 745
 location, A:6–7, 26, 458, 518, B:4, 79, 172,
 220, 230, 254, 453, 508, 521, 527,
 560, 592, 612, 722
 logging

- geophysical, A:59-60
 physical properties and, A:495-499
 Maestrichtian carbonate platform, A:465-467,
 B:940-941
 magnesium, pore water concentration,
 B:448-453
 magnetic basement, depth estimates,
 B:683-684
 magnetic properties, A:50-56, 179-181
 demagnetization, B:361-363
 inclination, B:367, 371
 inclination, shallow, B:371-374
 remanence disturbance, A:51
 susceptibility, A:491
 magnetostratigraphy, B:368
 chronozone boundaries, B:728
 Cretaceous-Paleogene, B:365, 367
 lithologic correlation, A:488-489, 491
 reversal boundaries, B:363
 mass accumulation rates, A:466-467
 Cretaceous-Paleogene fluxes, A:468
 navigation, A:63, 65, 69-70
 Neogene currents, A:469-470
 objectives, A:12-13, 457, B:721, 936
 organic carbon, A:493-494
 accumulation rate, B:470, 472, 475
 analytical methods, B:457
 source, B:477, 481
 organic matter, type of, B:472-473
 oxygen isotopes, pore water concentration,
 B:448-453
 paleoceanography, A:465-471, B:936
 paleodepth, B:724, 733, 746
 paleoenvironment, Paleogene, A:469
 paleolatitude, B:469, 724
 physical properties, A:57-59
 compaction effects, B:253-260
 lithification effects, B:267-268
 logging and, A:495-499
 units A-E, A:505, B:264-266
 physiography, B:468
 planktonic foraminifers, A:43, 483-484
 Cretaceous-Paleogene, B:81, 87, 94
 sedimentation rate, A:464, B:472, 735,
 737-740, 934, 936, 938-939
 seismic stratigraphy, B:80, 439
 onlapping sequences, A:81-84, B:746-747
 prograding downlapping wedge, A:75-81
 Sequence I, B:743-744
 Sequence II, B:744
 seismic velocity, B:682-683
 silica
 diagenesis, B:266, 521
 effects of chemical environment on, B:267
 mineralogy, B:521-522
 physical properties, B:521-522
 source of authigenic, B:267
 X-ray mineralogy, B:261-262, 264
 siliceous microfossils, preservation,
 B:262-264
 tephra, lithology, A:471-472
 uplift, A:87-89
 winnowing, pelagic oozes, A:469-470
 X-ray fluorescence, B:520
- Site 754
 acoustic impedance, seismic correlation, B:679
 alteration, pre- vs. post-rift, B:522
 anisotropy, B:376
 ash layers
 geochemistry, A:472-475, 480-482
 magnetic susceptibility, A:475, 482-483
 temporal variations, A:474-475
 Turonian-Santonian, A:463
- bathymetry, A:63, 72, 193, 460, B:262, 360,
 438, 458, 664, 682, 744, 764, 934
 benthic foraminifers, A:43-45, 485-487
 Cretaceous/Tertiary boundary, B:13, 19
 deep water associations, B:43-44
 diversity, B:9, 58
 faunal analysis, B:9-13, 16, 38-39, 59-60,
 62-65, 70
 Oligocene-Pleistocene, B:32, 34-39, 50-
 52
 Paleocene extinction event, B:19-21
 relative abundance, B:5-7, 56
 stratigraphic range, B:6-9, 17, 39, 69
 biomagnetostratigraphy, B:733-735
 biostratigraphy, A:476-488, B:369
 calcareous nannofossils, A:43, 478-483
 biostratigraphic summary, B:162
 Cretaceous, B:144-145, 148, 150-153
 zonation, Cretaceous, B:143
 calcareous sediments, lithification process,
 B:260
 calcium, pore water concentration, A:494,
 B:448-453
 calcium carbonate, A:497-498, B:81, 84
 carbon isotopes, B:232-234
 pelagic oozes, B:238
 carbon preference index, B:483
 carbonate, A:494-495, B:472
 accumulation rate, B:470
 chert layers, B:471
 Cretaceous-Paleogene, A:460, 462-463,
 B:937
 sedimentation, B:473, 475, 477
 chert, B:937
 chloride, pore water concentration, B:448-453
 clay mineralogy, B:521
 compaction, physical properties and,
 B:253-260
 Cretaceous/Tertiary boundary, depositional
 environment, B:941-943
 density, A:500, 504, B:472
 pelagic oozes, A:499
 diatoms, A:45, 487-488
 Eocene angular unconformity, A:460, 469
 eolian sediment flux, B:220-221
 geochemistry, A:56-57, 491-495
 carbonate-free residues, B:262, 266
 geological setting, A:71, B:460, 721
 geothermal gradient, B:522
 grain size, B:217
 pelagic oozes, A:462
 gravity anomaly, A:84-86, B:682
 topography and, B:684, 690-692
 heat flow, A:498, 506
 hydrocarbon gases, A:492-493
 hydrogen index, B:476
 igneous petrology, A:45-50
 interstitial-water chemistry, B:448-453
 isoprenoid hydrocarbons, B:483
 latitude and water depth, B:468
 lipid geochemistry, B:473
 lithologic column, B:723, 935
 lithologic summary, A:461, B:734
 lithology, B:369
 lithostratigraphy, A:194-195, B:520
 comparison, Kerguelen Plateau, A:463-
 464, B:938
 seismic correlation, A:165, 188, B:668-
 669, 745
 location, A:6-7, 26, 458, 518, B:4, 79, 172,
 220, 230, 254, 453, 508, 521, 527,
 560, 592, 612, 722
 logging
 geochemical, B:895-897, 902-903
- geochemical, vs. core-derived lithology,
 B:908
 geophysical, A:59-60
 physical properties and, A:495-499
 low-molecular-weight hydrocarbons
 analytical methods, B:459-460
 toluene contents, B:461
 yields, B:460-465
 Maestrichtian carbonate platform, A:465-467,
 B:940-941
 magnesium, pore water concentration,
 B:448-453
 magnetic basement, depth estimates,
 B:683-484
 magnetic properties, A:50-56
 demagnetization, B:361-363
 inclination, B:367, 371
 inclination, shallow, B:371-374
 remanence disturbance, A:51
 susceptibility, A:491-492
 magnetostratigraphy, B:369
 chronozone boundaries, B:728
 Cretaceous/Paleogene boundary, B:367
 lithologic correlation, A:488-489, 491
 reversal boundaries, B:363
 mass accumulation rates, A:466-467, B:224
 Cretaceous-Paleogene fluxes, A:468
 navigation, A:63, 65, 69-70
 Neogene currents, A:469-470
 objectives, A:12-13, 457, B:721, 936
 opal-CT-to-quartz transition, B:257, 522
 organic carbon, A:493-494, B:474
 accumulation rate, B:470, 472, 475
 analytical methods, B:457
 source, B:477, 481
 organic matter, type of, B:472-473
 oxides, logging data, B:904-905
 oxygen isotopes, B:226, 231-232
 pelagic oozes, B:213-215, 218, 238
 pore water concentration, B:448-453
 paleoceanography, A:465-471, B:936
 paleodepths, B:724, 735, 746
 paleoenvironment, Paleogene, A:469
 paleolatitude, B:469, 724
 pebble layer, A:462
 pelagic ooze, coarse fraction, B:218
 physical properties, A:57-59
 compaction effects, B:253-260
 laboratory vs. log measurements, B:257
 lithification effects, B:267-268
 logging and, A:495-499
 units A-E, A:505, B:264-266
 physiography, B:468
 planktonic foraminifers, A:43, 483-484
 Cretaceous-Paleogene, B:81, 88-89, 95-97
 saturated hydrocarbons, B:478-480, 484-486
 biological markers, B:483
 sediment classification
 chemical, A:42
 granular, A:40-42
 sedimentation rate, A:464, B:226, 472, 735,
 737-740, 934, 936, 938-939
 seismic stratigraphy, B:80, 439
 onlapping sequences, A:81-84, B:746-747
 prograding downlapping wedge, A:75-81
 Sequence I, B:743-744
 Sequence II, B:744
 seismic velocity, B:682-683
 silica
 diagenesis, B:266, 521
 effects of chemical environment on, B:267
 mineralogy, B:521-522
 phases, B:265
 physical properties, B:521-522

- source of authigenic, B:267
 X-ray mineralogy, B:261–262, 264
 siliceous microfossils, preservation,
 B:262–264
 tephra, lithology, A:471–472
 terrigenous organic particles, B:472, 476
 uplift, A:87–89
 velocity, laboratory vs. logged measurements,
 A:503
 winnowing, pelagic oozes, A:469–470
 X-ray fluorescence, B:520
 Site 755, A:237–258
 alteration, pre- vs. post-rift, B:522
 anisotropy, B:376
 ash layers
 geochemistry, A:472–475, 480–482
 magnetic susceptibility, A:475, 482–483
 temporal variations, A:474–475
 Turonian–Santonian, A:463
 bathymetry, A:63, 72, 94, 460, B:262, 360,
 438, 458, 664, 682, 764, 934
 benthic foraminifers, A:43–45, 485–487
 biomagnetostratigraphy, B:735–736
 biostratigraphy, A:476–488, B:370
 calcareous nannofossils, A:43, 478–483
 Campanian, B:154–156
 Cretaceous, B:143, 145
 calcium, pore water concentration, A:494,
 B:448–453
 calcium carbonate, A:498, B:81, 85
 carbon preference index, B:483
 carbonate, A:494–495, B:472
 accumulation rate, B:470
 chert layers, B:471
 Cretaceous–Paleogene, A:460, 462–463,
 B:937
 sedimentation, B:473, 475, 477
 chert, B:937
 chloride, pore water concentration, B:448–453
 clay mineralogy, B:521
 compaction, physical properties and
 B:253–260
 Cretaceous/Tertiary boundary, depositional
 environment, B:941–943
 density, A:500, 504, B:472
 pelagic oozes, A:499
 depositional environment, A:237
 diatoms, A:45, 487–488
 Eocene angular unconformity, A:460, 469
 formation
 depositional model, B:747, 753
 stratigraphic model, B:760
 geochemistry, A:57, 492–495
 carbonate-free residues, B:262, 266
 compositional end-members, B:439–440,
 442–444
 rifting mechanisms and, B:444
 geological setting, A:71, B:437–438, 460, 721
 geothermal gradient, B:522
 heat flow, A:498, 506
 hydrocarbon gases, A:492–493
 hydrogen index, B:476
 igneous petrology, A:45–50
 interstitial-water chemistry, B:448–453
 isoprenoid hydrocarbons, B:483
 latitude and water depth, B:468
 lipid geochemistry, B:473
 lithologic column, B:723, 935
 lithologic summary, A:461, B:736
 lithology, B:370
 lithostratigraphy, B:520
 comparison, Kerguelen Plateau, A:463–
 464, B:938
 seismic correlation, A: 165, 188, 235,
 B:668–669, 745
 location, A:6–7, 458, 518, B:4, 79, 172, 220,
 230, 254, 453, 508, 521, 527, 560,
 592, 612, 722
 logging
 geophysical, A:59–60
 physical properties and, A:495–499
 low-molecular-weight hydrocarbons
 analytical methods, B:459–460
 toluene contents, B:461, 465
 yields, B:460–465
 Maestrichtian carbonate platform, A:465–467,
 B:940–941
 magnesium, pore water concentration,
 B:448–453
 magnetic basement, depth estimates,
 B:683–484
 magnetic properties, A:50–56
 demagnetization, B:361–363
 inclination, B:367, 371
 remenance disturbance, A:51
 susceptibility, A:491
 magnetostratigraphy, B:370
 chronozone boundaries, B:728
 Cretaceous/Paleogene boundary, B:367
 lithologic correlation, A:488–489, 491
 reversal boundaries, B:363
 mass accumulation rates, A:466–467
 Cretaceous–Paleogene fluxes, A:468
 navigation, A:63, 65, 69–70
 Neogene currents, A:469–470
 objectives, A:12–13, 457, B:721, 936
 opal-CT-to-quartz transition, B:522
 organic carbon, A:493–494, B:474
 accumulation rate, B:470, 472, 475
 analytical methods, B:457
 source, B:477, 481
 organic geochemistry, A:56–57, 491–492
 organic matter, type of, B:472–473
 oxygen isotopes, pore water concentration,
 B:448–453
 paleoceanography, A:465–471, B:936
 paleodepths, B:724, 735, 746
 paleoenvironment, Paleogene, A:469
 paleolatitude, B:469, 724
 pebble layer, A:462
 pelagic cap, A:457–458, 460, B:936–937
 magnetostratigraphy, A:488
 physical properties, A:495, 497
 physical properties, A:57–59
 compaction effects, B:253–260
 lithification effects, B:267–268
 logging and, A:495–499
 units A–E, A:505, B:264–266
 physiography, B:468
 planktonic foraminifers, A:43, 483–484
 Cretaceous–Paleogene, B:81–82, 89, 98
 rare earth elements, B:443
 saturated hydrocarbons, B:478–480, 484–486
 biological markers, B:483
 sediment classification
 chemical, A:42
 granular, A:40–42
 sedimentation rate, A:464, B:472, 735,
 737–740, 934, 936, 938–939
 seismic stratigraphy, B:80, 439
 onlapping sequences, A:81–84, B:746–747
 pelagic cap, A:84–86
 prograding downlapping wedge, A:75–81
 Sequence I, B:743–744
 Sequence II, B:744
 seismic velocity, B:682–683
 siliceous microfossils, preservation,
 B:262–264
 tephra, lithology, A:471–472
 terrigenous organic particles, A:496, B:476
 tuff, B:937–938
 terrigenous plant material, B:747
 Turonian–Santonian tuffs, A:73
 uplift, A:87–89
 winnowing, pelagic oozes, A:469–470
 X-ray fluorescence, B:520
 Site 756, A:259–303
 alteration
 high-temperature, B:534–535, 541
 low-temperature, B:622
 post-magmatic, B:563–565, 614–615
 temperature of, B:453–455
 argon age dating, basalts, B:509–515
 basal sediments, depositional environmen-
 B:560–561
 basalts
 age determination, B:509–516
 geochemistry, B:561–562, 565, 567–569,
 574–582, 586–588
 isotopic values, B:568–569, 572–573, 578–
 579, 594–595, 607
 magnetic properties, B:526–533, 539, 541–
 543, 561
 major oxides, B:541–542
 petrography, B:509, 621–624
 primary oxides, B:535, 539
 subaerial eruption, B:229
 tholeiitic vs. alkaline, B:622
 basement
 age, B:447, 561
 lithology, B:562
 bathymetry, A:93–95, B:764
 benthic foraminifers, A:43–45
 deep water associations, B:43–44
 diversity, B:58
 faunal analysis, B:39–43, 61, 66–68
 Oligocene–Pleistocene, B:32–33, 38, 44–
 52
 relative abundance, B:56
 stratigraphic range, B:54, 69
 calcareous nannofossils, A:43
 strontium isotopes, B:928–929
 zonation, Cretaceous, B:143
 calcium, pore water concentration, B:448–453
 calcium carbonate, B:82–83, 86, 227
 calcium ion concentration, A:534
 carbon isotopes, B:232–234
 carbonate veins, B:453–455
 pelagic oozes, B:239
 carbonate, B:472
 accumulation rate, B:470
 sedimentation, B:473, 475, 477
 carbonate veins, formation temperatures,
 B:453–455
 chloride, pore water concentration, B:448–453
 clinopyroxene, B:623, 656–757
 compaction, physical properties and,
 B:253–260
 Cretaceous/Tertiary boundary, oceanic
 circulation, B:820–821
 density, B:472
 diatoms, A:45
 Dupal anomaly, basalts with, geochemical
 comparison, B:638–639
 eolian sediment flux, B:220–221
 feldspar, basalts, B:568–569
 geochemistry, A:56–57
 alteration effects, A:531–534
 intersite variation, B:633–638
 standard sediment sample, B:891–893

- geological setting, A:13, B:526
hot spots
 Cretaceous–Oligocene locations, B:769
 migration, B:776
 models, B:771–773
igneous mineralogy, B:567
igneous petrology, A:45–50, 523–531
 alteration effects, B:614–615
incompatible elements, basalts, B:564, 575, 577, 630–631, 640–645
India, northward motion, A:16
India-Asia convergence
 geological record, B:819
 paleolatitudes constraints, B:811–812, 816–830
interstitial-water chemistry, B:448–453
latitude and water depth, B:468
lavas
 comparison, Kerguelen–Heard and Indian Ocean lavas, B:598–401
 incompatible element-enriched, A:530
 petrology, B:616–620
lead isotopes, B:596–598, 600–602, 604–606
linear regression analysis, B:805, 809–810
lipid geochemistry, B:473
lithostratigraphy, A:517, 519–520
location, A:6–7, 26, 100, 458, 518, B:79, 220, 230, 254, 453, 508, 527, 560, 592, 612, 722, 922
logging, geophysical, A:59–60
magma mixing, B:633
magnesium, pore water concentration, B:448–453
magnesium oxide, B:626–628
magnetic anomalies, A:93
magnetic mineralogy, optical and microprobe analysis, B:534–540
magnetic properties, A:50–56, B:798–799
 demagnetization, B:784–786, 793–796, 800
 hysteresis loop measurements, B:533–534, 536–537
 natural remanent magnetization, B:526–533, 539, 541
 remanence disturbance, A:51
 susceptibility, B:799, 855
magnetic record, A:101
magnetostratigraphy, B:797–799
mass accumulation rates, B:224
mineralogy, alteration, B:659
nannofossil ooze, B:229
navigation, A:93, 96–97
neodymium isotopes, B:600–602
objectives, A:13–22, B:763
organic carbon, B:474
 accumulation rate, B:470, 472
 source, B:477, 481
organic matter, type of, B:472–473
oxygen isotopes, B:227, 231–234
 carbonate veins, B:453–455
 pelagic oozes, B:239
 pore water concentration, B:448–453
paleolatitude, B:469, 856–857, 867
 comparison, Indian Ocean sites, B:810
 evolution, B:778
 mean inclination and, B:799–803, 871
physical properties, A:57–59
 compaction effects, B:253–260
physiography, B:468
planktonic foraminifers, A:43
 Cretaceous–Paleogene, B:83, 89–90, 98, 100–102
 strontium isotopes, B:921, 924–925
rare earth elements, B:588
 basalts, B:576, 600–602
 chondrite-normalized, B:634–638
 ridge evolution
 geochemical record, A:525–530
 paleolatitude record, B:811
 volcanogenic record, A:524–525
 ridge formation, paleomagnetic record, B:766–768
 rocks and thin sections, B:648–655
 sediment classification
 chemical, A:42
 granular, A:40–42
 sedimentation rate, A:517, 519, B:472
 strontium isotopes, B:593, 596–598, 600–602
 calcareous nannofossils, B:928–929
 planktonic foraminifers, B:921, 924–925
 tephra, lithology, A:520–523
 titanomagnetite, B:658
Site 757, A:305–358
 alteration
 high-temperature, B:534–535, 541
 low-temperature, B:622
 oxygen isotope changes with, B:452–453
 post-magmatic, B:563–565, 614–615
 temperature of, B:453–455
 argon age dating, basalts, B:509–515
 basal sediments, depositional environment, B:560–561
 basalts
 age determination, B:509–516
 geochemistry, B:561–562, 565, 569–570, 574–582, 586–588
 isotopic values, B:570–573, 578–579, 594–595, 607
 magnetic properties, B:526–533, 539, 541–543, 561
 major oxides, B:541–542
 petrography, B:509, 624–629
 primary oxides, B:535, 539
 subaerial eruption, B:229
 tholeiitic vs. alkaline, B:622
 basement
 age, B:447, 561
 lithology, B:562
 bathymetry, A:93, 102, 105, B:764
 benthic foraminifers, A:43–45
 calcareous nannofossils, A:43
 strontium isotopes, B:928, 930
 zonation, Cretaceous, B:143
 calcareous sediments, lithification process, B:260
 calcium, pore water concentration, B:448–453
 calcium carbonate, B:83, 87
 calcium ion concentration, A:534
 carbon isotopes, B:232–234, 236
 carbonate veins, B:453–455
 pelagic oozes, B:240
 carbonate, B:472
 accumulation rate, B:470
 sedimentation, B:473, 475, 477
 carbonate veins, formation temperatures, B:453–455
 chloride, pore water concentration, B:448–453
 clinopyroxene, B:623, 656–757
 compaction, physical properties and, B:253–260
 consolidation tests, B:258–259
 Cretaceous/Tertiary boundary, oceanic circulation, B:820–821
 density, B:472
 diatoms, A:45
 feldspar
 basalts, B:568–569
 glassy inclusions, B:571
 geochemistry, A:56–57
 alteration effects, A:531–534
 intersite variation, B:633–638
 geological setting, A:13, B:462, 526
 hot spots
 Cretaceous–Oligocene locations, B:769
 migration, B:776
 models, B:771–773
 igneous mineralogy, B:567
 igneous petrology, A:45–50, 523–531
 alteration effects, B:614–615
 incompatible elements, basalts, B:575, 577, 579, 630–631, 640–645
 India, northward motion, A:16
 India-Asia convergence
 geological record, B:819
 paleolatitude constraints, B:811–812, 816–830
 interstitial-water chemistry, B:448–453
 volcanic alteration effects, B:448
 latitude and water depth, B:468
 lavas
 comparison, Kerguelen–Heard and Indian Ocean lavas, B:598–601
 incompatible element-enriched, A:530
 petrology, B:616–620
 lead isotopes, B:596–598, 600–602, 604–606
 linear regression analysis, B:805, 809–810
 lipid geochemistry, B:473
 lithologic summary, B:255
 lithostratigraphy, A:517, 519–520
 location, A:6–7, 26, 104, 458, 518, B:79, 220, 230, 254, 453, 592, 922
 logging, geophysical, A:59–60
 low-molecular-weight hydrocarbons
 analytical methods, B:459–460
 yields, B:462–465
 magma mixing, B:633
 magnesium, pore water concentration, B:448–453
 magnetic anomalies, A:93, B:769
 magnetic mineralogy, optical and microprobe analysis, B:534–540
 magnetic properties, A:50–56, B:806–809, 812–818, 820–822, 824–826
 demagnetization, B:784, 786–788, 792, 804–805, 810
 hysteresis loop measurements, B:533–534, 536–537
 natural remanent magnetization, B:526–533, 539, 541
 remanence disturbance, A:51
 susceptibility, B:799, 855
 magnetic record, A:101, 105
 magnetostratigraphy, B:797–799
 mineralogy, alteration, B:659
 morphology, B:560
 nannofossil ooze, B:229
 navigation, A:93, 96–97
 neodymium isotopes, B:600–602
 objectives, A:13–22, B:763
 organic carbon, B:474
 accumulation rate, B:470, 472
 analytical methods, B:457
 source, B:477, 481
 organic matter, type of, B:472–473
 oxygen isotopes, B:231–232, 236
 alteration effects, B:452–453
 carbonate veins, B:453–455
 pelagic oozes, B:240
 pore water concentration, B:448–453
 paleolatitude, B:469, 858–860, 867
 comparison, Indian Ocean sites, B:810
 evolution, B:778

- mean inclination and, B:799-803, 871
 physical properties, A:57-59
 compaction effects, B:253-260
 physiography, B:468
 Plagioclase, trapped melt, B:658
 planktonic foraminifers, A:43
 Cretaceous-Paleogene, B:83, 90, 92-93,
 95, 97
 strontium isotopes, B:921, 925
 porosity, B:257
 productivity, B:472
 rare earth elements, B:589
 basalts, B:576, 600-602
 ridge evolution
 geochemical record, A:525-530
 paleolatitude record, B:811
 volcanogenic record, A:524-525
 ridge formation
 geochemical record, B:639-645
 models, B:559-560, 578-581, 768-773
 paleomagnetic record, B:766-768
 rocks and thin sections, B:648-655
 sediment classification
 chemical, A:42
 granular, A:40-42
 sedimentation rate, A:517, 519, B:472
 strontium isotopes, B:593, 596-598, 600-602
 calcareous nannofossils, B:928, 930
 planktonic foraminifers, B:921, 925
 tephra, lithology, A:520-523
 titanomagnetite, B:568
 Site 758, A:359-453
 alteration
 basaltic flow units, B:549
 high-temperature, B:534-535, 541
 low-temperature, B:622
 oxygen isotope changes with, B:452-453
 post-magmatic, B:563-565, 614-615
 temperature of, B:453-455
 argon age dating, basalts, B:509-515
 ash layers
 age range, B:275
 magnetic properties, B:307-309, 796
 basal sediments, depositional environment,
 B:560-561
 basalt, magnetic properties, A:395-396
 basaltic flow units
 contacts, B:548-549
 grain size variation, B:549
 thickness of, B:548
 basalts
 age determination, B:509-516
 geochemistry, B:561-562, 565, 571, 574-
 582, 586-588
 isotopic values, B:571-573, 578-579, 594
 595, 607
 magnetic properties, B:526-533, 539, 541-
 543, 561
 major oxides, B:541-542
 petrography, B:509, 629-633
 primary oxides, B:535, 539
 basement
 age, B:447, 561
 lithology, B:562
 magnetic properties, B:796-797
 bathymetry, A:93, 106, B:764
 benthic foraminifers, A:43-45
 Brunhes Chronozone, B:245
 calcareous nannofossils, A:43
 Cretaceous, B:145, 149, 159-161
 zonation, Cretaceous, B:143
 calcium, pore water concentration, B:448-453
 calcium carbonate, B:84, 88, 298, 307,
 310-311, 315, 316-328, 340-355, 495
 ash layer, B:490
 foraminifers, B:491
 mid-Pleistocene shift, B:313
 carbon isotope stratigraphy, B:248-251,
 304-308, 310-311
 Chron 6 carbon shift, B:251
 carbon isotopes, B:329-355
 carbonate veins, B:453-455
 carbon preference index, B:483
 carbonate, B:472, 493-494
 accumulation rate, B:470
 sedimentation, B:473, 475, 477
 carbonate veins, formation temperatures,
 B:453-455
 chloride, pore water concentration, B:448-453
 clinopyroxene, B:623, 656-757
 coarse fraction, B:316-328, 340-355
 stratigraphy, B:307
 compaction, physical properties and,
 B:253-260
 composite depth section, B:300-302, 306,
 310, 316-328
 magnetic susceptibility, B:312
 oxygen isotope record, B:303
 post-coring expansion/stretching, B:302
 sediment recovery gaps, B:300-302
 consolidation tests, B:258
 Cretaceous/Tertiary boundary
 geochemistry, B:498
 oceanic circulation, B:820-821
 density, B:299, 316-328, 340-355, 472
 diatoms, A:45
 Pliocene-Quaternary, B:190-191, 195
 stratigraphic ranges, B:193
 zonation, Oligocene-Miocene, B:195-199
 zonation, Pliocene-Quaternary, B:191-194
 feldspar
 basalts, B:568-569
 tephra layers, B:284
 gadolinium, B:910-912
 geochemistry, A:56-57
 alteration effects, A:531-534
 intersite variation, B:633-638
 microtektites, B:490
 geological setting, A:13, B:273, 462, 526, 703
 glass
 basaltic unit contacts, B:549
 geochemistry, B:277-283
 grain size
 basaltic flow units, B:549
 mineralogy and, B:379, 382
 hot spots
 Cretaceous-Oligocene locations, B:769
 migration, B:776
 models, B:771-773
 hydrogen index, B:476
 igneous mineralogy, B:567
 igneous petrology, A:45-50, 523-531
 alteration effects, B:614-615
 incompatible elements, basalts, B:575, 577,
 580, 630-631, 640-645
 India, northward motion, A:16
 India-Asia convergence
 geological record, B:819
 Neogene sedimentary record, B:821-829
 paleolatitude constraints, B:811-812, 816-
 830
 interstitial-water chemistry, B:448-453
 isoprenoid hydrocarbons, B:483
 latitude and water depth, B:468
 lavas
 comparison, Kerguelen-Heard and Indian
 Ocean lavas, B:598-601
 incompatible element-enriched, A:530
 petrology, B:616-620
 lead isotopes, B:596-598, 600-602, 604-606
 linear regression analysis, B:805, 809-810,
 872-874
 lipid geochemistry, B:473
 lithologic summary, B:255
 lithology, B:297-298, 703
 lithostratigraphy, A:517, 519-520
 magnetic property correlation, B:882
 location, A:6-7, 26, 458, 518, B:79, 125, 190,
 220, 230, 242, 254, 298, 453, 508,
 527, 560, 592, 612, 698, 722
 logging
 geochemical, B:895-897, 906-908
 geochemical, vs. core-derived lithology,
 B:908
 geophysical, A:59-60
 low-molecular-weight hydrocarbons
 analytical methods, B:459-460
 toluene content, B:463, 465
 yields, B:462-465
 magma mixing, B:633
 magnesium, pore water concentration,
 B:448-453
 magnetic anomalies, B:769
 magnetic mineralogy, optical and microprobe
 analysis, B:534-540
 magnetic properties, A:50-56, B:827-829,
 842-855
 ash layers, B:796
 basement, B:796-797
 declinations, B:710-712
 demagnetization, B:379-380, 382-384,
 788-795, 827-828, 839-841, 847,
 850-851
 directional changes, B:382-383, 386
 hysteresis loop measurements, B:533-534,
 536-537
 lithostratigraphic correlation, B:882
 mid-Pleistocene shift, B:313
 natural remanent magnetization, B:526-
 533, 539, 541
 remanence disturbance, A:51
 susceptibility, B:299-300, 307-312, 340-
 355, 384, 799, 855, 881
 transition field, B:383-385
 viscous remanent magnetization, B:701,
 709
 magnetostratigraphy, B:797-799
 Jaramillo polarity transition, B:385-388,
 391
 reversal boundaries, B:855
 mass accumulation rates, B:299
 microtektites, B:489-493, 502-503
 geochemistry, B:490, 494
 morphology, B:495-496
 mineralogy
 alteration, B:659
 grain size variation and, B:379, 382
 morphology, B:560
 navigation, A:93, 96-97
 neodymium isotopes, B:600-602
 northward movement, B:868
 northward movement rate, B:868, 872-874
 objectives, B:763
 paleoclimatic, A:18-21
 petrologic, A:13-16
 organic carbon, B:474
 accumulation rate, B:470, 472
 analytical methods, B:459-460
 source, B:477, 481
 organic matter, type of, B:472-473
 oxides, logging data, B:910-912

- oxygen isotope stratigraphy, B:247–248, 250, 304–308, 310–311
 magnetostratigraphic correlation, B:306–307
- oxygen isotopes, B:329–355
 alteration effects, B:452–453
 carbonate veins, B:453–455
 mid-Pleistocene shift, B:313
 planktonic foraminifers, B:298–299
 pore water concentration, B:448–453
- paleoceanography
 high-frequency fluctuations, B:311–312
 long-term change, B:309–311
- paleoclimatic objectives, A:362–363
- paleolatitide, B:469, 861–867
 comparison, Indian Ocean sites, B:810
 evolution, B:778
 mean inclination and, B:385, 799–803, 871
- physical properties, A:57–59, B:704
 compaction effects, B:253–260
- physiography, B:468
- phytoliths, Pliocene–Quaternary, B:192
- planktonic foraminifers, A:43, B:492
 carbon isotopes, B:245
 carbonate and productivity of, B:134
 Cretaceous, B:106
 Cretaceous–Paleogene, B:84–85, 108–109
 dissolution effects, B:130, 133
 dissolution-resistant species, B:135–136
 ecological factors influencing, B:133–134, 136–137
- Oligocene, B:103
 oxygen isotopes, B:133, 245
 Paleocene–Eocene, B:103–104
 relative abundance, B:126–127
 resistant species ratio, B:127, 130
 spatial distribution, B:138
 strontium isotopes, B:921
 temporal variation, B:127–130
- porosity, stress and, B:702, 704
- productivity, calcium carbonate MAR and decreases in, B:309
- rare earth elements, B:590
 basalts, B:576, 600–602
- ridge evolution
 geochemical record, A:525–530
 paleolatitide record, B:811
 volcanogenic record, A:524–525
- rocks and thin sections, B:648–655
- saturated hydrocarbons, B:478–480, 484–486
 biological markers, B:483
- sediment classification
 chemical, A:42
 granular, A:40–42
- sediment recovery gaps, B:300–301
- sedimentation rate, A:517, 519, B:242, 302–304, 472
 subsidence and, B:309–310
- silicoflagellate
 Pliocene–Quaternary, B:192
 stratigraphic ranges, B:193
- stress, B:701–702, 704–706, 713–717
 deformation rate analysis, B:698–699
 measurement procedures, B:699–701
- strontium isotopes, B:593, 596–598, 600–602
- sulfides, B:659
- tephra
 dating of, B:284–287
 evolution, B:287
 formation mechanisms, B:287
 geochemistry, B:277, 284–285, 288–291
 lithology, A:520–523
 Neogene, B:275
- terrigenous clay, A:534
- terrigenous organic particles, B:476
- titanomagnetite, B:568
- Toba ash, B:489
 carbonate content, B:490
- trace elements, Cretaceous/Paleogene boundary, B:498
 winnowing, B:309
- Site NER-2C, seismic reflection profiling, A:103
- Slump scars, Broken Ridge, A:81, 84
- Slumping
 Broken Ridge, B:746
 tephra layer A, Site 758, B:276
- Smectite
 Broken Ridge, in altered glass, A:472
 Ninetyeast Ridge
 alteration phases, A:275
 in lavas, B:629
- Smectite, green, Ninetyeast Ridge, A:520
- Smectite spherules, Site 758, Cretaceous/Paleogene boundary, B:491–492
- Sodium oxide, Ninetyeast Ridge
 basalts, B:567
 mobility during alteration, B:615
 vs. magnesium oxide, basement basalts, A:401
- Southeast Indian Ridge
 post-rift oceanic crest, vs. pre-rift Broken Ridge crest, B:690
 reconstruction, Chron 26, B:768
 subsidence history, B:691
- Southwest Indian Ridge, spreading direction, change in, B:816
- St. Paul Fracture Zone, basalt, isotopic ratios, B:639
- St. Paul Island
 isotopic ratios, Ninetyeast Ridge basalts compared, B:599
 lanthanum/samarium ratio, vs. lead isotopes, B:599–600
- lava, isotopic ratios, A:526
- lavas
 formation of, B:600–601
 Ninetyeast Ridge compared, B:600–601
 lead isotopes, basalts, B:569
 paleolatitide, present day latitude correlation, B:802, 811
 tholeiitic basalt, A:526
- Stevens Klint, Cretaceous/Tertiary boundary
 clay origins, B:415, 419
 iridium, B:415
- Stress
 deformation rate analysis, Site 758, B:698–706
 geomagnetic measurements, Site 758, B:701
 Ninetyeast Ridge, breakouts, A:22
 Site 758
 breakout, A:363
 deformation rate analysis, B:698–701, 706
 horizontal, B:704–705, 714–715
in-situ memory, B:699, 705
 measurement procedures, B:699–701
 pore pressure effect, B:702, 704–705, 713
 predicted vs. *in-situ*, B:705, 716–717
 vertical, B:701–702, 705
- Strontium
 Ninetyeast Ridge
 basalts, B:571
 ferrobasalts, A:526
 lavas, B:600–602
 vs. calcium oxide/aluminum oxide ratio, B:639
 vs. plagioclase content, B:627
 zirconium correlation, B:565, 568–569
 Site 752, Cretaceous/Tertiary boundary, B:424
- Strontium isotopes
 Broken Ridge, A:14
- Indian Ocean, mid-ocean ridge basalts, A:15
- Ninetyeast Ridge, B:580, 924–925
 alteration effects, B:596–598
 basalts, B:569–573
 basement basalts, B:593
 lavas, A:529, B:596–598, 600–603
 nannofossil correlation, B:925, 928–930
 ocean-island and spreading-ridge basalt affinities, B:599
 vs. lanthanum/samarium ratio, B:599, 608
 vs. lead isotopes, A:529, B:607
 vs. neodymium isotopes, B:578, 603
 Site 756, B:924–925
- Strontium/neodymium ratio, Ninetyeast Ridge intersite differences, B:637
 vs. lanthanum content, B:640
- Stumpata Quartzarenite, deposition, correlation, Ninetyeast Ridge Eocene unconformity, B:819
- Subsidence
 Broken Ridge, B:229
 grain size correlation, B:214
 pelagic cap deposition during, B:753
 pre- vs. post-rift, A:87–89
 Turonian–Eocene, A:499
 Ninetyeast Ridge, A:525
- Suiko Seamount
 lanthanum content, B:643
 primary oxide composition, B:539
 sandstone, B:633
- Sulfate
 Broken Ridge, A:136
 calcite-gypsum supersaturation and, B:448
 Ninetyeast Ridge, A:398
 calcite-gypsum supersaturation and, B:448
 calcium and variations in, A:286, 399
 gradient, effect of volcanic alteration on, A:334
- Sulfides, Ninetyeast Ridge, B:659
 nickel and cobalt content, B:629
- Sunda Arc, eruptions, B:273
- Sunda Trench, bending lithosphere, B:705
- Surtseyan eruptions, Ninetyeast Ridge, lapilli creation, A:523
- Tahiti, volcanism, Jaramillo Subchronozone B:385
- Takena Formation
 magnetic properties, Albian–Cenomanian, B:816
 paleoposition, Tertiary, B:816
- Takli Formation (Nagpur), Intertrappean Beds, pelobatid frogs, B:818
- Tantalum, Ninetyeast Ridge
 vs. lanthanum/tantalum ratio, B:582
 vs. thorium and lanthanum, basalts, B:578
- Tasman Fold Belt, extensional tectonism, Cretaceous, A:491
- Tasman Sea, spreading, end of, B:816
- Tectonic history
 Broken Ridge, A:5–12, 457, B:933–934
 Campanian/Maestrichtian boundary event, A:491
 Ninetyeast Ridge, A:12, 534–535, B:765–766
- Temperature
 Broken Ridge, A:152
 blocking, B:361
 Miocene gradient, B:215
 opal-A to -CT conversion, B:522
 Site 758, isotopic effects of, B:304
- Tephra
 Broken Ridge
 hot spot association, A:471
 lithology, A:471–472

- Ninetyeast Ridge
 air-fall deposits, A:520-521
 lithology, A:520-523
 macroscopic characteristics, A:274
- Site 758
 evolution, B:287
 grading, B:276
 macroscopic characteristics, B:274, 276
 magma petrogenesis and residence time variation, B:287
 Neogene, B:275
 parental magma, B:287
- Tephra layers
 major oxide composition, B:284
- Site 758
 age and chemical correlation, B:277
 correlation, Toba Caldera eruptions, B:287
 dating of, B:284-287
 depositional environment, B:276
 feldspar composition, B:284
 formation mechanisms, B:287
 geochemistry, B:277-284
 layers A-D, B:276-277
 macroscopic characteristics, B:274-276
 major oxide geochemistry, B:288-291
 major oxide ratios and grouping of, B:277, 284
 oxygen isotopes, paleomagnetic dating correlation, B:285
 thickness of, B:276
- Tephrochronology, Site 758, B:273
- Terrigenous plant material, Broken Ridge, in Turonian-Santonian tuffs, A:89
- Terrigenous sediment
 Broken Ridge, in Turonian-Santonian tuffs, A:238, B:747
 high-frequency fluctuations, Site 758, B:311-312
 long-term changes, Site 758, B:309-310
- Ninetyeast Ridge
 Himalayan source, B:477
 Neogene influx, B:821-822
- Site 758
 after Indian-Asian collision, A:363
 calcium carbonate dissolution and increase in, B:311
 correlation, susceptibility breaks, B:822-823
 Himalayan source, A:519-520, B:310-311, 822
 magnetic properties, B:792-793
 mass-accumulation rate, A:517, 519
 Miocene-Holocene, A:375, 424
 Neogene influx, B:822
 riverine output and increase in, B:309-310
 sources, B:297-298
- Thermal conductivity
 Broken Ridge, A:58, 143-144, 147, 150-151, 183-185, 219, 229, 252, 254, 498
 Ninetyeast Ridge, A:58, 289-290, 298, 342, 352-353, 407, 436-437
 lithologic Unit I, A:343
 tuffs, A:344
- Thermal subsidence curve, Broken Ridge, Neogene, A:469
- Thorium, Ninetyeast Ridge
 lavas, B:600-602
 vs. tantalum and lanthanum, basalts, B:578
- Thorium/lanthanum ratio, Ninetyeast Ridge, B:639
 continental-related component, B:644-645
 oceanic island basalt characteristics, B:644
 vs. lanthanum content, B:640
- Thorium/tantalum ratio, Ninetyeast Ridge, B:639-640
- basalts, B:571
 continental-related component, B:644-645
 oceanic island basalt characteristics, B:644
 vs. lanthanum content, B:640
 vs. lanthanum/tantalum ratio, B:581, 584, 641-642, 644
- Thorium/uranium ratio, Ninetyeast Ridge
 alteration effects, B:624
 lavas, B:625
 vs. lanthanum content, B:640
- Tibetan Plateau
 extensional tectonism, Ninetyeast Ridge
 susceptibility breaks and, B:826
- uplift
 after Indian-Asian collision, A:363
 sedimentation rate impact, A:20
- Titanium dioxide, Ninetyeast Ridge, A:281, 327, 329
 plagioclase content and, A:329
 vs. zirconium, basement basalts, A:401
- Titanium/europium ratio, Ninetyeast Ridge
 intersite differences, B:637
 vs. lanthanum content, B:640
- Titanium oxide
 Broken Ridge, vs. susceptibility, ash layers, A:475, 482
 Site 757, basalt, A:331
- Titanium oxide/magnesium oxide ratio, Site 757, lavas, B:625-627
- Titanium/phosphorus ratio, Ninetyeast Ridge, basalts, B:577
- Titanium/zirconium ratio, Ninetyeast Ridge, basalts, B:568-570
- Titanomagnetite, Ninetyeast Ridge, B:658
 fractionation, B:576-577
 low-temperature alteration, B:539
 oxidation exsolution, B:534-535
 oxide relationship, A:333
 trellis pattern, B:535
- Toba ash, Site 758, A:368, 373
 carbonate content, B:489
- Toba Caldera
 ash distribution, B:273
 correlation, Ninetyeast Ridge tephra, B:287
 magma chamber, zoning, B:284
- Toba Lake event, planktonic foraminiferal changes, B:489, 492
- Toba tuff, Site 758, feldspar composition, B:284
- Toluene
 Broken Ridge, olefin content correlation, B:461-462
 Ninetyeast Ridge, B:462-463, 465
- Trace elements
 Broken Ridge
 ash layers, A:474, 476-479
 basalt, B:594-595
 Cretaceous/Paleogene boundary, B:492
- Ninetyeast Ridge
 basalt, B:572, 581, 608
 control of melting process on, B:579
 Cretaceous/Paleogene boundary, B:492, 498
 vs. isotopic ratios, B:599, 608
- Site 752, B:426-428
 Cretaceous/Tertiary boundary, B:499-500, 913-919
- Tristan da Cunha, lava, Dupal isotopic signature, B:599
- Tristan hot spot, predicted path, B:516
- Tropical-Subtropical Transition Zone, Indian Ocean, productivity, A:18
- Tuff
 Broken Ridge
 depositional environment, A:89
 terrigenous component, B:747
- Ninetyeast Ridge, A:311, 360, 374
 basalt contacts, A:391
 depositional environment, A:350
 physical properties, A:337, 344
 toluene content, B:462-465
- Site 755
 Cretaceous, B:735
 toluene content, B:462-465
 Turonian-Santonian, A:73, 239-240, 242, B:743-744, 937-938
- Tuff, calcareous, Site 758, magnetic susceptibility, A:394, 424
- Turbidites, glauconite, Broken Ridge, A:464-465, B:939-940
 in Turonian tuffs, A:464
- Turkmenia, geomagnetic polarity transition, B:385
- Ulvöspinel, Site 757, A:333
- Uplift
 Broken Ridge, A:87-89, 457
 duration, B:444
 Maestrichtian, A:467
 origin, A:500
 rift-related, A:5
- Kerguelen-Heard Plateau, unconformity resulting from, A:89
- Ninetyeast Ridge, sedimentary record, A:17, B:310-311
- Uplift, flexural, Broken Ridge, B:740, 934
 magnitude, A:9-10, 86
 rifting duration and, A:172
- Upwelling
 Broken Ridge, productivity increase with, B:941
 Ninetyeast Ridge, oxygen isotopes and rate of, B:452
- Uranium, Ninetyeast Ridge, lavas, B:600-602
- Vanadium, Ninetyeast Ridge, A:280, 282
 basement basalts, A:330, 333, 399, 402, 527
 vs. niobium, A:402
- Veins, carbonate, Ninetyeast Ridge
 isotopic values, B:453-455
 temperature of formation, B:453
- Veins, oceanic crust, Ninetyeast Ridge, lead isotopes, B:597
- Velocity
 Broken Ridge, A:58-59
 density and, B:684
 dipping and truncated sequence, A:497
 laboratory vs. logging data, A:503
 lithologic association, B:521
 pelagic cap, A:495, B:254
 truncated limestone-chert/prograding wedge boundary, A:73, 75
 laboratory vs. logged measurements, Site 752, A:160
 Ninetyeast Ridge, A:58-59
 calcareous chalks, B:256
- Site 752, A:141, 146-147, 163
 chert layer, A:149
 laboratory vs. logging data, A:149-150, 160
 opal-A to -CT transition, B:256-257
 sonobuoy solutions, A:151, 164
- Site 753, A:183
 laboratory vs. logging data, A:185
 sonobuoy solutions, A:187
- Site 754, A:217-219, 223-224
 across unconformity, A:217
 sonobuoy solutions, A:234
- Site 755, A:252, 254
 sonobuoy solutions, A:258

- Site 756, A:289, 292, 295–296
basalt, A:297
sonobuoy vs. refraction, A:293
- Site 757, A:339–340, 348
basement basalt, A:344
lithologic Unit II, A:344
- Site 758, A:404, 414, 432–434
- Velocity, seismic
Broken Ridge, B:682
laboratory vs. sonobuoy measurements, A:254
Santonian–Turonian section, B:683
sonobuoy solutions, B:686
Ninetyeast Ridge, A:347
basalts, A:393
vs. measured, A:414
- Velocity anisotropy
Broken Ridge, A:147–148, 217, 225–226, 252
horizontal vs. vertical, A:142, 218
Ninetyeast Ridge, A:288, 340, 349, 404
Site 754, A:225–226
Site 757, A:349
- Virtual geomagnetic pole (VGP)
Atlantic Ocean, equatorial, B:385, 388
Site 758
Matuyama/Brunhes transition, B:377
reliability criteria, B:385, 387
- Vitrite
Broken Ridge, A:496, B:472
random reflectance values, B:476
Ninetyeast Ridge, B:472
random reflectance values, B:476
- Volcanic centers, Ninetyeast Ridge, A:534
- Volcaniclastic sediment
Broken Ridge, classification, A:41–42
Ninetyeast Ridge, classification, A:41–42
- Volcanism
Broken Ridge
basal sediment record, B:507
iridium content and, B:417
magnetic susceptibility and intensity of, A:475
Turonian–Santonian, A:465, B:939–940
Ninetyeast Ridge
air-fall deposits, A:520–521
alkalic, A:530
characteristics, A:524
Hawaiian volcanoes compared, A:15–16, B:633
hot spot role, B:639
late-stage, B:768
phreatic eruptions, A:306, 520–521, 524, 531
source of, paleolatitudes, B:768
sources of excess, B:763
- Volcanism, constructional, A:499, B:559, 933–934
- Volcanism, explosive, Ninetyeast Ridge, Neogene, B:273
- Volcanogenic sediment, Site 752, B:519–520
diagenetic enrichment, B:429
inter-element ratios, B:425, 430
source, B:428–429
vs. carbonate content, B:427–428
- Walvis Ridge
lava, Dupal isotopic signature, B:599
winnowing, pelagic oozes, B:212
- Water content
Broken Ridge, A:140, 144, 182–183, 217, 250–251, 505
Ninetyeast Ridge, A:287–288, 292, 337, 402–404
Site 758, host sediment and interstitial water influence, B:277
- Wharton Basin
fossil spreading ridge, B:765
magnetostratigraphy, east-west anomalies, B:765
ridge jump, B:801
spreading center, A:13, 261
northward migration, B:765
tectonic chart, B:765–766
- Winnowing
Broken Ridge
Antarctic Circumpolar Current influence, A:90
pelagic ooze, A:458, 460, 469, 471, 497, 500, B:212, 753, 936, 943–944
Ninetyeast Ridge, effect on coarse fraction, B:309
- Woodside Creek
clay, Cretaceous/Tertiary boundary, origin, B:415, 419
iridium, Cretaceous/Tertiary boundary, B:415
- X-ray diffraction, Broken Ridge, A:140, 214–215, B:520–521
- X-ray fluorescence
Broken Ridge, A:47–48, 50
Ninetyeast Ridge, A:47–48, 50
- Ytterbium, Ninetyeast Ridge, vs. lanthanum, basalts, B:568
- Yttrium, Ninetyeast Ridge, A:280, 282
basement basalts, A:333, 399
as function of age, A:531
- Yttrium/niobium ratio, Ninetyeast Ridge
ash beds, A:393
basement basalts, A:330, 403–434
of incompatible elements, A:282
intersite differences, B:637
two-component magma mixing, B:641
vs. zirconium/niobium ratio, A:334–335, 403–404, 527–528, 530, B:644
- Zanskar region, Cretaceous/Tertiary boundary, sedimentary record, B:819
- Zeolite
Broken Ridge
burial diagenesis, B:521
mineralogy, B:521–522
Ninetyeast Ridge, A:324, B:629
non-oxidative alteration, B:563
- Zhob Valley, ophiolites, obduction, A:16
- Zinc, Site 757, vs. niobium, A:334
- Zirconium, Ninetyeast Ridge, A:280–281, 327
basement basalts, A:333, 399, 526, B:568
strontium correlation, B:565, 569
vs. aluminum oxide, basalts, B:577
vs. barium, basalts, B:565
vs. chromium, basalts, B:574
vs. modal plagioclase, basalts, B:580
vs. nickel, basalts, B:574
vs. niobium, basement basalts, A:400
vs. rubidium, basalts, B:564
vs. titanium dioxide, basement basalts, A:401
vs. titanium oxide and scandium, basalts, B:575
- Zirconium/niobium ratio
Broken Ridge, ash layers, A:474
Ninetyeast Ridge
ash beds, A:393
basalts, B:568, 571, 577, 581
basement basalts, A:330
of incompatible elements, A:282
intersite differences, A:393, B:637
post-magmatic alteration, A:329
source differences and variation in, A:526, 528
two-component magma mixing, B:641
vs. yttrium/niobium ratio, A:334–335, 403–404, 527–528, 530, B:644
- Zirconium/tantalum ratio, Ninetyeast Ridge, basalts, B:571
- Zirconium/yttrium ratio, Ninetyeast Ridge, basalts, B:568, 571, 577, 581
- Zones, fossil. *See* in Taxonomic Index

TAXONOMIC INDEX

- Abathomphalus mayaroensis*/*Globotruncana gansseri* boundary, Broken Ridge, A:489
- Abathomphalus mayaroensis* Zone
Broken Ridge, A:485
Site 752, A:128, B:86, 731
Site 754, A:210, B:356, 733
Site 758, A:384, B:97
- Acarinina* sp.
Site 752, B:122
Site 757, B:92
- acervulinoides*, *Planoglobulina*, B:104
- aculeatus*, *Hyalodiscus*, Site 752, B:183
- acuta*, *Gublerina*, B:102
- acuta*, *Morozovella*, B:113
Site 758, A:383
- acutiloba*, *Asterolampira*, Site 758, B:200
- acutus*, *Ceratolithus*
Site 752, first occurrence, A:123
Site 753, first occurrence, A:175
Site 754, first occurrence, A:199
Site 757, first occurrence, A:316
Site 758, A:376
- aequa*, *Morozovella*, B:113
Broken Ridge, A:484
- alabamensis*, *Hantkenina*, B:115
- alata*, *Rhizosolenia*, Site 758, B:207
- alazanensis*, *Cibicidoides*, Site 756, B:32
- albatross*, *Brizalina*, Site 756, B:71
- albeari*, *Morozovella*, B:113
- Allomorphina* sp., Site 752, B:25
- allomorphinoides*, *Quadriformina*, Site 752, B:25
- altiaperturus*, *Globigerinoides*, B:111
Site 757, B:121
- altispira*, *Dentoglobigerina*
Site 709
carbon isotopes, B:244, 248–249
oxygen isotopes, B:243, 248–249
Site 758
carbon isotopes, B:245–246
oxygen isotopes, B:245–246
- altispira globosa*, *Globoquadrina*, B:111
- altus*, *Chiasmolithus*, Site 754, A:200
- americana*, *Pullenia*, Site 752, B:29
- ammophila*, *Hanzawaia*
Site 742, B:26
Site 752, B:9, 26
Maestrichtian–Eocene, B:5–6
Site 754, B:9
Site 756, B:33
- Ammophaeroidina* sp., Site 752, B:29
- Amphistegina*, Site 754, A:205
- amphora*, *Hanzawaia*, Site 758, A:385
- ampliapertura*, *Globigerina*, B:109
Site 755, B:90
- angiporoides*, *Globigerina*, B:109–110
Site 756, B:120
- angulata*, *Globorotalia*, Site 752, A:128
- angulata*, *Globotruncanita*, B:107
- angulata*, *Morozovella*, B:113
Broken Ridge, A:484
- angulaturalis*, *Globigerina*, B:110
- angulosa*, *Acarinina*, B:111
- angustiumbilicata*, *Globigerina*, B:110
Site 758, B:120
- Anomalina capitatus*/*A. danicus* assemblage, Site 752, B:14
- Anomalinoidea capitatus*/*A. danicus* assemblage, Site 752, B:9
- anthophorus*, *Reinhardtites*, B:149
Site 754, A:200, B:145, 166
- antillea*, *Orthomorphina*, Site 756, relative abundance, B:56
- aragonensis*, *Aragonia*, Site 752, B:25
- aragonensis*, *Morozovella*, B:113
- arca*, *Globotruncana*, B:107
- Archaeoglobigerina*, Site 758, A:384
- Archaeoglobigerina* spp., Site 754, B:89
- architecturalis*, *Melosira*
Site 752, B:180
Site 758, A:386, B:205
last occurrence, B:195
- aries*, *Trinacria*, Site 752, B:184
last occurrence, B:727
- Arkhangelskiella cymbiformis*/*Nephrolithus frequens* Zone, Site 752, B:143
- arkhangelskiyi*, *Prediscosphaera*, Site 758, B:169
- asperus*, *Globigerinelloides*, B:109
- australiformis*, *Planorotalites*, B:114
Site 752, B:123
- australis*, *Archaeoglobigerina*, B:105
Site 758, B:119
- australis*, *Hornibrookina*, Broken Ridge, A:481
- baltica*, *Whiteinella*, B:108
- barbadiensis*, *Discoaster*, Site 757, last occurrence, A:316
- barnesae*, *Watznaueria*, Site 752, B:143
- barri*, *Globigerinatheka*, B:111
- barronii*, *Craspedodiscus*, Site 758, B:200, 206
- beccariiiformis*, *Stensioina*
Broken Ridge, A:487
Site 752, B:9, 25
carbon isotope values, B:16, 19
isotopic values, B:885
last occurrence, A:129, B:6
Maestrichtian–Eocene, B:5
Site 758, A:386
- belemnus*, *Sphenolithus*
Broken Ridge, A:479
Site 754, first occurrence, A:199
Site 756, A:268, 270
- bella*, *Pseudopodosira*, Site 752, B:183
- bijugatus*, *Zygrhablithus*
Broken Ridge, last occurrence, A:479
Site 752, last occurrence, A:123
- binaiensis*, *Globigerina*, B:110
- bipartita*, *Globigerina*, B:110
Site 758, B:120
- bipolaris*, *Stictodiscus*, n. sp., Site 752, B:180, 183, 187
- Biscutum coronum* Zone, Site 754, B:145
- Biscutum magnum* Zone
Site 752, B:144
Site 754, B:145
- Biscutum* sp., Site 752, B:413
- bisecta*, *Reticulofenestra*
Broken Ridge, last occurrence, A:479
Site 752, last occurrence, A:123
- blowi*, *Archaeoglobigerina*, B:105
- Bogorovia* sp., Site 758, B:207
- Bogorovia veniamini* Zone, Site 758, A:387, B:198
- bosquensis*, *Archaeoglobigerina*, B:106
Site 758, A:384
- Braarudosphaera*
Broken Ridge, A:477
Site 752
Cretaceous/Tertiary boundary, B:406
- shallow-water origin, A:480
Site 754, shallow-water origin, A:480
- brachiatum*, *Triceratium*, Site 752, B:184
- bradburyi*, *Bulimina*, Site 752, B:25
- bradyi*, *Cibicidoides*, Site 758, A:385
- bradyi*, *eggerella*
Site 754, Indian Deep Water association, B:43
Site 756, B:71
Indian Deep Water association, B:43
- bradyi*, *Karrieriella*, Site 754, B:32, 71
relative abundance, B:40
- bradyi*, *Trifarina*, Site 754, relative abundance, B:42
- bramlettei*, *Tribrachiatum*, Site 752, first occurrence, A:127, 132–133
- brevispira*, *Hantkenina*, B:115
- brevispira*, *Turrilina*, Site 752, B:25
stratigraphic range, B:9
- brittonensis*, *Whiteinella*, B:108
- broedermanni*, *Acarinina*, B:111
- brouweri*, *Discoaster*
Site 752, last occurrence, A:123
Site 754, last occurrence, A:198–199
Site 757, last occurrence, A:316
Site 758, last occurrence, A:376
- brunii*, *Baxteriopsis*, Site 758, B:206
- bukryi*, *Thalassiosira*, Site 758, A:386, B:201
- Buliminella* sp., Site 752, B:25
- bullbrookii*, *Acarinina*
Site 753, A:178
Site 754, A:204
Site 757, A:320
- bulletta*, *Dorothia*, Site 752, B:29
- bulloides*, *Globigerina*, B:110
Site 752, A:128
Site 753, A:177
Site 754, A:201
Site 758
abundance, B:132
coarse fraction association, B:130, 135
dissolution trend, B:133
ecological role, B:127
- bulloides*, *Pullenia*
Site 753, A:179
Site 754
Indian Deep Water association, B:43
relative abundance, B:41
Site 756, B:73
Indian Deep Water association, B:43
- bulloides*, *Sphaeroidina*
Site 754, B:32
relative abundance, B:42
Site 756, B:75
- bulloides tripartita*, *Globigerina*, B:111
- Burseolina pacifica*/*Cibicidoides mundulus* assemblage, Site 754, B:38, 62
correlation, Site 756 assemblages, B:51
Miocene decrease, B:50
- Calculites obscurus* Zone, Site 755, B:145
- canaliculata*, *Dicarinella*, B:107
- capita*, *Anomalinoidea*, Site 757, A:320
- capitatus*, *Anomalinoidea*, Site 752, B:6
- carinata*, *Ehrenbergina*, Site 754, B:32
relative abundance, B:40
- castrorum*, *Biscutum*, Site 752, B:412
Cretaceous/Tertiary boundary, B:405–406
- Catapsydrax* sp., Site 757, B:120
- caucasica*, *Morozovella*, B:113
Site 757, B:92

- cavus*, *Coccolithus*
Site 752, Cretaceous/Tertiary boundary,
B:405, 407
Site 758, A:380
- cerroazulensis*, *Turborotalia*, B:114
Site 757, A:320
- Cestodiscus* sp., Site 758, B:204-205
- chapmani*, *Planorotalites*, B:114
- cheneveri*, *Triceratium*, Site 752, B:184
- chipolensis*, *Cassidulina*, B:114
- chipolensis*, *Cassigerinella*, B:114
- Cibicidoides*, Site 752, A:129
- Cibicidoides hyphalus* assemblage, Site 752, B:9,
15
- Cibicidoides mundulus/Burseolina pacifica* as-
semblage, Site 756, B:43, 68
- Cibicidoides mundulus/Oridorsalis umbonatus* as-
semblage, Site 756, B:43, 66
Oligocene decrease, B:50
- Cibicidoides pseudoperlucidus* assemblage, Site
752, B:9, 14
- Cibicidoides* sp.
Site 709, oxygen isotopes, B:244, 246
Site 752, B:28
oxygen isotopes, B:231
Site 754, B:74
- Cibicidoides subspiratus* assemblage, Site 752,
B:10, 15
- ciesielskii*, *Hemiaulus*, Site 752, B:186
- ciperoensis*, *Globigerina*, B:110
- ciperoensis*, *Sphenolithus*
Site 754, last occurrence, B:733
Site 756, A:270
Site 757, last occurrence, A:316
Site 758, last occurrence, A:376
- citae*, *Globotruncanella*, Site 754, A:204
- clavicamerata*, *Protentella*, B:114
Site 757, B:92-93, 95
- coalingensis*, *Acarinina*, B:112
Site 752, B:121
Site 753, B:121
Site 757, B:121
- cocoaensis*, *Turborotalia*, B:114
- collactea*, *Truncorotaloides*, B:114
- columbiana*, *Guembelitra*, B:102
Site 753, B:87, 120
- compressa*, *Planorotalites*, B:114
- conicotruncata*, *Morozovella*, B:113
- conicus*, *Globorotalites*, Site 752, B:29
- conicus*, *Hemiaulus*, Site 752, B:186
- constans*, *Biscutum*, Site 752, B:413
- continua*, *Jenkinsella*, B:112
- convexa*, *Morozovella*, B:113
- convexa aspinosa*, *Thalassiosira*, Site 758, B:193
- coronata*, *Marginotruncana*, B:107
- coronum*, *Biscutum*, Site 758, B:164
- corpulenta*, *Globigerina*, B:110
- coryelli*, *Pullenia*, Site 752, B:28
Maestrichtian-Eocene, B:5
- corystus*, *Nephrolithus*, Site 752, B:165
last occurrence, B:144
- Coscinodiscus* spp., Site 752, B:180
- costa*, *Planulina*, Site 758, A:385
- costata*, *Planulina*, Site 754, B:74
- costulata*, *Heterohelix*, B:102
- costulata*, *Pseudoguembelina*, B:104-105
- Craspedodiscus elegans* Zone, Site 758, A:386,
B:195
- Craspedodiscus undulatus* Zone, Site 752, B:175
- crassa*, *Globocassidulina*, Site 754, A:204
- crassaformis*, *Globorotalia*, Site 753, first occur-
rence, A:177
- crassaformis hessi*, *Globorotalia*, Site 758, A:382
- crassata*, *Morozovella*, B:113
- creta*, *Alabama*, Site 752, B:25
- cretacea*, *Archaeoglobigerina*, B:106
Site 754, B:119
- cretacea*, *Guembelitra*, B:102
- cretacea*, *Rhizosolenia*, Site 758, B:208
- Cribrostomoides* sp., Site 758, A:386
- cruciatatus*, *Neocrepidolithus*, Site 752, B:412
- Cruciplacolithus* lineage, Site 752, Creta-
ceous/Tertiary boundary, B:407
- crux*, *Zygodolithus*, Site 752, B:407, 411
Cretaceous/Tertiary boundary, B:406
- cryptomphala*, *Globigerina*, B:110
- cubensis*, *Chiloguembelina*, B:99
Site 754, B:88
Site 756, B:120
- cuneata*, *Cassidulina*, Site 754, A:204
- cunialensis*, *Turborotalia*, B:114
- Cyclagelosphaera* sp., Site 752, Cretaceous/Terti-
ary boundary, B:407
- cymbiformis*, *Arkhangelskiella*
Site 752, B:413
Site 758, B:168
Cretaceous/Tertiary boundary, B:405-406
- daniae*, *Cribrosphaerella*
Site 752, B:413
Site 758, Cretaceous/Tertiary boundary, B:705
- danicus*, *Anomalinoidea*, Site 752, B:6, 26
isotopic values, B:885
- danicus*, *Chiasmolithus*, Site 752, A:127
first occurrence, A:133, B:365, 730
- daubjergensis*, *Globigerina*, B:110
- daubjergensis*, *Globoconusa*, Site 758, A:384
- dayi*, *Cibicidoides*
Site 752, B:28
stratigraphic range, B:6
Site 754, B:6
- decussata*, *Micula*
Site 752, B:413
Cretaceous/Tertiary boundary, B:405-406
Site 754, A:200
- deflandrei*, *Discoaster*, Broken Ridge, A:479
- dehiscens*, *Globoquadrina*, B:111
Site 754, B:733
last occurrence, A:202
Site 755, B:89
Site 756, A:271
Site 757, last occurrence, A:319
Site 758, last occurrence, A:382
- dehiscens*, *Sphaeroidinella*
Site 757, first occurrence, A:319
Site 758, A:382
- delicatulus*, *Bolivinoidea*
Site 752, B:25
Site 753, B:25
- densa*, *Acarinina*, B:112
Site 757, B:122
- dentata*, *Heterohelix*, B:104
- diastypus*, *Discoaster*, Site 752, A:127
- Discoaster binodosus* Subzone, Site 757, A:315
- Discoaster multiradiatus* Zone
Site 752, B:728
Site 757, A:315
- dissimilis*, *Biscutum*, Site 758, B:164
- dissimilis*, *Globigerina*, B:109
- dissonata*, *Alabama*, Site 752, B:9, 25, 726-727
- distentus*, *Sphenolithus*
Site 754, B:733
Site 756, A:270
- doliolus*, *Pseudoeunotia*, Site 758, A:386
first occurrence, B:191, 194
- draco*, *Bolivinoidea*
Site 752, B:25
Cretaceous/Tertiary boundary, B:6, 12
- Site 754, B:6
- dutertrei*, *Neogloboquadrina*
Indian Ocean N, spatial distribution, B:134,
137
- Site 758
abundance, B:132, 136
coarse fraction association, B:130, 135
ecological conditions favoring, B:126-127
ecological conditions influencing, B:134,
136
- ecclesiastica*, *Thiersteinia*, Site 755, A:483
- echolsi*, *Astrononion*
Site 754, B:32
comparison, Site 756, B:38
Indian Deep Water association, B:43
relative abundance, B:41
Site 756, B:33, 38, 73
comparison, Site 754, B:38
Indian Deep Water association, B:43
relative abundance, B:57
- edita*, *Globigerina*, B:110
- edwardsii*, *Hornibrookina*, Site 752, comparison,
Maud Rise species, A:512
- Eggerella* sp., Site 752, B:29
- ehrenbergi*, *Globigerinelloidea*, Broken Ridge,
A:485
- ehrenbergi*, *Planorotalites*, B:114
- Ehrenbergina carinata* assemblage, Site 754,
B:38-39
- Ehrenbergina* spp., Site 754, B:65
- elegans*, *Craspedodiscus*, n. var., Site 758, B:199,
206
- elegans*, *Pseudotextularia*, B:105
- elegans asymmetricus*, *Craspedodiscus*, n. var.,
Site 758, B:206
- Ellipsoidella* sp., Site 752, B:25
- ellipsoides*, *Ellipsoidina*, Site 752, B:29
- ellipticus*, *Actinocyclus*, Site 758, B:205
- eocaena*, *Globigerina*, B:110
- eocaenus*, *Cibicidoides*, Site 752, B:9, 27
- eocenica*, *Pullenia*
Site 752, B:29
stratigraphic range, B:9
Site 753, stratigraphic range, B:9
- Epistominella unbonifera/Cibicidoides mundulus*
assemblage, Site 756, B:43, 67
correlation, Site 754 assemblages, B:51
faunal changes, B:50
- Eprolithus* sp., B:149
Site 755, B:170
- esnaensis*, *Acarinina*, B:112
- euapertura*, *Globigerina*, B:110
Site 754, B:120
Site 756, B:120
Site 758, B:95
- Eunotogramma* sp., Site 758, B:201, 208
- excavata*, *Tribrachia*, Site 758, A:386
- excavata tetragona*, *Trinacria*, Site 752, B:184
- excertus*, *Gyroidinoidea*, Site 752, B:29
- exigua*, *Epistominella*, Site 756, B:73
- exilis*, *Discoaster*, Broken Ridge, first occurrence,
A:479
- eximius*, *Eiffellithus*, Site 758, B:167
- eximius*, *Eiffellithus*
Site 755, B:145
Site 758, last occurrence, B:149
- favus*, *Favocassidulina*, Site 757, A:320
- fennerae*, *Anaulus*, n. sp., Site 752, B:180, 186-
187
- finalis*, *Pulleniatina*, Site 758, first occurrence,
A:382

- fistulosus*, *Globigerinoides*, Site 758, last occurrence, A:382
- florealis*, *Nuttallinella*, Site 752, B:26
- floridanus*, *Cyclicargolithus*
Broken Ridge, A:479
Site 757, A:316
Site 758, A:376
- fohsi fohsi*, *Globorotalia*, Site 758, oxygen isotopes, B:247
- fossilis*, *Nitzschia*, Site 758, last occurrence, A:386, B:191
- fossus*, *Neorepidolithus*, Site 752, B:411
- fraga*, *Thalassiosira*, Site 758, B:201, 205
- frequens*, *Nephrolithus*
Broken Ridge, first occurrence, A:488
first occurrence, B:143
Site 752, B:165, 731
first occurrence, A:127, 134, B:144
Site 754, A:200
Site 758, Cretaceous/Tertiary boundary, B:705
- frequens frequens*, *Nephrolithus*, Site 752, B:413
- frequens miniporus*, *Nephrolithus*, Site 752, B:413
- fringa*, *Globigerina*, B:110
- frontosa*, *Globigerina*, B:110
Site 753, B:87
- fructicosa*, *Racemiguembelina*, B:105
- furcatus*, *Marthasterites*, Site 755, A:242, 244, 246, B:145
- galavisi*, *Globigerina*, B:110
Site 756, B:120
- Gansserina gansseri* Zone, Site 752, B:731
- gartneri*, *Quadrum*, Site 755, B:170
- Gavelinella* sp., Site 752, B:28
- Gavelinopsis lobatulus/Uvigerina proboscidea* assemblage, Site 754, B:38, 63
Pliocene development, B:50
- gelida*, *Rocella*, Site 758, absence of, B:195, 199
- gigas*, *Chiasmolithus*
Site 753, A:481
last occurrence, B:432
Site 757, A:316
- Globigerina calida calida* Subzone, Site 758, A:382
- Globigerina* sp., Site 757, B:120
- Globigerinatheka index* Zone, Site 753, A:178
- Globigerinatheka* spp., Site 757, A:320
- Globigerinelloides* spp., B:108-109
Site 752, Cretaceous/Tertiary boundary, A:507, 509
- Globigerinoides* spp., B:111
- Globigerinoides trilobus* Zone, Site 753, A:178
- Globocassidulina* sp.
Site 754, B:32, 74-75
comparison, Site 756, B:38
relative abundance, B:40
Site 756, B:33, 66-67, 75
comparison, Site 754, B:38
relative abundance, B:56
- Globocassidulina* sp. assemblage
Site 754, B:38, 62
Site 756, B:39, 43
- Globoquadrina*, Ninetyeast Ridge, conditions for, A:18
- Globorotalia conoidea/Globorotalia inflata* lineage
Broken Ridge, A:484
Site 754, A:201
- Globorotalia fohsi* evolutionary lineage
Site 753, A:178
Site 756, A:270
- Globorotalia inflata* Zone, Site 754, A:201
- Globorotalia merotumida/Globorotalia tumida* lineage, Site 757, A:319
- Globorotalia peripheroronda-Globorotalia peripheroacuta* Zone
Site 753, A:178
Site 754, A:202-203
- Globorotalia truncatulinoidea* Zone, Site 758, A:382
- globosa*, *Globocassidulina*
Site 752, B:29
Site 754, B:6, 75
Indian Deep Water association, B:43
Site 756, Indian Deep Water association, B:43
- globosus*, *Gyroidinoides*, Site 752, B:28
last occurrence, B:6
- Globotruncana coronata* group, Site 755, A:246
- Globotruncana coronata-Globotruncana linneiana* group, Broken Ridge, A:485
- Globotruncana gansseri* Zone, Site 752, A:121
- Globotruncana primitiva* group, Site 755, A:246
- Globotruncanella citae/Abathomphalus intermedius/Abathomphalus mayaroensis* evolutionary sequence, Site 754, A:204
- Globotruncanella citae/Abathomphalus intermedius/Abathomphalus mayaroensis* lineage, Site 752, A:129
- globulifera*, *Tritaxia*, Site 752, B:29
- globulosa*, *Heterohelix*, B:104
- glomerosa*, *Praeorbulina*, Site 754, A:203
- glomerosa curva*, *Praeorbulina*, Site 757, A:319
- glutinata*, *Globigerinita*, B:111
Site 758
abundance, B:132
coarse fraction association, B:130, 135
dissolution trend, B:133
ecological role, B:126
- gombosii*, *Triceratium*, Site 752, B:181, 184, 187
- gortanii*, *Globigerina*, B:110
- gothicus*, *Tetralithus*, Site 758, A:382
- graciliformis*, *Uvigerina*, Site 754, B:71
- gracilis*, *Pyxilla*, Site 752, B:186
- gracillima*, *Stilostomella*, Site 752, B:25
- gracillium*, *Triceratium*, Site 752, A:130
- grandis*, *Prediscosphaera*, Site 758, B:169
- grata*, *Buliminella*, Site 752, B:25
- Gravelinella* sp., Site 752, B:29
- gravelli*, *Acarinina*, B:112
Site 752, B:122
- grimsdalei*, *Cibicidoides*, Site 752, B:27
- Gyroidina orbicularis/Rectuvigerina striata* assemblage, Site 754, B:38, 64
- Gyroidinoides* sp., Site 752, B:28
- hagni*, *Dicarinella*, B:107
- harnatus*, *Discoaster*, Site 756, A:268
- Hantkenina*, B:115
- havanensis*, *Cibicidoides*
Site 756, B:72
Site 757, A:320
Site 758, A:385
- havanensis*, *Globotruncanella*, B:107
- havanensis*, *Nonion*
Site 752, B:27
Site 756, B:73
- hebetata*, *Rhizosolenia*, Site 752, B:186
- Hemiaulus incurvus* Zone, Site 752, A:130, 488, B:175-176, 179, 728
- Hemiaulus periperus* Zone, Site 752, B:179-180, 728
- Hemiaulus* sp.
Site 752, B:180, 185-186
Site 758, B:206, 208
- Heterohelix* spp., Broken Ridge, A:485
- heteromorphus*, *Sphenolithus*, Site 756, A:268
- hexacamerata*, *Rugoglobigerina*, B:107
Site 752, B:86
- Site 758, A:384
- hispidia*, *Uvigerina*, Site 756, B:33, 71
- holmdelensis*, *Hedbergella*, B:107
- horizontalis*, *Globocassidulina*, Site 754, B:75
- Hornibrookina*
Site 738, Cretaceous bloom, B:395
Site 750, Cretaceous bloom, B:395, 407
Site 752
Cretaceous bloom, B:407
Cretaceous/Tertiary boundary, B:408
- howei*, *Catapsydrax*, B:109
- hustedtii*, *Denticulopsis*, Site 758, A:386
- Huttonia* sp., Site 752, A:130
- huxleyi*, *Emiliana*, Site 758, A:376
- Hyalodiscus* sp., Site 752, B:183
- hyphalus*, *Cibicidoides*, Site 752, B:27
Maestrichtian-Eocene, B:5
- hystrix*, *Ehrenbergina*, Site 756, B:75
- imbricata*, *Dicarinella*, B:107
- imitata*, *Planorotalites*, B:114
- impedens*, *Bulimina*, Site 756, B:32, 71
- inaequilateralis*, *Hemiaulus*, Site 752, B:180, 185
- inaequispira*, *Globigerina*, B:110
- inconstans*, *Morozovella*, B:113
Site 758, A:383-384
- incrassata*, *Coryphostoma*, Site 752, B:25
Cretaceous/Tertiary boundary, B:12
last occurrence, B:6
- increbescens*, *Turborotalia*, B:114
- incurvus*, *Hemiaulus*, Site 752, B:186
- index*, *Globigerinatheka*, B:111
Site 752, B:121
Site 754, A:204, B:121
Site 756, B:83
- inflata*, *Cribrorhantkenina*, B:114
- inflata*, *Globorotalia*
Broken Ridge, A:483
Site 753, first occurrence, A:177
Site 754, A:201
- Inoceramus* shell fragments, Site 754, A:195
- intermedia*, *Pseudotextularia*, B:105
- intermedius*, *Abathomphalus*, B:105
Site 754, A:204
- inversus*, *Markalius*, Site 752, B:412
Cretaceous/Tertiary boundary, B:405-407
- jarvisi*, *Bulimina*
Site 756, B:32-33
Site 757, A:320
- jouseae*, *Nitzschia*, Site 758, A:386
- kleinpellii*, *Heliolithus*, Site 752, first occurrence, A:127, 133, B:728
- kolchidica*, *Morozovella*, B:113
- kugleri*, *Fohsella*, B:112
Site 757, B:122
- kugleri*, *Globorotalia*, Site 756, A:271
- labiacrassata*, *Zeaglobigerina*, B:111
- lacunosa*, *Pseudoemiliana*, Site 758, last occurrence, A:376
- Lapideacassis* sp., Site 752, Cretaceous/Tertiary boundary, B:407
- laurisae*, *Cibicidoides*
Site 752, B:27
Site 756, B:32, 72
- Lenticulina*, Site 757, A:320
- Lenticulina* spp.
Site 752, B:6
Site 753, B:6
Site 754, B:6, 32, 64
relative abundance, B:42
Site 756, B:38

- relative abundance, B:57
Site 758, A:386
- Lenticulina* spp. assemblage
Site 752, B:9, 15
Site 753, B:9
Site 754, B:9, 38
- lepidula*, *Stilostomella*, Site 756, B:75
relative abundance, B:56
- levis*, *Reinhardtites*
Site 752, B:731
last occurrence, A:127
Site 754, B:145, 166
- lewisianus rhomboides*, *Coscinodiscus*, Site 758, B:195
- linaperta*, *Globigerina*, B:110
- linneiana*, *Globotruncana*, B:107
- linneiana*, *Globotruncana*?, Site 758, A:384
- lobatulus*, *Cibicidoides*, Site 754, B:72
- lobatulus*, *Gavelinopsis*
Site 754, B:32
relative abundance, B:40
Site 756, B:73
- lodoensis*, *Discoaster*
Broken Ridge, first occurrence, A:488
Site 752, first occurrence, A:132
Site 757, A:318
- longicornis*, *Hemiaulus*, Site 758, B:206
- Lucianorhabdus cayeuxii* Zone, Site 755, B:145
- Lucianorhabdus maleformis* Zone, Site 755, B:145
- macellus*, *Ellipsolithus*, Site 752, A:127
- macilenta*, *Bulimina*
Site 752, B:9, 25
Site 756, B:71
- magna*, *Rossiella*, n. comb., Site 758, B:200, 207
- magnificus*, *Kamptmerius*
Site 755, A:246, B:145
first occurrence, A:244
Site 758, B:168
Cretaceous/Tertiary boundary, B:705
- magnum*, *Biscutum*, Site 758, B:164
- maleformis*, *Lucianorhabdus*, Site 755, B:145
- maleinterpretaria*, *Nitzschia*, Site 758, B:207
first occurrence, B:195
- margaritae*, *Globorotalia*, Site 754, first occurrence, A:202
- marginata*, *Marginotruncana*, B:107
- marginodentata*, *Morozovella*, B:113
Site 752, A:128
- Marthasterites furcatus* Zone, Site 755, B:145
- martini*, *Chiloguembelina*, B:99
- martinii*, *Prinsius*, Site 752, first occurrence, A:133, B:730
- marylandicus*, *Raphidodiscus*, Site 758, B:205
- mayaroensis*, *Abathomphalus*, B:105
Broken Ridge, A:484
Site 754, B:89
first occurrence, A:204
Site 758, A:384, B:97
- mayeri*, *Globorotalia*, Site 753, first occurrence, A:178
- mayeri*, *Jenkinsella*, B:112
- mckannai*, *Acarinina*, B:112
- mediaconvexa*, *Thalassiosira*, Site 758, A:387, B:205
- menardii*, *Globorotalia*, Site 758
abundance, B:132, 136
coarse fraction association, B:130, 135
Cretaceous/Paleogene boundary, B:489
dissolution trend, B:133
ecological conditions favoring, B:126-127
- mexicana*, *Bulimina*, Site 754, B:32, 71
relative abundance, B:41
- mexicana*, *Osangularia*, Site 756, B:74
- mexicanus*, *Cibicidoides*, Site 756, B:32
- micra*, *Pseudohastigerina*, B:115
Site 758, A:383
- Micula decussata* Zone, Site 755, B:145
- midwayensis*, *Chiloguembelina*, B:102
- midwayensis*, *Coryphostoma*, Site 752, B:25
last occurrence, B:6
- minuta*, *Paracassidulina*, Site 754, B:75
- miozea*, *Uvigerina*, Site 756, B:71
- mirabile*, *Triceratium*, Site 752, B:184
- mohleri*, *Discoaster*, Site 752, first occurrence, B:728
- monmouthensis*, *Hedbergella*, B:107
- moratus*, *Lithastrinus*, Site 755, B:167
- morenoensis*, *Coscinodiscus*, Site 758, B:208
- Morozovella*, B:113
- Morozovella trinidadensis* Zone, Site 758, A:384
- Morozovella uncinata* Zone, Site 758, A:384
- Morozovella velascoensis* Zone, Site 752, B:728
- multicamerata*, *Planoglobulina*, B:104
- multicostata*, *Rectuvigerina*, Site 756, B:71
- multiradiatus*, *Discoaster*
Site 752, first occurrence, A:133
Site 757, A:316
- multispinatus*, *Globigerinelloides*, B:109
Site 758, B:97
- munda*, *Tenuitella*, B:114
- mundulus*, *Cibicidoides*
Site 754, B:32
comparison, Site 756, B:38
relative abundance, B:40-41
Site 756, B:33, 72
comparison, Site 754, B:38
relative abundance, B:56-57
- murus*, *Micula*, first occurrence, B:143
- naguwichiensis*, *Pseudohastigerina*, B:115
Site 753, B:87
- nana*, *Jenkinsella*, B:112-113
- navarroensis*, *Praebulimina*, Site 752; B:25
- nazcaensis*, *Clavigerinella*, Site 757, B:92-93, 95
- neocrassus*, *Neorepidolithus*, Site 752, B:412
- Neorepidolithus* spp., Site 752, Cretaceous/Tertiary boundary, B:405, 407
- Nephrolithus frequens* Zone
Site 752, A:512
Site 758, B:149
- nicobarensis*, *Clavatorella*, B:112
Site 758, B:97, 122
- nitida*, *Acarinina*, B:112
- Nitzschia jouseae* Zone, Site 758, A:386
- Nitzschia reinholdii* Zone, Site 758, A:386, B:191
- nobilis*, *Discoaster*, Site 752, first occurrence, B:728
- noda*, *Gyroidina*, Site 752, B:29
- nuttalli*, *Pseudotextularia*, B:105
- Nuttallides truempyi* assemblage
Site 752, B:9, 15
Site 753, B:9
- obesa*, *Globigerinella*, B:112
- obliquiloculata*, *Pulleniatina*, Site 758
abundance, B:132, 136
coarse fraction association, B: 130, 135
Cretaceous/Paleogene boundary, B:489
dissolution trend, B:133
ecological conditions favoring, B:126-127
- obliquum*, *Gartnerago*, Site 755, A:246, B:145
- obscurus*, *Calculites*, Site 754, B:145
- occlusa*, *Morozovella*, B:113
- officinalis*, *Globigerina*, B:110
- okadaï*, *Discoaster*, Site 758, A:378
- oligocenica*, *Azpeitia*, Site 758, B:207
last occurrence, B:195
- operculata*, *Thoracosphaera*, Site 752, B:412
- opima*, *Jenkinsella*, B:113
- opima nana*, *Globorotalia*, Site 758, A:383
- orbicularis*, *Gyroidina*, Site 754, relative abundance, B:40
- orbiculatum*, *Triceratium*, Site 752, B:184
- Oridorsalis* sp., Site 752, B:29
- ornata*, *Lisitzinia*, Site 758, B:206
- orthostylus*, *Tribrachiatum*, Site 757, A:315
- Osangularia* sp., Site 752, Maestrichtian-Eocene, B:5-6
- Osangularia* sp./*Hanzawaia ammophila* assemblage, Site 752, B:9, 14
- ostoensis*, *Pullenia*, Site 756, B:73
- ouachitaensis*, *Globigerina*, B:110
- ovalis*, *Cestodiscus*, Site 758, B:205
- pacifica*, *Burseolina*
Site 754, A:204, B:32
relative abundance, B:41
Site 756, B:38, 75
relative abundance, B:57
- paleacea*, *Rossiella*, Site 758, B:207
first occurrence, B:195
- paleocenica*, *Tritaxia*, Site 752, B:29
- palpebra*, *Pseudoguembelina*, B:105
- parcus*, *Aspidolithus*, Site 758, A:381-382
- parcus constrictus*, *Aspidolithus*
Site 754, B:145
Site 758, B:168
reworked specimens, A:381-382
- pauperata*, *Laticarinina*, Site 756, B:71
- pectinatus*, *Monomarginatus*, Site 758, B:163
- pelagicus*, *Coccolithus*, Broken Ridge, A:479
- pennyi*, *Rugoglobigerina*, B:108
Site 752, B:119
- pentacamerata*, *Acarinina*, B:112
Site 752, B:122
- pentaradiatus*, *Discoaster*, Broken Ridge, first occurrence, A:479
- peregrina*, *Uvigerina*, Site 754, B:71
- peripheroacuta*, *Globorotalia*
Site 753, A:178
Site 758, oxygen isotopes, B:247
- peripheroronda*, *Globorotalia*, Site 753, A:178
- peripterus longispinus*, *Hemiaulus*, n. var., Site 752, B:180, 185
- peripterus peripterus*, *Hemiaulus*, Site 752, B:185
- petaloidea*, *Globotruncanella*, B:107
- petalosa*, *Futyania*
Site 750, Cretaceous/Tertiary boundary, B:407
Site 752, B:411
Cretaceous/Tertiary boundary, B:407
- petalosis*, *Toweius*, Site 752, comparison, Maud Rise species, A:512
- phacelosus*, *Tranolithus*
Site 754, B:145
Site 758, B:167
last occurrence, A:381
- pileolus*, *Trinacria*, Site 758, B:208
- planata*, *Heterohelix*, B:104
- planispira*, *Hedbergella*, B:107
Site 752, B:119
- planoconica*, *Planorotalites*, B:114
- Planorotalites pseudomenardii* Zone, Site 758, A:383
- planulatus*, *Gyroidinoides*, Site 752, B:27
- Planulina wuellerstorfi*, Site 754, B:63
- Planulina wuellerstorfi* assemblage, Site 754, B:38
bottom water cooling and Miocene decrease in, B:51

- Miocene decrease, B:50
pleniporus, *Misceomarginatus*, Site 758, B:163
polycystinorum, *Hemiaulus*, Site 752, B:185
polymorphus, *Eunotogramma*, Site 758, B:208
pomeroli, *Turborotalia*, B:114
postcretacea, *Globorotalia*, B:112
 Site 756, B:122
praeacuta, *Anomalinoidea*, Site 752, B:26
praebergonii, *Rhizosolenia*, Site 758, last occurrence, B:191
praebergonii praebergonii, *Rhizosolenia*, Site 758, A:386
 last occurrence, B:191
praebergonii robusta, *Rhizosolenia*, Site 758, A:386
 last occurrence, B:191
Praebulimina sp., Site 752, B:25
praebulloides, *Globigerina*, B:110
 Site 756, B:120
praecursoria, *Morozovella*, B:113
praedeheiscens, *Globoquadrina*, B:111
 Site 754, A:203, B:121
 Site 755, B:89
praequadratus, *Lithraphidites*, first occurrence, B:143
primalabiata, *Thalassiosira*, Site 758, last occurrence, B:195
primitiva, *Acarinina*, B:112
 Site 753, A:178
 Site 757, A:320
primus, *Amaurolithus*
 Broken Ridge, first occurrence, A:479
 Site 753, last occurrence, A:175
 Site 754, A:199
primus, *Cruciplacolithus*, Site 752, B:411
 Cretaceous/Tertiary boundary, B:405, 407
princeps, *Rocella*, Site 758, B:207
 last occurrence, B:199
Prinsius sp., Site 752, Cretaceous/Tertiary boundary, B:405, 407
proboscidea, *Uvigerina*
 Site 754, B:32
 relative abundance, B:41
 Site 756, B:38, 71
 relative abundance, B:56
prolixa, *Protentella*, B:114
 Site 758, B:97, 123
pschadae, *Globotruncanella*, B:107
pseudobulloides, *Morozovella*, B:113
 Site 758, A:383–384
Pseudoeumotia doliolus Zone, Site 758, A:386, B:191
pseudogrosserugosus, *Anomalinoidea*, Site 752, B:26
pseudolinneiana, *Marginotruncana*, B:107
pseudomenardii, *Planorotalites*, B:114
 Site 758, B:95
pseudoperlucidus, *Cibicoides*, Site 752, B:27
 Maestrichtian–Eocene, B:5
pseudoplicata, *Brizalina*, Site 756, B:71
Pseudopodosira sp., Site 752, B:180, 183
pseudotopilensis, *Acarinina*, B:112
pseudoumbilica, *Reticulofenestra*, Site 753, last occurrence, A:175
punctata, *Globocassidulina*, Site 754, B:9
puncticulata, *Globorotalia*
 Site 753, first occurrence, A:177
 Site 754, first occurrence, A:202
punctulata, *Heterohelix*, B:104
 Site 754, B:89, 367
pusilla, *Brizalina*, Site 756, B:71
pusilla, *Morozovella*, B:113
pyramidata, *Gaudryina*, Site 752, B:29
Pyramidina sp., Site 752, B:25
- Pyxilla gracilis* Zone, Site 752, A:130, B:175, 727
- quadrangula*, *Mesocena*, Site 758, last occurrence, B:191
quadrangula, *Rouxia*, Site 758, B:207
quadratas, *Gyroidinoidea*
 Site 752, B:28
 Site 754, B:6
quadratus, *Lithraphidites*
 first occurrence, B:143
 Site 752, B:144, 413
 Site 758, first occurrence, A:381
quadripunctata, *Prediscosphaera*
 Brazos River, Texas, A:512
 El Kef, Tunisia, A:512
 Site 752, Cretaceous/Tertiary boundary, B:406
Quadrium trifidum Zone, Site 758, B:149
quaternarius, *Monomarginatus*, Site 758, B:163
quinqueloba, *Globigerina*, B:110–111
 Site 758, B:121
quinqueloba, *Pullenia*, Site 756, B:73
- radiata*, *Trochosira*, Site 752, B:183
rajagopalani, *Heterohelix*, B:104
recurvus, *Isthmolithus*
 Site 754, first occurrence, B:733
 Site 757, A:318
reflexa, *Globocassidulina*, Site 756, B:75
reinhartii, *Cyclagelosphaera*
 Site 752, B:412
 Site 758, Cretaceous/Tertiary boundary, B:405
Reinhardtites anthophorus Zone, Site 755, B:145
Reinhardtites levis Zone, Site 758, B:149
reinholdii, *Nitzschia*, Site 758, A:386
 last occurrence, B:191
renzi, *Planulina*
 Site 754, B:74
 Site 756, B:33
resurgens, *Uvigerina*, Site 754, B:6
reticulata, *Stephanopyxis*, Site 758, B:208
Rhizosolenia praebergonii Zone, Site 758, A:386, B:191, 194
ribbingi, *Cibicides*, Site 752, B:27
riedellii, *Heliolithus*, Site 752, first occurrence, A:133
riograndensis, *Planoglobulina*, B:104
robertsonianus, *Cibicoides*, Site 754, B:72
Rocella gelida Zone, Site 758, A:387
Rocella vigilans Zone, Site 758, B:198
rohri, *Truncorotaloides*, B:114
romeinii, *Biscutum*, Site 752, comparison, Maud Rise species, A:512
rosetta, *Globotruncana*, B:107
Rossiella paleacea Zone, Site 758, A:386, B:195
rostrata, *Bulimina*, Site 756, B:71
ruber, *Globigerinoidea*
 Site 754, A:201
 Site 756, A:270
 Site 758
 abundance, B:132
 coarse fraction association, B:130, 135
 Cretaceous/Paleogene boundary, B:489
 dissolution trend, B:133
 productivity of, B:126
Rugoglobigerina spp.
 Site 752, Cretaceous/Tertiary boundary, A:507, 509
 Site 754, A:204
rugosa, *Globotruncana*, B:107
rugosa, *Rugoglobigerina*, B:108
 Site 752, B:86–87
 Site 758, B:119
 last occurrence, A:384
Rzehakina sp., Site 758, A:386
- sacculifer*, *Globigerinoidea*, B:111
 Site 709
 carbon isotopes, B:244, 248–249
 oxygen isotopes, B:243, 248–249
 Site 758
 abundance, B:132
 carbon isotopes, B:245–246, 329–339
 coarse fraction association, B:130, 135
 Cretaceous/Paleogene boundary, B:489
 dissolution trend, B:133, 309
 oxygen isotopes, B:245–246, 329–339
 productivity of, B:126
saepes, *Neochiastozygus*, Site 752, first occurrence, A:133
saipanensis, *Discoaster*
 Site 754, last occurrence, B:733
 Site 756, A:270
 Site 757, A:318
 last occurrence, A:316
saturalis, *Orbulina*, Site 753, first occurrence, A:178
sawamurae, *Coscinodiscus*, Site 758, B:205
Sceptroneis spp., Site 752, B:180
schencki, *Uvigerina*, Site 756, B:71
schulzii, *Stephanopyxis*, Site 758, B:208
schulzii, *Triceratium*, Site 758, B:201, 208
scoticus, *Hyalodiscus*, Site 752, B:183
scotus, *Acuturris*, Site 752, B:413
sellii, *Globigerina*, B:111
 Site 758, A:383
semicostata, *Heterohelix*, B:104
semicibratus, *Anomalinoidea*
 Site 752, B:26
 Site 754, B:72
 Site 756, B:33
senni, *Globigerina*, B:111
septenarius, *Lithastrinus*, Site 758, B:169
sicana, *Praeorbulina*, Site 757, A:319
sigmoides, *Zygodiscus*, Site 752, A:512, B:412
 Cretaceous/Tertiary boundary, B:405–406
simulatilis, *Morozovella*, B:113
sissinghii, *Quadrum*
 first occurrence, B:143
 Site 754, B:166
 Site 758, last occurrence, B:149
Skeletonema sp., Site 752, B:180, 183
soldadoensis, *Acarinina*, B:112
 Site 757, A:320
soldanii, *Gyroidina*
 Site 754, B:74
 Site 756, relative abundance, B:57
sparsus, *Biantholithus*, Site 752, B:405
 Cretaceous/Tertiary boundary, B:407, 731
 first occurrence, A:127, 133, 509, 511, B:402
 stratigraphic range, B:396
speciosus, *Gladius*, Site 758, B:208
spectabilis, *Spiroplectammina*, Site 752, B:25
spinosa, *Prediscosphaera*, Site 758, B:169
spinosa, *Thalassiosira*, Site 758, B:205
spinosa, *Vulvulina*, Site 756, B:74
spinulosa, *Uvigerina*, Site 756, B:32
spiralis, *Globigerina*, B:111
spissiformis, *Anomalinoidea*, Site 752, B:26
spumellaroides, *Thalassiosira*, Site 758, B:205
stella, *Macrora*, Site 758, B:196
Stellarima sp., Site 752, B:180, 183
Stensioina beccariiiformis assemblage, Site 752, B:9, 14
 Paleocene decrease, B:19
Stephanopyxis, Site 752, B:264
Stilostomella, Site 752, A:129
stokesianus, *Cestodiscus*, Site 758, B:205
stoveri, *Prediscosphaera*
 Maud Rise, A:512

- Site 750, B:395
 Site 752, B:413
 comparison, Site 690, B:408
 Cretaceous/Tertiary boundary, B:405-406
striata, *Rectuvigerina*, Site 754, B:71
 relative abundance, B:42
stuarti, *Globotruncanita*, B: 107
stuartiformis, *Globotruncanita*, B:107
subacutus, *Hemiaulus*, Site 758, B:206
subbotinae, *Morozovella*, B:113
 Site 752, A:128
subcarinata, *Pullenia*, Site 756, B:73
subcarinatus, *Globigerinelloides*, B:109
subcircummodifer, *Rugotruncana*, B:108
 Site 752, B:86, 119
 Site 754, B:119
subconglobata, *Globigerinathea*, B:111
subglobosa, *Globocassidulina*
 Site 753, A:179
 Site 754, B:32, 75
 Site 756, relative abundance, B:56
subspinosa, *Stilostomella*, Site 754, B:75
subspiratus, *Cibicidoides*, Site 752, B:27
 stratigraphic range, B:9
subtriangularis, *Chiloguembelina*, B:102
 Site 753, B:120
sulcata, *Paracassidulina*, Site 754, B:75
surculus, *Discoaster*, Site 753, first occurrence,
 A:175
suteri, *Globorotaloides*, B:112
 Site 758, B:122
suturalis, *Orbulina*, Site 754, first occurrence,
 A:202
symmetrica, *Rossiella*, Site 758, B:207
- tamalis*, *Discoaster*, Site 756, A:268
tapuriensis, *Globigerina*, B:111
teniculum, *Prinsius*, Site 752, first occurrence,
 A:512
tenuicarinata, *Siphonina*, Site 756, B:32
teniculum, *Prinsius*, Site 752, B:411
 Cretaceous/Tertiary boundary, B:407
tenuis, *Cruciplacolithus*
 Site 752, B:411
 Cretaceous/Tertiary boundary, B:405, 407
 first occurrence, A:128, 133, 512, B:730
 Site 758, A:380
tesselatum, *Triceratium*, Site 752, B:181, 184
teuriensis, *Hornibrookina*
 Broken Ridge, A:481
 Site 752, B:411
Thalassiosira grunowii group, Site 758, A:386
Thalassiosira sp., Site 752, B:183
Thalassiosira yabei group, Site 758, A:386
Thalassiosiropsis sp., Site 752, B:181
thalmanni, *Brizalina*, Site 756, B:71
Thoracosphaera, Site 752, A:512
Thoracosphaera spp., Site 752, Cretaceous/Tertiary
 boundary, B:405-407
topilensis, *Truncorotaloides*, B:114
tosaensis, *Globorotalia*
 Site 753, first occurrence, A:177
 Site 754, A:201
Towieus sp., Broken Ridge, A:481
trabeculatus, *Helicolithus*, Site 755, B:145, 167
Tranolithus phacelosus Zone
 Site 754, B:145
 Site 758, B:149
triangularis, *Bulimina*, Site 752, B:25
triangularis, *Globigerina*, B:111
Triceratium sp.
 Site 752, B:184
 Site 758, B:208
Triceratium tessellatum Zone, Site 752, A:130, 488
- Trifarina* sp., Site 756, B:71
trifidum, *Quadrum*
 first occurrence, B:143
 Site 758, B:166
trigonula, *Guttulina*, Site 752, B:29
trilobus, *Globigerinoides*, B:111
 Site 753, A:178
 Site 757, B:121
 Site 758, Cretaceous/Paleogene boundary,
 B:489
triloculinoides, *Globigerina*, B:111
 Site 758, B:95
trinidadensis, *Morozovella*, B:113
 Site 758, A:383-384
trinitatensis, *Bulimina*, Site 752, B:6, 25
tripartite, *Globigerina*, B:111
 Site 758, A:383
Trochosira sp., Site 752, B:183
trochus, *Cestodiscus*, Site 758, B:204
truepyi, *Nuttallides*, Site 752, A:129, B:9, 26
 Maestrichtian–Eocene, B:5
truncanus, *Cibicidoides*, Site 752, B:9, 26
truncatulinoides, *Globorotalia*, Site 754, first oc-
 currence, A:201
tumida, *Globorotalia*
 Site 757, first occurrence, A:319
 Site 758, A:382
 abundance, B:132
 coarse fraction association, B: 130, 135
 Cretaceous/Paleogene boundary, B:489
 dissolution trend, B:127, 130, 133
Turborotalia, B:114
turris, *Stephanopyxis*, Site 752, B:271
turrieffelii, *Eiffellithus*, Site 758, B:167
tuxpamensis, *Bulimina*
 Site 754, B:71
 relative abundance, B:41
 Site 756, B:71
tympaniformis, *Fasciculithus*, Site 752, first oc-
 currence, A:133
- umbilica*, *Reticulofenestra*, Site 756, A:270
umbonatus, *Oridorsalis*
 Site 752, B:6
 Site 756, B:73
 relative abundance, B:57
umbonifera, *Epistominella*
 Site 754
 Antarctic Bottom Water association, B:50
 corrosive water effects, B:51
 Site 756, B:33, 73
 Antarctic Bottom Water association, B:50
 corrosive water effects, B:51
 relative abundance, B:56
uncinata, *Morozovella*, B:113
unicavus, *Catapsydrax*, B:109
universa, *Orbulina*, Site 758, A:382
Uvigerina proboscidea assemblage, Site 756,
 B:43, 68
 Miocene development, B:50
Uvigerina sp.
 Site 752, B:25
 Site 754, B:71
 Indian Deep Water association, B:43
 Site 756
 Indian Deep Water association, B:43
 relative abundance, B:56
uwula, *Globigerinita*, B:111
- varians*, *Remesella*, Site 752, B:29
variospira, *Globorotalia*, B:113–114
variospira, *Morozovella*, B:113–114
 Site 758, B:95, 113, 123
velascoensis, *Bulimina*, Site 758, A:386
- velascoensis*, *Cibicidoides*
 Site 752, B:27
 Site 758, A:386
velascoensis, *Globigerina*, B:111
 Site 752, last occurrence, B:727
velascoensis, *Morozovella*, B:114
 Site 757, A:320
 Site 758, A:383
velascoensis, *Osangularia*, Site 752, last occur-
 rence, B:6
velatus, *Hemiaulus*, Site 752, B:186
venezuelana, *Globigerina*, B:111
venezuelana, *Globoquadrina*
 Site 709
 carbon isotopes, B:244, 248-249
 oxygen isotopes, B:243, 246, 248–249
 Site 758
 carbon isotopes, B:245
 oxygen isotopes, B:245-246
veniamini, *Bogorovia*
 first occurrence, B:198
 Site 758, B:207
 last occurrence, B:195
ventricosa, *Globotruncana*, B:107
vigilans, *Rocella*, Site 758, B:207
volutus, *Globigerinelloides*, B:109
 Site 754, B:119
 Site 758, B:119
- walfischensis*, *Rosita*, B:107
walli, *Cibicidina*
 Site 752, B:6, 27
 Site 754, A:205
wilcoxensis, *Acarinina*, B:112
 Site 752, B:122
wilcoxensis, *Chiloguembelina*, B:102
 Site 752, B:120
wilcoxensis, *Pseudohastigerina*, B:115
wilcoxensis, *Trifarina*, Site 752, B:25
woodi, *Zeaglobigerina*, B:111
wuellerstorfi, *Planulina*
 Site 753, A:179
 Site 754, B:32, 74
 Indian Deep Water association, B:43
 relative abundance, B:42
 Site 756, B:33
 Indian Deep Water association, B:43
 Site 758, A:385
- yaucensis*, *Globigerinelloides*, B:109
 Site 758, B:119
- Zones
 CC14, Site 755, A:244
 CC14–CC16, Site 755, B:735
 CC15–CC16, Site 755, A:244
 CC19–CC22, Site 758, A:382
 CC21–CC22, Site 758, B:199
 CC22–CC23 boundary, Site 754, A:200
 CC22–C25, Site 758, A:381
 CC23–CC24, Site 754, B:733
 CC23–CC25, Site 754, A:200
 CC24, Site 752, A:121, 127
 CC24–CC23b, Site 754, A:195
 CC24–CC25, Site 754, B:367
 CC24–CC26, Site 752, B:731
 CC25, Site 754, B:6
 CC25–CC26, Site 752, A:121, 127
 CN1
 Site 753, B:732
 Site 754, A:200
 Site 756, A:270
 CN1/CN2 boundary, Site 754, A:199
 CN2–CN4, Site 709, B:242

TAXONOMIC INDEX

- CN2–CN9, Broken Ridge, A:479
 CN3–N4, Site 754, A:199
 CN3–N12, Site 756, A:268, 270
 CN4, Site 753, A:175
 CN5, Site 757, A:316
 CN5/CN6, Site 755, A:242
 CN5–N6, Site 754, A:199
 CN8–N11, Site 753, A:175
 CN9b–CN10a, Site 754, A:199
 CN13
 Site 755, A:242
 Site 756, A:268
 CN19a, Site 754, A:195
 CP1, Site 752, A:512
 CP1–CP3, Site 758, A:380
 CP1–CP8, Site 752, B:728
 CP1–CP10, Site 752, A:481
 CP2/CP1 boundary, Site 752, A:127
 CP2/CP3 boundary, Site 752, B:730
 CP3, Site 752, A:127
 CP3–P10, Site 752, A:126
 CP4–CP8
 Site 752, B:5
 Site 758, A:376, 378
 CP8–CP14, Site 757, A:316
 CP10–CP12, Site 752, B:727
 CP13–15, Broken Ridge, A:481
 CP13–CP15
 Site 753, B:732
 Site 754, B:733
 CP13c, Site 753, A:176
 CP14–CP15, Site 754, A:200
 CP15
 Site 752, A:480, B:725–726
 Site 754, A:480
 CP15–CP18
 Site 754, B:733
 Site 756, A:270
 CP15b, Site 754, A:195
 CP17–CP19, Site 758, A:376
 CP18, Site 754, B:733
 CP19
 Site 752, A:123, 480
 Site 754, A:200, 480
 N4
 Site 755, B:89
 Site 757, A:319
 N8–N12, Site 757, A:319
 N18–N21, Site 758, A:382
 P1, Site 758, B:97
 P1–P4, Site 758, A:383–384
 P1–P5
 Site 752, B:86
 Site 758, B:95
 P6, Site 757, A:320
 P6–P8, Site 757, B:92
 P10–P14
 Site 752, B:85–86
 Site 753, B:87
 P10–P17, Site 757, B:92
 P12–P13, Site 752, B:89
 P12–P14, Site 753, B:732
 P13–P14
 Site 752, B:725–726
 Site 754, A:204
 P13–P19, Site 754, B:733
 P14, Broken Ridge, A:477
 P17, Site 757, A:320
 P17–P20, Site 755, B:90
 P18–P19, Site 757, A:319–320
 P18–P21, Site 757, B:90, 92
 P19, Site 758, A:383
 P19–P22, Site 758, B:95
 P21
 Site 754, B:88
 Site 755, B:89–90
 P22
 Site 753, B:87, 732
 Site 755, B:89–90
 P22/N4
 Site 752, B:85
 Site 754, B:87–88, 733
 Site 757, B:90
 Site 758, B:95