

INDEX TO VOLUMES 127 AND 128

This index provides coverage for the *Initial Reports* portions of Volumes 127 and 128 of the *Proceedings of the Ocean Drilling Program* (published as separate leg-specific books) and for the *Scientific Results* portion of those *Proceedings* (published as Volume 127/128, Parts 1 and 2). The reports in Parts 1 and 2 of the *Scientific Results* are not leg-specific but are arranged by subject matter. References to page numbers in the *Initial Reports* are preceded by "A," followed by the appropriate leg reference in parentheses, and then a colon: "A(127)"; those in the *Scientific Results* are preceded by "B" with a colon.

The index was prepared by Wm. J. Richardson Associates, Inc., under subcontract to the Ocean Drilling Program. The index contains two hierarchies of entries: (1) a main entry, defined as a keyword or concept followed by a reference to the page on which that word or concept appears, and (2) a subentry, defined as a further elaboration on the main entry followed by a page reference.

The index is presented in two parts: (1) a Subject Index and (2) a Taxonomic Index. Both parts cover text figures and tables but not core-description forms ("barrel sheets") or core photographs. Also excluded are bibliographic references, names of individuals, and routine front and back matter.

The Subject Index follows a standard format. Geographic, geologic, and other terms are referenced only if they are subjects of discussion. This index also includes taxonomic entries above the generic level, as well as broad fossil groups such as foraminifers and radiolarians. The notation "ff" following a page listing indicates that reference to a topic continues beyond the last page given but is not sequential. This would be the case where one or more figures or tables follow a principal topic of discussion that makes up a major section. A site chapter in the *Initial Reports* is considered the principal reference for that site and is indicated on the first line of the site's listing in the index. A parenthetical reference is made to the appropriate leg. Such a reference to Site 798, for example, is given as "Site 798, A(128):121-236."

The Taxonomic Index is an index relating to significant findings and/or substantive discussions, not of species names *per se*. This index covers three varieties of information: (1) individual genera and species that have been erected or emended formally, (2) biostratigraphic zones, and (3) fossils depicted in illustrations. A taxonomic entry consisting of both genus and species is listed alphabetically by genus and also by species. Biostratigraphic zones are listed alphabetically by genus; zones with letter prefixes are listed under "zones."

For further information, including available electronic formats, contact the Chief Production Editor, Ocean Drilling Program, 1000 Discovery Drive, College Station, Texas 77845-9547, U.S.A.

SUBJECT INDEX

- AACT. *See* aluminum, logging; geochemical logging
- acanthostyles
Site 795, B:542
- acanthoxeas
Site 795, B:542
- acanthstrongyles
Site 795, B:542
- accelerometer, three-axis feedback-type. *See* downhole seismic experiment
- acoustic impedance
opal-A to opal-CT transition, B:1146
Site 799, B:1137-1138, 1140
- acoustic velocity. *See* velocity
- ACT. *See* aluminum, logging
- AF demagnetization. *See* magnetic properties
- age dating. *See* biostratigraphic zonations; biostratigraphy; radiometric age
- age spectra
Site 794, B:820, 828-832
Site 795, B:820-821, 832-833
Site 797, B:820, 828-829
- Aikawa section, diatom abundance, B:316
- air-gun profiling experiment, B:1109-1114, 1117-1118
- Aira-Tn, tephra marker, B:796
- Akita Basin, lithostratigraphy, A(128):17
- Akita oil field, benthic foraminifers, B:381
- Akita-Yamagata oil field basalt
Site 794 basalt compared, B:918, 1339
Site 797 basalt compared, B:921
- albitization. *See also* feldspar
and calcium concentration, B:137
occurrence modes, B:136
provenance information destroyed, B:148
of sandstone, B:136-137
silica source, B:148
and sodium in interstitial water, B:137
and sulfate formation, B:143-144
temperature, B:136
- alginite
Site 798, B:429
Sites 798-799, B:670
- alkaline feldspar. *See* feldspar, alkaline
- alkaline glass, B:1378, 1387
- alkalinity
and carbonate precipitation, A(127):204
and diagenetic carbonate, B:94
and dolomitization, A(127):362
in interstitial water, B:1264
at Japan Sea sites, B:1265
and organic carbon decomposition, B:1262
and organic matter degradation, A(127):362
Site 794, A(127):108
Site 795, A(127):204
Site 796, A(127):279
Site 797, A(127):362, 368
Site 798, A(128):173, 181
Site 799, A(128):317-318, 328; B:610
and sulfate reduction, A(127):279
vs. calcium, A(128):183
vs. sulfate, A(128):183
- alkane-alkene doublets, Site 799, B:627
- n*-alkanes
Site 799, A(128):324, 343
Sites 798 and 799, B:669-671
- alkenones, Sites 798 and 799, B:669-670
- alteration
ash layers, A(127):109-110, A(128):151, 288; B:1267, 1375-1378
- augite clinopyroxene, B:885
- basalt, B:838, 884-888, 899, 901
- basement, A(127):122-125, A(128):88-91, 321; B:849, 883-889, 891, 908-916, 1265-1267, 1282, 1338
- basement rock and adjacent pore water elemental variations correlated, B:911
and element composition, B:799
and element mobility, B:838, 892, 908-916
glass shards, A(127):95
groundmass, B:885-886
hydroclastic shards, A(128):89-90
and liquid line of descent calculations, B:911
and magnesium enrichment, B:842
olivine, B:885
plagioclase feldspar, B:884-885
and rare earth element mobility, B:911-916
secondary minerals, B:838
Site 794, B:908-911, 916
Site 795, B:918
Site 797, B:911
temperature, B:839
timing, B:838-839
and XRF loss on ignition, B:892-893, 909
- alteration halo, around fractures, A(128):91
- alternating field demagnetization. *See* magnetic properties
- Altyn-Tagh Fault, offset along, B:1325
- aluminum
in ash layers, B:1380
in augite, B:851-853, 856
in dark/light cycles, B:569
in Japan Sea sediment, B:1236-1237
and parent magma, B:856
Site 794, B:654, 656
Site 795, B:706
Site 796, B:654, 656
Site 798, B:654, 656, 722
Site 799, B:654, 656, 722
vs. chromium, B:733
vs. iron, B:732
vs. iron/magnesium ratio, B:894
vs. lanthanum, B:737
vs. ytterbium, B:737
XRF and geochemical log analyses compared, B:1028-1029, 1031-1035
- aluminum, logging
clay, A(127):306-307
core data correlated, B:1030
diatom ooze, A(127):306-307
diatom vs. clay content, B:1417
opal-A to opal-CT transition, B:22
sand, A(127):306
Site 794, B:1416, 1420-1421
Site 796, B:1416, 1423
Site 797, B:7, 1417, 1426-1427
Site 798, A(128):212, 215; B:1030, 1401-1402
Site 799, A(128):362; B:1404, 1408-1409
vs. potassium, A(128):215
- aluminum activation clay tool (AACT). *See* aluminum, logging; geochemical logging
- aluminum clay tool (ACT). *See* aluminum, logging
- aluminum oxide. *See* aluminum
- aluminum/silica ratio, vs. terrigenous content, B:1035
- ammonia
in clay minerals, A(127):204, 362
and organic matter degradation, A(127):204, 362
Site 794, A(127):108
- Site 795, A(127):204
Site 797, A(127):362, 368
Site 799, B:610
- ammonium
in clay minerals, A(127):279-280
in interstitial water, B:1264
and organic carbon decomposition, B:1262
and organic matter degradation, A(127):279-280
Site 796, A(127):279-280
Site 798, A(128):173, 181
Site 799, A(128):317-318, 328
- amorphous silica. *See* opal-A; silica, biogenic
- amphibole, in ash layers, B:1379
- Amur River, and Japan Sea, B:69-70, 442
- Amurian microplate, A(128):9
- anaerobic bacterial degradation. *See* methanogenesis; sulfate reduction
- analcime, Site 797, B:139, 143
- andesite
ash layers, B:793
microphotograph, B:859
rare earth elements, B:924
Site 795, B:850, 854, 1338
spidergrams, B:924
spinel composition, B:844
- andesite, basaltic
brecciated, A(127):221-222
plagioclase pyroxene phyrlic, A(127):174, 217
rare earth elements, B:924
Site 795, B:918, 1338
spidergrams, B:924
- anhydrite
replacing plagioclase, B:148
sandstone replacement mineral, B:143
- Aniai-type flora, Japan Sea, B:249, 486
- anisocheles
Site 795, B:543
- anisotropy
in oceanic crust, B:1110-1115
in velocity measurements, A(127):229
- ankerite
and Mossbauer parameters, B:741
Site 796, A(127):266
Site 799, B:82
- anomalies. *See* free-air gravity anomalies; magnetic anomalies
- anorthite
Sites 794 and 796, B:126
vs. orthoclase, B:122
- anoxic conditions. *See* oxygenation conditions
- anoxic-nonsulfidic sediments, Site 799, B:612
- antimony
Site 795, B:711, 714
Site 798, B:1370-1371
Site 799, B:724-725
- aperture, vertical, in logs, B:396-397
- Ar/Ar age. *See* radiometric age
- arc tholeiite suite, Site 795, B:1338
- arenite. *See* felspathic arenite; lithic arenite; sandstone
- argon/argon age. *See* radiometric age
- argon isotopes. *See* isotopes, argon
- aridity signal, in logs, A(128):188-192
- arkose. *See* sandstone
- aromatic species, Site 799, B:627
- arsenic
Site 794, B:1362
Site 795, B:711, 714, 1365
Site 797, B:1366

- Site 798, B:1368-1369
 Site 799, B:724
 ash. *See* ash layers
 ash beds. *See* ash layers
 ash layers, B:1373-1393. *See also* glass, volcanic; tephrochronology; volcanic activity; volcanic phases; volcanic pulses; *specific volcanic rocks*
 acidic, A(128):151, 158, 287, B:793
 age, A(128):123, 155-156, 288, 300; B:963
 alteration, A(127):22, 109-110, 363-364, A(128):151, 288; B:701, 1267, 1375-1378
 alteration effects, B:641, 643, 646, 701
 aluminum vs. silica oxide composition, B:1380
 areal distribution, B:791
 Aso-4 ash, B:581
 barium anomalies, B:1377-1378
 basic, A(128):151, 285-286; B:793
 and bioturbation, A(128):153
 boron content, B:639-640
 in Brunhes Chron sediments, B:963
 calcium, A(127):280; B:1383
 color, A(128):151
 core/log correlation, B:398, 401, 1025
 and core recovery, B:1319-1320
 crystal fraction, A(128):150; B:791
 in dark/light cycles, B:581
 definition, A(128):150
 erosion, A(128):153
 feldspar composition, B:1388
 in FMS logs, A(128):153, 184; B:398, 400
 frequency, A(127):95-97, 190, 193, 346, 350, A(128):157, 301; B:1313
 frequency underestimation with depth, B:1375
 frequency vs. age, B:1319-1320
 frequency vs. thickness, A(128):159, 300
 frequency vs. time, B:794, 798
 glass fraction, A(128):150
 graded, A(127):186, A(128):151-153, 285; B:793, 802
 grain size, A(128):150; B:791
 heterogeneous, A(128):153, 285; B:793, 802
 homogeneous, A(128):151-152, 285; B:793, 801
 intermediate composition, B:793
 and interstitial water chemistry, B:1265
 iron vs. silica oxide composition, B:1381
 Japan Sea, A(128):32-33; B:1316, 1319-1321
 lanthanum vs. ytterbium, B:1386
 lithology, B:791
 in logs, A(128):288, 339-340
 magmatic composition, B:793
 magnesium content, A(127):280; B:1265, 1382
 magnetic susceptibility correlated, A(127):105, 199
 mineral fragment composition, B:1379
 neodymium vs. zirconium, B:1386
 origin, A(128): 156-157, 291-292
 and oxygen isotopes, B:52
 petrography, A(128):150-153, 284-288
 potassium vs. silica oxide composition, B:1385
 pyroxene composition, B:1388
 secondary mineralogy, B:1375-1378
 sedimentary types, B:793
 sedimentation processes, A(128):151-153, 284-288
 seismic stratigraphy, A(128):122, 125, 153-155, 344-351, 379
 silica diagenetic evolution, B:1375-1377
 Site 794, A(127):90; B:1374
 Site 795, A(127):186; B:1374
 Site 796, A(127):251, 264; B:1374
- Site 797, A(127):341; B:1373-1374
 Site 798, A(128):33, 35, 122-124, 137, 139-140, 150-157, 196-198, B:397-398, 791-793, 963, 1374
 Site 799, A(128):33, 240, 256, 284-291, 296-297, B:605, 791-793, 963, 1373
 Sites 798 and 799 compared, A(128):291-292, 301
 slumps, A(128):285
 sodium vs. silica oxide composition, B:1384
 sources, A(128):122, 198; B:791
 strontium, A(127):280; B:607
 synsedimentary/post-sedimentary processes, A(128):153
 on synthetic seismogram, A(128):184
 tephra markers, A(128):157, 291-292; B:793-796, 798-799
 thickness, A(128):157, 301; B:791, 1319
 thickness underestimation with depth, B:1375
 thickness vs. deposition age, B:963
 thickness vs. time, B:795, 799
 thorium/uranium ratio, A(128):339
 thorium vs. zirconium, B:1386
 and turbidity flows/slumps, A(128):153-154; B:793
 volcanic activity, B:800
 volcanic pulses, B:796-797
 XRF analyses, A(128):151
 Asia, sandstone provenance, B:110-111
 Aso-4, B:581, 796
 Aso volcano, ash layers from, A(128):157
 asthenosphere, injection of, B:903
 Atlantic II Fracture Zone, cation exchange capacity, B:1282
 Atlantic Ocean N, calcareous nannofossils, B:173
 Atlantic Ocean NE, paleoenvironment, B:338
 Atterberg limits, Sites 798 and 799, B:1124, 1126
 augite
 altered to clay, B:885
 anhydrous pressure, B:871
 augite-liquid relations, B:870-872
 chemical composition, B:851-853, 867-868
 and cooling rate, B:852, 870-871
 crystallization, B:870-872
 experimental chemical compositions, B:867-868
 hypersthene-normative, B:864
 in melting experiments, B:861-862
 Mg/(Mg+Fe) histograms, B:855
 minor element content, B:854
 natural vs. experimental augite compared, B:870-871, 878-879
 and parent magma, B:854, 856
 Site 794, B:851-852
 Site 795, B:852
 Site 797, B:852-853
 sodium content, B:858
 titanium, B:855-856
 back-arc rifting
 Japan Basin N, A(127):238
 Japan Sea, A(127):16, 258, A(128):11
 mechanisms for, A(127):5
 Site 794, A(127):84, A(128):82-83
 Site 795, A(127):179-180
 Site 796, A(127):258
 Site 797, A(127):336
 Yamato Basin S, A(127):336-337
 back-arc tholeiite. *See* tholeiite, back-arc
 bacteria. *See also* microbiological studies
 acetogens, B:766-768
 aerobic ammonifiers, B:766-768
 anaerobic, B:767, 770-771
 anaerobic ammonifiers, B:766-768
- anaerobic viable count, B:772
 and bishomohopanoid acid, B:669
 chemistry, B:765
 contamination concerns, B:757-758
 in dark/light cycles, B:772
 in deep aquifers, B:755, 759
 depth distribution of dividing and divided cells, B:766
 and diagenetic processes, A(128):178-179
 direct counts, B:756, 765-766
 distribution, B:769-771
 dormant, B:759
 filamentous, B:757-758
 frequency of dividing and divided cells (FDDC), B:757-758, 765-766, 769, 772
 growth, B:755-760
 H-thymidine incorporation, B:757, 759, 767, 769, 771-773
 heterotrophs, B:766-768
 importance, B:761
 incubation, B:756, 764
 under metabolic stress, B:759
 and methanogenesis, B:767-769, 772-773
 methods of study, B:755-756, 761-765
 nitrate reducers, B:766-768
 Peru margin, B:755, 757, 759, 761, 769, 773
 potential activity, B:763, 767-769
 presence at depth, B:761
 productivity, B:758-759
 rates of bacterial activity, B:771-772
 sampling times, B:756
 ship-shore transportation, B:762
 shipboard handling, B:761-762
 shore lab handling, B:762-765
 Site 798, A(128):125-126; B:755-776
 sulfate reducers, B:766-768, 770, 772-773
 and temperature, B:761
 total anaerobic, B:767
 total count, B:758, 766-767, 772
 viability, B:759, 763-767, 773
 Bagnold effect, and vein fillings, B:1180
 Baja California, diagenetic carbonates, B:75
 Barbados forearc, veins observed, B:1180
 Barbados Ridge Complex, and chloride, A(128):172
 barite
 in ash layer, B:1392
 authigenic precipitation, B:665
 crystals, B:664
 energy dispersive spectra, B:1378
 estimated ion activity product (IAP) nodules, B:637-638
 replacement mineral in sandstone, B:143
 Site 795, B:640
 and sulfur isotopes, B:637-638
 barium, B:651-665
 abundance, B:663
 accumulation rate, B:660
 authigenic precipitation, B:654
 barium front, B:637, 663
 basement, A(128):99
 biogenic, B:653, 656, 659
 burial rates, B:652
 C_{org} barium ratio, B:653-654, 661
 detrital vs. biogenic component, B:657
 diagenetic redistribution, B:652
 dissolution, B:659, 661
 in interstitial water, B:657-659
 mass accumulation rates, B:665
 paleoproductivity indicator, B:651-656
 preservation, B:652
 and productivity, B:651-656, 661-662

SUBJECT INDEX

- remobilization, B:654, 656-659
schematic geochemical diagram, B:664
and sediment accumulation rate, B:660-661
Site 794, B:653-656, 1362
Site 795, B:710, 1365
Site 796, B:654, 656
Site 797, B:1366
Site 798, B:654, 656, 1368-1369
Site 799, B:654, 656, 724-725
and volcanic glass alteration, B:1377-1378
vs. sedimentation rate, B:661
vs. yttrium, A(128):100
vs. zirconium, A(128):100
barium accumulation rate. *See* barium
basalt. *See also* basement; *specific minerals in basaltic rocks*
AFM diagram, B:874
alteration, B:669-701, 838, 883-889, 891, 899, 901, 908-916
altered veins, B:886-887
augite-liquid relations, B:870-872
calc-alkalic evolution trend, B:870, 875
chemistry, B:839-842
and chromium spinel, B:837-847
crystallization history, B:853-854
experimental and core data compared, B:870
fractionation models, B:872-874
grain size, A(127):370
lava flows, B:819
liquid line of descent, B:869-882
major elements, B:869-870
neodymium isotopes, B:901-903
oxide variations, B:869-870, 873
parent magma, B:855-856
petrography, B:892
phase equilibria, B:870
pseudoternary normative projections, B:875
sills, B:819
Site 794, B:869
Site 795, B:1338
Site 797, B:869, 899-900, 911
strontium isotopes, B:901-903
thickness of altered layer, B:699-701
tholeiitic evolution trend, B:870, 875
vesicles, B:887-888
basalt, alkali, Site 797, B:920, 1339-1340
basalt, aphanitic, Site 797, A(127):345-346
basalt, aphyric
alteration, B:885, 887
microphotograph, B:859
radiometric age, B:822-823, 825
Site 794, A(128):68-69, 90-91; B:853, 1339
Site 797, A(127):369-370, 375-380; B:850, 911
basalt, augite phyrlic, Site 795, B:854
basalt, back-arc, isotopes, B:805
basalt, calc-alkaline
Site 795, B:919
Site 797, B:1339-1340
basalt, doleritic, Site 794, A(128):68, 86, 88
basalt, high-Al, B:893
Site 794, B:861
Site 797, B:861, 900, 920, 1339-1340
basalt, island-arc, isotopes, B:805
basalt, mid-ocean ridge (MORB), and helium isotopes, B:747
basalt, olivine phyrlic
Site 794, B:849, 853
Site 797, B:850, 854-857
basalt, olivine plagioclase phyrlic, Site 794, B:908
basalt, olivine-pyroxene microphyric, Site 794, A(128):68, 90
basalt, olivine tholeiitic, liquid line of descent, B:864
basalt, phyrlic pyroxene plagioclase, Site 795, B:918
basalt, plagioclase olivine phyrlic, Site 797, A(127):369
basalt, plagioclase phyrlic
crystal accumulation, B:853-854
radiometric age, B:823, 826
Site 794, B:849, 853
Site 795, A(127):174, 217; B:850, 854
Site 797, A(127):370-371, 380; B:850, 854
basalt, pyroxene phyrlic, Site 795, B:850
basalt, pyroxene plagioclase phyrlic
radiometric age, B:826-827
Site 795, A(127):174, 217
basalt, tholeiitic
Site 794, B:899, 916
Site 797, B:900, 1339-1340
basement. *See also* fractionation models; mantle source; parent magma; petrology, experimental; radiometric age; X-ray fluorescence (XRF) analyses; *specific minerals in basement rocks*; *specific rock types*
age, A(127):22, 107, 192-193, 195, 361, A(128):69; B:821, 906-908, 1333-1337
age, determination methods, B:1303-1304
age, lithospheric stretching models, B:1304-1305
alteration, A(127):23, 122-125, 217-219, 280, 369-380, A(128):68-69, 321; B:779, 849, 883-889, 891, 908-916, 1265-1267, 1282, 1338
alteration effects, B:641-643, 646
alteration mineralogy, B:884-888
arc tholeiite suite, B:1338
baked contacts, A(127):369-380
boron as alteration product, B:639
brecciation, B:1296
calc-alkaline suite, B:1338
chemistry, A(127):24-25, 123, 126, 174, 219, 223, 325, 381-383, A(128):68-69, 96ff, 98; B:779-789, 805-817, 839, 872, 892-894, 905-916, 1338-1340
chilled margins, A(127):369-380
compared to MORBS and BABBS, A(127):126, 219
crustal anisotropy, B:1107-1121
depth, B:1120
drilling rubble, Site 794, A(127):125
electrical resistivity, B:1354
emplacement, A(127):381
eruptive history, A(127):174
fractionation models, B:872-874
fractures, A(127):217, A(128):69
grain size, A(128):90-91
groundmass, A(128):88-91, B:849-859, 892
igneous event sequence, A(128):95-97
interlayered sediments/igneous rocks, A(127):369
isotopes, B:779-789, 805-817, 884-888, 1338-1341
Japan Basin N, A(127):169, 176-177
Japan Sea, A(127):24-25, 29, A(128):67, 32-33
Kita-Oki Bank, A(128):127-129
Kita-Yamato Bank, A(128):247-249
Legs 127/128 Site 794 rocks correlated, A(128):93-95
logging and logs, A(127):28, A(128):86, 93-95, 103-104; B:1282
Lower Volcanic Complex (LVC), Site 794, B:780
magma genesis, B:1340-1341
magma suites recognized, B:869-870
magmatic affinity, A(128):96ff
magnetic intensity, A(127):174
magnetic polarity, A(128):86; B:936-938
magnetic properties, B:933-945
magnetic susceptibility, A(127):174
magnetics, A(127):203, 209, 358-359, A(128):101-102
major elements, B:869-870, 893
Middle Volcanic Unit (MVU), Site 794, B:780
mineral composition, B:850-853
N-MORB normalized composition, B:1338
nature of unit contacts, A(128):68, 88-91
neodymium isotopes, B:786-787
normalized composition, A(127):25, 126, 223, 383
Oki Islands, A(128):127-129
Oki Ridge, A(128):127-129
oxide variation, B:869-870
parent magma, B:855-856, 894
petrography, A(128):86-91; B:849-850, 891-892
phenocrysts, B:849-859, 892
physical properties, A(127):26, 135, 139-140, 227-228, A(128):102ff; B:1282-1284, 1289-1291
primary mineralogy, B:891-892, 908
radiogenic isotopes, B:807
radiometric age, B:819-836
secondary mineralogy, B:891, 908
seismic expression, A(127):28-29, 143, 234-236, 312-313
sills and flows, A(127):126-127; B:853-855
Site 794, A(127):73, 120-127, A(128):32-33, 36, 67-69, 86ff, B:849, 883-889, 891-894, 906, 927, 933, 1112-1113, 1336, 1339, 1354
Site 795, A(127):174, 217ff; B:849, 906, 933, 1336, 1338-1339
Site 797, A(127):325, 369-383; B:849, 891-894, 906-908, 933, 1336, 1339-1340
spidergram, B:923-924, 926
stratigraphy, B:871, 901
strontium isotopes, B:786-787
structural features, A(128):91-93; B:1181-1184
Takuyo Bank, A(128):248-249
tectonic significance, B:1337-1341
temperature, A(127):137
thermal conductivity, B:1281
thickness of altered layer, B:699-701
trace elements, B:807
unit boundaries correlated, A(128):93-95
units, Site 794, A(127):73, 121-125, A(128):68-69, 75, 86-91; B:779-781, 884, 907-908
units, Site 795, A(127):174, 217-219, 221; B:909
units, Site 797, A(127):369-380; B:910
Upper Volcanic Complex' (UVC), Site 794, B:780
velocity vs. porosity, A(127):140
X-ray fluorescence analyses, B:892-894
Yamato Bank, A(128):247-249
Yamato Basin, A(127):72, A(128):23-24, 32-33
Yamato Basin N, A(127):77-81, A(128):73-75, 117
Yamato Basin S, A(127):331-334
Yamato Rise, A(128):247-249

- basin. *See* pull-apart basin; *particular basins*
- bedding
 in blue tuff, B:117
 color-banded, A(127):186, 189
 dips, Site 799, B:1176, 1178
- Bering Sea, opal flux, B:439
- beryllium isotopes. *See* isotopes, beryllium
- BHTV. *See* borehole televiewer
- bicarbonate, and alkalinity, A(128):173
- biochronology synthesis, B:1219–1228
- bioclastic sand. *See* sand, bioclastic
- biosilica. *See* silica, biogenic
- biosiliceous sediments
 laminated, B:439–455
 permeability, B:1128
- biostratigraphic zonations, B:1225–1226. *See also* biostratigraphy; microfossils; *names of specific microfossils*
- foraminiferal and diatom zones correlated, B:222–223
- foraminifers, B:200–201
 Japan Sea, A(127):20
 of onshore exposures, A(128):17
 Site 794, A(127):98, 100; B:344–345
 Site 795, A(127):196–197; B:346–347
 Site 796, A(127):271–272
 Site 797, A(127):353; B:348–349
 Site 798, A(128):161
 Site 799, A(128):304–305
- biostratigraphy, B:1219–1228. *See also* biostratigraphic zonations; microfossils; *names of specific fossil groups*
- age of oldest sediments, B:1335–1337
- datum levels used, B:1222
- high-resolution study, B:193–200
- Japan Sea, A(127):19–22, A(128):28–30
- Leg 127, A(127):19–22
- Leg 128, A(128):28–30
- Legs 127 and 128 correlated, A(128):29
- Neogene foraminifers, B:188–189
- problems with, B:1219
- radiometric ages compared, B:821–822, 1337
- Site 794, A(127):72, 96–103, A(128):99–100; B:187–224, 1221–1223, 1225
- Site 795, A(127):174, 192–199; B:187–224, 1223, 1225
- Site 796, A(127):250–251, 269–274; B:187–224, 1223, 1225
- Site 797, A(127):324, 351–357; B:187–224, 1223, 1225
- Site 798, A(128):28–30, 124–125, 158–166; B:155–169, 1223–1226
- Site 799, A(128):28–30, 244, 298–312; B:155–169, 1224, 1226
- biotite, in ash layers, B:1379, 1392
- bioturbated sediments. *See* sediments, bioturbated
- bishomohopanoid acid, Sites 798 and 799, B:669, 671
- Black Sea
 oxygenation conditions, B:705
 pyrite, B:712
 trace metals, B:1361
- Blake Outer Ridge, carbon isotopes, B:88
- blooms, Site 798, A(128):165
- blue tuff
 age of Yamato seamount chain samples compared, B:126
 clay minerals, B:119
 deposition, B:119, 123–126
 lithology, B:117
 petrography, B:117–119
 plagioclase, B:124–126
- pyrogenic crystal chemistry, B:116, 121–123, 126
- radiometric age, B:116, 123, 126–127
- sedimentology, B:115–120
- Site 794, B:120
- Site 796, B:121
- Sites 794 and 796, B:115–130
- stratification, B:129
- strontium isotopes, B:116, 123, 127
- Yamato Basin and Japan Basin samples compared, B:127–128
- boninite, spinel composition, B:844
- borehole diameter, and geochemical logging errors, B:1029–1030
- borehole televiewer
 fractures, A(127):307, 399
 hole ellipticity, A(127):307
 Site 796, A(127):307
 Site 797, A(127):399, 409
 stress field determination, B:1047
- boron
 adsorption/desorption processes, B:640–641
- and ash layer alteration, B:639–640
- and basement alteration, B:639, 1266
- and gas hydrate, B:639
- in interstitial water, B:638–642
- and organic matter, B:640
- and sand transport, B:640
- Site 794, B:638–641
- Site 795, B:638–641
- Site 796, B:638–641
- Site 797, B:638–641
- vs. chloride, B:642
- vs. magnesium, B:642
- boron isotopes. *See* isotopes, boron
- Boso Peninsula, magnetostratigraphy, A(127):199
- bottom simulating reflector (BSR)
 age, B:1155–1156
- causes, B:1145–1147
- and heat flow, B:1147–1148
- and opal-A to opal-CT transition, B:1149
- polarity, B:1147
- and sedimentation rate, B:1147–1148
- Yamato Basin, B:1155
- Bottom Water, B:577
- bottom water oxygenation conditions. *See* oxygenation conditions
- breccia, tuff. *See also* tuff, rhyolitic
- in logs, A(128):287
- Site 799, A(128):243, 287; B:34
- breunnerite, Site 799, B:82
- brittle structures
 Site 794, B:1181–1183
- Site 798, B:1181–1182
- Site 799, B:1182–1183
- broadband seismometer. *See* downhole seismic experiment
- bromine
 Site 798, B:1368–1369
- Site 799, B:724–725
- Brunhes Chron
 and ash layers, B:963
- and diatom occurrence, A(128):159
- and inclination variations, B:960
- and sediment accumulation rate, B:468
- and sedimentation rate, B:961
- silicoflagellate and ebridian zones correlated, A(128):162
- Site 796, A(127):275
- Site 797, B:973
- Site 798, A(128):124
- warm/cold intervals in, B:459, 464
- Brunhes/Matuyama boundary
- and diatom occurrence, A(128):161
- field behavior during polarity transition, B:970, 972, 974, 978
- and foraminiferal coiling direction, B:208
- and hematite, B:972
- inclination variations, A(128):172
- multiple reversals at, A(127):22
- sedimentation rate, B:961
- silicoflagellate and ebridian zones correlated, A(128):305
- Site 794, B:218, 224, 970, 1221
- Site 795, A(127):199; B:972, 1223
- Site 797, B:218, 224, 973
- Site 798, A(128):30, 156, 165, 167; B:164, 563, 973, 1224
- Site 799, A(128):314; B:974, 1224
- BSEM images, faunal/floral morphotaxa recognized, B:548
- bulk density. *See* density
- Burma earthquake, and downhole seismic experiment, B:1162, 1168
- burrows
 in cyclic lithofacies, A(128):140–141
- in dark/light cycles, B:564–565, 569
- with organic carbon fillings, B:38
- and oxygenation conditions, A(128):141; B:1232
- Site 795, A(127):189
- Site 796, A(127):265–266
- Site 797, A(127):344, 346, 348–349
- Site 798, A(128):143–145
- Site 799, A(128):262
- iso-butane
 Site 798, A(128):125, 176, 187
- Site 799, A(128):244–245, 322
- n*-butane
 Site 798, A(128):125, 176, 187
- Site 799, A(128):244–245, 322
- C/N ratio
 in color-banded beds, A(127):365–367
- in dark/light cycles, A(127):212; B:431
- Site 794, A(127):113–114, 119
- Site 795, A(127):211–212, 216; B:708–709
- Site 796, A(127):283–285, 287
- Site 797, A(127):365–367, 375
- Site 798, A(128):189–193; B:425–426, 435–437
- Site 799, A(128):324, 334–338, 341; B:427, 429
- C/S ratio
 and oxygenation condition, B:1241, 1249
- Site 794, B:1246–1247, 1255
- Site 796, A(127):283
- Site 798, A(128):189–192
- Site 799, A(128):324, 334–338, 342; B:623–624
- Ca/K atomic ratio, and radiometric dating of blue tuff, B:123
- calc-alkalic evolution trend
 in basalt, B:870, 875
- and suppression of plagioclase crystallization temperature, B:873
- calc-alkaline glass, B:1378, 1387
- calc-alkaline suite, Site 795, B:1338
- calcareous nannofossils. *See* nannofossils, calcareous
- calcite
 abundance in Japan Sea sediments, B:1235–1241
- in dark/light cycles, B:588
- isotope data, B:888
- Site 795, A(127):188
- Site 799, A(128):289; B:611

- in veins, B:887
- calcite, diagenetic
 chemical composition, B:81
 formation depth, B:85
 isotopic composition, B:83
 occurrence, B:80
 replacing orthoclase, B:147
 in sandstone, B:141–143, 146
 shallowest occurrence, A(128):273
 Site 798, A(128):137
 Site 799, A(128):272; B:75–98
- calcium
 and albitization, B:137
 alteration and, B:909, 911
 and ash alteration, A(127):280, 363–364
 in ash layers, B:1383
 and basement, A(127):280, B:892
 in dark/light cycles, B:569
 and dolomite formation, A(127):363
 in interstitial water, B:607, 1263, 1268
 Japan Sea, B:1236, 1265
 mobility during alteration, B:838
 Site 794, A(127):109–110
 Site 795, A(127):174, 205
 Site 796, A(127):280
 Site 797, A(127):363–364, 370
 Site 798, A(128):173–174, 182; B:1368–1369
 Site 799, A(128):318, 329; B:611
 sources of, B:143
 vs. alkalinity, A(128):183
 vs. magnesium, A(127):115
 vs. strontium isotopes, B:645
 XRF and geochemical log analyses compared,
 B:1028–1029, 1031–1035
- calcium, logging
 Site 794, B:1420–1421
 Site 796, B:1423
 Site 797, B:1426–1427
 Site 798, B:1024
 Site 799, A(128):364; B:1408–1409
- calcium/aluminum ratio, Site 794, B:1236
- calcium carbonate (CaCO₃). *See* carbon, inorganic; carbonate content
- calcium oxide. *See* calcium
- California
 planktonic foraminifers, A(128):165
 silicoflagellate, B:237
- caliper logging
 in dolomite, A(128):332–333
 and lithology, A(127):306
 Site 794, A(127):142, 147
 Site 796, A(127):303–306
 Site 797, A(127):395–396, 403
- calthrops
 Site 795, B:543
- carbon, inorganic
 Japan Sea, A(127):23
 Site 794, A(127):113
 Site 795, A(127):209–212
 Site 796, A(127):283–285
 Site 797, A(127):365–367
 Site 798, A(128):125, 176–177, 189–192
 Site 799, A(128):323–324, 334–338
- carbon, organic. *See also* organic matter; Rock-Eval analyses; sediments, organic-rich
 accumulation rate, B:425, 428, 660
 and alkalinity, B:1262
 and ammonium, B:1262
 in burrows, B:38
 and carbon isotopes, B:1262–1263
 C_{org}/barium ratio, B:653–654, 661
 and cyclic lithofacies, A(128):125, 177
 in dark-colored beds, A(127):283, 365
- in dark/light cycles, B:431–432, 568–569, 579, 586–588, 594, 671
 decomposition, B:1262–1263
 and diagenetic carbonate, B:92
 environment needed for accumulation, B:429–430
 flux rate, B:423–437
 glacial/interglacial variation, B:589
 and gray value, B:589
 and increased productivity, A(127):283
 Japan Sea, A(127):24
 mass accumulation rate, B:425, 428
 and oceanographic interpretations,
 A(127):368–369
 and phosphate, B:1262
 and productivity, B:430
 and sedimentation rate, B:628
 shipboard vs. shorebased analyses compared,
 B:623, 628
 Site 794, A(127):113–114, 119; B:656, 1246–1247, 1255
 Site 795, A(127):209–212, 216; B:706–710, 714
 Site 796, A(127):251, 283–285, 287; B:656
 Site 797, A(127):325, 365–367, 375; B:592
 Site 798, A(128):32, 125, 176–177, 189–194, 196; B:423–426, 435–437, 656, 668–669, 776
 Site 799, A(128):32, 238–239, 245, 323–324, 334–338, 340; B:38, 423–437, 623–627, 629, 656, 668–669
- and sulfate, B:1262
 vs. biogenic barium, B:659
 vs. free hydrocarbons (S_i), B:631
 vs. nitrogen, A(128):343–344
 vs. sulfur, A(128):343–344; B:594, 628, 712–713, 1241, 1254
- carbon, total organic (TOC). *See* carbon, organic; Rock-Eval analyses
- carbon isotopic ratios. *See* isotopes, carbon
- carbon/nitrogen ratio. *See* C/N ratio
- carbon/sulfur ratio. *See* C/S ratio
- carbonaceous siltstone. *See* siltstone, carbonaceous
- carbonate. *See also* carbonate, biogenic; carbonate, diagenetic
 in dark/light cycles, B:584
 diagenesis, B:1263–1265, 1267
 glacial/interglacial variation, B:590
 in Japan Sea sediments, B:1235–1241
 logged thickness vs. cored thickness, B:79
 in logs, B:78–79
 mass accumulation rate, B:428
 nodules, B:611–612, 615, 1232
 occurrence, B:78–79
- carbonate, authigenic. *See* calcite, diagenetic; carbonate, diagenetic; dolomite; siderite
- carbonate, biogenic
 diagenetic phases, A(127):267
 preservation, A(127):169
 recrystallization, A(128):318
 Site 799, B:78
- carbonate, diagenetic. *See also* carbonate diagenesis
 abundance and distribution, B:80, 82
 and calcareous nannofossils, B:166
 carbon isotopes vs. depth, B:85–90
 chemical composition, B:81–85
 compositional zoning, B:83
 disseminated rhombs, B:78
 equilibrium oxygen isotope values, B:83–85, 89
- formation conditions, A(128):277; B:83–87, 89–90, B:888
 and inorganic geochemical data, A(128):276–277
 interstitial water chemistry and, B:646
 isotopic composition, B:83
 layers, B:78
 in logs, B:77
 nodules, B:78
 occurrence, B:78
 oxygen isotopes, B:86–88
 paleoenvironmental implications, B:93–95
 petrography, B:81
 provenance, B:148
 replacing quartz, B:147
 and revised lithostratigraphy, B:1232–1233
 in sandstone, B:141–143
 SEM photographs, B:97–98
 Site 796, A(127):266–267
 Site 798, A(128):147–150; B:701–702
 Site 799, A(128):240, 260, 272–278, 292, 354; B:38, 75–98, 697, 701–702
 stratigraphically controlled distribution, B:76
 textural relations, B:81
 thickness, B:77
 X-ray diffraction analyses, B:79–80, 86–87
- carbonate-compensation depth (CCD)
 and benthic foraminifers, B:504–505
 and calcareous nannofossils, B:1220
 Japan Sea, A(127):24, A(128):20; B:155, 208, 440, 1201
 and microfossil preservation, A(127):19
 Oki Ridge, B:409
 and sedimentation rate, B: 1228
 Site 794, A(127):103, 113, 147
 Site 795, A(127):169, 199, 211
 Site 796, A(127):247, 283, 315
 Site 797, A(127):324, 356, 365, 409
 Site 798, A(128):122, 158; B:168
 Site 799, A(128):292
- Yamato Basin, A(127):103
- carbonate content
 core analyses, B:404
 in dark/light cycles, B:431–432, 579, 586–588
 logging estimates, B:402, 404
 Site 794, A(127):113–114, 119
 Site 795, A(127):209–212, 216
 Site 796, A(127):283–285, 287
 Site 797, A(127):365–367, 375; B:592
 Site 798, A(128):32, 189–194; B:401, 435–437, 445–446, 668–669
 Site 799, A(128):32, 245, 307, 323–324, 334–338, 340; B:668–669
- carbonate diagenesis. *See also* carbonate, diagenetic and interstitial water oxygen isotopes, B:701–702
 and magnesium, A(128):174
 material balance calculation of carbonate precipitation, B:702
- carbonate minerals
 formation, A(127):22–23
 and iron, B:722–723
- carbonate nodules. *See* nodules, carbonate
- carbonate preservation
 Site 794, B:188
 Site 795, A(127):211
 Site 797, B:188
- carboxylic acids
 Site 798, B:674
 Sites 798 and 799, B:669–671
- Cariaco Basin, oxygenation conditions, B:705
- cation exchange capacity (CEC), and clay alteration minerals. B:1282

- CCD. *See* carbonate-compensation depth
- Ce anomaly
and diagenetic fractionation, B:692-693
and dolomite, B:92
and hydrogenous precipitation, B:731
and hydrothermal activity, B:719, 729
and lanthanum/ytterbium ratios, B:693-694
and light rare earth elements, B:693-694
and oxygenation conditions, B:692
and sediment contamination of mantle source, B:815
Site 794, B:682, 693
Site 795, B:683
Site 797, B:688-691, 693
Site 798, B:726, 735
Site 799, B:90-91, 729, 736
vs. Eu anomaly, B:737
vs. lanthanum/ytterbium ratio, B:737
vs. manganese content, B:737
- Ce/Ce. *See* Ce anomaly
- celadonite
in vesicles, A(127):217
and volcanic glass alteration, B:1375
- cerium. *See also* Ce anomaly
distribution, B:692
and hydrogenous manganese precipitation, B:726-727
Site 794, A(128):101
Site 795, B:707
Site 798, B:1368-1369
Site 799, B:90-91
- cesium
Site 795, B:707
Site 798, B:1368-1369
Site 799, B:724
- Chenopodiaceae
Site 794, B:491
- chert
cycles in FMS logs, B:1041-1044
in FMS logs, A(127):307; B:1039-1040, 1232
in logs, A(127):27-28, 302, 394
and revised lithostratigraphy, B:1232
in sandstone, B:144-148
seismic expression, A(127):407-408
Site 795, A(127):187-188
Site 796, A(127):265
Site 797, A(127):343-344, 349
- China
dust storms, B:394
loess deposits, B:394
subcontinental lithosphere, B:811-812
- chloride
and clay minerals, B:608-610
distribution, B:605
and opal-A, B:608-610
Site 795, A(127):205
Site 796, A(127):278-279
Site 798, A(128):172, 179
vs. boron, B:642
- chlorine
and geochemical logging, B:1023-1024
Site 794, A(127):107
Site 797, A(127):364, 374
Site 799, A(128):317, 328; B:610
- chlorite
in Japan Sea sediments, B:1235-1241
in sandstone, B:139
as sandstone cement, B:134
Site 798, B:411, 416, 418
- chlorite/saponite mixed-layer clay
from altered materials, B:884-886, 888
chemical composition, B:885-887
and isotopes, B:888
- in sandstone, B:139
Site 794, B:885
in veins, B:886-887
X-ray diffraction spectra, B:886
- chlorophyllinite, Sites 798 and 799, B:669-670
- chromatography. *See* pyrolysis-gas chromatography
- chromite, B:840. *See also* spinel; spinel, chromium
- chromium
basement, A(128):99
Site 795, B:706
Site 797, B:920
Site 798, B:1368-1369
from terrigenous sources, B:723
vs. aluminum, B:733
vs. yttrium, A(128):101
- chromium/barium ratio, basement, A(128):99
- chromium spinel. *See* spinel, chromium
- Chron 5, Site 798, A(128):30
- Chron 6
Site 796, A(127):277
Site 798, A(128):30
- Chugoku Backbone Range, uplift, B:526, 528
- Cichorioideae
Site 795, B:491
- clathrate. *See also* gas hydrate
Site 796, A(127):251, 278, 281, 288-289
- clay
aluminum, logging, A(127):306-307
and revised lithostratigraphy, B:1229-1230
Site 794, A(127):90-92, A(128):77, 81
Site 795, A(127):186
Site 797, A(127):340-341
Site 798, A(128):124, 137
Site 799, A(128):256
- clay, coatings of, B:57
- clay, diatom
and revised lithostratigraphy, B:1230-1232
Site 794, A(127):90
Site 795, A(127):186
Site 796, A(127):261-264
Site 797, A(127):341-343
Site 798, A(128):124, 137-138
Site 799, A(128):256, 260
- clay, siliceous silty, and revised lithostratigraphy, B:1232
- clay, silty
in dark/light cycles, B:568
and revised lithostratigraphy, B:1229-1232
Site 794, A(127):90, A(128):77
Site 795, A(127):186
Site 796, A(127):261-264
Site 797, A(127):340-341
Site 798, A(128):124, 137-138
Site 799, A(128):256
- clay content
and density, B:1278-1279
and gamma ray logging, B:6
- clay minerals. *See also* clay minerals, diagenetic; *names of specific clay minerals*
abundance in Japan Sea sediments, B:1235-1244
in altered materials, B:885-888
and ammonium, logging, A(127):279
and basement alteration, B:908-909, 918
in blue tuff, B:119
chemical composition, B:885-887
compaction structures, B:39
crystallization, A(128):320-321
D/H ratios, interstitial water, B:642
in dark/light cycles, B:569
in diagenetic sequence, B:39
- fabric, B:35-36, 38
and gamma-ray logging, A(127):393
glacial/interglacial variation, B:588
and low-chloride interstitial fluids, B:608-610
major element molar ratios, B:611
and rare earth elements, B:691
as sandstone cement, B:134-135
SEM photographs, B:30
Site 794, A(127):94
Site 798, B:411, 416, 418
Site 799, B:611
- clay minerals, diagenetic. *See also* clay minerals; *names of specific clay minerals*
in sandstone, B:139-141
Site 799, B:35-36
- clayey diatom ooze. *See* ooze, clayey diatom
- claystone. *See also* pelite
and revised lithostratigraphy, B:1232-1233
sediment accumulation rate, A(127):191
Site 794, A(127):92-93, A(128):81
Site 795, A(127):189
Site 796, A(127):264; B:1260
Site 797, A(127):343-344
Site 798, A(128):124, 138
Site 799, A(128):265; B:41-43
- claystone, dark, deposition, A(127):267
- claystone, diatom
Site 795, A(127):186-187
Site 796, A(127):264
- claystone, pebbly, Site 796, A(127):264-265
- claystone, siliceous
grain shape, B:38
grain size, B:38
and revised lithostratigraphy, B:1230-1232
sediment accumulation rate, A(127):191
Site 794, B:1258
Site 795, A(127):187-189
Site 796, A(127):264-265
Site 797, A(127):343-344
Site 798, A(128):124, 138
Site 799, A(128):260, 264-265; B:39-43
in vitric tuff, B:34
- claystone, silty
and revised lithostratigraphy, B:1232-1233
Site 794, A(127):93-94
Site 797, A(127):343, 345-346
Site 799, B:43
- claystone, silty dolomitic, Site 799, B:41
- claystone, silty phosphatic, Site 799, B:43
- claystone, silty siliceous, Site 799, B:40-41
- claystone, spicular silty, Site 797, A(127):343
- claystone, tuffaceous, Site 794, A(128):81
- climate
Japan Sea, B:180, 337
pollen and, B:320
precipitation and, B:326
radiolarians and, B:299
Site 796, A(127):274
Site 797, A(127):356
vegetation and, B:325-326
- clinoptilolite
abundance, B:1235-1244
in ash layers, B:1390
and opal-A to opal-CT transition, B:10
SEM photographs, B:30, 1294
Site 796, B:139
Site 797, A(127):344
- clinopyroxene
in ash layers, B:1379
in melting experiment, B:895
- cluster analysis, Japan Sea, B:525-526, 531, 535
- CNT-G. *See* porosity, logging
- cobalt

- Site 795, B:710
 Site 798, B:1368-1369
 Site 799, B:724
 coccolithophores
 Site 799, A(128):310
 coccoliths
 Site 799, A(128):307, 309
 Cochiti Subchron, Site 797, B:1223
 coiling direction. *See also* coiling ratio; foraminifers, planktonic
 of planktonic foraminifers, A(128):165, 310-311
 as surface-water temperature indicator, B:1220
 coiling dominance (CD) curve, B:459. *See also* coiling direction; coiling ratio; foraminifers, planktonic
 Site 798 coiling ratios correlated, B:460
 Site 798 warm/cold events correlated, B:463-465
 coiling ratio. *See also* coiling direction; coiling dominance (CD) curve; foraminifers, planktonic
 of N Pacific foraminifers, B:459
 oxygen isotopes correlated, B:467
 as paleoclimate indicator, B:463, 467-468
 of planktonic foraminifers, B:188, 193, 208, 210-215, 218-219
 collosphaerids
 Japan Sea, B:296
 color-banded bedding. *See also* dark/light cycles; lithofacies, cyclic
 in clay and silty clay, A(127):341
 in diatom-bearing claystone, A(127):261
 and organic carbon content, A(127):365
 origin, A(127):19, 24, 349
 Site 797, A(127):324, 343
 color-banding, in dark/light cycles, B:581
 compaction structures, Site 799, B:38, 39, 48
 compensated neutron tool (CNT-G). *See* porosity, logging
 composition, sediment. *See* sediment composition
 compression
 along Hokkaido W-Honshu N thrust belt, A(128):76-77
 and fracture formation, B:1181
 compressional stress field, Japan Sea and, B:529-531
 compressional velocity. *See* velocity
 conductivity model, Japan Arc, A(128):85
 conglomerate, of porcellanite pebbles, A(127):344-345
 conifers
 Japan Sea, B:327-328
 Japanese archipelago, B:334
 Site 798, B:318-319
 connate brackish water, and interstitial water analyses, B:607-608
 consolidation
 and Atterberg limits, B:1124
 in debris flows and slumps, B:1125
 definitions used, B:1123
 determined from in situ pore water pressure, B:1124
 excess pore-water pressure, B:1125-1127, 1129
 and gassy sediments, B:1129
 and leakage from artesian water or gas pressure source, B:1129-1130
 normally consolidated/underconsolidated sediment boundary, B:1124
 from oedometer tests, B:1125-1127
 overburden pressure, B:1125-1127, 1129
 overconsolidation ratio, B:1125-1127, 1129
 preconsolidation pressure, B:1125-1127, 1129
 processes causing, B:1123
 and sedimentation B:1124-1125, 1129-1130, 1277
 shear strength/overburden pressure ratio, B:1125-1126
 Site 798, B:1125-1127
 Site 799, B:1125, 1127
 underconsolidation, B:1129-1130
 void ratio vs. effective pressure, B:1126, 1128
 convergence
 along Hokkaido W-Honshu N thrust belt, A(128):76
 Japan Basin E, A(127):258
 copper
 Site 794, B:1362
 Site 795, B:1365
 Site 797, B:1366
 in sulfide deposits, A(128):21
 core/log correlation
 and ash layers, B:398, 401
 and resistivity logging, B:398-400, 402
 Site 798, B:1024-1025
 correlation coefficients
 for core XRF and geochemical log oxides, B:1029
 in FMS data, B:1040-1041, 1046
 cristobalite. *See* opal-CT
 cross correlation, of FMS logs, B:1046
 crest, lower, structure, B:1342-1343
 crust, oceanic
 electrical resistivity structure, B:1351
 Japan Basin, A(127):73
 Yamato Basin, A(127):73, B:928
 crustal anisotropy, and regional stress field, B:1114-1115
 crustal extension, in Yamato Basin, B:1345-1346
 crustal structure
 3.5 km/s layer, meaning of, B:1342
 4.5 km/s layer thickness variation, B:1120
 anisotropy, B:1110-1115, 1343
 electrical resistivity, B:1343
 and heat flow, B:1302-1303
 Honshu N shelf, A(128):73
 Japan Arc, A(128):74
 Japan Basin, A(127):9, 79, 176, A(128):9-11, 70-72, 74; B:924, 1075, 1083, 1086, 1104, 1107, 1304-1305, 1311-1314, 1341-1346
 Japan Basin E, A(127):251-252
 Japan Basin N, A(127):176
 Japan Sea, A(128):6-7, 9-11; B:1075, 1083, 1104, 1107, 1311-1314, 1333, 1341-1346
 Japan Sea E, A(128):9-11
 Japan Trench, A(128):74
 Kita-Yamato Bank, A(128):248
 Kita-Yamato Trough, A(128):245-249
 lateral heterogeneity, B:1110-1112
 lower crest, B:1342-1343
 and ocean bottom seismometer data, B:1075
 Oki Ridge, A(128):127-128
 Oki Trough, A(128):130
 Okushiri Ridge, A(127):251-252
 and paleobathymetry, B:1201
 and plate reconstruction geometrical constraints, B:1311-1314
 Takuyo Bank, A(128):249
 and tectonic history, B:1318-1319
 and temperature, B:1343
 Tsushima Basin, B:1311-1314
 velocity structure, B:1078-1083
 Yamato Bank, A(127):330, A(128):248
 Yamato Basin, A(127):9, 79, 176, A(128):9-11, 24, 70-73, 127-128, 130, 248; B:837, 899, 1083, 1086, 1104, 1107, 1304-1305, 1311-1314, 1341-1346
 Yamato Basin N, B:1075-1106
 Yamato Basin S, A(127):329-330, 333; B:1083, 1104-1106
 Yamato Rise, A(127):9, 329-330, 333, A(128):9-11, 245-249; B:1075
 Yamato Seamount Chain, A(128):128
 crystallization age, Site 794, B:821
 Cupressaceae
 Site 797, B:490
 cut fluorescence. *See* fluorescence, cut
 cycladophorids
 Site 794, B:307
 cyclopentane, Site 799, A(128):321-322
 D-phosphate, Site 799, B:38-39
 dacite, ash layers, B:793
 Daijima-type flora, Japan Sea, B:249, 486
 dark/light cycles, B:1221. *See also* lithofacies, cyclic
 and bacteria, B:772
 biogenic component and detrital grain size compared, B:584, 588
 biogenic opal abundance, B:432
 burrows, B:564-565, 569
 C/N ratio, B:431
 carbon and sulfur analyses, B:579
 carbonate content, B:431-432, 586-588
 characterization, B:563
 clay minerals, B:569
 color, B:581
 core continuity and depth corrections, B:578
 correlations at Site 798, B:562-563
 dark layers, B:580, 584, 591-594, 597-601
 density, B:568
 depositional environment, B:590-594
 and dropstones, B:581
 excess silica content, B:1241, 1253
 first-order rhythms, B:446-447, 559, 563
 geochemistry of second-order cycles, B:569
 gray value, B:570-571, 579, 581-584, 586-587
 hydrogen index, B:431-432
 and interbedded ash layers, B:581
 isotopic analyses, B:569-571
 and Japan Sea circulation, B:574-575
 layer thickness and organic carbon content correlated, B:594
 major and minor elements, B:568, 570-571
 microfossils, B:568
 and Milankovitch cycles, B:446-447, 571-575
 and molecular organic fossils, B:670, 675
 and monsoons, B:574-575
 nature of basal boundaries, B:569
 nitrogen content, B:431
 occurrence, B:562-563
 opal content, B:446, 569
 organic carbon, B:430-432, 568, 586-588, 671
 organic matter, B:446, 564, 668-669
 origin of cyclicity, B:577-601
 origin of detrital component, B:594-595
 oxygen index, B:431-432
 oxygen isotopes, B:431, 588
 and oxygenation conditions, B:432, 590-591
 paleoceanographic significance, B:574-575
 paleoenvironmental interpretation, B:419-421
 productivity, B:431-432, 591-594
 and revised lithostratigraphy, B:1229-1232
 second-order rhythms, B:446, 559, 563-569
 sediment composition, B:579, 582-584
 sedimentological observations, B:569

- Site 795, B:1257
 Site 797, B:574
 Site 798, B:393, 421, 446, 458, 559-576
 Site 799, B:571-574
 Sites 794, 795, and 797 cycles correlated, B:577-601
 Sites 797-799 compared, B:574
 structure/composition, B:564-565
 sulfur, B:586-588
 synthetic sequence, Site 798 drillholes, B:566-567
 thickness, B:581
 third-order rhythms, B:559, 563
 trace metals, B:1361
 X-ray diffraction analyses, B:579, 584-588
- dark/light sediment cycles. *See* dark/light cycles; lithofacies, cyclic
- debris flow
 consolidation, B: 1125
 seismic expression, B:1140-1141, 1143
 Site 799, A(128):267
- declination. *See* magnetic properties
- deformation, soft-sediment
 and lithologic boundaries, A(128):142ff
 Site 798, A(128):141ff
 Site 799, A(128):241
- deformation rate analysis (DRA), and stress field, B:1048-1049
- density. *See also* physical properties
 acoustic impedance, B:1137-1138
 basement, A(127):227
 and clay content, B:1278-1279
 corrected analyses, B:990-1015
 in dark/light cycles, B:568
 data correction methods, B:987
 data quality, A(127):127, 228-229
 and diatom content, B:1278-1279
 errors in wet volume measurement, B:985-986
 Japan Sea, B:1278-1281
 and loss on ignition, B:1283
 and opal-A to opal-CT transition, B:18-19, 21, 1146-1147, 1278-1279, 1288
 Site 794, A(127):127-134, A(128):102-103, 105; B:1050, 1280-1281
 Site 795, A(127):222-228, 230, 233; B:1280-1281
 Site 796, A(127):290-295; B:1280-1281
 Site 797, A(127):383-390; B:1280-1281
 Site 798, A(128):180-181, 197-198, 202-205, 208-210; B:1280-1281
 Site 799, A(128):325-326, 345-349, 354-356; B:1137-1138, 1140, 1280-1281
 vs. loss on ignition, B:1291
 vs. opal abundance, B:402
 vs. porosity, B:25, 27, 986, 989, 1281, 1290
 vs. water content, A(127):233, 395
- density, logging
 Site 794, A(127):139-142, 144-146
 Site 795, A(127):232, 237
 Site 796, A(127):302-305, 319-320
 Site 798, A(128):185-187; 212, 229-232; B:399, 401
 Site 799, A(128):332-338, 362-364, 383-385, 389-392
 vs. gamma ray, logging, A(128):215
- density flow, of blue tuff, B:123-126
- deposition rate. *See* opal flux; sediment accumulation rate; sedimentation rate; terrigenous flux
- depositional cycles. *See* lithofacies, cyclic
- desiccation, as problem for visual microanalysis, B:57
- detrital accumulation rate. *See* terrigenous flux
- detrital minerals
 eolian origin, B:1238-1239
 in Japan Sea sediments, B:1241-1249
 deuterium isotopes. *See* isotopes, deuterium
- dewatering
 Site 798, B:1176-1178
 Site 799, B:1178
- diachrony, species, of calcareous nannofossils, B:164
- diactinal monaxons
 Site 795, B:542
- diatom-bearing clay. *See* clay, diatom-bearing
- diatom clayey siltstone. *See* siltstone, diatom clayey
- diatom claystone. *See* claystone, diatom
- diatom number. *See also* diatoms
 in dark/light cycles, B:579, 588
 Site 797, B:593
- diatom ooze. *See* ooze, diatom
- diatomaceous clay. *See* clay, diatom
- diatoms. *See also* biostratigraphic zonations; biostratigraphy; microfossils, siliceous; opal-A; silica, biogenic
 abundance and opal-A to opal-CT transition, B:1230-1232
 abundance and preservation, A(127):97, 192-193, 271-272, 351, A(128):28, 159-161, 299-303; B:312, 314, 344-349, 352, 593, 1219
 abundance and Tsushima Current inflow correlated, B:591
 age constraints, A(127):271, A(128):69; B:1335
 BSEM images, B:554
 in carbonate concretion, A(127):351
 in dark/light cycles, B:568, 584
 datum levels, B:251, 258-259
 and density, B:1278-1279
 diachronous occurrences, B:309
 dissolution, B:309-316, 341-342, 353, 1220
 foraminifers correlated, B:188-189, 222-223
 freshwater, A(128):249
 frustule structure, A(127):137
 internal porosity, B:1277
 Japan Sea, B:363
 Kita-Yamato Bank, A(128):249
 Leg 127, A(127):19-20
 Leg 128, A(128):28
 Miocene/Pliocene boundary, A(127):97, 193, 272, 351
 morphotaxa recognized in BSEM images, B:548
 and opal-A to opal-CT transition, B:3
 as paleoenvironmental indicator, B:341-342
 Pliocene/Pleistocene boundary, A(127):97, 351
 productivity, A(127):274, A(128):122; B:342
 Quaternary/Pliocene boundary, A(127):192, 271
 reworked, A(127):97, 192-193, 273; B:250
 and sediment velocity, A(127):133
 SEM photograph, B:28-29, 1292-1293
 and silica abundance, A(127):362-363
 Site 794, A(127):97, A(128):99; B:264-269, 284, 344, 1221
 Site 795, A(127):192-193; B:276-281, 286, 344-345
 Site 796, A(127):271-273; B:270-275, 285
 Site 797, A(127):351; B:252-258, 283, 345, 593, 1223
 Site 798, A(128):124, 159-161; B:361
 Site 799, A(128):299-303; B:34-35, 361
 as stratigraphic markers, B:1220
- and Tsushima Strait closure, B:1219-1220
 upwelling-specific, A(127):267, 271, 349
 zonations, A(127):19-20, 97, 192-193, 271-273, 351, A(128):159-161, 299-303; B:249-251, 282, 361-362, 1219, 1225-1226
- digital downhole seismometer. *See* downhole seismic experiment
- dikes
 and crustal heterogeneity, B:1113-1114
 Site 797, B:942-943
- discoasters. *See also* nannofossils, calcareous
 Site 797, A(127):354
- dissolution
 of biogenic silica, B:344
 of calcareous nannofossils, B:166
 and geologic age, B:310
 Leg 128, A(128):28, 30
 and opal abundance, B:310-311
 of planktonic foraminifers, B:465
 of radiolarians, A(127):198-199; B:295-297
 of siliceous microfossils, A(127):169; B:353
 Site 798, A(128):162, 165
 Site 799, A(128):310
 and temperature, B:310
- dissolved oxygen. *See* oxygen, dissolved
- DIT. *See* resistivity, logging
- dolerite
 alteration, B:883-889
 isotopic data, B:884-888
 petrography, B:892
 sills, A(128):86
 Site 794, B:849, 869
 vesicles, B:887-888
- dolerite, aphyric, A(128):75
 radiometric age, B:822-826
 Site 794, A(127):122-125, A(128):68, 88-90; B:908, 1339
 Site 797, A(127):371-375, 380-381
 zoned vein alteration, B:886
- dolerite, olivine, Site 794, A(128):69, 91
- dolerite, olivine microphyric, Site 794, A(128):68, 89
- dolerite, plagioclase phyric
 Site 794, A(127):122-125, A(128):75
 Site 797, A(127):371
- dolerite, plagioclase-pyroxene, phyric, leucocratic, Site 794, A(128):68, 88
- Dolgorae-1 Well
 paleobathymetry, B:1210
 subsidence plot, B:1208
- dolomiticrite
 Site 795, A(127):186
 Site 797, A(127):343
- dolomite. *See also* ankerite; carbonate, diagenetic; proto-dolomite; siderite
 beds, A(128):275-276
 calcium and magnesium abundances, A(127):363-364
 and Ce anomaly, B:92
 cementing sandstone, A(128):276
 and cerium concentration, B:90-91
 composition, B:82-85
 in FMS, A(127):307, A(128):184, 368
 formation, A(127):23; B:85, 93
 Funakawa Formation, B:75
 in geochemical logs, B:1399
 and interstitial water chemistry, B:1265
 isotopic composition, B:83
 layers, A(128):274-275
 in logs, A(127):302, A(128):34, 245, 332-333, 340-341
 major elements, B:1241

- Monterey Formation, B:75
 and Mossbauer parameters, B:740-741
 nodules, A(127):113, 186-187, 199, 343,
 A(128):275
 occurrence, B:80
 Onnagawa Formation, B:75
 oxygen isotopes vs. degree of recrystalliza-
 tion, B:83, 89
 paleoenvironmental implications, B:94
 patches, A(128):274
 petrography, B:81
 precipitation, B:92-93
 recrystallization categories, B:81, 92-93
 rhombs, A(128):273-274
 in sandstone, B:141-143
 and sedimentation rate, A(128):148
 seismic stratigraphy, A(128):34, 125; B:1141,
 1143
 SEM photographs, B:98
 shallowest recurrence, A(128):273
 Site 795, A(127):187-188
 Site 796, A(127):266
 Site 798, A(128):137-138, 147-150, 174
 Site 799, A(128):240, 260, 272-278, 289,
 354; B:38, 40-41, 75-98, 611-612
 on synthetic seismogram, A(128):184
 texture, B:81
 X-ray diffraction analyses, B:79-80
 dolomite, siliceous, Site 799, B:43
 dolomite nodules. *See* dolomite
 dolomitization
 and alkalinity, A(127):362
 and interstitial water analyses, B:607
 Site 799, A(128):320-321
 downhole seismic experiment. *See also* air-gun
 profiling experiment; ocean bottom seis-
 mometer
 air-gun profile locations, A(128):112; B:1070,
 1077, 1109, 1159
 air-gun reflection records, B:1076, 1079
 air-gun shooting, A(128):113-114; B:1162-
 1163, 1167
 and anisotropic models, B:1117
 assignment of cable conductors, A(128):109
 battery specifications/wiring, B:1067
 broadband digital telemetry package, B:1064
 and crustal structure, B:1342
 data/power link, B:1064-1065
 data selection, B:1067
 digital telemetry of data, B:1063
 equipment used, A(128):69-70
 first arrival traveltimes, B:1089
 frequency response of sensors, B:1158-1159,
 1161
 HIG and NORDA experiments, B:1061
 horizontal sensor azimuth, B:1159
 installation, A(128):69-70, 110; B:1067-1068
 local earthquakes observed, B:1161, 1165-
 1166
 logging cable, A(128):106-109; B:1063
 navigation during, B:1107-1108
 noise levels, B:1162-1163
 objectives, A(128):18, 22-23, 83, 106;
 B:1061, 1157
 OBS-array locations, B:1070, 1077, 1107,
 1109, 1159
 OBS records, B:1077-1078, 1084, 1160
 OBSs deployed during real-time experiment,
 A(128):111, 113; B:1076, 1159
 off-line experiment, B:1158-1162
 operations, A(128):68, 85, 113
 power supply and consumption, B:1066,
 1159-1160
 pressure housing, B:1063-1064
 ray tracing results, B:1090-1103
 real-time experiment, A(128):110-111;
 B:1068, 1077, 1157-1158
 recorder unit, A(128):109-110
 recovery unit, B:1066-1068, 1073
 seafloor data recorder, B:1065-1066
 seafloor package deployment, A(128):111;
 B:1068, 1070-1071
 seafloor recording/recovery unit specifica-
 tions, A(128):109
 seismometer records obtained, B:1077-1078,
 1080-1083
 seismometer specifications, A(128):109
 seismometer system layout, A(128):84, 114;
 B:1063, 1161
 sensors, A(128):106-109; B:1063-1064,
 1072, 1158-1160
 ship positioning during experiment, B:1076-
 1077
 Site 794, A(128):36, 68-70, 106ff, 117;
 B:1061-1073, 1075-1078, 1107-
 1121, 1157-1171
 system design, A(128):106-110; B:1061-1063
 technological difficulties, A(128):106; B:1157
 teleseismic observations, B:1162, 1167-1168
 temperature change during, B:1160-1161
 tilting during, B:1160-1161
 traveltime, offset distance and basement depth
 correlated, B:1112-1113
 velocity power spectrum plot, A(128):115
 velocity structure, one-dimensional, B:1078-
 1081, 1085-1086
 velocity structure, two-dimensional, B:1081-
 1083, 1086, 1089
 very long-period data (VLP), B:1159-1161,
 1164
 weather during, A(128): 113; B:1163, 1171
 downslope transport
 Site 794, A(127):103
 Site 795, A(127):199
 dropstones
 in dark/light cycles, B:581
 Japan Sea, B:1229-1230
 Site 795, B:1256
 dry-bulk density. *See* density
 dust, eolian
 Japan Sea, B:394, 401, 1238-1239
 in logs, B:403
 Oki Ridge, B:419
 origin of dark/light cycle detrital component,
 B:594-595
 in Pacific Ocean, B:394, 403
 dust storms, in China and Mongolia, B:394
 dysaerobia. *See* oxygenation conditions
 dysaerobic conditions. *See* oxygenation conditions
 earthquakes
 Burma, B:1162
 during downhole seismic experiment, B:1161-
 1163, 1165-1166
 Japan Basin, A(128):74
 Japan Basin E, A(127):255
 Japan Sea, A(128):76
 Japan Sea E, A(128):11
 Japan Sea quake (1983), B:1047, 1062, 1161
 and sediment consolidation, B:1130
 and stress field, B:1054, 1120
 and vein formation, B:1180-1181
 East Pacific Rise, sulfide deposits, A(128):21
 ebridians, B:1220
 Site 798, A(128):162; B:241-244
 Site 799, A(128):305; B:241, 246-247
 zonal data, B:1225-1226
 zonation, B:237, 239-241
 eccentricity frequencies, in dark/light cycles,
 B:446-447, 571, 574-575
 electrical resistivity, crustal, B:1343, 1351
 electrical resistivity experiment, B:1351-1359
 cable used, A(128):114
 current used, A(128):113, 114, 116; B:1352
 data collection/operations, B:1352-1354
 electrodes used, A(128):114, B:1352
 logging resistivity compared, B:1354
 methods, A(128):111-115
 noise source, B:1352
 objectives, A(128):18, 23, 83, 111
 operations, A(128):68, 70, 85, 115-116
 schematic illustration of, A(128):115
 sensor cable configuration, A(128):116;
 B:1351-1353
 shooting ship positions, B:1352
 signal amplitude, A(128):116; B:1352, 1359
 signal forms observed, B:1355-1358
 Site 794, A(128):36, 68, 70, 111-117; B:1343
 source current time variation, B:1354
 vertical electrical field (Ez) observed, B:1352
 element mobility, Site 794, B:916
 ellipticity, of borehole, Site 799, A(128):339-340,
 368
 eolian dust. *See* dust, eolian
 epidote, in ash layers, B:1379
 ethane
 in gas hydrate, A(127):288-289
 Site 794, A(127):119
 Site 795, A(127):174, 213-216, 220
 Site 796, A(127):251, 287
 Site 797, A(127):368
 Site 798, A(128):125, 175-176, 187
 Site 799, A(128):244-245, 321-322, 339
 ethylene
 Site 794, A(127):119
 Site 796, A(127):287
 Eu anomaly
 in crest-derived materials, B:731
 and rhyolitic volcanic activity, B:731-732
 and siliceous input, B:691
 Site 794, B:682-683, 916-917
 Site 795, B:683, 918
 Site 797, B:688-691, 920
 Site 798, B:729, 735
 Site 799, B:729, 736
 vs. Ce anomaly, B:737
 Eu/Eu*. *See* Eu anomaly
 Eurasian Plate, A(127):5-6, A(128):9
 europium. *See also* Eu anomaly
 Site 798, B:1368-1369
 euxinic conditions. *See* oxygenation condition
 excess pore-water pressure. *See* consolidation
 excess silica. *See* silica, excess
 experimental petrology. *See* petrology, experimen-
 tal
 extension, and fracture formation, B:1181
 F (emplacement mode parameter). *See* magnetic
 properties
 F-phosphate, Site 799, B:36-39
 fabric, of clay minerals, B:35-36, 38
 n- fatty acids, Sites 798 and 799, B:669-671
 faults. *See also* microfaults; normal faults; reverse
 faults; thrust faults
 Site 794, B:1182-1183
 with slickensides, A(128):92; B:1185, 1190
 and slumps, B:1176
 and stress field analysis, B:1052
 feldspar. *See also* albitization
 albitization, B:107, 136-137
 in ash layers, B:1379, 1388, 1392

- chemical composition, B:106, 136, 139
and sandstone, B:105-107
Site 798, B:411-415, 417
twinning characteristics, B:107
- feldspar, alkali, in ash layers, B:1379
- feldspar, alkaline, in ash layers, B:1392
- feldspar, calcic plagioclase, B:872
- feldspar, orthoclase, B:147
- feldspar, plagioclase
abundance, B:1235-1244
albitized, B:105-106, 109, 137-138
altered, B:141, 144-145, 884-885
anorthite, B:121-123, 126
in ash layers, B:1379
in blue tuff, B:117, 121-126
in dark/light cycles, B:584
in dolerite, A(128):88
effect of water-undersaturated conditions,
B:856
experimental chemical compositions, B:867
glacial/interglacial variation, B:588
K(0-8) vs. mole percent An, B:851
in melting experiment, B:861-862, 895
microprobe analyses, B:124-126
and parent magma, B:856
and potassium, B:856, 869
and radiometric age, B:785-786
replaced by anhydrite, B:148
in sandstone, B:104
Site 794, B:850-851
Site 795, B:850-851
Site 796, B:107, 136
Site 797, B:136, 850-851
Site 799, B:137, 140
textures, B:136
vs. illite, B:1245
vs. quartz, B:1245, 1252
water-undersaturated conditions, composition
effect, B:872
- feldspar, potassium. *See also* feldspar, orthoclase
altered to kaolinite, B:141
in ash layers, B:1379, 1390
in sandstone, B:104
Site 797, B:105, 137
Site 799, B:137
- feldspar, sodic, B:1379
- feldspar accumulation rate, Site 798, B:411-420
- feldspathic arenite. *See also* arenite, feldspathic;
sandstone
Site 799, B:105
- ferrobasalt, Site 797, B:861
- fissility
Site 798, A(128):146ff
Site 799, A(128):272
- flame structures, Site 799, A(128):268-269;
B:1178
- flora, brackish-water, Japan Sea, B:249
- fluid flow
along thrust faults, A(127):247
evidence for, A(127):295
heat transport by, A(127):301
Site 799, A(128):321
- fluorescence, cut, Site 799, A(128):323; B:623
- fluoride
and francolite, B:67
in interstitial water, B:64, 66
uptake rates, B:65-66
- FMS. *See* formation microscanner
- folds
along Okushiri Ridge, A(127):255-256
Site 799, A(128):268
- foliation, clay mineral fabric, B:38
- Foram Sharp Line (FSL)
Japan Sea, B:493, 506
timing of, B:526, 528-529
- foraminifers
abundance and preservation, A(127):99-100,
195-198, 273-274, 355-356,
A(128):258; B:188
assemblages, A(127):20, 100-101, 195-198,
273-274, 355-356
BSEM images, B:548, 557
and climate fluctuation, A(127):195-197
diatom zones correlated, B:188-189, 222-223
morphotaxa recognized in BSEM images,
B:548
paleodepth, A(127):198, 274, 355
paleoenvironment, A(127):199
reworked, A(127):195, 273, 355
Site 794, A(127):99-101, A(128):99; B:187-
193
Site 795, A(127):195-198; B:187-197
Site 796, A(127):273-274; B:187-200
Site 797, A(127):355-356; B:187-207, 1223
Site 799, B:78
textularid taxa, A(127):195-198, 356
and water temperature interpretation,
A(127):355
zonations, B:200-201
- foraminifers, agglutinated
assemblages, B:188-189
Site 794, B:500, 528
Site 795, B:501-502
Site 797, A(127):355-356; B:503-504
Site 798, B:373
- foraminifers, benthic, B:1220-1222
abundance and preservation, A(128):30, 166,
311-312
assemblages, A(128):166, 311-312; B:188-
189
in dark/light cycles, B:568
deposition depth, A(128):354
depth distribution, B:379
Japan Basin, B:208
Japan Sea, B:365, 1208
in laminated sediments, B:551
Leg 128, A(128):30
oxygen isotopes, B:443-444
oxygenation, A(128):166, B:1221
and paleobathymetric analysis, B:1201
as paleodepth indicator, A(128):166; B:208,
1220-1221
preservation, B:366
Site 794, A(127):101; B:210-215, 495-496,
500, 506-507, 516-517
Site 795, A(127):197-198; B:500-505, 518-
521
Site 796, A(127):273-274
Site 797, A(127):355-356; B:210-215, 502-
504, 508-515, 522-528
Site 798, A(128):166; B:367-369, 373-375,
382, 443
Site 799, A(128):311-312; B:370-373, 376-
378
species distribution, B:208
Yamato Basin, B:189, 208
zonation, B:383, 517, 521, 524
- foraminifers, planktonic, B:1220
abundance and preservation, A(128):30, 165-
166, 310-311; B:193, 201, 221, 460
assemblages, A(128):310-311; B:188-189,
462-463, 466
and basal sediment age, B:1335
BSEM images, B:553
coiling direction, A(128):165, 310-311;
B:1220
coiling ratios, B:188, 193, 208, 210-215, 218-
219, 459, 463
in dark/light cycles, B:568, 584
dissolution, A(128):165; B:465
faunal provinces, B:459-462
gyre-margin fauna, B:459
Honsu exposures, B:459
Leg 127, A(127):20
Leg 128, A(128):30
monospecific assemblages, A(128):165
and oxygen isotopes, B:442-443
paleoclimatic interpretations, B:465-468
polar-subpolar fauna, B:459, 462
preservation in diatom-rich sediments,
A(128):165
Q-mode cluster analysis, B:188, 193-200,
208-209, 220, 224
Site 794, A(127):99-101; B:193-200, 210-215
Site 795, A(127):195-197
Site 796, A(127):273-274
Site 797, A(127):355-356; B:193-200
Site 798, A(128):124, 165-166; B:442-443,
457-470
Site 799, A(128):310-311
subtropical fauna, B:459
and surface-water temperature, B:187
transitional fauna, B:459
upwelling evidence, B:469
- forceps
Site 795, B:543
- forests
Japanese archipelago, B:320, 325, 334
Site 798, B:317
- formation factor
data quality, A(127):127, 222, 289-290
Site 794, A(127):128-133, 137
Site 795, A(127):224-228, 231
Site 796, A(127):290-295
Site 797, A(127):383-389, 391
- formation microscanner (FMS)
ash-flow tuff, B:793
ash layers, A(128):153, 339-340; B:398, 400,
1025
chert, A:307; B:1039-1040, 1232
and core recovery, B:1038-1040
cross correlations, B:1040-1041, 1045
cumulate layering in sill, A(128):104
cyclicality in chert layers, B:1041-1044
data processing, B:1038-1043
depth alignments, B:1038-1040
diagrams, B:1038-1039
dolomite, A(127):307, A(128):368
Fourier transform analysis, B:1043, 1046
hole parameters, A(127):307, 399
in interbedded sediments, A(127):399
and intersite lithologic correlation, B:1233
and Milankovitch cycles, B:1043-1044
operations, B:1037-1038
porcellanite, B:1040
qualitative comparison of Site 794 and 797
logs, B:1039-1040
signal processing, B:1040-1043
of siliceous shale, B:1232
Site 794, A(127):142, A(128):104ff, 109;
B:1037-1046
Site 796, A(127):307, 310
Site 797, A(127):399, 408; B:1037-1046
Site 798, A(128):184-185
Site 799, A(128):339-341
Sites 794 and 797 correlated, B:1042-1044
veins and fractures, A(128):104-106
Fourier transform analysis, of FMS data, B:1043,
1046

- fractionation models, B:872-874, 881
- fractures
- ages, A(128):92-93
 - basement, A(127):217, A(128):69, 88-92; B:1181-1184
 - in BHTV data, A(127):307, 399, 409
 - calcite-filled, A(127):188
 - fillings, A(128):91-92; B:1181
 - geometry, B:1181
 - origin, A(128):91-93
 - relative ages, A(128):93
 - Site 794, B:1182-1184
 - Site 798, A(128):143ff
 - and stress regime, B:1181
- fractures, calcite-filled, and normal faults, A(127):188
- framboids, of pyrite, B:36
- francolite, occurrence, B:65, 67
- free-air gravity anomalies
- Japan Basin, A(128):74
 - Japan Basin E, A(127):253, 255-256
 - Japan Basin N, A(127):176, 179
 - Japan Sea E, A(128):252
 - Kita-Yamato Trough, A(128):247
 - Oki Ridge, A(128):127, 133
 - Oki Trough, A(128):127
 - Site 794, A(127):81
 - Yamato Basin, A(128):127
 - Yamato Basin N, A(127):77, A(128):73
 - Yamato Basin S, A(127):330-331, 335
 - Yamato Rise, A(128):247
- frictional heating. *See* heating, frictional
- frontal boundary, oceanographic, Japan Sea, B:155-169
- Funakawa Formation, A(127):198, A(128):13
- deposition, A(128):17
 - dolomite, B:75
 - foraminifers, B:187
 - lithostratigraphy, A(128):354
- fungi spores, Japan Sea, B:485
- gadolinium, logging
- Site 794, B:1416, 1420-1421
 - Site 796, B:1423
 - Site 797, B:1426-1427
- gamma ray logging
- of ash-flow tuff, B:793
 - calibrated by strip samples, B:1367
 - and clay content, B:6
 - of clay minerals, A(127):393
 - and core-log depth correlation, B:1024-1025
 - and depth alignments, B:1038-1040
 - lithostratigraphy correlated, B:1416-1417
 - of Olduvai Subchron, B:403
 - of opal-A to opal-CT transition, B:22
 - and oxygen isotopes, B:403
 - paleoclimatic significance, A(128):188-192
 - power spectra, A(128):217, 369; B:407
 - responses, A(128):103-104
 - Site 607, oxygen isotopes correlated, B:405
 - Site 794, A(127):142, 147, 156-159, 164-167, A(128):103-104, 107-108, 119-120; B:1069, 1416, 1418-1419
 - Site 795, A(127):244-245
 - Site 796, A(127):303-306, 319-322; B:1416, 1422
 - Site 797, A(127):393-395, 400-402, 416-421; B:7, 1417, 1424-1425
 - Site 798, A(128):185-187, 212, 215-216, 225-232; B:399, 401, 1398-1400
 - Site 799, A(128):332-338, 362-364, 367, 369, 380-385, 389-397; B:1399, 1403, 1406-1407
- terrigenous content compared, B:1024-1025, 1028
- used for depth shifting of logs, B:1398, 1415
 - vs. density logging, A(128):215
- gamma-ray spectroscopy tool (GST). *See* geochemical logging
- gas
- and magnetic properties, A(127):275
 - sediment consolidation and, B:1129
- gas, thermogenic, Site 799, A(128):322
- gas chromatography. *See* pyrolysis-gas chromatography
- gas hydrate. *See also* clathrate
- and boron content, B:639
 - bottom simulating reflector (BSR), B:1145, 1147
 - Japan Sea, A(127):24
 - and oxygen isotopes, B:52
 - Site 796, A(127):260, 288-289, 315
 - temperature stability, A(127):288, 299-300
- Gauss Chron
- and diatom occurrence, A(128):160
 - and N Pacific diatoms, A(128):302
 - silicoflagellate and ebridian zones correlated, A(128):305
 - Site 795, B:970, 1223
 - Site 796, A(127):275
 - Site 797, B:973
 - Site 798, A(128):30, 124-125, 170; B:974
 - Site 799, A(128):303, 314; B:1224
- Gauss/Gilbert boundary
- Site 794, A(127):109; B:970
 - Site 797, B:973, 1223
 - Site 798, A(128):30; B:1224
 - Site 799, A(128):314; B:975, 1224
- geochemical logging, B:1411-1427. *See also specific elements analyzed*
- aluminum concentration calculation, B:1398, 1415
 - of ash layers, A(128):339
 - borehole diameter effect, B:1029-1030
 - with boron sleeve, B:1023, 1030
 - calibrated by strip samples, B:1367
 - core comparison, B:1398-1399
 - and core-log depth correlation, B:1024-1025
 - data reduction, B:1395-1399, 1415-1416
 - depth shifting, B:1398, 1415
 - drillpipe iron effect, B:1415
 - elemental weight fraction calculation, B:1398, 1415-1416
 - elements not measured, B:1395, 1415
 - errors in, B:1028-1030
 - industry and ODP data compared, B:1021
 - K, Th, and U calculation, B:1398, 1415
 - Kleiner-Hartigan diagrams, B:1030
 - normalization factors, B: 1029-1030, 1034-1035
 - in opal-A to opal-CT transition, A(127):396-399
 - operations, B:1411-1412
 - and orbital obliquity cycles, B:1021
 - oxide factors for closure normalization, B:1402
 - oxide percentage calculation, B:1398, 1416
 - porosity effect, B:1022-1024, 1029
 - post-cruise processing schematic, B:1397
 - problems in, B:1022-1024
 - relative elemental yield reconstruction, B:1396, 1415
 - Site 794, A(127):142, 150-153, 160-163; B:1416, 1420-1421
 - Site 796, A(127):306-309, 317-318; B:1416, 1423
- Site 797, A(127):396-399, 404-407, 412-415; B:1417, 1426-1427
- Site 798, A(128):187, 233-236; B:1021-1035, 1398-1399, 1401-1402
- Site 799, A(128):366, 386-388, 398-402; B:1399, 1404, 1408-1409
- tool components, B:1022, 1395-1396, 1412-1415
- uses of, B:1021
- X-ray fluorescence analyses compared, B:1025-1029, 1402, 1416-1417, 1420-1421, 1426
- geochemistry, inorganic, B:1261-1274. *See also* geochemical logging; geochemistry, sediment; interstitial water; *specific inorganic geochemical measurements*
- and ash alteration, A(127):22; B:1265
 - and authigenic carbonates, A(128):276-277
 - and bacterial degradation of organic matter, A(127):22
 - and basement alteration, A(127):23, 205; B:1265-1267
 - and biogenic silica diagenesis, B:1263, 1267
 - and carbonate chemistry, B:1263-1265, 1267
 - and carbonate formation, A(127):22-23, 205
 - and clathrates, A(127):278
 - of dark/light cycles, B:569
 - geochemical intervals, Site 799, A(128):244, 318-321
 - Japan Sea, A(127):22-23, A(128):31-32
 - new metal-free squeezer, B:635, 648-650
 - and organic matter diagenesis, B:1262-1263, 1267
 - safety considerations, A(128):322-323
 - shorebased analyses of Leg 127 samples, B:635-650
 - and silica diagenesis, A(127):23
 - Site 794, A(127):72-73, 107ff
 - Site 795, A(127):174, 204ff
 - Site 796, A(127):251, 278-285
 - Site 797, A(127):324-325, 362ff
 - Site 798, A(128):31-32, 125, 171ff; B:605-621
 - Site 799, A(128):31-32, 244, 316ff, 326-327; B:605-621
- geochemistry, organic. *See also* hydrocarbons, volatile; interstitial gas; molecular organic fossils; Rock-Eval analyses; *specific organic geochemical analyses*
- of hydrocarbons, A(128):175-176, 321-322
 - Japan Sea, A(127):23-24, A(128):32
 - Rock-Eval analyses, A(127):114-119, 213, 285, 367, A(128):177
 - Site 794, A(127):72-73, 111-119, 121
 - Site 795, A(127):174, 209-217
 - Site 796, A(127):251, 283-290
 - Site 797, A(127):324-325, 365ff
 - Site 798, A(128):32, 125, 175ff; B:423-437, 667-675
 - Site 799, A(128):32, 244-245, 321ff; B:423-437, 623-633, 667-675
 - van Krevelen plot, A(127):121, 219, 288, 367, A(128):196; B:629
- geochemistry, sediment. *See also* geochemical logging; major elements; minor elements; rare earth elements (REE); trace elements
- of bulk sediment, B:610-611, 613, 615-619
 - of carbonates, B:611-612, 615
 - of clay-size fraction, B:611, 613, 620-621
 - major element molar ratios, B:616-619
 - Site 795, B:705-717
 - Site 798, B:610-612, 719-737
 - Site 799, B:610-612, 719-737

- Sites 794-797, B:677-695
 trace element molar ratios, B:616-619
 geophysical experiments. *See* downhole seismic experiment; electrical resistivity experiment
 geosyncline, Japan Sea, A(128):11
 Gilbert Chron
 Site 796, A(127):275
 Site 798, A(128):30
 Site 799, A(128):303, 314, 316; B:975, 1224
 glacial/interglacial cycles
 and biogenic silica variation, B:591
 carbonate variation, B:590
 grain size variation, B:589
 isotopic variations, B:442
 and Japan Sea circulation, B:457-458, 608
 and oceanographic front oscillations, B:167-168
 and opal flux variation, B:446
 and organic carbon variation, B:589
 and oxygenation condition, B:590-591
 and salinity variation, B:591
 and sediment composition variation, B:588-590
 sulfur variation, B:589-590
 and terrigenous flux variation, B:446
 glacial periods
 and biogenic opal supply, B:404
 interglacial periods, B:403-404
 and loess deposition, B:403
 glaciation
 and Japan Sea, B:338, 342-343, 419-421
 and opal, B:350, 404
 and oxygen isotopes, B:403
 and Pacific Ocean, B:528
 and productivity, B:430
 Site 796, A(127):274
 Site 797, A(127):356
 Site 798, A(128):164
 Site 799, A(128):312
 and Tsugaru Strait sill depth, B:351
 and upwelling, A(127):271, 274
 vegetation and, B:320
 glass. *See* alkaline glass; calc-alkaline glass; glass, volcanic; peralkaline glass
 glass, volcanic
 andesitic magma series, B:1378
 chemistries, B:1374-1379
 dissolution, B:1375, 1389
 and original magmatic composition, B:1377
 in sandstone, B:104
 shoshonite series, B:1378
 glass shards. *See* shards, glass
 glauconite
 allochthonous, B:67, 74
 autochthonous, B:38, 67, 74
 in diagenetic sequence, B:39
 formation, B:71
 microprobe analyses, B:67, 70
 and revised lithostratigraphy, B:1232
 Site 794, B:1259
 Site 795, A(127):189
 Site 796, A(127):265
 Site 797, A(127):344
 Site 798, B:63-74
 Site 799, A(128):260, 264; B:38, 63-74
 glauconitic sand. *See* sand, glauconitic
 gneiss
 Oki Islands, A(128):127
 Goban Spur, inorganic nitrogen, A(127):285
 gold
 Site 798, B:1368-1369
 in sulfide deposits, A(128):21
 GPIT. *See* formation microscanner
 grain density. *See* density
 grain shape, of siliceous claystone, B:38
 grain size
 of ash layers, A(128):150; B:791
 of blue tuff, B:117
 of dolerite, A(128):90-91
 glacial/interglacial variation, B:589
 of magnetic minerals, B:949-951
 magnetic properties affected, B:962
 and permeability, B:1127-1128
 sandstone, B:100, 104
 of siliceous claystone, B:38
 Site 794, A(127):94, 96, 127
 Site 795, A(127):189, 192
 Site 797, A(127):346; B:593
 Site 798, B:1132
 Site 799, B:1132
 grainstone, foraminifers, Site 798, B:551
 Gramineae
 Site 794, B:491
 granite
 Kita-Oki Bank, A(128):127
 Korea, B:111
 Oki Islands, A(128):127
 and sandstone provenance, B:108
 Yamato Rise, A(128):248-249
 GRAPE density. *See* density
 gravity gliding. *See* gravity-induced structures; slumps
 gravity-induced structures, Sites 798 and 799, B:1175-1176
 gray analyses. *See* gray value
 gray value
 of dark/light cycles, B:570-571, 579, 581-587
 and organic carbon content, B:589
 power spectra, B:574
 Site 797, B:592
 time-series analyses, B:571
 Green Tuff, A(128):9, 73
 and abiogenic origin of methane, B:749
 deposition, A(128):17
 Honshu, A(128):127-128
 Oki Islands, A(128):128
 Oki Ridge, A(128):127-128
 and strontium isotopes, B:644
 Yamato Rise, A(128):249
 groundmass
 altered to clay, B:885-886
 in basalt and dolerite, A(128):88-91
 in basaltic basement, B:892
 chemical composition, B:886
 microphotograph, B:859
 GST. *See* geochemical logging
 Guaymas Basin, basalt emplacement, B:1337
 Gulf of Aden, calcareous nannofossils, B:173
 Gulf of California, diagenetic carbonates, B:75
 Gulf of Mexico, microbiological studies, A(128):178
 hafnium
 mobility during alteration, B:911
 Site 795, B:707
 Site 798, B:1368-1369
 Site 799, B:724
 Hannuoba basalt, and subcontinental lithosphere, B:811
 harmotome
 in ash layer, B:1393
 and volcanic glass alteration, B:1377-1378
 headspace analyses. *See* hydrocarbons, volatile
 heat flow
 BSR-derived and probe data compared, B:1148, 1154
 and crustal radiogenic heat, B:1302-1303
 estimated from opal-A/opal-CT BSR, B:1156
 and fluid flux through Toyama Deepsea Fan, B:1148
 helium isotope ratios correlated, B:747
 Japan Basin, A(128):71-74, 246; B:1302, 1345
 Japan Basin E, A(127):252-253
 Japan Basin N, A(127):174
 Japan Sea, A(127):26-27, A(128):14; B:1297-1298, 1301-1304, 1375
 Kita-Oki Bank, A(128):127, 131
 Kita-Yamato Trough, A(128):246
 and lithospheric stretching models, B:1304-1305
 Oki Ridge, A(128):125, 131; B:1302
 Oki Trough, A(128):131
 Okushiri Ridge, A(127):30, 300-301; B:1303, 1307
 and opal-A/opal-CT bottom simulating reflector, B:1147-1148, 1150
 and sedimentation rate, B:1147-1148, 1302
 Site 794, A(127):73, 137
 Site 795, A(127):174, 230-232
 Site 796, A(127):247, 251, 299-300, 314
 Site 797, A(127):325, 390
 Site 798, A(128):125, 213-214
 Site 799, A(128):338, 365
 sources, A(127):251
 topography effect, A(127):300
 vs. basement age, B:1305
 vs. opal-A/opal-CT BSR 2-way traveltime, B:1152-1153
 Yamato Basin, A(127):73, 82, 330, A(128):127, 138, 246, 250; B:1147-1148, 1150, 1302, 1345
 Yamato Basin N, A(128):71-73, 75
 Yamato Basin S, A(127):332
 Yamato Rise, A(128):246, 250; B:1302-1303
 heating, frictional, along thrust faults, A(127):247
 heavy minerals, and rare earth elements, B:691
 heavy rare earth elements (HREE), and elemental ratios, B:693, 730-732
 helium, types of, B:748-749
 helium isotopes. *See* isotopes, helium
 hematite
 at Brunhes/Matuyama boundary, B:972
 in sandstone, B:144
n-heptane, Site 799, A(128):321-322
 Hess Rise, diatoms, B:360
 heulandite, Site 797, B:139
n-hexane, Site 799, A(128):321-322
 HI. *See* hydrogen index
 Hidaka Shear Zone, and Japan Sea tectonic evolution, B:1314-1315
 high-field-strength elements (HFSE)
 alteration and, B:911
 mobility, B:911
 Site 795, A(127):213
 Hokkaido, A(127):14, 251
 basement, A(127):177-178
 depositional history, A(127):349
 rock outcrops, A(127):254
 sandstone provenance, B:110
 tectonic setting, A(128):75-77
 terrigenous sediment source, A(127):307
 Hokkaido W-Honshu N thrust belt, convergence and compression along, A(128):76
 Honshu, A(127):325, A(128):70
 basement, A(128):73
 crustal structure of shelf, A(128):73
 diatom abundance, B:316
 Green Tuff, A(128):73, 127-128

- Kuroko sulfide deposits, A(128):239, 251
 lithostratigraphy, A(127):1415, A(128):19
 paleobathymetry, B:1200
 planktonic foraminifers, B:459
 rock outcrops, A(127):333-334
 rotation, A(128):9
 tectonic setting, A(128):75-77; B:1316
 uplift, A(128):158
 volcanic events, B:1316
- Honshu arc
 compressional stress field and, B:529-531
 diatoms, B:360
 geologic events and, B:536
 movement of, B:528
 paleodepths, B:526
- hopanoid acid, Sites 798 and 799, B:669, 671
 hornblende, in ash layers, B:1379
 hotspot activity, in Yamato Basin, B:1345
 HREE. *See* heavy rare earth elements
 Huang Ho River, and freshening of Japan Sea,
 B:442
- hydrate. *See* gas hydrate
 hydrocarbons, free (S.)
 chromatograms, B:633
 Site 799, B:625, 633
 vs. organic carbon, B:631
- hydrocarbons, generatable (SZ), Site 799, B:625
 hydrocarbons, heavy, Site 799, B:628-629
 hydrocarbons, high-molecular-weight, Site 799,
 A(128):324
 hydrocarbons, liquid, Site 799, A(128):323
 hydrocarbons, nonindigenous, Site 799, B:627
 hydrocarbons, total pyrolytic yields (S,+ S.), Site
 799, B:624
 hydrocarbons, volatile. *See also names of specific
 volatile hydrocarbons*
 and opal-A to opal-CT transition, A(128):323
 Site 794, A(127):119
 Site 795, A(127):213-216
 Site 796, A(127):285-289
 Site 797, A(127):368, 378
 Site 798, A(128):32, 125, 175-176, 185-189
 Site 799, A(128):32, 244-245, 321-322, 330-
 331, 333
 thermogenic origin, A(128):322
- hydroclast, vesicular
 in blue tuff, B:117, 130
 and pressure compensation level, B:119
- hydroclastic shards, in tuffaceous clayey siltstone,
 A(128):89-90
- hydrogen
 and geochemical logging, B:1023-1024
 organic, Site 799, B:629
- hydrogen index (HI)
 chromatograms compared, B:627
 in dark/light cycles, B:431-432
 and organic matter preservation, B:628
 Site 794, A(127):115-118
 Site 795, A(127):213, 219
 Site 796, A(127):285
 Site 797, A(127):367
 Site 798, A(128):177, 196; B:425-426, 668-
 669
 Site 799, B:427, 429, 624, 668-669
 vs. marine maceral content, B:670
- hydrogen isotopes. *See* isotopes, hydrogen
 hydrogen sulfide, shallow presence, A(127):259-
 260
- hydrothermal activity, and Ce anomaly, B:729
 hygromagmaphile elements, spidergram, B:783
 igneous complex. *See* basement
 illite
 glacial/interglacial variation, B:588
- in Japan Sea sediments, B:1235-1244
 in sandstone, B:139
 Site 797, B:592
 Site 798, B:411, 416, 418
 vs. kaolinite + chlorite, B:1245
 vs. Plagioclase, B:1245
 vs. smectite, B:139, 1245
- ilmeneite, in ash layers, B:1379
 Inazumi Formation, A(128):129
 inclination. *See* magnetic properties
 inclination tool (GPIT). *See* formation microscan-
 ner
- incompatible elements. *See* rare earth elements;
 trace elements; X-ray fluorescence analy-
 ses
- index properties. *See* physical properties; *specific
 index properties*
- Indian Ocean
 benthic foraminifers, B:505
 calcareous nannofossils, B:173
 heat flow anomalies, A(127):301
 induction tool. *See* resistivity, logging
 inertinite, Sites 798 and 799, B:670
 intensity. *See* magnetic properties
 interglacial periods. *See* glacial periods
 interstitial gas
 and helium isotope ratios, B:747-751
 methane, B:749-750
 methane vs. helium content, B:750
 Site 798, B:748
 Site 799, B:748
- interstitial water
 ash layer alteration and, B:646
 barium content, B:657-659
 basement alteration and, B:646
 basement rock and adjacent interstitial water
 elemental variations correlated, B:911
 boron content, B:642
 carbonate formation and, B:646
 D/H ratios, B:642
 and diagenetic reactions, B:699
 and fluid/solid exchange reactions, B:1261
 hydrogen isotopes, B:639, 641-643
 isotopic studies, B:49-56
 organic matter degradation and, B:646
 and oxygen isotope depletion, B:697-703
 oxygen isotopes, B:51-52, 54, 84-85, 639,
 641-643, 699-702
 pH, B:94
 silica diagenesis and, B:646
 Site 798, B:607-610
 Site 799, B:607-610
 stable and radiogenic isotopic content, B:635-
 650
 strontium, B:643-646
 sulfate content, B:639, 657-659
- Inyo dike, and stress field, B:1323
- iron. *See also* Mossbauer spectroscopy
 alteration and, B:911
 in ash layers, B:1381
 in augite, B:851-853
 in basalt, B:870
 in carbonate minerals, B:722-723
 in dark/light cycles, B:569
 depth profiles, B:742
 in experimental augite, B:862-863
 high-spin Fe³⁺ in aluminosilicates and carbon-
 ates, B:740-741, 744-746
 in Japanese volcanic outcrops, B:874
 in Japan Sea sediment, B:1236
 paramagnetic Fe³⁺, B:739, 741, 745-746
 silicate Fe³⁺ content, B:744-745
 Site 795, B:706
- Site 798, B:722, 1368-1369
 Site 799, B:90-91, 722
 in veins, B:1178
 vs. aluminum, B:732
 XRF and geochemical log analyses compared,
 B:1025-1029, 1031-1035
- iron, logging
 Site 794, B:1416, 1420-1421
 Site 796, B:1416, 1423
 Site 797, B:1426-1427
 Site 798, B:1401-1402
 Site 799, B:1404, 1408-1409
- iron/magnesium ratio, in basaltic basement rocks,
 B:893-894
- iron oxide. *See* iron
 iron sulfide, and magnetic properties, B:947
 island-arc tholeiite. *See* tholeiite, island-arc
 isochelae
 Site 795, B:543
- isochron plots
 Site 794, B:828-832
 Site 795, B:832-833
 Site 797, B:828-829
- isopentanes, Site 799, A(128):321-322
 isoprenoids, Site 799, B:628-629
 isothermal remanent magnetization (IRM). *See*
 magnetic properties
- isotherms, silica transitions, B:53
 isotopes, argon, in basement rocks, B:822-833
 isotopes, beryllium, and mantle source, B:815
 isotopes, boron
 adsorption/desorption processes, B:640-641
 basement alteration, B:1266
 in interstitial water, B:638-642
 Site 794, B:638-641
 Site 795, B:638-641
 Site 796, B:638-641
 Site 797, B:638-641
- isotopes, carbon
 and carbonate diagenesis, B:85-90
 in chlorite/saponite mixed-layer clays, B:888
 in dark/light cycles, B:569-571
 in interstitial water, B:607
 and methanogenesis, B:88-90
 and organic carbon decomposition, B:1262-
 1263
 in saponite clays, B:888
 in secondary calcite, B:888
 Site 799, B:614, 1265
 and sulfate reduction, B:88
 vs. oxygen isotopes, B:88
- isotopes, deuterium
 in interstitial water, B:607
 Site 799, B:614
 vs. oxygen isotopes, B:615
- isotopes, helium
 heat flow correlated, B:747
 in interstitial gas, B:747-751
 Japan Sea, B:748
 Japan Trench, B:748-749
 in mid-ocean ridge basalt (MORB), B:747
 Nankai Trough, B:748-749
 Site 799, B:614
 vs. helium to neon ratio, B:749-750
 vs. total carbon to helium ratio, B:750
- isotopes, hydrogen
 in chlorite/saponite mixed-layer clays, B:888
 D/H ratios, B:642
 in interstitial water, B:639, 641-643
 in saponite clays, B:888
 in secondary calcite, B:888
 Site 794, B:641-643
 Site 795, B:641-643

- Site 796, B:641-643
 Site 797, B:641-643
 vs. isotopes, oxygen, B:643
 vs. potassium, B:1273
 vs. rubidium, B:1273
- isotopes, lead
 in basement, B:807, 1340-1341
 Pb-Pb plots, B:808, 813
 vs. neodymium isotopes, B:810, 813
 vs. strontium isotopes, B:810
- isotopes, neodymium
 in basalt, B:901-903
 in basement, B:786-787, 807, 893
 and crustal contamination, B:809
 and mantle sources, B:903
 secular variation, B:899
 Site 794, B:1339-1340
 Site 795, B:1338-1340
 Site 797, B:1339-1340
 vs. lanthanum/samarium ratio, B:811
 vs. lead isotopes, B:810, 813
 vs. strontium isotopes, B:809, 812
- isotopes, oxygen
 and ash layer alteration, B:52, 641, 701
 and basalt alteration, B:699-702
 and basement alteration, B:641-642
 and benthic foraminifers, B:443-444
 and carbonate diagenesis, B:701-702
 in chlorite/saponite mixed-layer clays, B:888
 coiling ratio correlated, B:467
 correlated with diatom number in dark/light
 cycles, B:588
 D/H ratios, interstitial water, B:642
 in dark/light cycles, B:431, 569-571
 in diagenetic carbonate, B:83, 86-88
 and diagenetic reactions in interstitial water,
 B:699
 equilibrium values vs. depth of carbonate for-
 mation, B:89
 gamma-ray logs correlated, B:405
 and gas hydrate, B:52
 glacial/interglacial variability, B:442-443
 gradients with dolomite layers, B:89
 in interstitial water, B:51-52, 54, 84-85, 607,
 639, 641-643, 697-703
 isotopic stages, B:442-443
 and Japan Sea, B:442-443, 447
 loess correlated, B:395
 and Northern Hemisphere glaciation, B:403
 oceanographic front oscillations correlated,
 B:168
 Oki Ridge stratigraphy, B:444
 and opal-A to opal-CT transition, B:641-642
 opal-CT, B:51-54
 in planktonic foraminifers, B:442-443
 quartz, B:51-54
 in saponite clays, B:888
 in secondary calcite, B:888
 silica diagenesis and, B:701-702
 and silica transformation temperature, B:49
 Site 794, B:641-643
 Site 795, B:641-643
 Site 796, B:641-643
 Site 797, B:593, 641-643
 Site 798, B:442-443, 698-699
 Site 799, B:614, 698-700
 stage boundaries, B:468
 stratigraphy, B:1221
 surface-water temperature correlated, B:464-
 465, 468
 vs. carbon isotopes, B:88
 vs. deuterium isotopes, B:615
 vs. hydrogen isotopes, B:643
- vs. potassium, B:1273
 vs. rubidium, B:1273
- isotopes, radiogenic
 in basement rocks, B:805-817
 in interstitial water, B:635-650
- isotopes, stable, B:635-650
- isotopes, strontium
 ash layer/basement alteration and, B:643
 in ash layers, B:607
 in basalt, B:901-903
 in basement, B:786-787, 807, 893
 and basement alteration, B:1266
 in blue tuff, B:123, 127
 and crustal contamination, B:809
 in Green Tuff, B:644
 in interstitial water, B:643-646
 isotopic ratios vs. ^{137}Sr , B:646
 and seawater alteration of basalt, B:901
 secular variation, B:899
 Site 794, B:901-903, 1339-1340
 Site 795, B:901-903, 1338-1340
 Site 797, B:901-903, 1339-1340
 Site 799, B:614
 and thermal regime, B:644
 vs. calcium, B:645
 vs. lead isotopes, B:810
 vs. magnesium, B:645
 vs. neodymium isotopes, B:809, 812
 in Yamato Seamount Chain lavas, B:127, 812
- isotopes, sulfur
 in interstitial water, B:637-639
 and organic carbon decomposition, B:1262
 in pyrite, B:711-713
 Site 794, B:637-638
 Site 795, B:637-638, 640, 708-709, 715
 Site 796, B:637-638
 Site 797, B:637-638
 vs. sulfur content, B:715
- isotopic stages, Site 798, B:442-443, 468
- Itoigawa-Shizuoka Tectonic Line, and Japan Sea
 stress field, B:1185
- Izu-Bonin Arc
 basement isotopic composition, B:807
 pressure compensation level, B:119
 sediment contamination of mantle source,
 B:805
 strontium isotopes vs. neodymium isotopes,
 B:809
- Japan. *See also* Japanese archipelago
 Tertiary system faunal changes, B:493
- Japan Arc, A(128):5
 ash, A(128):122, 156-157; B:791
 conductivity model, A(128):85
 crust, A(128):74; B:1345
 geologic history, A(128):1617, 157
 iron enrichment, B:874
 paleoposition, B:1331
 Pb-Pb plots of arc volcanic rocks, B:808
 rotation, A(127):10, A(128):9, 11-12; B:187,
 899, 1054, 1185, 1315, 1317-1318,
 1334
 sandstone provenance, B:110
 strontium isotopes vs. neodymium isotopes,
 B:809
 subcontinental lithosphere, B:811-812
 tectonic evolution, B:1328-1329
 volcanic history, B:1311-1331
 volcanic outcrops, B:874, 882
 Yamato Basin basalt correlated, B:903
- Japan archipelago
 electrical resistivity structure, B:1351
 radiolarians, B:296
 separation from Asia, B:531
- vegetation, B:320, 325, 334
- Japan Basin, A(128):70
 age, A(127):6, 29, A(128):7; B:179, 906
 ash layers, areal distribution, B:791
 basement, A(127):6, 29, B:906
 benthic foraminifers, B:208, 500-505, 518-
 521
 biostratigraphy, B:1223
 blue tuff, B:127-128
 calcareous nannofossils, B:176, 178
 crustal structure, A(127):9, 79, 176, A(128):9-
 11, 71-72, 74; B:924, 1075, 1083,
 1086, 1104, 1107, 1304-1305, 1311-
 1314, 1318-1319, 1341-1346
 crustal structure and tectonic history, B:1318-
 1319
 diatoms, B:276-281, 286
 earthquakes, A(128):74
 Foram Sharp Line and, B:506, 517
 formation age, B:830-831
 free-air gravity anomalies, A(128):74
 heat flow, A(127):27, A(128):71-74, 246;
 B:1302, 1345
 Layer 1, A(128):71-72
 Layer 2, A(128):71-72; B:1342
 Layer 3, A(128):71-72; B:1342-1343
 and lithospheric stretching models, B:1304-
 1305, 1307
 lithostratigraphy, B:496-497
 magnetic anomalies, A(127):6, 10, A(128):7;
 B:942, 1314, 1345
 magnetic properties, B:941-942
 magnetostratigraphy, A(127):22; B:970-972,
 1223
 oceanic crest, A(127):73
 oceanographic setting, B:342
 opal-A to opal-CT transition, B:50
 opal-CT to quartz transition, B:50
 oxygenation conditions, B:201-208
 paleodepth, B:201, 1220-1221
 pollen, B:328, 471, 480, 484
 radiolarians, B:295
 seafloor spreading, B:1345-1346
 sedimentation rate, B:288, 1227
 sponge spicules, B:541-543
 subsidence, A(127):20; B:209, 1208-1211,
 1214
 surface-water temperature, B:209
 tectonic history, B:1317
 Yamato Basin, B:189, 1337
- Japan Basin E
 crustal structure, A(127):251-252
 dredge samples, A(127):253
 earthquakes, A(127):255
 free-air gravity anomalies, A(127):253, 255-
 256
 heat flow, A(127):252-253
 initiation of convergence, A(127):258
 magnetic anomalies, A(127):252-254
 oceanographic history, A(127):259
 sedimentation, A(127):259
 stress field, A(127):258
 tectonic setting, A(127):254-255, 257
 thrust faults, A(127):255
- Japan Basin N
 age, A(127):169, 176
 back-arc rifting, A(127):238
 basement, A(127):169
 bathymetry, A(127):174-176
 crustal structure, A(127):176
 depositional history, A(127):190-192
 dredge samples, A(127):176-178
 free-air gravity anomalies, A(127):176, 179

- geologic history, A(127):169
 heat flow, A(127):174, 176-177
 Layer 1, A(127):176
 Layer 2, A(127):176
 Layer 3, A(127):176
 magnetic anomalies, A(127):176, 178
 oxic conditions, A(127):199
 paleoenvironment, A(127):199
 rifting history, A(127):179
 sedimentation, A(127):179, 181, 238
 seismic stratigraphy, A(127):179, 233-234
 tectonic setting, A(127):178-180
 upwelling, A(127):199
- Japan Sea
 age, A(127):29; B:297, 327, 362, 1208, 1335-1336
 back-arc rifting, A(127):16
 bacteria, B:755-776
 basaltic submarine volcanism, B:874
 basement, A(127):22, 24-25, A(128):6-7; B:749, 1265-1266, 1303-1304
 basin subsidence and volcanic activity, B:796-797
 bathymetry, A(127):6, A(128):6, 10; B:1199, 1276
 benthic environments, B:379
 biosiliceous productivity, B:313
 biostratigraphy, A(127):19; B:187, 1219
 carbonate chemistry, B:1263-1265
 carbonate-compensation depth, B:155, 168, 208, 440, 1201
 climate, B:320
 consolidation/permeability studies, B:1123-1133
 continental crust, B:809
 continental fragments, A(128):5, 10-11
 crustal structure, A(128):6-7, 9-11; B: 1075, 1083, 1104, 1107, 1311-1314, 1333, 1341-1346
 current systems, B:228
 deposition, A(128):15-18; B:1261
 development, B:536
 diatoms, B:309, 363
 dissolved oxygen, A(128):20
 downhole seismic experiment, B:1061-1073, 1157-1171
 earthquakes, A(128):76
 eolian dust, B:394
 faunal change, B:506
 formation, A(127):5, 9-10
 formation age, B:819-836, 1220, 1228
 formation models, A(128):11-13; B:1183-1187
 gateways to, B:362-363
 geochemistry, A(127):22-23
 geographic setting, B:365-366
 geologic history, A(128):16-17
 geologic setting, B:651, 677
 heat flow, A(127):26-27, A(128):14; B:1297-1298, 1301-1303, 1375
 helium isotope ratios, B:748-749
 inorganic geochemistry, B:1261-1274
 lithofacies, B:635
 lithospheric thickness, A(127):255
 lithostratigraphy, A(127):17-19, A(128):13-15; B:679, 1229-1233, 1275, 1315-1316
 logging, A(127):27-28
 magma genesis, B:1340-1341
 magmatic history, B:787-788
 magnetic anomalies, A(127):11; B:1334
 magnetics, A(127):22
 magnetostratigraphy, B:969-982, 1219
- mantle, anisotropic velocity model, B:1075
 models for backarc spreading, B:903
 neodymium/strontium isotopes, B:899-904
 oceanographic front, B:155-169
 oceanographic history, A(127):16, 19-22, 24, 30-31, A(128):20-22; B:678-679
 oceanographic setting, A(128):20; B:166-167, 342-343, 394-395, 423, 440, 457-458, 559-560, 577, 605, 651, 705
 onshore exposures, A(128):15-18
 opal-A to opal-CT transition, A(127):31; B:49, 341, 1145-1156
 opal-CT to quartz transition, A(127):31; B:49
 opening of, B:171, 179, 529-531
 and organic matter, A(127):31; B:1262-1263
 origin, B:249
 oxygenation conditions, B:216, 342-343, 423, 430, 678-679, 705, 713-715, 1221
 paleobathymetry, B:1209-1210
 paleoceanography, B:179-180, 298-300, 419-421, 526, 528-529
 paleoclimate, B:403, 459
 paleodepth, B:1228
 paleoenvironment, B:187, 493
 paleogeography, A(128):22
 paleotemperature fluctuations, B:457-470
 phosphorus, B:68-71
 physical properties, A(127):25-26; B:1275-1296
 phytoplankton pigment concentration, B:167
 plate reconstructions, B:1321-1324
 pollen, B:328-336, 395, 471-478
 pre-ODP data available, B:1333
 productivity, A(127):20; B:167, 430, 651
 radiometric ages, B:1336
 rifting, A(127):16; B:891, 899, 1334
 and sea ice formation, A(127):20-22; B:167
 sediment composition, B:1233-1249
 sediment isopachs, A(127):13, A(128):12
 sediment structures, A(128):13-15
 sedimentation, A(127): 10-16, 30-31; B:635, 1361
 sedimentation rate, A(127):22, 24
 seismic reflection profile, A(127):14
 seismic stratigraphy, A(127):28-29
 silica, B:447, 1263
 sill depths, B:299-300
 sites drilled, B:172, 226, 238, 250, 318, 326-327, 479, 494, 543
 slumping, A(128):14
 stress field, A(127):17
 structure, A(127):6; B:780, 792, 1175-1193
 submarine volcanism, A(127):95
 subsidence, A(127):16, 19-20; B:1197-1218, 1343-1344
 surface currents, A(128):21
 surface-water temperatures, B:201, 463-465
 tectonic events and microfossil assemblages, B:300, 526
 tectonic evolution, B:903, 1183-1187, 1311-1331, 1343-1347
 tectonic setting, A(127):5-10, A(128):9; B:1186, 1312
 temperature distribution, A(128):20
 tephra markers, A(128):291-292; B:796
 thermal conductivity, B:1301-1303
 thermal gradient, A(127):26-27; B:1301-1303, 1375
 thermal history, B:49-56, 1297-1309
 thermocline, B:167
 thrust faults, A(127):81-82
 volcanic activity, B:796-797, 829-832, 1319, 1324
- Wadati-Benioff zone, B:1062, 1158
 water circulation, A(128):21; B:69-71, 423-424
 water convection, B:394
- Japan Sea Bottom Water, isotope ratios, B:748
- Japan Sea E
 bathymetry, A(127):12
 compression, A(127):16-17, 268, A(128):9
 crustal structure, A(128):9-11
 deformation, B:1314-1315
 earthquakes, A(127):9, A(128):11
 free-air gravity anomalies, A(128):252
 initiation of convergence, A(127):314
 magnetic anomalies, A(128):251
 plate reconstruction, B:1314-1315
 rift propagation, B:1346
 stress field, B:1059
 and strike-slip shear, B:1314-1315
 tectonic history, B:1316-1317
 tectonic map, A(127):83
 thrust faults, A(127):9
- Japan Sea N
 heat flow, A(127):231-232
 stress field, A(127):180-181
- Japan Sea Plate, electrical resistivity structure, B:1351
- Japan Sea Proper Water (JSPW), B:577
 benthic foraminifers and, B:365-366
- Japan Trench, A(127):6, A(128):9
 basement age, B:749
 carbon isotopes, B:88
 crustal structure, A(128):74
 diagenetic carbonates, B:75
 helium isotope ratios, B:748-749
 helium to neon ratios, B:748
 interstitial water oxygen isotopes, B:698
 rare earth elements, B:719
- Jaramillo Event. *See* Jaramillo Subchron
- Jaramillo Subchron
 and diatom occurrence, A(128):159
 inclination variations, A(128):172
 and magnetic intensity variations, B:962
 and sedimentation rate, B:959-960
 silicoflagellate and ebridian zones correlated, A(128):162
 Site 795, B:1223
 Site 797, A(127):359; B:973, 1223
 Site 798, A(128):30, 125, 168; B:164, 564, 973, 1224
 Site 799, A(128):314; B:974, 1224
- joints, Site 794, B:1182-1183
- Juan de Fuca Ridge, basalt emplacement, B:1337
- K/Ar age, Site 794, B:784-786
- Kaena Subchron
 Site 798, B:1224
 Site 799, A(128):314; B:1224
- kaolinite
 abundance, B:1235-1241
 in sandstone, B:139-141
 as sandstone cement, B:134
 Site 798, B:411, 416, 418
 Site 799, B:144
- kaolinite/illite ratio, Site 798, B:411, 416
- Kita-Okii Bank, A(127):325
 basement, A(127):333, A(128):127-129
 bathymetry, A(128):126-127
 heat flow, A(127):330, A(128):127, 131
 seismic stratigraphy, A(127):399
- Kita-Yamato Bank, A(127):325
 basement, A(128):247-249
 bathymetry, A(128):245-246
 crustal structure, A(128):248
 diatoms, A(128):249, 303

- freshwater sediments, A(128):249
magnetic anomalies, A(128):247
phosphate source, B:67-68
phosphatic sediments, A(128):249
sandstone provenance, B:110-111
subsidence, A(128):295
- Kita-Yamato Trough, A(127):13, 325. *See also*
Yamato Trough
n-alkanes, B:669
bathymetry, A(128):245-246, 374; B:76
benthic foraminifers, B:370-373, 376-378
biostratigraphy, B:1224
consolidation/permeability studies, B:1123-1133
crustal structure, A(128):245-249
depositional history, A(128):252-253, 292-298, 353-355
diagenesis, B:35-39
diatoms, B:361
ebridians, B:241, 246-247
free-air gravity anomalies, A(128):247
geologic history, A(128):21-22
heat flow, A(128):246
Layer 1, A(128):246
Layer 3, A(128):246
magnetostratigraphy, B:974-975, 1224
normal faults, A(127):334
oceanographic conditions, A(128):312, 355-356
oceanographic drilling objectives, A(128):19
oceanographic history, A(128):252-253
organic matter, B:668
paleodepth, B:373, 379
paleoenvironment, B:381
radiolarians, B:225-226, 233-234
sediment thickness, A(128):237, 353-354
sedimentation, A(128):250; B:1227
seismic stratigraphy, A(128):34, 245, 250, 344-351; B:1135-1143
silicoflagellate, B:241, 246-247
Site 799, B:425, 560, 605, 623, 667, 721, 739, 1135
stress field, A(128):293-294
structural cross-section, A(128):303
structural setting, B:1135
subsidence, A(128):239, 295-298, 355
surface-water productivity, A(128):324
tectonic evolution, B:1187
tectonic setting, A(128):249-250; B:156
- Kitaura Formation
deposition, A(128):18
foraminifers, B:187, 459
submarine fan deposition, B:1202
- Kleiner-Hartigan diagrams, for core XRF and geochemical log oxides, B:1030
- Koenigsberger ratio (Q)
in basement, A(127):203
Site 795, A(127):199
- Korea, granite exposures, B:111
Korea Strait, A(127):355
Korean Current, B:424
Korean Peninsula, A(128):15
Korean Plateau, basement, A(127):9
Kuril Basin, formation, A(127):10
Kuroko sulfide deposits, A(128):21-22, 24, 239, 251
- Kuroshio Current, A(127):355, A(128):20-21, 164
climate and, B:326
and diatom assemblages, A(128):159
and foraminiferal coiling direction, A(128):165, 310-311
Japan Sea and, A(128):30
Site 798, A(128):164-166
- Site 799, A(128):312
and Tsushima Strait, B:166-167
- Kyushu, ash layers from, A(128):156-157
- laminated sediments. *See* laminations; sediments, laminated
- laminations
annual or seasonal, B:548, 551
in BSEM images, B:549-557
calcareous/biosiliceous alternations, B:549
Chaetoceros resting-spore laminae, B:549-551, 555-556
in cyclic lithofacies, A(128):140
in dark/light cycles, B:581
detrital-rich/biosiliceous alternations, B:549
in dolomite, B:92
foraminifer-rich laminae, B:551
mixed-biosiliceous laminae, B:549-551
thickness, B:548, 550
and upwelling events, B:551
- La/Yb values. *See* lanthanum/ytterbium ratio
- lansfordite
and alkalinity, B:92
changed to magnesite, B:80, 91
chemical composition, B:82
CuK- α vs. depth, B:81
formation, B:91-92
isotopic composition, B:83
occurrence, B:80
and organic carbon content, B:92
paleoenvironmental implications of, B:94
Site 799, B:75-98
X-ray diffraction analyses, B:80
- lanthanum
in ash layers, B:1386
Site 795, B:707
Site 798, B:1370-1371
vs. aluminum, B:737
- lanthanum/samarium ratio
in basement rocks, B:807
vs. neodymium isotopes, B:811
- lanthanum/ytterbium ratio
and Ce anomaly, B:693-694
and manganese accumulation, B:729
and migration of heavy rare earth elements (HREE), B:731-732
Site 794, B:683
Site 795, B:683
Site 797, B:688-691
Site 798, B:729, 735
Site 799, B:729, 736
vs. Ce anomaly, B:737
- large-ion lithophile elements (LILE)
mobility during alteration, B:779, 838
Site 795, A(127):219
and slab-derived fluids, B:809
- Last Glacial Maximum
and Japan Sea oceanographic conditions, B:394-395
and siliceous microfossil abundance, B:311
- Lau Basin
formation model, B:928
tectonic history, B:1347
- lava flows
eruptive depth, A(128):95
Site 794, B:906, 1339
Site 795, B:850, 906, 943
Site 797, B:838, 906-908
suites observed, B:853-855
- lava flows, basaltic, Site 795, B:819
- Layer 1
Japan Basin N, A(127):176
Kita-Yamato Trough, A(128):246
Oki Ridge, A(128):127
- Oki Trough, A(128):127
Yamato Basin, A(128):127
Yamato Basin S, A(127):330
- Layer 2
Japan Basin, B:1342
Japan Basin N, A(127):176
Oki Ridge, A(128):127-128
seismic anisotropy, B:1107
and thickness of altered basalt, B:701
Yamato Basin, A(127):73, A(128):9, 127-128; B:1342, 1345
Yamato Basin S, A(127):330
Yamato Rise, A(128):246
- Layer 3
Japan Basin, B:1342-1343, 1345
Japan Basin N, A(127):176
Kita-Yamato Trough, A(128):246
Oki Ridge, A(128):127-128
Yamato Bank, A(128):246
Yamato Basin, A(127):73, A(128):127-128; B:1342-1343, 1345
Yamato Basin S, A(127):330
Yamato Rise, A(128):246
- LDT. *See* density, logging
- lead
in basement rocks, B:807
Site 794, B:1362
Site 795, B:1365
Site 797, B:1366
in sulfide deposits, A(128):21
- lead isotopes. *See* isotopes, lead
- Leg 13, trace metal composition, B:1361
- Leg 19, silicoflagellate, B:237
- Leg 31
biostratigraphic zones used, A(128):162
consolidation tests, B:1123
diatoms, B:250
drilling history, A(127):16, A(128):5, 82; B:1197
foraminifers, A(127):273
Japan Sea formation age, B:1334
lithostratigraphy, A(128):13
organic carbon, B:430
poor carbonate preservation, B:457
radiolarians, A(128):162; B:225, 291, 297
silicoflagellate, A(128):28; B:237
site locations, B:156
Toyama Deepsea Fan, B:1148
- Leg 42B, trace metal composition, B:1361
- Leg 57
ash layers, A(127):95-96; B:1319-1320
silicoflagellate, B:237
volcanic activity, B:796
- Leg 64, basalt emplacement, B:1337
- Leg 65
downhole seismic experiment, B:1061
seismometer emplacement, A(128):106
- Leg 67
downhole seismic experiment, B:1061
seismometer emplacement, A(128):106
- Leg 73, magnetic properties, B:949
- Leg 78A
downhole seismic experiment, B:1061
seismometer emplacement, A(128):106
- Leg 78B, seismometer emplacement, A(128):106
- Leg 80, inorganic nitrogen, A(127):285
- Leg 86, silicoflagellate, B:237
- Leg 87
ash layers, B:1319-1320
volcanic activity, B:796
- Leg 88
downhole seismic experiment, B:1061
seismometer emplacement, A(128):106

- Leg 91
 downhole seismic experiment, B:1061
 seismometer emplacement, A(128):106
- Leg 93, sponge spicules, B:541
- Leg 112, microbiological studies, A(128):178
- Leg 113, sponge spicules, B:541
- Leg 120, sponge spicules, B:541
- Leg 124, Zijderveld plots, A(127):277
- Leg 135
 Lau Basin formation, B:928
 tectonic history, B:1347
- Leg 139, basalt emplacement, B:1337
- lerite, plagioclase phyrlic, A(128):75
- light rare earth elements (LREE), B:730
 and Ce anomaly, B:693-694
- Liman Current, B:342, 423
 cold surface water, B:440, 457
 in Japan Sea, A(128):20-21; B:70
- lipids
 in organic matter, A(128):324
 Sites 798 and 799, B:669-671
- lipid fraction, in organic matter, A(128):324
- liptinite, and thermal maturity levels, B:669
- liptodetrinite, Sites 798 and 799, B:670
- liquid limit. *See* Atterberg limits
- liquid line of descent
 in melting experiments, B:863
 mid-ocean ridge basalts and Site 797 sills correlated, B:870
 for olivine tholeiitic basalts, B:864
 problems with, caused by alteration, B:911
 Site 794, B:869-882, 1339
 Site 795, B:918, 1338
 Site 797, B:869-882, 920
- lithium
 and basement alteration, A(127):364; B:1266
 and biogenic silica content, A(127):205
 and diatom dissolution, A(127):364
 in interstitial water, B:1272
 and opal-A to opal-CT transition, B:1266
 Site 794, A(127):109
 Site 795, A(127):205
 Site 796, A(127):280-281
 Site 797, A(127):364, 371
 Site 798, A(128):174-175, 184
 Site 799, A(128):318, 332; B:611
 vs. silica, A(128):185
- lithodensity tool (LDT). *See* density, logging
- lithofacies, cyclic, A(128):27. *See also* color-banded bedding; dark/light cycles
 and bottom water conditions, A(128):141
 burrows, A(128):140-141
 compositional variability, A(128):140
 couplet thickness, A(128):140
 cycle boundary characteristics, A(128):140-142
 and foraminifers Q-mode cluster analysis, B:200
 formation, A(128):283
 Japan Sea, A(128):14
 laminations, A(128):140
 in logs, A(128):34, 125, 185
 orbital obliquity, A(128):35
 and organic carbon, A(128):125, 177
 organic matter, A(128):140
 origin, A(128):158
 and paleoceanographic changes, A(128):121-122
 Site 798, A(128):35, 121-122, 124, 137ff, 195-196; B:156
 Site 799, A(128):259, 282-284, 293
- lithospheric stretching models, heat flow and basement ages compared, B:1304-1305
- lithostratigraphy. *See also specific sediment types*
 basement, B:781, 884, 901
 and biogenic silica dissolution, B:344
 blue tuff, B:120-121
 gamma ray logs correlated, B:1416-1417
 Honshu, A(127):15
 intersite lithologic correlation, B:1233
 Japan Basin, B:679
 Japan Sea, A(127):17-19, A(128):25-28; B:1229-1233, 1275, 1315-1316
 Oga Peninsula, B:1203
 opal dissolution transition zone (ODTZ) correlated, B:347-349
 physical properties correlated, A(127):383-389, A(128):182, 327-329; B:1143
 Pohang Basin, B:1205
 problems with shipboard identifications, B:1229
 revised unit boundaries defined, B:1230-1233
 Sakhalin Island, B:1206
 and sedimentary structures, B:1177
 seismic intervals correlated, B:1140-1143
 seismic stratigraphy correlated, A(127):155, 174, 234-238, 243, 251, 312-313, 325, 403-408, 410, A(128):125, 193-194, 224, 351-353, 379
 Site 794, A(127):72, 74-77, 91, A(128):25-26, 77-81; B:5, 118, 494, 496, 655, 1049, 1277, 1313, 1336, 1377, 1413
 Site 795, A(127):170-174, 186-192, A(128):26; B:5, 118, 655, 706, 1277, 1313, 1336, 1377, 1413
 Site 796, A(127):248-250, 261-269, A(128):26; B:5, 105, 118, 133, 655, 1277, 1313, 1336, 1377, 1413
 Site 797, A(127):324, 326-330, 340-351, A(128):26; B:5, 105, 118, 133, 496, 498-499, 655, 1277, 1313, 1336, 1376, 1413
 Site 798, A(128):25, 27-28, 122-124, 136 158; B:5, 157, 655, 697, 800, 1277, 1336, 1376
 Site 799, A(128):25, 27-28, 35-36, 238-244, 255ff; B:5, 33-48, 76-77, 105, 133, 157, 655, 697, 800, 1137, 1277, 1313, 1336, 1376
 Sites 794-796 compared, A(127):269
 Sites 794 and 797 correlated, B:1038
 units, revised, B:1229-1233, 1250
 units, Site 794, A(127):90-94; B:679
 units, Site 795, A(127):186-190; B:4, 679, 706
 units, Site 796, A(127):261-266
 units, Site 797, A(127):340-346; B:4-5, 679
 units, Site 798, A(128):124, 137-138; B:156, 697, 973
 units, Site 799, A(128):240-244, 256-265; B:76-77, 158-160, 697, 974
 and vein abundance, B:1180
 vitric tuff, Site 799, B:36
 Yamato Basin, B:679
- loess
 in China, B:394
 clay fraction, B:401
 deposition at glacial maxima, B:403
 and Northern Hemisphere glaciation, B:421
 origin of dark/light cycle detrital component, B:594-595
 oxygen isotopes correlated, B:395
 and quartz abundance, B:401
 source and deposition regions, B:411
 logging, B:1395-1409, 1411-1427. *See also specific logging measurements*
 aridity signal, A(128):188-192
- of ash layers, A(128):288
 and Asian dust supply, B:403
 of basement, A(128):69, 86
 of breccia tuff, A(128):287
 of chert and porcellanite layers, A(127):394
 correlated with basement recovery, A(128):93-95
 of cyclic lithofacies, A(128):125
 cyclicity observed, B:398, 405
 and intersite correlation, B:1234
 Japan Sea, A(127):27-28, A(128):33-34
 lithostratigraphy correlated, A(127):393-395
 log characteristics, A(128):103-104
 log quality and processing, A(127):139, 232, 301-302, 392-393, A(128):103, 183, 329-332
 of opal-A to opal-CT transition, B:15-16, 19, 21-22
 and opal content variations, B:398-400
 of opal-CT to quartz transition, B:23
 operations, A(127):27, 139, 142, 183-186, 230, 232, 236, 260, 296, 301-302, 325, 338-339, 389-390, 392, 398, A(128):85, 103, 106, 211, 253-255, 329, 361; B:395-396, 1411-1412
 orbital obliquity observed, B:403
 paleoclimate, A(128):33-34, 187-192, 341-342; B:393-407
 periodicity resolution, B:396-397
 physical properties correlated, A(127):139, A(128):338
 and sediment composition estimates, B:402-403
 seismic stratigraphy correlated, A(128):224, 351-353, 379
 of silica diagenetic transitions, B:3
 Site 794, A(127):73, 135-142, 144-167, A(128):34, 69; B:1037-1046, 1069, 103ff
 Site 795, A(127):174, 230ff
 Site 796, A(127):251, 296ff
 Site 797, A(127):325, 389ff; B:7, 1037-1046
 Site 798, A(128):33-34, 125, 182ff; B:393-407, 1021-1035
 Site 799, A(128):34, 245, A:329ff
 Sites 798 and 799 compared, A(128):342
 stratigraphy, B:1221
 temporal resolution, B:396, 404-405
 in terrigenous/biogenic opal cycles, B:398
 tool string schematic diagrams, B:396
 units, Site 799, A(128):332-338
 used to determine alteration extent, B:1282
 vertical aperture, B:396
 of vitric tuff, B:37
- logging, geochemical. *See* geochemical logging
- long-space sonic tool (LSS). *See* velocity, logging
- loss on ignition (LOI)
 as alteration proxy, B:909
 and hydrous mineral content, B:1282-1284
 and physical properties, B:1283
 vs. density, B:1291
 vs. elemental data, B:918-921
 vs. magnesium, B:839
 vs. porosity, B:1290
 vs. velocity, B:1291
- LREE. *See* light rare earth elements
- LSS. *See* velocity, logging
- lutetium, Site 798, B:1370-1371
- lysocline, and nannofossil preservation, A(128):164
- macerals, Sites 798 and 799, B:668-670
- magma. *See* mantle composition; mantle source; parent magma

magma batches, Site 794, B:917
 magma composition, and trace elements, B:1378-1379
 magma genesis, in Japan Sea, B:1340-1341
 magma intrusion, and crustal deformation, B:1323
 magma source
 depleted asthenospheric mantle, B:1341
 enriched subcontinental upper mantle, B:1341
 and neodymium isotopes and titanium correlated, B:902
 Site 794, B:780-782
 Yamato Basin, B:782-784, 902
 magma suites
 in basement rocks, B:869-870
 Site 794, B:871, 875
 Site 797, B:871, 875
 magmatic affinity, of basement, A(128):96ff
 magmatic eruption, and blue tuff, B:119, 126
 magmatic history, Japan Sea, B:787-788
 magnesiochromite, B:840. *See also* spinel; spinel, chromium
 magnesite. *See also* breunnerite
 chemical composition, B:82, 84-85
 CuK- α vs. depth, B:81
 derived from lansfordite, B:80, 91
 formation, B:91-92
 formation depth, B:85
 isotopic composition, B:83
 occurrence, B:80
 organic carbon content, B:92
 origin, B:76
 paleoenvironmental implications, B:94
 petrography and textural relations, B:81
 SEM photographs, B:97
 Site 799, B:75-98
 X-ray diffraction analyses, B:80
 magnesite, ferroan, B:745-746
 magnesium
 alteration and, B:909, 911
 and ash alteration, A(127):280, 363-364
 in ash layers, B:1382
 and basement alteration, A(127):280; B:892-893, 1265-1266
 and dolomite formation, A(127):363-364
 enrichment during alteration, B:842
 and geochemical logging, B:1395, 1415
 in interstitial water, B:1263, 1268
 Japan Sea, B:1236, 1265
 mobility during alteration, B:838
 Site 794, A(127):109-110
 Site 795, A(127):174, 205
 Site 796, A(127):280
 Site 797, A(127):363-364, 370
 Site 798, A(128):173-174, 182
 Site 799, A(128):318, 329; B:611
 vs. boron, B:642
 vs. calcium, A(127):115
 vs. loss of ignition, B:839
 vs. smectite, B:1250
 vs. strontium isotopes, B:645
 magnesium/aluminum ratio
 in Japan Sea sediment, B:1236
 Site 794, B:1249
 as volcanic input indicator, B:1239
 magnesium/calcium ratio
 in interstitial water, B:1263, 1269
 Japan Sea, B:1265
 Site 799, A(128):329; B:611
 magnesium number
 basement, A(128):98
 Site 794, B:839
 Site 797, B:839
 spinel, B:840-842, 844

magnesium oxide. *See* magnesium
 magnetic anomalies
 Japan Basin, A(128):7; B:1314, 1345
 Japan Basin E, A(127):252-254
 Japan Basin N, A(127):176, 178
 Japan Sea, B:1334
 Japan Sea E, A(128):15, 251
 Kita-Yamato Bank, A(128):247
 Oki Ridge, A(128):127, 132
 Site 794, A(127):80
 Tsushima Basin, B:1314
 Yamato and Japan basins compared, B:942
 Yamato Bank, A(128):247
 Yamato Basin, A(128):7; B:1314, 1345
 Yamato Basin N, A(127):77, A(128):73
 Yamato Basin S, A(127):330, 334
 Yamato Rise, A(128):247, B:1314
 magnetic declination. *See* magnetic properties
 magnetic inclination. *See* magnetic properties
 magnetic intensity. *See* magnetic properties
 magnetic minerals. *See also specific magnetic minerals*
 diagenetic dissolution, B:951
 formation by thermal demagnetization, B:949, 951
 grain size, B:949-951
 and magnetic properties, B:947, 962
 Site 797, B:949, 951
 magnetic polarity. *See* magnetostratigraphy
 magnetic properties. *See also* magnetic minerals; magnetostratigraphy
 AF demagnetization, B:949, 951
 in basalt flows vs. dolerite sills, A(128):102
 in basement, A(127):209, 358-359
 and basement alteration, A(127):203
 basement VRM, intersite comparisons, B:941
 and bioturbation, A(127):275
 data quality variations, B:975
 declination, A(127):106-107, 200-202, 276, 358-359, 363, A(128):101-102, 167-170, 313-315; B:960, 977, 979, 1052-1053, 1317
 demagnetization, A(128):169, 314-315
 emplacement mode parameter (F), B:934, 936-937, 942
 equal-area plots, A(127):104-105, 206-207, 278-279, 360-361, A(128):104-105, 173, 316, 318-320, 322-323; B:976
 frequency dependence susceptibility, B:934, 936-937, 942
 and gas, A(127):275
 and geomagnetic field intensity variations, B:962-963, 977
 inclination, A(127):106-107, 200-202, 276, 358-359, 363, A(128):101-103, 105, 167-170, 173-174, 313-316; B:960, 974, 977-981
 intensity, A(127):72, 103, 106-107, 174, 200-202, 251, 275-276, 324, 357-359, 363, A(128):101-103, 105, 167-170, 173, 313-315; B:934, 937, 957, 969-973, 975-981
 intensity decay plots, A(127):104-105, 206-207, 278-279, 360-361, A(128):104-105, 173, 316, 318-320, 322-323
 isothermal remanent magnetization (IRM), A(127):357, 359; B:972-973, 975
 Japan Sea, A(127):22, A(128):30-31
 Koenigsberger ratio (Q), A(127):199, B:934-938
 magnetic intensity, B:959-967
 and magnetic minerals, B:951
 in RCB cores, B:975, 981

remagnetization, A(127):203, 359; B:938
 remanence, A(128):101
 and rotation of Japan Arc, B:1317-1318
 and rust contamination, A(127):357; B:973
 Site 794, A(127):103-105, A(128):30-31, 69, 100-105; B:934-935
 Site 795, A(127):174, 199ff; B:934, 937
 Site 796, A(127):251, 275ff
 Site 797, A(127):324, 357ff; B:934, 937-938, 947-957
 Site 798, A(128):30-31, 166ff
 Site 799, A(128):30-31, 312ff
 soft vs. stable components of remanence, A(128):101
 and stress field analysis, B:1050
 susceptibility, A(127):72, 105, 110-111, 174, 199, 203-204, 275, 277, 324, 358, 362, A(128):101-102, 167-171; B:934, 937, 954, 959, 961, 973, 975-977, 979, 981
 thermal demagnetization, B:938-939, 949, 951, 954
 vector demagnetization diagrams, B:935, 940, 950, 952-953
 viscous remanent magnetization (VRM), B:934, 936-937, 939-942, 944-945
 of volcanic rocks, B:933-945
 in XCB cores, B:975, 978, 980
 Yamato and Japan basins compared, B:941-942
 Zijderveld plots, A(127):104-105, 203, 206-207, 277-279, 360-361, A(128):104-105, 173, 316, 318-320, 322-323; B:971
 magnetic susceptibility. *See* magnetic properties
 magnetite, B:853
 in basaltic andesite, A(127):217
 magnetostratigraphic analyses and, B:976
 produced by thermal demagnetization, B:947
 and sedimentation rate, B:976
 Site 797, B:949, 951
 magnetostratigraphy, B:1219-1228. *See also* magnetic minerals; magnetic properties
 basement polarity, A(128):86; B:936-938
 chrons and subchrons, A(127):108
 core orientation, A(127):103, 199, A(128):167
 data quality, B:969, 975
 datums used, B:163, 1222
 depths to polarity boundaries, B:973
 diatoms and, B:361-362
 and drilling disturbance, A(127):103
 ebriidians and, B:240
 in igneous rocks, A(127):104, 203
 Japan Sea, A(127):22, A(128):30-31; B:969-982
 magnetic polarity, B:943
 problems with, B:1219
 reference time scale correlated, A(127):22-23, 103-105, 108, 110, 199-203, 208, 275, 280, 359, 364, A(128):170, 174-176, 317; B:970, 972
 silicoflagellate and ebriid zones correlated, A(128):162; B:240
 Site 794, A(127):72, 103-105, A(128):69, 100-105; B:969-970, 972, 1221-1223
 Site 795, A(127):174, 199ff; B:970-972, 1223
 Site 796, A(127):251, 275ff; B:1223
 Site 797, A(127):324, 357ff; B:972-973, 1223
 Site 798, A(128):30-31, 124-125, 166ff; B:399, 417, 972-974, 1223-1224
 Site 799, A(128):30-31, 244, 312ff, B:972, 974-975, 1224

- major element molar ratios
Site 798, B:616-617
Site 799, B:611, 618-621
- major elements, B:677-695. *See also specific major elements*
Al-normalized concentrations, B:687
alteration and, B:799, 901, 909, 911
in augite, B:852-854
in basalt, B:839, 869-870
basement, A(128):98; B:782, 893
in dark/light cycles, B:568, 570-571
and fractionation models, B:881
in ilmenite, B:856-857
interelement comparisons, B:690
of Japan Sea sediment, B:1235-1237, 1246-1248
in Japanese volcanic outcrops, B:874
in magnetite, B:856-857
mobility during alteration, B:838
in olivine, B:850
in Plagioclase, B:852-853
silica, B:682
Site 794, B:680, 682, 686-687, 782, 839, 858, 869-870, 893, 912-913, 918
Site 795, B:680-681, 683, 686-687, 913-914
Site 796, B:681
Site 797, B:681, 686-688, 839, 858, 869-870, 893, 914-916, 920
Site 798, B:720-722
Site 799, B:722-723, 726-727
in spinel, B:840-842, 856-857
in volcanic glass, B: 1378
vs. loss on ignition, B:918, 920
XRF and geochemical log analyses compared, B:1025-1029, 1031-1035
- Mamiya Strait
depth, B:440
Japan Sea and, B:362-363
sill depth, B:457
- Mammoth/Gauss boundary, Site 798, A(128):156
Mammoth Subchron, Site 798, B:1224
- manganese
accumulation, B:90-91
and cerium, B:726-727
in interstitial water, B:607
Japan Sea, B:1236
Site 795, B:710-711, 714
Site 798, A(128):172-173, 180; B:722, 1263
Site 799, A(128):317, 328; B:90-91, 610, 722, 727-729
vs. Ce anomaly, B:737
- manganese, hydrogenous, rare earth element source, B:729
- manganese nodules, and cerium, B:726-727
- manganese oxide. *See* manganese
- mantle
anisotropic velocities, B:1075
composition, B:786-787, 927
depleted, B:807-815, 903
heterogeneity, B:927
processes, B:927-928
and rare earth mixing model, B:817
Yamato Basin, B:927
- mantle source
and beryllium isotopes, B:815
and crustal contamination, B:807, 809-810
depleted asthenospheric mantle (AM), B:903
depleted mantle and pelagic sediment mixing plots, B:814
end-member compositions for mixing calculation, B:815
enriched mantle II (EMII) component, B:786-787
- enriched subcontinental upper mantle (SCUM), B:903
and Japan Sea basalt, B:903, 1341
and lithospheric contamination, B:810-812
N-MORB source depleted mantle (DM) component, B:786-787
and sediment contamination, B:805
Site 794, B:917
and slab-derived fluid contamination, B:809
and subducted sediment contamination, B:812-815
Yamato Basin, B:786-787
for Yamato Seamount Chain, B:812
marcasite, in vesicle, A(128):90
marginal basins, Pacific Ocean W, B:1311, 1333-1334
- marginal sea
formation, A(127):5
Japan Sea as, B:365-366
- Mariana Arc
Pb-Pb plots of arc volcanic rocks, B:808
sediment contamination of mantle source, B:805, 812
strontium isotopes vs. neodymium isotopes, B:809
- Mariana Trough
basement isotopic composition, B:807
lava flows, B:874
lava flows, chemical composition, B:882
Pb-Pb plots of arc volcanic rocks, B:808
- Maritime province, and sea ice, B:167
- mass accumulation rate. *See also* sediment accumulation rate
of barium, B:665
of carbonate, B:428
of organic carbon, B:425, 428
Site 798, B:428
- matrix sieving, and vein fillings, B:1179-1180
- Matuyama/Brunhes boundary, and inclination variations, B:974
- Matuyama Chron
and diatom occurrence, A(128):159
Site 796, A(127):275
Site 797, B:973
Site 798, A(128):124, 170
- Matuyama/Gauss boundary
silicoflagellate and ebridian zones correlated, A(128):162, 305
Site 794, B:970
Site 795, B:970
Site 796, A(127):275
Site 797, B:973, 1223
Site 798, A(128):30, 170; B:1224
Site 799, A(128):314; B:974, 1224
- Matuyama/Jaramillo boundary, inclination variations, A(128):172
- Matuyama/Olduvai boundary
Site 798, A(128):164
Site 799, A(128):307
- Mediterranean E, and trace metal composition, B:1361
- Meiyo-Daisan Seamount, A(128):70
- melt distribution, and electrical resistivity, A(128):111
- mesostasis
altered to clay minerals, A(127):217
Site 795, B:821
- methane
biogenic, A(127):251, 288
concentrations, A(127):119
in gas hydrate, A(127):288-290
Japan Sea, A(127):24
and methanogenesis, A(127):216
- nitrogen correlated, B:750
origin, B:749-750
potential production rates, B:771
shallow presence, A(127):259-260
Site 794, A(127):119
Site 795, A(127):174, 213-216, 220
Site 796, A(127):287-288, 290, 315
Site 797, A(127):368
Site 798, A(128):175-176, 180, 187; B:771-772, 776, 1263
Site 799, A(128):244, 321-322, 339
and sulfate abundance, A(127):368
thermogenic, A(127):251, 288
vs. depth, A(127):220
- methane-clathrate. *See* clathrate
- methane/ethane ratio (C_2/C_1)
in gas hydrate, A(127):288, 290
Site 795, A(127):220
Site 796, A(127):287-288, 290
Site 798, A(128):176, 188
Site 799, A(128):339
- methane/propane ratio (C_2/C_3)
Site 798, A(128):176, 188
Site 799, A(128):339
- methanogenesis
and bacteria, B:767-769, 771
and carbon isotopes, B:88-90
and interstitial water analyses, B:607
and methane content, A(127):216
and organic carbon decomposition, B:1262-1263
and organic matter degradation, A(127):207
Site 796, A(127):267
Site 797, A(127):368
Site 798, B:772-773, 776
- methylcyclohexane, Site 799, A(128):321-322
- methylcyclopentane, Site 799, A(128):321-322
- Mg#. *See* magnesium number
- Mg/Ca ratio. *See* magnesium/calcium ratio
- Mg-calcite, Site 796, A(127):266
- microbiological studies
bacteria, B:755-760, 1262
methods, B:755-756, 761-765
objectives, A(128):18, 24-27, 133
shipboard sampling procedures, A(128):179
shorebased laboratory studies, A(128):179-180
Site 798, A(128):24-27, 125-126, 178-180; B:755-776
- microfaults
and Japan Sea stress field, B:1187
Site 798, A(128):143ff
Site 799, A(128):269-272
and veins, B:1180
- microfossils. *See also* biostratigraphic zonations; biostratigraphy; *names of individual fossil groups*
abundance and preservation, A(127):19, 102, 192, 194-195, 269-271, 351-352, 354, A(128):28-30, 99, 158-159, 163, 298, 308-309
paleoenvironmental interpretation, A(127):20-22, 102-103, 199, 274, 356, A(128):99-100, 166, 312
phosphatized, B:64
preservation, B:1219
reworked, A(127):270, 356
Site 799, B:34-35
in tuffaceous clayey siltstone, A(128):68, 89
microfossils, calcareous, Leg 128, A(128):28
microfossils, siliceous
abundance and preservation, B:341-357

- abundance below shallowest opal-CT, A(127):19
dissolution, A(127):169; B:353
Leg 127, A(127):19
Leg 128, A(128):28
and opal-A to opal-CT transition, B:341
and opal dissolution transition zone (ODTZ), B:342
preservation, A(128):137
as productivity indicators, B:429-430
Site 794, B:341-357
Site 795, B:341-357
Site 797, B:341-357
micropumice, in ash-flow tuff, B:793
microquartz, in sandstone, B:144-148
microstructure, Site 799, B:33-48
mid-mean ridge basalt. *See* basalt, mid-ocean ridge
Milankovitch cycles. *See also* eccentricity frequencies; obliquity cycles
in FMS logs, B:1043-1044
mineral-liquid-disequilibrium, and spinel zoning, B:844
minerals, magnetic. *See* magnetic minerals
minor elements
in augite, B:854
in basement, B:912-916
in dark/light cycles, B:568, 570-571
Site 794, B:912-913
Site 795, B:913-914
Site 797, B:914-916
Site 798, B:720-722
Site 799, B:722-723, 726-727
Miocene, early/middle boundary, Site 799, A(128):299; B:1224
Miocene, middle/late boundary, Site 799, A(128):299; B:796, 1224
Miocene/Pliocene boundary
Site 794, A(127):97
Site 795, A(127):193; B:1223
Site 796, A(127):272
Site 797, A(127):351; B:1223
Site 798, B:1224
Site 799, A(128):298, 301-302, 305; B:1224
Mitoku-type flora, Japan Sea, B:486
Miura Group, veins observed, B:1180
mixed sediment
Site 795, A(127):186-187
Site 799, A(128):260
mixed sediment, diatomaceous clayey, Site 799, A(128):256
Mogami Channel, A(128):70
Moho
Japan Basin, A(128):246
Yamato Basin, A(128):246
Yamato Rise, A(128):245-246
molecular organic fossils
in dark/light cycles, B:670, 675
low-molecular-weight compound concentration, B:673
Site 798, B:669-670
Site 799, B:669-670
molybdenum
Site 794, B:1362
Site 795, B:1365
Site 797, B:1366
Site 798, B:1370-1371
monactinal monaxons
Site 795, B:542-543
monaxons
Site 795, B:541-543
Mongolia, dust storms, B:394
monsoons
and dark/light cycles, B:574-575
Japan Sea, B:320-321, 326
Monterey Formation
dolomite, B:75
lithostratigraphy, A(128):354
veins observed, B:1180
Monzen Formation
and age of rifting, B:1202
deposition of, A(128):17
MORB. *See* basalt, mid-ocean ridge
mordenite, Site 797, A(127):344
morphotaxa, faunal/flora, recognized in BSEM images, B:548
Mossbauer spectroscopy, B:739-746
absorption areas and sediment chemical compositions compared, B:741-742
iron species depth profiles, B:742-746
Mossbauer parameters, B:740-742
and partial dissolution of sediment samples, B:741
typical spectra, B:743
Mout St. Helens, stress field, B:1323
mud clasts, Site 799, A(128):267
mud plumes, Site 799, A(128):268-269; B:1178
mudstone, Site 799, B:42
Musasi Basin, dredge sample, A(127):176-177
Nakanami Formation, A(128):129
Nankai Trough, A(127):5-6, A(128):9
basement age, B:749
fissility, A(128):146
helium isotope ratios, B:748-749
interstitial water oxygen isotopes, B:698
veins, A(128):145-146; B:1180
nanofossils, calcareous
abundance and preservation, A(127):98-99, 193-195, 273, 351-355, A(128):28-30, 162-165, 305-310; B:161
and authigenic carbonates, B:166
and basal sediment age, B:1335
and carbonate-compensation depth fluctuations, B:1220
datums, Site 798, B:163-166
dissolution, A(128):162; B:166
diversity, B:175-177
events, B:172-174, 178
and Japan Sea formation age, B:1220
Leg 127, A(127):19-20
Leg 128, A(128):28, 30
magnetostratigraphy, B:163-164
monospecific assemblages, A(128):307
and oceanographic front fluctuations, B:167-168, 1220
paleoenvironmental interpretation, A(128):164-165
preservation determinations, B:161-162
reworked, A(127):99, 194
and sedimentation rate, B:1043-1044
Site 794, A(127):98-99; B:174-176, 1221
Site 795, A(127):193-195; B:176, 178
Site 796, A(127):273; B:176, 178
Site 797, A(127):351-355; B:177-179, 1223
Site 798, A(128):162-165; B:155-161, 164-166
Site 799, A(128):305-310; B:78, 155-163, 166
species diachrony, B:164
stratigraphic standards used, B:162-164
and surface-water temperature, A(127):273
units, Site 798, B:165-166
zonations, A(127):20, 98-99, 193-195, 273, 351-355, A(128):162-165, 305-310; B:172-174
naphthenic envelope, Site 799, B:627-628
natural gamma-ray tool (NGT). *See* gamma ray, logging
natural remanent magnetization (NRM). *See* magnetic properties
navigation, during downhole seismic experiment, B:1107-1108
neodymium
in ash layers, B:1386
in basement rocks, B:807
Site 798, B:1370-1371
vs. strontium, B:902
vs. titanium, B:902
neodymium isotopes. *See* isotopes, neodymium
neopentane, Site 799, A(128):321-322
nesquehonite, and alkalinity, B:92
neutron activation (INAA) analyses
of basement, B:912-916
Site 794, B:912-913
Site 795, B:913-914
Site 797, B:914-916
neutron porosity. *See* porosity, logging
NGT. *See* gamma-ray logging
nickel
and basalt fractionation models, B:873
in dark/light cycles, B:569
and fractionation models, B:881
Site 794, B:893, 1362
Site 795, B:1365
Site 797, B:893, 920, 1366
Niigata Basin, lithostratigraphy, A(128):17
niobium
basement, A(128):99
vs. zirconium, A(128):100; B:783
Nishikurosawa Formation
deposition, A(128):17
and subsidence history of Oga Peninsula, B:1202
nitrate, and nitrate-reducing bacteria, B:770
nitrogen
in dark/light cycles, B:431
methane correlated, B:750
Site 794, A(127):113-114
Site 795, A(127):209-212; B:708-709
Site 796, A(127):283-285
Site 797, A(127):365-367
Site 798, A(128):176-177, 189-192
Site 799, A(128):323-324, 334-338, 341
vs. carbon, organic, A(128):343-344
nodules, carbonate
with diatoms, A(127):193
Site 795, A(127):186
Site 797, A(127):343
nodules, phosphatic, Site 797, A(127):344, 346
non-euxinic-anoxic condition. *See* oxygenation condition
normal faults
in basement, A(128):91-93
and fractures, A(127):188
Kita-Yamato Trough, A(127):334
Oki Ridge, A(128):130
Oki Trough, A(128):130
Site 795, A(127):188
Site 798, A(128):145; B:1181-1182
Site 799, A(128):263, 267, 269-272, 294-295; B:1182-1183, 1189, 1190
and stress field, B:1182-1183
and veins, B:1178
Yamato Basin, A(128):130
Yamato Rise, A(128):76
Yamato Rise E, A(127):82-83
normalization factors, and geochemical logs, B:1029-1030, 1034-1035
North American Plate, A(127):5, A(128):9

- North Atlantic Deep Water, and biosiliceous sediments, B:439
- North Korean Current, A(128):21; B:343, 440
- northern hemisphere glaciation. *See* glaciation
- Noto Peninsula
- diatom abundance, B:311
 - and Japan Sea stress field, B:1185
 - rock exposures, A(128):129
 - uplift, A(128):158
- NRM intensity. *See* magnetic properties
- Nunivak Subchron
- Site 797, B:1223
 - Site 799, A(128):314; B:1224
- nutrient supply, and formation of color-banded bedding, A(127):349-351
- oblique electrical resistivity experiment. *See* electrical resistivity experiment
- obliquity cycles
- in biogenic opal, B:447
 - in dark/light cycles, B:446-447, 571, 574-575
 - in FMS logs, B:1043-1044
 - in geochemical logs, B:1021
 - in logs, B:403, 1221
 - of surface-water temperatures, B:467-468
- obliquity frequency. *See* obliquity cycles
- ocean bottom seismometer (OBS). *See also* air-gun profiling experiment; downhole seismic experiment
- air-gun signals in downhole seismic experiment, B:1162
 - anisotropic models, B:1117
 - array location, B:1070, 1077, 1109, 1159
 - and crustal structure, B:1075, 1342
 - data variance, B:1115
 - downhole seismometers compared, B:1061, 1078, 1084
 - first arrival traveltimes, B:1089
 - frequency response, B:1159, 1162
 - and navigation, B:1107-1108
 - ray tracing results, B:1092-1103
 - record sections obtained, B:1080-1083
 - slowness vectors, B:1118
 - specifications, for downhole seismic experiment, B:1160
 - traveltime, offset distance, and basement depth correlated, B:1112
 - velocity structure, one-dimensional, B:1078-1081, 1085-1086
 - velocity structure, two-dimensional, B:1081-1083, 1086, 1089
- oceanic crust. *See* crust, oceanic
- oceanization, Japan Sea, A(128):11
- oceanographic front
- and calcareous nannofossils, B:167-168, 1220
 - in Japan Sea, B:155, 167, 458
 - oscillations, B:167-168
 - oxygen isotope curve correlated, B:168
 - and productivity, B:167
 - and sea surface isotherms, B:167
- oceanographic frontal boundary. *See* oceanographic front
- ODTZ. *See* opal dissolution transition zone
- Oga Peninsula, A(128):70
- age, B:1202
 - basement, B:1202
 - benthic foraminifers, A(127):198; B:379, 381
 - depositional environment, B:1203
 - lithostratigraphy, A(128):17-18; B:1203
 - paleobathymetry, B:1203-1204, 1210
 - planktonic foraminifers exposed, B:459
 - sediment accumulation rate, B:1204
 - stress field, B:1204-1205
 - subsidence/uplift history, B:1202-1205
- uplift, A(128):158; B:1202-1205
- OI. *See* oxygen index
- Okhotsk microplate, A(128):9
- Okhotsk Sea
- diatoms, B:309-310
 - formation, A(128):13
 - and opal flux, B:439
- Oki-Dogo Island, ash layers from, A(128):156
- Oki Islands, A(128):126-127
- basement, A(128):127-129
 - Green Tuff rocks, A(128):128
 - and low-chlorinity pore fluids, A(128):172
- Oki Ridge
- ash layers, A(128):122
 - basement, A(128):127-129
 - bathymetry, A(128):126-127
 - benthic foraminifers, B:367-369, 373-375, 382
 - biostratigraphy, B:1223-1224
 - bottom water conditions, A(128):125
 - carbonate-compensation depth, A(128):122, 158; B:409
 - chronostratigraphy, B:317-318
 - consolidation/permeability studies, B:1123-1133
 - crustal structure, A(128):127-128
 - cyclic lithofacies, A(128):195-196
 - depositional history, A(128):157-158, 194-195
 - diatoms, B:361
 - ebriidians, B:241-244
 - eoian supply, B:409, 419
 - faulting, A(128):194-195
 - forests, B:317
 - free-air gravity anomalies, A(128):127, 133
 - Green Tuff rocks, A(128):127-128
 - heat flow, A(128):125, 127, 131; B:1302
 - Layer 1, A(128):127
 - Layer 2, A(128):127-128
 - Layer 3, A(128):127-128
 - magnetic anomalies, A(128):127, 132
 - magnetostratigraphy, B:973-974, 1223-1224
 - oceanographic conditions, A(128):166, 195-196
 - oceanographic drilling objectives, A(128):18-19
 - oxygen isotope stratigraphy, B:444
 - paleoclimatic conditions, B:419-421
 - paleodepth, B:373
 - paleoenvironment, B:379-380
 - pollen, B:318-324
 - radiolarians, B:225, 227, 229-230
 - sediment, B:409-422
 - sedimentary rocks dredged, A(128):129
 - sedimentation, A(128):131-132; B:1227
 - seismic stratigraphy, A(128):34, 125
 - silicoflagellate, B:241-244
 - Site 798, B:393, 409, 423-425, 440, 458-459, 560, 605, 667, 720, 739
 - tectonic setting, A(128):130-131, 134
 - uplift, A(128):35, 122, 157, 194-195; B:155, 373, 419, 1208
- Oki Ridge Basin, seismic stratigraphy, A(128):122
- Oki Trough
- bathymetry, A(128):126-127
 - crustal structure, A(128):130
 - free-air gravity anomalies, A(128):127
 - heat flow, A(128):131
 - Layer 1, A(128):127
- Okinawa Trough, sulfide deposits, A(128):21
- Okushiri Island, A(127):251, 254, A(128):76
- Okushiri Ridge
- anomalous heat flow, A(127):300-301
- basement, A(127):176, 312
 - bathymetry, A(127):251
 - biostratigraphy, B:1223
 - calcareous nannofossils, B:176, 178
 - compression, A(127):255
 - crustal structure, A(127):251-252
 - diatoms, B:270-275, 285
 - dredge samples, A(127):254
 - earthquakes along, A(127):178
 - fluid flow along faults, A(127):275
 - folding, A(127):255-256
 - frictional heating along thrust faults, A(127):247
 - heat flow, A(127):27, 30, 300-301, 314; B:1303, 1307
 - interstitial water chemistry, B:1265
 - magnetostratigraphy, B:1223
 - opal-A to opal-CT transition, B:50
 - opal-CT to quartz transition, B:50
 - physical properties, B:1275
 - pollen, B:471, 481, 484-485
 - sediment thickness, A(127):313-314
 - sedimentation, A(127):256-258; B:288, 1227
 - seismic stratigraphy, A(127):256-258, 307, 312-313
 - slumping, A(127):268
 - tectonic history, A(127):176
 - thermal model, A(127):16
 - uplift, A(127):19, 30, 176, 247, 258, 268, 314; B:1208
- Olduvai Event. *See* Olduvai Subchron
- Olduvai Subchron
- gamma-ray logging, B:403
 - and sedimentation rate, B:975
 - silicoflagellate and ebridian zones correlated, A(128):162
 - Site 794, B:218, 224, 970
 - Site 795, A(127):174, 199; B:970, 1223
 - Site 797, B:224
 - Site 798, A(128):30, 125, 170; B:164, 564, 974, 1224
 - Site 799, A(128):314; B:974, 1224
- olivine
- altered to clay, B:885-886
 - experimental chemical compositions, B:866
 - forsterite and, B:850
 - importance in basalt fractionation models, B:873-874
 - in melting experiment, B:861-862, 895
 - and parent magma, B:855-856
 - Site 794, B:850
 - Site 797, B:850
- Oman margin, and chloride, A(128):172
- Onnagawa Formation, A(128):13, 265
- ash-flow tuffs correlated, B:796
 - deposition, A(128):17
 - dolomite, B:75
 - foraminifers, B:187
 - lithostratigraphy, A(128):354
 - opal flux, B:439, 446
 - paleoenvironmental interpretation, B:1202
- Ontong Java Plateau, magnetic properties, B:951
- ooze, clayey diatom
- and revised lithostratigraphy, B:1230-1232
 - Site 794, A(127):90
 - Site 796, A(127):264
 - Site 797, A(127):341-343
 - Site 799, A(128):260
- ooze, diatom
- aluminum, logging, A(127):306-307
 - and enhanced productivity, A(127):267
 - and oxygenation conditions, B:350
 - and revised lithostratigraphy, B:1230-1232

- sedimentation rate, A(127):204
 Site 794, A(127):90, A(128):77
 Site 795, A(127):186
 Site 797, A(127):341-343
 Site 798, A(128):124, 137-138
 Site 799, A(128):256, 260
- opal
 abundance variations seen in logs, B:398-400
 and geochemical logging, B:1022-1024
 physical properties correlated, B:400
 stratigraphy, B:1221
- opal, biogenic. *See also* opal-A; opal flux; silica, biogenic
 abundance, B:443, 446-447, 449-455
 cycles, B:444-446
 in dark/light cycles, B:432, 446, 569
 obliquity cycles, B:447
 seismic expression, B:1140-1141, 1143
 Site 798, B:443-446, 449-455
 variations in supply, B:404
 vs. age, B:445
 vs. depth, B:445
- opal-A. *See also* opal; opal, biogenic; silica, biogenic
 abundance, A(127):94; B:1235-1241
 deepest occurrence, A(127):187
 and lithium, A(128):174-175
 and low-chloride interstitial fluids, B:608-610
 physical properties interrelationships, B:25-27
 vs. excess silica, B:1253
- opal-A to opal-CT transition, A(128):28; B:6-17, 1145-1156. *See also* silica diagenesis and acoustic impedance, B:1146
 age, A(127): 19; B:251, 1145-1147, 1305-1306
 as barrier to interstitial water, A(128):244
 bottom simulating reflector, B:1145-1147, 1149, 1152-1153
 and chloride concentration, B:609-610
 and deepest diatom occurrence, A(127):264
 and density, B:18, 1146-1147, 1288
 depth, B:1224-1227
 diatoms, B:3, 249, 313, 361, 1230-1232
 and dissolution patterns, B:311
 and dissolved silica maximum, A(127):174
 factors controlling phase change, B:1145
 in geochemical logs, B:1417
 and heat flow, B:1147-1148, 1150, 1156
 and inorganic geochemical data, A(127):23, 187, 204-209, 362-364, A(128):32, 321; B:607
 Japan Sea, A(127):31; B:49, 341
 Leg 127, A(127):19
 Leg 128, B:359
 and lithium content, B:1266
 lithology and, B:17
 in logs, A(127):139, 251, 302, 393-394, A(128):186-187, 333; B:3, 15-16, 19, 21-22
 and opal-CT abundance, B:10
 oxygen isotopes and, B:641-642
 and physical properties, A(127):26, 127, 133-134, 187, 223, 290-295, 388-389, A(128):33; B:10-18, 1145, 1277-1278, 1284, 1305
 and porosity, B:3, 6, 16, 18
 and radiolarian preservation, A(127):198, 355, A(128):303
 and radiolarians, B:226, 291-292, 295
 and rate of opal-CT precipitation, B:1306
 and sedimentation rate, B:1147-1148
 seismic expression, A(127):28, 143, 169, 179, 234-236, 251, 307-314, 403, 407, 410, A(128):13, 193; B:3, 1141ff, 1305
 and silica, A(128):125, B:1263
 and siliceous microfossils, A(128):28; B:341
 Site 794, A(127):111
 Site 795, A(127):169, 198; B:4, 50
 Site 796, A(127):247; B:50
 Site 797, A(127):343, 351; B:5, 50
 Site 798, A(128):149, 240
 Site 799, A(128):238, 260, 278
 and slumping of slope, A(127):268
 in synthetic seismograms, B:1146-1148
 temperature, A(127):19, 137, 232, 347, 392, A(128):150, 280; B:14, 1145-1147, 1152, 1263, 1305-1307, 1375
 and thermal conductivity, B:1147
 and thermal gradient, A(127):268, 314
 and thermal history, Japan Sea, B:1305-1307
 as thermal marker, B:3
 thickness, B:16, 1240, 1281-1282
 time-temperature plot, B:1266, 1288
 transition temperature and sedimentation rate correlated, B:1306
 and velocity, B:18
 and volatile hydrocarbons, A(128):323
 X-ray diffraction analyses, A(127):347; B:10, 17-18
 and zeolites, B:10
- opal abundance
 core analyses vs. log estimates, B:404
 and dissolution patterns, B:310-311
 logging estimates, B:402
 and productivity, B:311
 Site 798, B:401-402
 vs. density, B:402
 vs. porosity, B:402
- opal accumulation rate. *See* opal flux
- opal-CT
 abundance, A(127):94, A(128):141; B:53, 1235-1241
 deepest occurrence, A(127):189, A(128):280
 formation, A(127):109; B:49
 isotopic temperatures, B:55-56
 and opal-A to opal-CT transition, B:10
 oxygen isotopes, B:51-54
 permeability, B:1263
 physical properties, B:25-27
 porosity, B:1263
 shallowest occurrence, A(127):187, 343, 347, A(128):124, 138, 149, 174, 278
 Site 795, A(127):190
 and thermal gradient, B:49
 time-temperature plot for opal-CT formation, B:1266
 vs. excess silica, B:1253
- opal-CT to quartz transition, B:16-17. *See also* silica diagenesis
 age, A(127):19
 and core recovery, B:1038
 and inorganic geochemical data, A(127):204
 Japan Sea, A(127):31, B:49
 in logs, A(127):303, A(128):34, 333; B:3, 23
 and physical properties, A(127):223-227
 and porosity, B:3
 resistivity logging data and, B:1040
 SEM photographs, B:31
 and silica content, B:1263
 silicon content, A(127):307
 Site 794, B:50
 Site 795, A(127):169; B:4, 50
 Site 796, A(127):265; B:50
 Site 797, A(127):347; B:4, 50
 Site 799, A(128):238, 242, 260, 280
- temperature, A(127):19, 137, A(128):280; B:17, 1263, 1375
 thickness, B:17, 1240, 1281-1282
 time-temperature plot, B:1288
 X-ray diffraction analyses, B:24
- opal dissolution transition zone (ODTZ)
 age, B:350
 Japan Sea, B:342
 lithostratigraphy correlated, B:347-349
 and Northern Hemisphere glaciation, B:350
 photomicrographs, B:355-357
 and radiolarians, B:345-347
 Site 794, B:344
 Site 795, B:345
 Site 797, B:345
 thickness, B:347, 350
- opal flux. *See also* opal, biogenic; opal-A; silica, biogenic
 in Bering Sea, B:439
 in Onnagawa Formation, B:439, 446
 in Santa Barbara Basin, B:439
 in Sea of Okhotsk, B:439
 Site 798, B:439-455
- operations. *See particular site*
- orbital obliquity, in logs, A(128):34, 342
- orbital obliquity cycles. *See* obliquity cycles
- orbital precession cycles. *See* precession cycles
- organic carbon accumulation rate. *See* carbon, organic
- organic fossils. *See* molecular organic fossils
- organic matter. *See also* carbon, organic; molecular organic fossils; organic matter decomposition
 autochthonous, B:627-628
 bacterial degradation, A(127):22
 and barium remobilization, B:654
 and boron content, B:640
 composition, B:423-429
 in cyclic lithofacies, A(128):140
 in dark/light cycles, B:446, 564, 584, 668-669
 degradation, A(128):172-173, 318; B:646
 diagenesis, B:1262-1263, 1267
 and geochemical logs, A(128):187
 and hydrogen index (HI), B:628
 and inorganic geochemical data, A(127):204
 lipid fraction, A(128):324
 maturity, B:429
 origin, A(128): 125, 324
 and paleoenvironmental conditions, B:670-672
 recycled, B:625
 and sedimentation rate, B:1265
 Site 798, A(128):125; B:446, 667-669
 Site 799, A(128):245, 323-324; B:624, 667-669
 and thermal maturity, B:671-672
- organic matter decomposition
 and high ammonium values, A(127):279-280
 and interstitial water analyses, B:607
 Japan Sea, A(127):31
 and salinity, A(128):172
 Site 799, A(128):318
- organic petrology. *See* organic matter
- organic-rich sediments. *See* sediments, organic-rich
- orthoclase, vs. anorthite, B:122
- orthopyroxene
 in ash layers, A(128):151; B:1379
 in melting experiment, B:895
- Oshima Island, basement, A(128):73
- Oshima Plateau, dredge samples, A(127):78
- overburden pressure. *See* consolidation
 and stress field analysis, B:1050-1051

- overconsolidation. *See* consolidation
overconsolidation ratio (OCR). *See* consolidation
oxeas
Site 795, B:542
oxic conditions. *See* oxygenation conditions
oxides. *See* specific elements
oxides, elemental. *See* specific element
oxygen
benthic foraminifers and, B:528
fugacity, inferred from spinel analyses. B:842-844
in Japan Sea, A(128):20
oxygen index (OI)
in dark/light cycles, B:431-432
Site 794, A(127):115-118
Site 795, A(127):213, 219
Site 796, A(127):285
Site 797, A(127):367
Site 798, A(128):177, 196; B:425, 668-669
Site 799, B:429, 624, 626-627, 668-669
oxygen isotopes. *See* isotopes, oxygen
oxygen level, and formation of color-banded bedding, A(127):349
oxygenation conditions
and bioturbation, A(127):267, 315
and burrows, B:1232
and carbon/sulfur ratio, B:1249
and Ce anomaly, B:692
and dark/light cycles, B:432, 574-575, 590-591, 595
determined by sulfur vs. organic carbon plots, B:708-709
and diagenetic carbonate, B:93-94
and dolomite formation, B:92
glacial/interglacial variation, B:590-591
Japan Basin, B:201-208
Japan Basin N, A(127):199
Japan Sea, B:216, 342-343, 430, 678-679, 705, 713-715, 1221
Kita-Yamato Trough, A(128):354-355
and manganese concentration, B:90
and organic carbon content, A(127):283
and organic geochemical data, A(127):267
and oxygen isotopes, B:442-443
Site 794, A(127):96; B:209
Site 795, A(127):212; B:708-709, 713-715
Site 796, A(127):247, 251, 267
Site 797, A(127):349; B:209, 593
Site 799, B:76-77, 91-92
and wind velocity, B:447
Yamato Basin, B:189, 201-208
Oyashio Current, B:343
climate and, B:326
cold surface water, B:457
Japan Sea and, A(128):30
Site 799, A(128):312
p-wave velocity. *See* velocity
Pacific Bottom Water, helium, B:748
Pacific Deep Water, A(127):16
Japan Sea and, B:365
Pacific Ocean
benthic foraminifers, B:505
deep-water inflow, B:528
eolian dust, B:394
Japan Sea and, A(128):312; B:180, 299-300
tectonic setting, A(128):8
Pacific Ocean, equatorial, C_{org} /barium ratio, B:653-654
Pacific Ocean N
biosiliceous sediments, B:439
diatoms, A(128):160, 302, 315-316; B:258, 341, 358
foraminifer coiling ratio, B:459
paleoclimate studies, B:459
pollen, B:320
radiolarians, B:228
silicoflagellate, B:237
Site 794, A(127):96
Pacific Ocean NE, planktonic foraminifers, B:465
Pacific Ocean NW, diatoms, B:309, 360
Pacific Ocean SW, paleoenvironment, B:338
Pacific Ocean W
eolian dust, B:403
marginal basins, B:1311, 1333-1334
subduction zones, B:1333-1334
tectonic setting, A(127):7
Pacific Plate, A(127):5, A(128):9
electrical resistivity structure, B:1351
Pacific Plate oceanic model (PPOM), B:1304
paleodepth
benthic foraminifers and, B:504-506
Japan Basin, B:201
Japan Sea, B:526, 528-529
Site 797, A(127):357
Site 798, B:373
Site 799, B:373, 379
Yamato Basin, B:201
paleoecology, Japanese archipelago, B:320, 334
paleoenvironment. *See also* specific regions and topics
pollen and, B:320
Site 794, A(127):103, A(128):99-100
Site 795, A(127):199
Site 796, A(127):274
Site 797, A(127):357
Site 798, A(128):166; B:379-380
Site 799, A(128):312; B:381
paleomagnetism. *See* magnetic properties; magnetostratigraphy
paleoproductivity. *See* productivity
palyinology, Japan Sea, B:317-321, 325-338
parent magma
and augite, B:854
of basalt, B:855-856
of basement, B:893-894
estimates, B:894
Japan Basin, B:927-928
Japan Sea, B:1341
origin, B:897
production depth, B:897
Site 794, B:844-845, 917
Yamato Basin, B:927-928
pebbly claystone. *See* claystone, pebbly
PEF. *See* photoelectric factor
pelite, Site 799, B:33
n-pentane, Site 799, A(128):321-322
peralkaline glass, B:1378
periodicity, orbital. *See also* obliquity cycles
in logs, B:396-398
permeability
in biosiliceous sediments, B:1128
and grain size, B:1127-1128
and opal-CT formation, B:1263
porosity effect, B:1128-1129
Site 798, B:1127-1128, 1131-1132
Site 799, B:39, 1128, 1131-1132
and temperature measurements, A(127):299
vs. porosity, B:1132
Peru/chile Current, diatoms, A(128):16
Peru margin
bacteria, B:755, 757, 759, 761, 769, 773
chloride, A(128):172
D-phosphate nodules, B:38
fluoride, B:66
sediment consolidation, B:1277
vein fillings, B:1178
petrology, experimental
AMF diagram for experimental glasses, B:865
An mole percent, Plagioclase vs. Fo mole percent, B:867
anhydrous melting-phase relations, B:896
augite, Cr and Ti vs. Mg/(Mg + Fe), B:867
augite, Fe/Mg vs. Fe/Mg for coexisting liquid, B:868
of basaltic basement, B:861-868, 891-898
chemical compositions of experimental materials, B:863-864, 866-868, 894
crystallization, B:864-866
and differential processes of basement basaltic rocks, B:895-897
equilibrium materials, B:862-863
exchange coefficients, B:864
experimental conditions, B:862
experimental liquid line of descent, B:863
experimental phase proportions vs. liquid percentage, B:862
experimental pyroxene-liquid relations and natural augite compared, B:878-879
experimental results and Site 794 basalt compared, B:870
Fe/Mg olivine vs. Fe/Mg coexisting liquid, B:866
melting experiments, B:861, 894-897
mineral-liquid equilibria, B:862-863
minerals crystallized, B:861-862, 895-896
normative projections, B:868
precision of microprobe analyses, B:862
and primary magma production, B:897
procedures used, B:861, 895
starting materials, B:861-862, 894-895
pH
in interstitial water, B:94
Site 794, A(127):107
Site 795, A(127):205
Site 796, A(127):283
Site 797, A(127):364, 374
Site 799, A(128):317-318, 328; B:610
phase equilibria, in basalt, B:870
phasor dual induction tool (DIT). *See* resistivity, logging
phenocrysts
augite, B:851-853
in basaltic basement, B:892
ilmenite, B:853
olivine, B:850
plagioclase, B:850-851
Site 794, B:849
Site 795, B:850
Site 797, B:850
spinel, B:853
Philippine Basin, rare earth elements, B:719
Philippine Sea
basement isotope geochemistry, B:805
Pb-Pb plots of arc volcanic rocks, B:808
Philippine Sea Plate, A(127):5-6, A(128):9
electrical resistivity structure, B:1351
phillipsite, Site 797, B:139
phonolite, ash layers, B:793
phosphate. *See also* D-phosphate; F-phosphate; phosphorus
abundance, B:36
allochthonous, B:64, 73
chemistry, B:64-65
in diagenetic sequence, B:39
in interstitial water, B:36-38, 64-66, 607, 1264
microprobe analyses, B:64-67
and organic carbon decomposition, B:1262
pelagic, B:36

- peloidal, B:36
 pristine, B:63-64, 73
 and sedimentation rate, B:67
 and siliceous veins, B:64
 Site 794, A(127):108
 Site 795, A(127):205
 Site 796, A(127):279-280
 Site 797, A(127):362, 368
 Site 798, A(128):173, 181; B:63-74
 Site 799, A(128):317-318, 328; B:36-38, 47, 63-74, 610
 source of allochthonous particles, B:67-68
 total oxide values, B:65
 X-ray fluorescence analyses, B:64-65
 phosphate, Sites 798 and 799, B:64-66
 phosphatic nodules. *See* nodules, phosphatic
 phosphatic sediments, Kita-Yamato Brink, A(128):249
 phosphorus. *See also* phosphate
 Japan Sea, B:68-69, 1237
 P_{org}/P_{inorg} ratio, B:71
 phosphorus oxide. *See* phosphorus
 photoelectric factor (PEF), and magnesium and sodium content, B:1395, 1415
 phreatomagmatic eruption, submarine, B:119, 123-126
 physical properties. *See also* specific physical properties analyses; *specific physical properties logs*
 basement, A(127):26, 135, 227-228, 389, A(128):102ff; B:1282-1284, 1289-1291
 basin and rise sites compared, Japan Sea, B:1275-1296
 consolidation studies, B:1123-1133
 corrected analyses, B:990-1015
 data quality, A(127):127, 220-222, 289-290, 382-383; B:987-988
 determination of wet volumes, B:988-989
 and diatom abundance, A(127):222
 error identification, B:985-986
 index properties, B:985-1015
 of interbedded basement/sediments, A(127):394; B:1283-1284
 interrelations at opal-A to opal-CT transition, B:17-18
 Japan Sea, A(127):25-26, A(128):33
 lithostratigraphy correlated, A(127):127-135, 222-227, 290-295, 383-389, A(128):182, 327-329; B:1143
 logs correlated, A(128):338
 opal-A interrelations, B:25-27
 and opal-A to opal-CT transition, B:10-16, 1145, 1277-1278, 1281-1282, 1305
 opal content correlated, B:400
 opal-CT interrelations, B:25-27
 permeability studies, B:1123-1133
 quartz interrelations, B:25-27
 seismic stratigraphy correlated, A(128):224, 379; B:1143
 and silica diagenesis, A(128):280; B:3-31, 1281-1282
 Site 794, A(127):73, 127-136, 138-140, A(128):102ff
 Site 795, A(127):174, 220ff, 231
 Site 796, A(127):251, 289-297, B:1275
 Site 797, A(127):325, 382ff, 392-393, 395
 Site 798, A(128):33, 180ff
 Site 799, A(128):33, 324ff; B:1137-1140
 Sites 794 and 795 compared, A(127):228
 Sites 794 and 797 compared, A(127):389, 396
 systematic errors corrected, B:985-1015
 water loss discrepancy, B:988
 X-ray fluorescence basement analyses compared, B:1283
 phytane, Site 799, A(128):343; B:628-629, 633
 phytoplankton, Japan Sea, B:167
 Pisco Formation, veins, B:1180
 placoliths
 Site 799, A(128):309
 plagioclase. *See* feldspar, plagioclase
 plagioclase feldspar. *See* feldspar, plagioclase
 plankton blooms. *See* blooms, plankton
 plastic limit. *See* Atterberg limits
 plasticity index. *See* Atterberg limits
 plate reconstructions
 geometrical constraints, B:1311-1315
 Japan Sea, B:1321-1324
 kinematic model used, B:1315
 Miocene, B:1322-1324
 Pliocene-Quaternary, B:1321-1322
 Present time, B:1321
 reconstruction poles, B:1325
 tectonic events, B:1316-1317, 1324-1325
 plateau ages
 Site 794, B:820
 Site 795, B:820-821
 Site 797, B:820, 829
 Pliocene/Miocene boundary, Site 794, A(127):96
 Pliocene/Pleistocene boundary
 and biosiliceous productivity, B:313
 and diatom assemblages, A(128):160
 diatoms and, B:250
 and foraminiferal fauna, B:208
 and microfossil dissolution, A(128):28
 Site 794, A(127):97; B:1221
 Site 795, B:1223
 Site 796, B:1223
 Site 797, A(127):351, 355-356; B:1223
 Site 798, A(128):162, 166; B:1224
 Site 799, A(128):301; B:1224
 Pliocene/Quaternary boundary, Site 796, A(127):251
 Pohang Basin, subsidence/uplift history, B:1205-1206
 polarity. *See* magnetostratigraphy
 pollen, B:1221
 and basal sediment age, B:1335
 distribution, B:489
 flora represented, B:479, 484-486
 and glacial/interglacial periods, B:395
 Japan Sea, B:328-336, 395
 marine, B:320-321, 337-338
 Site 794, B:328, 471, 482, 485
 Site 795, B:328, 471, 480, 484
 Site 796, B:471, 481, 484-485
 Site 797, B:328, 479, 483, 485-486, 1223
 Site 798, B:318-324
 Site 799, A(128):302-303
 sources, B:332-333
 stratigraphy, B:486, 488
 zones, B:479, 484
 polyaxons
 Site 795, B:543
 porcellanite
 conglomerate pebbles, A(127):344-345
 elastic moduli, A(127):223
 in logs, A(127):27-28, 394; B:1040
 and revised lithostratigraphy, B:1230-1232
 seismic expression, A(127):407-408
 Site 795, A(127):187
 Site 799, A(128):260, 264; B:33, 40, 41, 44
 pore gas. *See* interstitial gas
 pore space, and clathrate formation, A(127):281
 pore water. *See* interstitial water
 porosity. *See also* physical properties
 basement, B:1289
 corrected analyses, B:987, 990-1015
 data quality, A(127):127, 228-229; B:987
 and diatom internal porosity, B:1277
 errors in wet volume measurement, B:985-986
 and geochemical logging, B:1022-1024, 1029
 and horizontal stress, B:1053
 in interbedded sediment/igneous rocks, B:1278
 Japan Sea, B:1275-1278
 and loss on ignition, B:1283
 and normalization factors, B:1035
 and opal-A to opal-CT transition, B:3, 6, 16, 18, 21, 1277-1278, 1282, 1298
 and opal-CT formation, B:1263
 and opal-CT to quartz transition, B:3, 1282
 permeability and, B:1128-1129
 Site 794, A(127):127-133, A(128):103, 105; B:6, 1050, 1278-1279
 Site 795, A(127):223-227, 230; B:6, 1278-1279
 Site 796, A(127):290-295; B:6, 1278-1279
 Site 797, A(127):383-389, 391; B:6, 1278-1279, 1300
 Site 798, A(128):181, 197-198, 202-205, 209; B:6, 1278-1279
 Site 799, A(128):325, 345-349, 355; B:6, 39, 700, 1278-1279
 thermal conductivity correlated, B:987
 velocity and, B:1280
 vs. density, B:25, 27, 986, 989, 1281, 1290
 vs. loss on ignition, B:1290
 vs. opal abundance, B:402
 vs. permeability, B:1132
 vs. thermal conductivity, A(127):135, 232, 395; B:25, 27, 989, 1287
 vs. velocity, A(127):140, 395; B:25, 27, 1005, 1286, 1290
 vs. water content, A(127):234
 porosity, logging
 and geochemical logs compared, B:1023
 Site 794, A(127):142, 147, 164-167
 Site 795, A(127):244-245
 Site 796, A(127):303-306, 319-320
 Site 797, A(127):395-396, 403, 419-421; B:7
 Site 798, A(128):185-187, 212
 Site 799, A(128):383-385, 389-392
 vs. velocity, B:26
 potassium
 alteration and, B:909,911
 in ash layers, B:1385
 and basement alteration, B:892, 1265-1266
 in clay minerals, A(127):205
 in igneous rocks, A(128):103
 in interstitial water, B:1270
 in Japan Sea sediment, B:1237
 and magma suite distinction, B:869
 partitioned into plagioclase, B:856
 Site 794, A(127):109, A(128):108
 Site 795, A(127):205
 Site 796, A(127):280-281
 Site 797, A(127):364, 371
 Site 798, A(128):174-175, 184
 Site 799, A(128):318, 332; B:611, 723
 vs. hydrogen isotopes, B:1273
 vs. oxygen isotopes, B:1273
 vs. sodium, B:732, 858
 XRF and geochemical log analyses compared, B:1028-1029, 1031-1035
 potassium, logging
 and diatom vs. clay content, B:1417
 Site 794, B:1416, 1418-1421
 Site 796, B:1416, 1422-1423
 Site 797, B:1424-1427

- Site 798, B:1400-1402
 Site 799, B:1403-1404, 1406-1409
 vs. aluminum, logging, A(128):215
 potassium/aluminum ratio, Site 794, B:1249
 potassium/argon age. *See* K/Ar age; radiometric age
 potassium feldspar. *See* feldspar, potassium
 potassium oxide. *See* potassium
 precession cycles
 and dark/light cycles, B:571, 574-575
 in FMS logs, B:1043-1044
 precipitation
 Japan Sea, B:337, 442-443
 of silica, A(127):174
 vegetation and, B:325-326
 preconsolidation pressure. *See* consolidation
 pressure, hydrostatic, thermal conductivity and,
 B:1017-1019
 pressure compensation level (PCL), and vesicular
 hydroclasts, B:119
 primary magma. *See* parent magma
 primary productivity. *See* productivity
 pristane, Site 799, A(128):343; B:628-629, 633
 pristane/phytane ratio, Site 799, B:628
 productivity
 and barium content, B:652, 661-662
 in dark/light cycles, B:431-432, 591-594
 and diatoms, A(128):122; B:342
 estimate methods, B:425
 Japan Sea, A(128):21, B:167, 296, 299, 651
 Kita-Yamato Trough, A(128):355
 and opal abundance, B:311
 and organic carbon, A(127):283; B:430
 and sea ice, B:311
 and sedimentation rate, B:1228
 and silica content, B:694
 and siliceous microfossil abundance, B:430
 Site 796, A(127):274
 and Tsushima Current, B:311
 propane
 Site 795, A(127):174, 213-216, 220
 Site 797, A(127):368
 Site 798, A(128):125, 176, 187
 Site 799, A(128):244-245, 321-322, 339
 proto-dolomite, Site 799, B:611
 provenance diagrams, and tectonic setting, B:109
 pull-apart basin, Japan Sea, A(128):12-13
 pumice, long-tube, B:117, 130
 pycnometer, and wet sample volume analysis,
 B:985-986, 988-989
 pyrite
 abundance, B:1235-1241
 in ash layers, B:1392
 in Black Sea, B:712
 in dark claystone, A(127):267
 in dark-colored sediments, A(127):211
 in dark/light cycles, B:584, 588
 in diagenetic sequence, B:39
 in diatom-bearing claystone, A(127):261
 in dolomite, A(128):148
 energy dispersive spectra, B:1378
 framboidal, B:711
 and high organic matter, A(127):283
 and magnetic properties, B:947
 and Mossbauer parameters, B:740
 and oxygenation conditions, B:712
 in sandstone, B:144
 and sediment demagnetization, A(127):203,
 275-276
 sedimentary, B:711-713
 Site 795, A(127):186-187, 189; B:717
 Site 797, A(127):344
 Site 799, A(128):243, 265; B:36, 46, 79, 745-
 746
 Sites 798 and 799, B:670, 771, 775
 and sulfur, A(127):212-213, 705-717;
 B:1249, 1254
 pyrite, framboidal, B:706-709
 pyrite-plus-sulfur (PVS), Site 798, B:771, 775
 pyroclastic component, sediment gravity flow
 processes, A(127):267
 pyroclastic deposits
 sandstone source rocks, B:107
 Site 796, A(127):265
 pyroclastic flows, of blue tuff, B:119
 pyrogenic crystal chemistry, of blue tuff, B:126
 pyrolysates, adsorption, A(127):285
 pyrolysis-gas chromatography, Site 799, B:627-
 628, 632-633
 pyroxene. *See also* clinopyroxene; orthopyroxene;
specific pyroxene minerals
 altered to clays, B:885
 in ash layers, B:1379, 1388
 chemical composition of alteration clay miner-
 als, B:886
 in dolerite, A(128):88
 fractionation models, B:874
 non-quadrilateral end-member molecules,
 B:858
 Site 794, B:851-852
 Site 795, B:852
 Site 797, B:852-853
 pyrrhotite, B:978
 and magnetic properties, B:947
 and magnetization, A(128):169
 Site 797, B:949, 951
 Q (Koenigsberger ratio). *See* magnetic properties
 Q-mode cluster analysis, and planktonic foraminif-
 ers, B:188, 193-200, 208-209, 220, 224
 QFR diagram, sandstone classification, B:106
 quartz
 abundance, A(127):94, 189; B:53, 1235-1244
 and Chinese loess deposits, B:401
 in dark/light cycles, B:584
 glacial/interglacial variation, B:588
 isotopic temperatures, B:55-56
 origin, B:51
 oxygen isotopes, B:51-54
 physical properties interrelations, B:25-27
 polycrystalline, B:107-108
 replaced by carbonate, B:147
 rutile inclusions, B:107
 in sandstone, B:34, 104, 135
 Site 795, A(127):190
 Site 797, B:109, 592
 Site 798, B:401, 411-415, 417
 tourmaline inclusions, B:108
 undulatory extinction, B:107
 vs. excess silica, B:1253
 vs. plagioclase, B:1245, 1252
 vs. terrigenous component abundance, B:403
 quartz, normative component, vs. iron/magnesium
 ratio, B:894
 quartz accumulation rate, Site 798, B:411-420
 quartz/feldspar ratio, Site 798, B:411-415, 417
 quartz sand. *See* sand, quartz
 Quaternary/Pliocene boundary
 Site 795, A(127):192
 Site 796, A(127):271
 radiogenic isotopes. *See* isotopes, radiogenic
 radiolarians. *See also* microfossils, siliceous
 abundance and preservation, A(127):101-102,
 198-199, 274, 356-357, A(128):28,
 161-162, 303-305; B:1220 and cli-
 mate fluctuation, A(127):274, 356
 cycladophorid, B:299
 dilution by detrital components, A(127):198
 dissolution, A(127):198-199, B:353
 dissolution-resistant, B:295-297
 Leg 127, A(127):19-20
 Leg 128, A(128):28
 and opal dissolution transition zone, B:345-
 347
 Pliocene/Pleistocene boundary, A(127):355
 preservation, B:292-296, 1220
 productivity, B:296, 299
 pyritized, A(127):198
 refractive index, B:296
 reworked, A(127):198-199, 274; B:296
 Site 794, A(127):101-102, A(128):99; B:293,
 1221
 Site 795, A(127):198-199; B:294
 Site 796, A(127):274
 Site 797, A(127):356-357; B:295-296
 Site 798, A(128):161-162; B:225, 227, 229-
 230
 Site 799, A(128):303-305; B:225-226, 233-
 234
 species diversity, A(127):101-102,
 A(128):162
 and water temperature, A(128):162
 zonations, A(127):20, 101-102, 198-199,
 274, 356-357, A(128):161-162, 303-
 305; B:1220, 1225-1226
 radiometric age, B:819-836
 age spectra, B:820-821, 828-836
 argon isotopes, B:822-827
 basement, B:1336-1337
 biostratigraphic age compared, B:1337
 of blue tuff, B:123, 126-127
 crystallization age, B:821
 evaluation of ages obtained, B:821
 formation age, B:820-821
 isochron plots, B:828-833
 of Japan Sea rock samples, B:1334
 plateau ages, B:820-821
 radiometric and paleontologic ages compared,
 B:821-822
 samples, B:819-820
 Site 794, B:784-786
 and timing of volcanic activity in Japan Sea,
 B:829-832
 vs. sedimentation rate, B:126
 radiometric dating. *See* radiometric age
 rare earth elements (REE), B:677-695, 719-737.
See also specific rare earth elements
 associated with detrital fraction, B:691
 basalt, B:900
 basement, B:780, 912-916
 and biogenic siliceous input, B:692
 complexation with organic and inorganic li-
 gands, B:730
 correlation coefficients, B:691
 and heavy minerals, B:691
 LREE enrichment relative to HREE, B:725
 mixing model of pelagic sediment into de-
 pleted mantle, B:817
 mobility during alteration, B:911-916
 and parent magma, B:927-928
 processes controlling distribution, B:692
 and silica content, B:682
 Site 794, B:682-684, 688, 912-913, 916-919,
 922, 1339
 Site 795, B:683-685, 688, 913-914, 918, 924,
 1338-1339
 Site 796, B:685
 Site 797, B:685, 688-691, 914-916, 920-921,
 925, 1339-1340

- Site 799, B:734
sources in sediment column, B:729
spidergram, B:783
terminology, B:681-682
and terrigenous sedimentation, B:691, 732
in volcanic glass, B:1387
vs. loss on ignition, B:919, 921
reconstructions. *See* plate reconstructions
recrystallization
Site 794, A(128):99
Site 799, A(128):303, (128):307
Red River Fault, offset along, B:1325
redox conditions. *See* oxygenation conditions
REE. *See* rare earth elements
reflection coefficient, Site 799, B:1138-1141
refractory elements. *See also* names of specific refractory elements
Site 798, B:706
remagnetization, and pyrite, A(127):203
resistivity, logging
core depths correlated, B:1040
core/fog correlation, B:398-400
electrical resistivity experiment compared, B:1354
of opal-A to opal-CT transition, B:22
opal-CT to quartz transition and, B:1040
Site 794, A(127):139-142, 144-146, 156-159, A(128):104, 107, 119-120; B:1069
Site 795, A(127):232, 237
Site 796, A(127):302-305, 321-322
Site 797, A(127):393-395, 400-402, 416-418; B:7
Site 798, A(128):185-187, 212, 225-228; B:399, 401-402
Site 799, A(128):332-338, 362-364, 380-382, 393-397
Sites 794 and 797 correlated, B:1039, 1041
resolution, in logs, B:396-397, 404-405
reverse faults
Site 799, A(128):263, 271-272; B:1183, 1189
and stress field, B:1182-1183
Yamato Basin N, A(128):75-76
rhodochrosite
and cerium concentration, B:90-91
chemical composition, B:81-82, 84-85
formation, B:85, 90-91
and iron distribution, B:742
isotopic composition, B:83
and Mossbauer parameters, B:741
occurrence, B:80
paleoenvironmental implication, B:94
petrography and textural relations, B:81
SEM photographs, B:97
Site 799, B:75-98, 744-745
X-ray diffraction analyses, B:79-80
rhyodacite, volcanic glass, B:1378
rhyolite
ash layers, B:793
tuff, B:731
volcanic glass, B:1378
rhythmic sedimentation. *See* lithofacies, cyclic
rift propagation, in Japan Sea E, B: 1346
rifting. *See also* back-arc rifting
ripup clasts, in sandstone, B:104
rivers, pollen and, B:333
Rock-Eval analyses. *See also* hydrogen index (HI); oxygen index (OI); Rock-Eval pyrolysis; T_{max}
carbon, organic, A(127):367
carbon, total organic (TOC), A(127):115, 213, 285
methods, A(127):114
 S_1/S_2 vs. depth, A(127):121
Site 794, A(127):114-120
Site 795, A(127):218-219
Site 796, A(127):285, 288
Site 797, A(127):367, 376-378
Site 798, A(128):177, 195; B:425
Site 799, B:429
source character, A(127):115-118, 205, 285, 367
thermal maturity, A(127):118, 285, 367
 T_{max} vs. depth, A(127):121
Rock-Eval pyrolysis. *See also* hydrocarbon; hydrogen index (HI); organic matter; oxygen index (OI); Rock-Eval analyses; T_{max} ; total pyrolytic yields; transformation ratio (TR); van Krevelen plot
Site 798, B:669
Site 799, B:627, 669
rotalids
Site 794, A(127):101
rotation event, Japan Sea, B:179, 528
rubidium
and ash alteration, B:1266
and basement, B:807, 1266
in interstitial water, B:1271
Site 794, A(127):109, B:1363
Site 795, A(127):205, B:1365
Site 796, A(127):280-281
Site 797, A(127):364, 371; B:1366
Site 798, B:1370-1371
vs. hydrogen isotopes, B:1273
vs. oxygen isotopes, B:1273
rutile, as quartz inclusions, B:107
 S_1 . *See* hydrocarbons, free
 $S_1 + S_2$. *See* hydrocarbons, total pyrolytic yield
 $S_1/(S_1 + S_2)$. *See* transformation ratio
 S_2 . *See* hydrocarbons, generatable
Sado Island
and Japan Sea stress field, B:1185
uplift, A(128):158
Sado Ridge, A(128):70, 76
basement, A(128):73
Sakhalin Island, A(128):15; B:1206-1207, 1210
salinity
and clathrate formation, A(127):281
glacial/interglacial cycles, B:591, 608
and organic matter decomposition, A(128):318
and oxygen isotopes, B:442-443, 447
Site 794, A(127):107
Site 795, A(127):205
Site 796, A(127):278-279
Site 797, A(127):364, 374
Site 798, A(128):172, 179
Site 799, A(128):317, 328; B:610
and sulfate removal, A(127):205
samarium
in basement rocks, B:807
Site 798, B:1370-1371
sand. *See also* sandstone; terrigenous component
and clathrate formation, A(127):281
composition, A(127):268, B:216
differential compaction, B:39
and fluid migration, A(128):321
and foraminifer abundance, B:193
Kita-Yamato Trough, A(128):354
in logs, A(127):251
microturbidite, B:44
provenance, A(127):268
sedimentary structures, A(127):268, A(128):243, 265
seismic expression, A(127):313
shallowest occurrence, A(127):272
Site 796, A(127):250, 261-264
Site 799, A(128):258, 265
temperature measurements in, A(127):299
Toyama Fan, A(128):14
sand, bioclastic, Site 797, A(127):343
sand, glauconitic
origin, A(128):157
Site 798, A(128):124, 138, 141
Site 799, A(128):241
sand, quartz
origin, A(128):157
Site 798, A(128):124, 138
Site 799, A(128):264
sandstone. *See also* sand
alteration, A(127):345
carbonate cementation, B:39, 141
cementation, B:132, 134-135
classification, B:106
compaction, B:132
composition, A(127):345
deposition, A(127):19
diagenesis, B:131-151
dolomite-cemented, A(128):276
feldspar albitization, B:131-151
feldspar composition, B:105-106, 136, 139
framework composition, B:104-105, 113
grain size, B:100, 104
interbedded with basalt, A(127):345-346
laminated, A(127):3347-349
lithic, B:104
mineral replacement in, B:131-148
normally graded, A(127):264-265, 346, 348
oldest deposited, A(127):268-269
provenance, B:99-111, 148
and revised lithostratigraphy, B:1233
sedimentary structures, B:100, 104, 1233
Site 796, B:99-113
Site 797, A(127):344-346; B:99-113
Site 799, A(128):265, 292; B:33-34, 42-43, 99-113
source area location, B:110-111
source of carbonate, B:141-143
source-rock lithology, B:106-109
tectonic provenance, B:109-110
thickness, B:100, 104
zeolitization, B:131-139
sandstone, bioclastic, Site 797, A(127):346
sandstone, feldspathic, Site 799, B:131
sandstone, tuffaceous, Site 796, A(127):265
sandstone, volcanoclastic
Site 796, B:102, 131
Site 797, B:103, 131
Santa Barbara Basin, and opal flux, B:439
saponite, B:885-888
saturation vapor pressure, and index property data errors, B:986, 989
scandium
Site 795, B:707
Site 798, B:1370-1371
scanning electron microscope (SEM), and visual microanalysis, B:57-58
sea ice
in Japan Sea, A(127):20-22; B:167
Leg 127, A(127):20
and productivity, B:311
sea level
Japan Sea, B:180
Site 798, A(128):166
Sea of Okhotsk. *See* Okhotsk Sea
sea-surface temperature. *See* temperature, surface-water
seafloor spreading, in Japan Basin, B:1345-1346
Second Spring Layer, benthic foraminifers and, B:365

- sediment accumulation rate. *See also* feldspar accumulation rate; mass accumulation rate; quartz accumulation rate; sedimentation rate
- and barium distribution, B:660-661
- of biogenic silica, A(128):122, 158
- Brunhes Chron, B:468
- of claystone, A(127):191
- Japan Sea, A(127):22, 24, A(128):31
- Oga Peninsula, B:1204
- of siliceous claystone, A(127):191
- Site 794, A(127):72, 105-107
- Site 795, A(127):174, 203-204
- Site 796, A(127):250-251, 277-278, 281-282
- Site 797, A(127):324, 359-362
- Site 798, A(128):31, 124-125, 171ff, 176-177; B:409-420, 446, 563-564
- Site 799, A(128):31, 244, 259-260, 315-316, 324
- and sulfate content, B:637, 646
- sediment composition
- carbonate components, B:1241
- of dark/light cycles, B:579, 582-584
- detrital components, B:1237-1239
- of fine-grained rocks, B:1367
- glacial/interglacial variations, B:588-590
- Japan Sea, B:1233-1249
- logging estimates, B:402-403
- major elements, B:1235-1237, 1246-1248
- of sand, B:216
- of sandstone, B:104-105, 113
- siliceous components, B:1239-1241
- Site 794, A(127):92; B:1235-1237, 1242
- Site 795, A(127):188; B:8-10, 1237-1240, 1243
- Site 796, A(127):264
- Site 797, A(127):345; B:11-14, 1240-1241, 1244
- Site 798, A(128):138; B:398-400, 409-422
- Site 799, A(128):258
- sulfide components, B:1241-1249
- variations seen in logs, B:398-400
- sediment geochemistry. *See* geochemistry, sediment
- sediment grain size. *See* grain size
- sediment gravity flow deposits
- and formation of color-banded bedding, A(127):349
- Site 796, A(127):247, 268, 315; B:100, 110
- Site 797, A(127):347; B:100
- sources, A(127):267
- sediment loading, sediment consolidation and, B:1130
- sediment texture, Site 799, B:33-48
- sediment thanatocoenosis, and diatom dissolution, B:309-316
- sedimentary structures. *See also specific sedimentary structures*
- in blue tuff, B:117
- brittle structures, A(128):269-272; B:1181-1183
- caused by compaction, B:38
- color-banded bedding, A(127):186
- dewatering structures, B:1176-1178
- faults, A(128):143ff
- gravitational compaction structures, A(128):146ff, 272
- gravity-induced structures, B:1175-1176
- Japan Sea, A(128):13-15; B:1183-1187
- laminations, A(127):95-96, 347-349
- and lithostratigraphy, B:1177
- in sandstone, B:100, 104, 1233
- in sandstone and siltstone, A(127):324, 345-346
- in sediment gravity flow deposits, A(127):268
- in siltstone, B:1233
- Site 798, A(128):141ff
- Site 799, A(128):265-272; B:48
- soft-sediment deformation, A(127):261-264, A(128):141ff, 241, 266-269
- thin bedding, A(127):96
- in tuff, A(128):81
- veins, A(128):143ff; B:1176-1181
- sedimentation, rhythmic. *See* dark/light cycles; lithofacies, cyclic
- sedimentation rate. *See also* sediment accumulation rate
- age control points, B:351
- in Brunhes Chron, B:961
- at Brunhes/Matuyama boundary, B:961
- and carbonate compensation depth, B:1228
- and consolidation, B:1277
- datum levels used, A(127):111, 209, 281, 365
- and dolomite, A(128):148
- error sources, A(128):171
- and heat flow, B:1302
- heat flow estimates from opal-A/opal-CT BSR and, B:1147-1148
- Japan Sea, A(127):22, 24, A(128):31
- and Jaramillo Subchron, B:959-960
- and logging resolution, B:396
- and magnetic intensity, B:966
- and magnetite degradation, B:976
- during Olduvai Subchron, B:975
- and opal-A to opal-CT transition, B:1306
- and orbital periodicity, B:397-398
- and organic carbon, B:628
- and organic matter preservation, B:1265
- and phosphate formation, B:67
- Pliocene/Quaternary boundary, A(127):251
- and productivity, B:1228
- related to upwelling, A(127):204
- Site 794, A(127):96, 105-107, 112, A(128):31, 69; B:287, 351, 653, 975, 982, 1043-1044, 1224, 1227
- Site 795, A(127):174, 203-204, 210; B:288, 351, 653, 975, 982, 1226-1227
- Site 796, A(127):250-251, 277-278, 281-282; B:288, 653, 1226-1227
- Site 797, A(127):324, 351, 357, 359-362, 366; B:287, 297, 351, 594, 975, 982, 1043-1044, 1226-1227
- Site 798, A(128):31, 124-125, 171ff, 176-177; B:407, 443, 446, 548, 563-564, 653, 959, 961, 974-975, 982, 1226-1227
- Site 799, A(128):31, 244, 259-260, 315-316, 324-325; B:653, 959, 961, 974-975, 982, 1227
- Sites 794 and 795 compared, A(127):204
- Sites 798 and 799 correlated, B:961, 965
- and sulfate, A(127):362; B:1262
- and turbidites, B:959
- vs. barium, B:661
- vs. depth of phosphate maximum, B:64-66
- vs. sulfate, B:661
- Yamato Basin S, A(127):356
- sediments, biosiliceous, Site 798, B:439-455
- sediments, bioturbated, in cyclic lithofacies, A(128):124
- sediments, laminated. *See also* laminations
- BSEM images, B:553-557
- in cyclic lithofacies, A(128):124
- lamination thickness, B:548, 550
- and low oxygen environment, B:189
- paleoceanographic significance, B:547-557
- Site 794, A(127):147
- Site 799, B:44
- visual microanalysis methods, B:57-62
- sediments, organic-rich, formation, A(127):212, 365
- sediments, subducted, and contamination of mantle source, B:805-817
- seismic experiment. *See* downhole seismic experiment
- seismic expression
- of biogenic opal, B:1140-1141, 1143
- of debris flows, B:1140-1141, 1143
- of dolomite, B:1141, 1143
- of opal-A to opal-CT transition, B:1141, 1143
- seismic reflection profile
- data collection and processing, A(127):61-63
- Japan Sea, A(127):14, A(128):13
- Kita-Yamato Trough, B:1137-1139
- reflector origin, B:1140-1142
- Site 794, A(127):68, 84, 86-87, 153, A(128):83; B:111, 1079
- Site 795, A(127):181, 183-184, 241-243
- Site 796, A(127):258, 260-261, 311-312
- Site 797, A(127):337-338, 340-341, 409
- Site 798, A(128):135, 218-223; B:395, 410
- Site 799, A(128):23, 254-255, 260, 375-379; B:1137-1139
- synthetic seismogram compared, B:1142
- Yamato Basin, B:1149
- seismic reflection survey, pre-cruise surveys, A(127):142, 233, 307, 399, A(128):192, 343-344
- seismic refraction profiles, Yamato Basin N, A(128):112
- seismic stratigraphy
- ash layers correlated, A(128):122, 153-155, 344-351, 379
- of basement, B:779
- of interbedded sediments/basalts, A(127):408
- Japan Sea, A(127):28-29, A(128):34
- Kita-Yamato Trough, A(128):245; B:1135-1143
- lithostratigraphy correlated, A(127):143, 155, 174, 234-238, 243, 251, 312-313, 325, 403-408, 410, A(128):125, 193-194, 224, 351-353, 379; B:1140-1143
- logs correlated, A(128):224, 351-353, 379
- Okushiri Ridge, A(127):256-258, 307
- physical properties correlated, A(128):224, 379; B:1143
- seismic intervals, Site 799, B:1136
- Site 794, A(127):73, 142-143
- Site 795, A(127):174, 233-238
- Site 796, A(127):251, 307-314
- Site 797, A(127):325, 399-410
- Site 798, A(128):34, 125, 192-194
- Site 799, A(128):34, 343-353; B:1135-1143
- Yamato Basin, A(127):83-84, 143
- seismology. *See* downhole seismic experiment
- seismometer. *See* downhole seismic experiment
- SFL. *See* resistivity, logging
- shards, glass
- altered, A(127):95
- in ash layers, B:791
- in blue tuff, B:117, 119, 130
- replaced by zeolites, B:139
- Site 795, A(127):190
- shards, vitric, in ash-flow tuff, B:793
- shear joints, Site 794, A(128):92
- shear strength
- and consolidation characteristics, B:1124-1125

- overburden pressure ratio, B:1125–1126
 Site 798, A(128):182, 207, 211
 Site 799, A(128):327, 353, 360
- shear zone, Site 799, A(128):270
- sheet flows, of doleritic basalt, A(128):86
- Shibikawa Formation, and planktonic foraminifers zonation, B:459
- Shikoku Basin, rare earth elements, B:719
- Shikoku–Parece Vela Basin, formation, A(127):10
- Shiribeshi Basin, A(127):268
- Shiribeshi Trough
 bathymetry, A(127):251
 free-air gravity anomalies, A(127):253
 sedimentation, A(127):256–257
 seismic stratigraphy, A(127):307
- siderite
 and Mossbauer parameters, B:741
 shallowest occurrence, A(128):273
 Site 799, A(128):260, 272, 275–276, 289;
 B:611–612
- sigmas
 Site 795, B:543
- Sikhote-Alin, basement, A(127):177–178
- silica
 in ash layers, B:1380–1385
 basement, A(128):98
 and biogenic silica diagenesis, B:1263
 in dark/light cycles, B:569
 diagenesis, A(127):190
 and diatom dissolution, A(127):362–363
 dissolution, B:350
 formation, B:52–54
 in interstitial water, B:1266
 Japan Sea, B:1235–1237
 precipitation, A(127):174
 and rare earth elements, B:682
 removal rates in Japan Sea, B:447
 and silicification front, A(127):205
 Site 794, A(127):108–109; B:682
 Site 795, A(127):205; B:683
 Site 796, A(127):280
 Site 797, A(127):362–363, 369; B:688
 Site 798, A(128):174–175, 184; B:722
 Site 799, A(128):318, 320–321, 332; B:611,
 722
 sources, B:148
 transitions, B:53
 vs. lithium, A(128):185
 XRD analyses, A(128):292
 XRF analyses, A(128):281; B:1025–1029,
 1031–1035
- silica, amorphous. *See* opal-A
- silica, biogenic. *See also* opal, biogenic; opal-A;
 opal flux
 abundance, A(128):150, 259
 accumulation rate, A(128):122, 158
 cycles, B:439
 in dark/light cycles, B:584
 diagenesis, B:1263, 1267
 dissolution, A(128):320–321
 fragmentation degree, B:549
 glacial/interglacial variation, B:591
 Japan Sea, B:1239–1241
 and productivity, B:694
 and rare earth elements, B:692
 XRF analyses, A(128):151
 XRF vs. smear slide analyses, A(128):281
- silica, diagenetic
 isotopic studies, B:49–56
 Japan Sea, B:1239–1241
 Site 799, B:36, 46
 XRF vs. smear slide analyses, A(128):281
- silica, dissolved, in Japan Sea, B:350–351
- silica, excess, B:1240–1244, 1253
- silica, logging
 and diatom vs. clay content, B:1417
 Site 794, B:1416, 1420–1421
 Site 796, B:1416, 1423
 Site 797, B:1417, 1426–1427
 Site 798, B:1401–1402
 Site 799, A(128):364; B:1404, 1408–1409
- silica/aluminum ratio (Si/Al)
 Site 795, B:683
 Site 797, B:688
- silica diagenesis. *See also* opal-A to opal-CT tran-
 sition; opal-CT to quartz transition
 and Ce anomaly, B:682
 effect of temperature, B:3
 effects on interstitial water chemistry, B:646
 and Eu anomaly, B:682
 and inorganic geochemical data, A(127):23
 and interstitial water oxygen isotopes, B:85,
 701
 Japan Sea, B:1263, 1267
 material balance calculation for, B:701
 and physical properties, A(128):280; B:3–31,
 1281–1282
 and silica content, B:682, 688
 Site 795, A(127):190
 Site 797, A(127):347
 Site 798, A(128):149–150
 Site 799, A(128):278–281
 and thermal history, Japan Sea, B:1305–1307
- silica phase change. *See* opal-A to opal-CT transi-
 tion; opal-CT to quartz transition
- silicate, alteration reactions, A(127):110–111
- siliceous claystone. *See* claystone, siliceous
- siliceous veins. *See* veins, siliceous
- silicification front, and silica content, A(127):205
- silicoflagellate, B:1220
 Site 798, A(128):162; B:241–244
 Site 799, A(128):305; B:241, 246–247
 zonal data, B:1225–1226
 zonation, B:237, 239–241
- silicon, at opal-CT to quartz transition A(127):307
- silicon oxide. *See* silica
- sill, Japan Sea, B:299–300, 365
- sills
 basaltic, B:819
 bathymetric, A(128):20, 35
 and crustal heterogeneity, B:1113–1114
 crystallization history, B:864–865
 dolerite, A(128):86
 emplacement, B:894, 927, 938
 intrusion, A(128):95; B:829–830
 demagnetization, B:938
 Site 794, B:837–838, 849, 906, 927, 1339
 Site 797, B:837–838, 850, 907–908
 suites, B:853–855
- silt
 Site 796, A(127):261–264
 Site 799, A(128):256, 264; B:43
- siltstone
 and revised lithostratigraphy, B:1233
 sedimentary structures, B:1233
- siltstone, carbonaceous, Site 797, A(127):345–346
- siltstone, diatom clayey, Site 795, A(127):187
- siltstone, siliceous, Site 799, B:39–40
- siltstone, tuffaceous clayey, A(128):68, 89–90
- silty clay. *See* clay, silty
- silver
 Site 798, B:1368–1369
 in sulfide deposits, A(128):21
 Site 79, physical properties, B:1010–1015
 Site 116, interstitial water boron content, B:638
 Site 125, trace metals, B:1361
- Site 126, trace metals, B:1361
- Site 127, trace metals, B:1361
- Site 128, trace metals, B:1361
- Site 130, trace metals, B:1361
- Site 141, paleoenvironment, B:338
- Site 148, siderite occurrences, B:612
- Site 173
 foraminifers, A(128):165
 paleoclimate, B:459
 silicoflagellate, B:237
- Site 290, basement isotope geochemistry, B:805
- Site 294, rare earth elements, B:719
- Site 296, rare earth elements, B:719
- Site 299, A(127):337
 bathymetry, A(127):6, 78, A(128):72
 consolidation tests, B:1123
 oceanic crust, A(127):73
 planktonic foraminifers, B:459
 radiolarians, A(128):162, B:225
 sediment thickness, A(127):84
 seismic stratigraphy, A(127):179
 Site 797 sandstone compared, B:111
 turbidite sands, B:99
- Site 300
 bathymetry, A(127):6, A(128):72
 nanofossils, B:168
 sand recovered, A(128):14
- Site 301
 bathymetry, A(127):6, A(128):72
 drilling history, A(127):180
 lithostratigraphy, A(128):13, 249; B:1141
 nanofossils, B:168, 174
 sand recovered, A(128):14
 sediment thickness, A(127):84
 seismic stratigraphy, A(127):179
 Site 799 sandstone compared, B:111
 turbidite sands, B:99
- Site 302
 bathymetry, A(127):6, 78, A(128):72
 consolidation tests, B:1123
 diatom clay, A(127):84
 faulting age, A(127):334
 nanofossils, B:168, 174
 normal faulting, A(127):83
 radiolarians, A(128):303, B:226
 sediment thickness, A(127):84
 tectonic history, A(128):130
- Site 303, mantle source contamination, B:815
- Site 304, mantle source contamination, B:815
- Site 366, paleoenvironment, B:338
- Site 395, temperature change during downhole
 seismic experiment, B:1160
- Site 397, paleoenvironment, B:338
- Site 417, alteration and rare earth elements corre-
 lated, B:911
- Site 433, diatom abundance, B:316
- Site 436
 ash layers, B:1319–1320
 basement age, B:749
 helium isotope ratios, B:749
 rare earth elements, B:719
 tectonic setting, B:1312
- Site 438
 ash layers, B:1319–1320
 basement age, B:749
 diatom abundance, B:311
 forearc subsidence, B:1213
 helium isotope ratios, B:749
 paleobathymetric history, B:1214
 silicoflagellates, B:237
 subsidence of Ishikari-Hidaka Basin, B:1199
 tectonic setting, B:1312
- Site 439

- ash layers, B:1319-1320
forearc subsidence, B:1213
paleobathymetric history, B:1214
subsidence of Ishikari-Hidaka Basin, B:1199
tectonic setting, B:1312
and volcanic front position, B:1324
- Site 440
ash layers, B:1319-1320
tectonic setting, B:1312
- Site 504
formation factor and cation exchange capacity correlated, B:1282
isotopic analyses, B:888
oxygen isotopes, B:699
stress field, B:1047
- Site 533, nannofossils, B:173
- Site 544, paleoenvironment, B:338
- Site 578, diatoms, B:350
- Site 579
biosiliceous productivity, B:311-313
diatoms, B:350
- Site 580
biosiliceous productivity, B:311-313
diatoms, B:350
radiolarian faunal change, B:447
- Site 581
diatoms, B:311, 316
temperature change during downhole seismic experiment, B:1160
- Site 582
interstitial water oxygen isotopes, B:698
veins observed, B:1180
- Site 583
basement age, B:749
helium isotope ratios, B:748-749
interstitial water oxygen isotopes, B:698
- Site 584
ash layers, B:1319-1320
basement age, B:749
helium isotope ratios, B:749
interstitial water oxygen isotopes, B:698
tectonic setting, B:1312
- Site 588, paleoenvironment, B:338
- Site 590, paleoenvironment, B:338
- Site 591, paleoenvironment, B:338
- Site 607, oxygen isotopes, B:403
- Site 672, veins, B:1180
- Site 684, helium isotope ratios, B:748
- Site 747, ash layer logging, A(128):339
- Site 752, ash layer logging, A(128):339
- Site 792, Brunhes/Matuyama boundary, A(127):199
- Site 794, A(127):71-167, A(128):67-120
aluminum, B:654, 656
ash layer chemistry, B:1373-1393
augite-liquid relations, B:870-872
barium, B:653-657
barium as productivity indicator, B:653-656
basalt alteration, B:883-889
basalt melting experiment, B:891-898
basement, B:849-850, 885, 906, 1339
basement chemistry, B:779-789, 839, 892-894, 908-911, 916-918
basement isotopes, B:786-787, 807-814, 888, 901-902
basement magnetic properties, B:934
basement mineralogy, B:850-855
basement structures, B:1181
biostratigraphy, B:1221-1223
blue tuff, B:115-130
boron, B:638-641
crustal anisotropy, B:1107-1121
crustal structure, B:1075-1081
- diatoms, B:344
downhole seismic experiment, B:1061-1073, 1075-1081, 1107-1121, 1157-1171
electrical resistivity experiment, B:1351-1359
FMS logs, B:1039-1046
foraminifers, B:190-193, 210-213
geochemical logs, B:1420-1421
geothermal analyses, B:1301-1307
gray value, B:585
hydrogen isotopes, B:641-642
inorganic geochemistry, B:637-646, 1261-1274
liquid lines of descent, B:870
lithostratigraphy, revised, B:1231
logs, B:1416, 1418-1419
magma suites, B:869
magmatic characterization, B:780-784
magnetostratigraphy, B:969-970
mantle source, B:807-815
Olduvai Subchron, B:218
operations, A(127):85-90, A(128):83-85
organic carbon, B:656
oxygen isotopes, B:641-643
parent magma, B:855-857, 894
physical properties, B:990-993, 1275-1296
radiometric age, B:780-784, 820, 824-826
sediment composition, B:1236-1237, 1242
sediment geochemistry, B:680, 682-684, 691-695, 1246-1249
sedimentation rate, B:1224-1225
spinel chemistry, B:840-843
stress measurements, B:1050-1059
strontium isotopes, B:643-646
subsidence history, B:1207-1211
sulfate, B:637-638, 656
sulfur isotopes, B:637-638
trace metals, B:1362-1363
Yamato Basin tectonics, B:787-788
- Site 795, A(127):169-245
ash layer chemistry, B:1373-1393
augite-liquid relations, B:870-872
basement, B:906, 1338-1339
basement chemistry, B:913-914, 918-919
basement isotopes, B:807-814
basement magnetic properties, B:934, 937
basement mineralogy, B:850-855
basement petrography, B:849-850
biostratigraphy, B:1223
boron, B:638-641
diatoms, B:344-345
foraminifers, B:194-197
geothermal analyses, B:1301-1307
gray value, B:585
hydrogen isotopes, B:641-642
inorganic geochemistry, B:637-646, 1261-1274
lithostratigraphy, B:496-497, 1231
magnetostratigraphy, B:970-972
mantle source, B:807-815
operations, A(127):182-186
oxygen isotopes, B:641-643
oxygenation conditions, B:711-713
parent magma, B:855-857
physical properties, B:993-997, 1275-1296
physical properties and silica diagenesis, B:3-31
pyrite, B:711-713
radiometric age, B:820-821, 826-827
reworking, A(127):198
sediment composition, B:8, 1237-1240, 1243
sediment geochemistry, B:680-681, 683-685, 691-695, 706-717
sedimentation rate, B:1225
- strontium, B:643-646
subsidence history, B:1207-1211
sulfate, B:637-638
sulfur, B:637-638
trace metals, B:1365
- Site 796, A(127):247-322
aluminum, B:654, 656
ash layer chemistry, B:1373-1393
barium, B:654, 654-657
biostratigraphy, B:1223
blue tuff, B:115-130
boron, B:638-641
clay mineral authigenesis, B:139
feldspar albitization, B:136
foraminifers, B:198-200
geochemical logs, B:1423
geothermal analyses, B:1301-1307
hydrogen, B:641-642
inorganic geochemistry, B:637-646, 1261-1274
lithostratigraphy, B:1231
logs, B:1416, 1422
operations, A(127):259-261
organic carbon, B:656
oxygen isotopes, B:641-643
physical properties, B:997-999, 1275-1288
reworking, A(127):273
sandstone provenance, B:99-113
sediment geochemistry, B:681, 685, 691-695
sedimentation rate, B:1225
strontium, B:643-646
subsidence history, B:1207-1211
sulfate, B:637-638, 656
sulfur, B:637-638
zeolitization, B:138-139
- Site 797, A(127):323-421
ash layer chemistry, B:1373-1393
augite-liquid relations, B:870-872
basalt melting experiment, B:861-868, 891-898
basement, B:906-907, 1339-1340
basement chemistry, B:839, 892-894, 911, 914-916, 919-921
basement isotopes, B:807-814, 901-902
basement magnetic properties, B:934, 937-939
basement mineralogy, B:850-855
basement petrography, B:850, 891-892
biostratigraphy, B:1223
boron, B:638-641
boron isotopes, B:638-641
clay mineral authigenesis, B:139
dark/light cycles, B:574, 577-601
diatoms, B:345, 348-349
feldspar albitization, B:136-137
FMS logs, B:1039-1046
foraminifers, B:202-207, 213-215
fractionation models, B:872-874
geochemical logs, B:1426-1427
geothermal analyses, B:1301-1307
gray value, B:585, 592
hydrogen isotopes, B:641-642
inorganic geochemistry, B:637-646, 1261-1274
liquid lines of descent, B:870
lithostratigraphy, B:4, 1231
logs, B:7, 1417, 1424-1425
magma suites, B:869
magnetic minerals, B:949-951
magnetostratigraphy, B:973
mantle source, B:807-815
opal-A to opal-CT transition, B:19
operations, A(127):337-340
organic carbon, B:586-587, 592

- oxygen isotopes, B:641-643
parent magma, B:855-857, 894
physical properties, B:3-31, 999-1104, 1275-1296
radiometric age, B:820, 822-824
reworking, A(127):357
sandstone provenance, B:99-113
sediment composition, B:8-10, 582-584, 586-587, 592, 1240-1241, 1244
sediment geochemistry, B:681, 685, 688-695
sedimentation rate, B:1225
spinel chemistry, B:840-843
strontium isotopes, B:643-646
subsidence history, B:1207-1211
sulfate, B:637-638
sulfur, B:586-587, 592
sulfur isotopes, B:637-638
trace metals, B:1366
zeolitization, B:138-139
- Site 798, A(128):121-236
accumulation rates, B:411-419
aluminum, B:654, 656
ash layers, B:699-701, 791-793, 1373-1393
bacteria, B:755-776, 755-769
barium, B:654, 656-659
basalt alteration, B:699-701
biostratigraphy, B:1223-1224
C/N ratio, B:435-437
carbonate content, B:435-437
carbonate diagenesis, B:701-702
chronostratigraphy, B:317-318
clay mineral composition, B:416
consolidation, B:1124-1127
dark/light cycles, B:430-432, 446-447, 559-576
feldspar, B:411-419
foraminifers, B:460-463
geochemical logs, B:1021-1035, 1398, 1400-1402
geologic setting, B:425, 440, 560
geothermal analyses, B:1301-1307
glauconite, B:63-74
inorganic geochemistry, B:607-608, 697-703, 1261-1274
interstitial gas helium isotopes, B:748-749
laminated biosiliceous sediments, B:547-557
lithostratigraphy, B:697
logs, B:398-400
magnetic properties, B:959-967, 975-978
magnetostratigraphy, B:973-974
methane origin, B:749-750
microbiology, B:761-776
Mossbauer characterization of sediments, B:739-746
opal content, B:443-446, 449-455
operations, A(128):133-136
organic carbon, B:425, 435-437, 656
organic geochemistry, B:667-675
organic matter, B:425, 667-675
oxygen isotopes, B:442-443, 698-699
permeability, B:1127-1129, 1131-1132
phosphate, B:63-74
physical properties, B:1006-1010, 1275-1296
quartz, B:411-419
sediment composition, B:398-400, 411
sediment geochemistry, B:610-612, 615-617, 720-737, 1022-1027, 1402
sedimentary structures, B:1175-1183
sedimentation, B:419-421, 1225
seismic stratigraphy, B:1135-1143
silica diagenesis, B:701
strip sample trace elements, B:1367-1372
subsidence history, B:1207-1211
- sulfate, B:656
surface-water temperature fluctuations, B:463-465
synthetic seismogram, B:1142
tephrochronology, B:793-797, 800
- Site 799, A(128):237-402
aluminum, B:654, 656
ash-flow tuff, B:793
ash layers, B:699-701, 791-793, 1373-1393
barium, B:654, 654-657
basalt alteration, B:699-701
biostratigraphy, B:34-35, 1224
carbonate diagenesis, B:701-702
clay mineral authigenesis, B:139-141
consolidation, B:1124-1127
dark/light cycles, B:571-574
diagenetic carbonates, B:75-98
diagenetic mineralogy, B:35-38
feldspar albitization, B:137
geochemical logs, B:1399, 1403-1408
geologic setting, B:425, 560
geothermal analyses, B:1301-1307
glauconite, B:63-74
inorganic geochemistry, B:608-610, 697-703, 1261-1274
interstitial gas helium isotopes, B:748-749
lithostratigraphy, B:33-34, 76-77, 697
magnetic intensity variations, B:959-967
magnetic properties, B:975-981
magnetostratigraphy, B:974-975
methane, B:749-750
Mossbauer characterization of sediments, B:739-746
operations, A(128):253-255
organic carbon, B:425, 427, 623, 626, 656
organic geochemistry, B:623-633, 667-675
organic matter, B:425, 667-675
oxygen isotopes, B:698-699
permeability, B:1127-1129, 1131-1132
phosphate, B:63-74
physical properties, B:1010-1015, 1275-1296
pyrolysis-gas chromatography, B:627-628, 632-633
Rock-Eval pyrolysis, B:624-627
sandstone provenance, B:99-113
sediment geochemistry, B:610-612, 618-620, 720-737, 1402
sediment structures, B:38-39, 1175-1183
sedimentation rate, B:1225-1227
silica diagenesis, B:701
subsidence history, B:1207-1211
sulfate, B:656
sulfur, B:623-624
tephrochronology, B:793-799
thermal conductivity, B:1017-1019
slickensides, in tuffaceous clayey siltstone, A(128):90
- slump folds
Site 798, A(128):142
Site 799, A(128):261-263, 266-267
- slumps
of ash layers, A(128):153-154, 285
and compositional banding, B:1176
and consolidation, B:1125
extent and magnitude, A(127):268
and faults, B:1176
Japan Sea, A(128):14
on Okushiri Ridge, A(127):268
and opal-A to opal-CT transition, A(127):268
Site 798, B:1175
Site 799, A(128):259, 355; B:1175-1176, 1189
smear slide analyses. *See* sediment composition
smectite
in blue tuff, B:119
in dark/light cycles, B:584-588
diagenetic origin, B:1238
glacial/interglacial variation, B:588
from Japanese Islands, B:595
in Japan Sea sediments, B:1235-1244
as sandstone cement, B:135
Site 797, B:592
Site 798, B:411, 416, 418
and volcanic glass alteration, B:1375
vs. illite, B:1245
vs. magnesium, B:1250
smectite/illite ratio, Site 798, B:411, 416
sodic feldspar. *See* feldspar, sodic sodium
in ash layers, B:1384
in augite, B:851-853, 858
and basement alteration, B:1266
and feldspar albitization, B:137
and geochemical logging, B:1395, 1415
Japan Sea, B:1237
and parent magma, B:856
Site 794, A(127):107
Site 795, A(127):205
Site 796, A(127):278-279
Site 797, A(127):364, 374
Site 798, B:1370-1371
Site 799, B:723
sources, B:137
vs. potassium, B:732, 858
sodium/aluminum ratio
Site 794, B:1249
Site 795, B:683
Site 798, B:682
sodium/chlorine ratio, in interstitial water, B:1274
sodium oxide. *See* sodium
sodium/potassium ratio, Site 799, B:723
soft-sediment deformation. *See* deformation, soft-sediment
soil horizons, formed during interglacials, B:403
sonic core monitor (SCM), Site 799, A(128):253, 255
sonic velocity. *See* velocity, logging
Soya Strait, B:423
depth, B:342, 440
Japan Sea and, B:70, 362-363, 365
sill depth, B:457
sphenoliths
Site 799, A(128):307, 309
spherasters
Site 795, B:543
spherically focused log (SFL). *See* resistivity, logging
spinel. *See also* spinel, chromium
chemistry, B:840-842
crystal chemical model, B:840-841
magmatic magnesium number inferred from composition, B:840-842
magmatic oxygen fugacity from spinel analyses, B:842-844
and mineral-liquid disequilibrium, B:844
photomicrographs, B:847
relict magmatic phase, B:838
spinel in other igneous rocks compared, B:840
zoned, B:840, 844
spinel, chromium, B:853
magnesium number vs. Cr (Cr + Al), B:843
magnesium number vs. Fe³⁺/(Cr + Al + Fe³⁺), B:843
as petrogenic indicator, B:837-847
spirasters
Site 795, B:543

- sponge spicules. *See also* microfossils, siliceous abundance, B:344-349
Leg 127, B:292
Site 794, B:344
Site 795, B:304, 345, 541-543
Site 797, A(127):343; B:303, 345
Site 799, A(128):260
- sponges, Site 799, B:35, 45
- spongoliscids
Site 795, B:542
- spores
BSEM images, B:554-556
formation and recycling of, B:550-551
resting-spore laminae, B:549-551
- sporinite, Sites 798 and 799, B:670
- squalene, Site 799, B:627, 632
- stable isotopes. *See* isotopes, carbon; isotopes, oxygen
- steranes, in Kita-Yamato Trough, B:669
- stilbite, and ash alteration, A(128):288
- strain difference functions, Site 794, B:1054, 1056-1057
- stratal disruption, Site 798, A(128):142
- stress field, B:1323-1324
azimuthal distribution of, B:1058
crack behavior, B:1121
crustal anisotropy, B:1114-1115, 1343
deformation rate analysis (DRA), B:1048-1049
determination methods, B:1047
from earthquake analysis, B:1054
and faults and fractures, B:1181-1183
horizontal stress, B:1051-1052
and Japan Arc rotation, B:1054
Japan Sea, A(127):17, B:1185-1187
Japan Sea E, B:1059
Kita-Yamato Trough, A(128):293-294
and magma ascent, B:1319
and Oga Peninsula tectonic history, B:1204-1205
porosity effect on horizontal stress, B:1053
Site 794, A(127):84; B:1047-1059, 1120
Site 795, A(127):180-181
Site 796, A(127):258
Site 797, A(127):337
Site 799, A(128):245
strain difference functions, B:1054, 1056-1057
stress measurement procedure, B:1049-1050
stress-strain curves, B:1051
vertical stress, B:1050-1051
and volcanic activity, B:796, 1319, 1321, 1325
Yamato Rise, A(128):245
Yamato Trough, A(128):340
- strongyles
Site 795, B:542
- strontium
and ash layer alteration, A(127):280
ash layer/basement alteration and, B:643
basement, B:807
in interstitial water, B:607, 643-646, 1263
isotopic ratios vs. $1/\text{Sr}$, B:646
Site 794, A(127):110; B:1363
Site 795, A(127):205-207; B:1365
Site 796, A(127):280
Site 797, A(127):363-364, 370; B:1366
Site 798, A(128):173-174, 182
Site 799, A(128):318, 329; B:611
vs. neodymium, B:902
- strontium isotopes. *See* isotopes, strontium structures. *See* sedimentary structures styles
Site 795, B:542
- subduction zones, in Pacific Ocean W, B:1333-1334
- submarine fans. *See also* Toyama Deep Sea Fan progradation of, A(127):16
- submarine volcanism
Japan Sea, A(127):95
Site 795, A(127):219-220
- subsidence
fault-controlled, B:1344
in forearcs, B:1213
geohistory and backstripping analysis methods, B:1199-1201
Japan Sea, B:1209-1214, 1343-1344
Oga Peninsula, B:1202-1205
onshore sections and offshore wells analyzed, B:1201-1202
paleobathymetric analysis methods, B:1200-1201
Pohang Basin, B:1205-1206
rates, B:1211-1212
Sakhalin Island, B:1206-1207
and sediment loading effects, B:1211
subsidence, B:1200, 1212-1213
thermal, B:1344
Tsushima Basin, B:1207
- Sugata Formation, A(128):129
- sulfate
and barite front development, B:637
and clathrate formation, A(127):281
in interstitial water, B:637-639, 657+59, 1264
and methane abundance, A(127):368
and organic carbon decomposition, B:1262
and organic matter degradation, A(127):362
in sandstone, B:143-144
and sediment accumulation rate, B:637, 646
and sedimentation rate, A(127):362; B:1262
Site 794, A(127):108; B:637-638
Site 795, A(127):204; B:637-638, 640
Site 796, A(127):279; B:637-638
Site 797, A(127):325, 362, 368; B:637-638
Site 798, A(128):172-173, 180; B:770, 775-776, 1263
Site 799, A(128):317, 328; B:610
and source of methane, A(127):288
and sulfate-reducing bacteria, B:770
vs. alkalinity, A(128):183
vs. sedimentation rate, B:661
- sulfate reduction. *See also* sulfate and bacteria, B:767, 771
and barium dissolution, B:656, 661
and carbon isotopes, B:88
and high alkalinity values, A(127):279
and interstitial water analyses, B:607
and organic carbon decomposition, B:1262-1263
and organic matter, A(127):207, B:637
potential rates, B:771
and sedimentation rate, A(127):207
Site 794, A(127):108, 111, 119
Site 795, A(127):174, 204
Site 796, A(127):251, 267
Site 797, A(127):362
Site 798, B:770, 772-773, 775-776
Site 799, A(128):318
and sulfate-reducing bacteria, B:770
- sulfide, acid-volatile (AVS), Site 798, B:771, 775
- sulfide deposits
in failed rifts, A(128):251-252
in Japan Sea sediments, B:1241-1249
Kuroko, A(128):24, 239, 251
occurrence of, A(128):21-22
- sulfur
in dark/light cycles, B:579, 586-588
as framboidal pyrite, B:706-709
glacial/interglacial variations, B:589-590
and organic carbon, B:1241
Site 794, A(127):113; B:1246-1247, 1255, 1363
Site 795, B:706-710, 714, 209-213
Site 796, A(127):283-285
Site 797, A(127):365-367; B:592
Site 798, A(128):176-177, 189-192
Site 799, A(128):323-324, 334-338, 342; B:623-624, 626-627
vs. carbon, organic, A(128):343-344
vs. organic carbon, B:594, 628, 712-713, 1254
vs. pyrite, B:1254
vs. sulfur isotopes, B:715
- sulfur, logging, Site 798, B:1024
- sulfur/iron ratio
and pyrite formation, B:1249
Site 794, B:1255
- sulfur isotopes. *See* isotopes, sulfur
- Sumisu Rift, Pb-Pb plots of arc volcanic rocks, B:808
- Sunda Arc, sediment contamination of mantle source, B:805
- surface-water temperature. *See* temperature, surface-water
- susceptibility. *See* magnetic properties
- Suttu Canyon, bathymetry, A(127):251
- synthetic seismogram
and opal-A to opal-CT transition, B:1146-1148
seismic field record compared, B:1142
Site 794, A(127):143, 154
Site 798, A(128):184, 214
Site 799, B:1138-1140, 1142
- Takuyo Bank
basement, A(128):248-249
bathymetry, A(128):245-246
crustal structure, A(128):249
- tantalum
Site 798, B:1370-1371
Site 799, B:724
- Tartar Strait
depth, B:342
Japan Sea and, B:69-70, 365
sea ice, B:167
sill depth, B:457
- tau- p mapping, and one-dimensional velocity structure, B:1078-1080
- Taxodiaceae
Site 795, B:490
- tectonic provenance. *See* provenance, tectonic tectonic reconstructions. *See* plate reconstructions
- teleseismic events, and downhole seismic experiment, B:1162, 1167-1168
- temperature
of alteration, B:839
and clathrate stability, A(127):281
contaminated measurements, A(127):136
of diagenetic reactions, B:148-149
and electrical resistivity, A(128):111
expected decay curve, A(127):298-299
and gas hydrate stability, A(127):288, 299-300
increase with depth, A(127):135-136
Japan Sea, A(127):26-27, A(128):20
and lower crust structure, B:1343
of opal-A to opal-CT transition, A(128):150, 280; B:14, 1145-1147, 1263, 1305-1306
of opal-CT formation, B:49
of opal-CT to quartz transition, A(128):280; B:17, 1263

- and opal dissolution, B:310
in permeable sand, A(127):299
Site 794, A(127):73, 140-142
Site 795, A(127):174, 230-232
Site 796, A(127):251, 296-301
Site 797, A(127):390ff, 396-397
Site 798, A(128):183-184, 213-214
Site 799, A(128):338-339, 365
Sites 794-797, B:51, 56
and strontium isotopes, B:644
typical temperature record, WSTP tool,
B:1299
vs. opal-A/opal-CT BSR 2-way traveltime,
B:1152
vs. thermal resistance, A(127):142, 235, 300,
397; B:1302
vs. time, A(127):235
WSTP measurement reliability, B:1308-1309
- temperature, isotopic
opal-CT, B:55-56
quartz, B:55-56
- temperature, logging
Site 798, A(128):125, 183-184, 213
Site 799, A(128):338-339
- temperature, silica transformation, and oxygen iso-
topes, B:49
- temperature, surface-water
calcareous nannofossils and, B:180
fluctuations, B:463-465
and foraminifers coiling ratio, B:193
Japan Basin, B:209
Japan Sea, B:201, 337
orbital cyclicities of, B:467-468
oxygen isotope record correlated, B:464-465,
468
and planktonic foraminifers, B:187
Site 794, A(127):102-103
Site 795, A(127):199
Site 796, A(127):273
Site 797, A(127):357
Site 798, B:457-470
Site 799, A(128):310, 312
- temperature gradient. *See* thermal gradient
- tephra markers
Japan Sea, A(128):157, 291-292; B:796
- tephrochronology
Site 798, A(128):153-157, 196-198; B:793-
796
Site 799, A(128):284-291; B:793-796, 798-
799
- terbium, Site 798, B:1370-1371
- terrigenous accumulation rate. *See* terrigenous flux
- terrigenous component. *See also* sand
core analyses, B:404
logging estimates, B:402, 404
sediment gravity flow deposits, A(127):267
Site 796, A(127):250
Site 798, B:401
vs. quartz abundance, B:403
- terrigenous flux
glacial/interglacial variations, B:446
to Japan Sea, B:409
in logs, B:403
Site 798, B:419, 446
- terrigenous sediment
and chromium, B:723
influx of, A(127):16
and major and minor element chemistry,
B:722-723
and rare earth elements, B:729, 732
X-ray diffraction analyses, B:400-401
- tetraxons
Site 795, B:543
- Th/Sc ratio. *See* thorium/scandium ratio
- thanatoecoenosis, and diatom dissolution, B:309-
316
- thermal conductivity
above opal-A to opal-CT boundary, B:1147
in basement, B:1281
correction factors applied, B:1018
data quality, A(127):127
hydrostatic pressure and, B:1017-1019
Japan Sea, B:1280-1281, 1287, 1298, 1301-
1303
measuring system schematic, B:1018
porosity correlated, B:987
in sediments, B:1281
Site 794, A(127):127ff, A(128):102, 106;
B:1287
Site 795, A(127):222ff; B:1287
Site 796, A(127):290ff; B:1287
Site 797, A(127):383ff, 397; B:1287, 1300
Site 798, A(128):181-182, 206-207, 210;
B:1287
Site 799, A(128):326-327, 338, 352-353,
360, 365; B:1287
typical temperature vs. log time plot, B:1018
vs. porosity, A(127):135, 232, 395; B:25, 27,
989, 1287
- thermal demagnetization. *See* magnetic properties
- thermal gradient
isotopic, B:56
Japan Sea, A(127):26-27; B:1301-1303, 1375
and opal-A to opal-CT transition, A(127):268
sediment, B:56
and silica formation, B:52-54
Site 796, A(127):247, 251, 297, 299
Site 797, A(127):390
Site 798, A(128):125, 183-184, 213-214
Site 799, A(128):338-339, 365
Sites 794-797, B:51
and thickness of opal-CT zone, B:49
- thermal history, Japan Sea, B:49-56
- thermal maturity
and *n*-alkanes, A(128):324
of organic matter, B:671-672
Site 798, B:669
Site 799, B:624, 669-670
- thermal resistance
Site 796, A(127):299
Site 797, A(127):390
Site 799, A(128):338
vs. temperature, A(127):235, 300, 397; B:1302
- thermocline, in Japan Sea N, B:167
- thermogenic gas. *See* gas, thermogenic
- thermohaline circulation, Japan Sea, B:528-529
- tholeiite
in basalt, B:870, 875
chemistry, B:839
rare earth elements, B:922, 925
Site 794, A(128):33, 68-69, 96-99, 117;
B:779-780
Site 795, B:919
Site 797, B:920
spidergrams, B:923, 926
- thorium
in ash layers, B:1386
in basement rocks, B:807
Site 795, B:707
Site 798, B:1370-1371
Site 799, B:724
- thorium, logging
Site 794, A(128):103, 108; B:1418-1419
Site 796, B:1422
Site 797, B:1424-1425
Site 798, B:1400
- Site 799, A(128):366-367; B:1403, 1406-1407
- thorium/scandium ratio
and volcanic activity, B:723, 732
Site 798, B:723, 733
Site 799, B:723, 734
- thorium/uranium ratio, logging
in ash layers, A(128):339
Site 799, A(128):366
- thrust faults
fluid flow along, A(127):247, 295, 301
frictional heating along, A(127):247, 301
Japan Basin E, A(127):255
Japan Sea, A(127):81-82
Japan Sea E, A(127):9
Yamato Basin N, A(128):75-76
- Tibetan Plateau, uplift, B:338
- titanium
in augite, B:851-853, 856, 862-863
in Japan Sea sediment, B:1236
and parental magma, B:856
Site 798, B:722
Site 799, B:722
vs. Mg/(Mg + Fe), B:858
vs. neodymium, B:902
vs. zirconium, A(128):101
XRF and geochemical log analyses compared,
B:1028-1029; 1031-1035
- titanium, logging
Site 794, B:1420-1421
Site 796, B:1423
Site 797, B:1426-1427
Site 798, B:1401-1402
Site 799, B:1404, 1408-1409
- titanium/aluminum ratio
in augite, B:851-853
as indicator of eolian contribution, B:1239
Site 794, B:1249
- titanium oxide. *See* titanium
- titanomagnetite, Site 797, B:949, 951
- T_{max}
Site 798, B:668-669
Site 799, B:624-625, 668-669
vs. transformation ratio (TR), B:630
- tomogram, X-ray radiograph compared, B:1133
- Tonga Arc
Pb-Pb plots of arc volcanic rocks, B:808
and sediment contamination of mantle source,
B:812
- topography, and heat flow, A(127):300-301
- tourmaline, as quartz inclusions, B:108
- toxons
Site 795, B:542
- Toyama Channel, pollen and, B:333
- Toyama Deep Sea Fan, A(127):11
distribution of deposits, B:1146
and heat flow, B:1148
sands, A(128):14
seismic expression, B:1148
- Toyama Trough, A(128):70
- TR. *See* transformation yield
- trace element molar ratios
Site 798, B:616-617
Site 799, B:618-621
- trace elements, B:677-695, 719-737. *See also* X-
ray fluorescence (XRF) analyses; *specific*
trace elements
Al-normalized concentrations, B:689
and ash layer magmatic distinctions, B:1378-
1379
of basalt, B:839
and basaltic basement rocks, B:893
basement, A(128):98; B:781-783, 807, 912-
916

- detection limits, INAA analyses, B:1372
 interelement comparisons, B:690
 mobility during alteration, B:911-916
 N-MORB normalized spidergram, B:923-924, 926
 Site 794, B:682-684, 689, 782-783, 839, 912-913, 918-919, 923
 Site 795, B:683-685, 689, 705-717, 913-914, 924
 Site 796, B:685
 Site 797, B:685, 688-691, 839, 914-916, 920-921, 926
 Site 798, B:722-725, 1367-1372
 Site 799, B:723-725, 728-731
 in spinel, B:840-842
 in strip samples, B:1367-1372
 vs. loss on ignition, B:918-921
- trace metals, B:1361-1366
 Site 794, B:1362-1363
 Site 795, B:1365
 Site 797, B:1366
- transformation ratio (TR)
 Site 799, B:624-625
 vs. T_{max} , B:630
- trees, broadleaf, Site 798, B:319
- triaxons
 Site 795, B:543
- trissocyclids
 Japan Sea, B:294, 296
 Site 797, B:303
- Tsugaru Strait, B:423
 depth, B:342, 440
 Japan Sea and, B:70, 362-363, 365
 sill depth, B:351, 457
- Tsushima Basin
 crustal structure, B:1311-1314, 1318-1319
 magnetic anomalies, B:1314
 paleobathymetry, B:1210
 subsidence history, B:1207-1211, 1214
- Tsushima Current, B:342, 423-424
 benthic foraminifers and, B:365
 calcareous nannofossils and, B:180
 and foraminiferal coiling direction, A(128):165, 310-311
 inflow and diatom abundance correlated, B:591
 and Japan Sea, A(128):20-21; B:365
 and onset of glaciation, B:466
 and productivity, B:311
 radiolarians and, B:298-299
 Site 799, A(128):310, 312
 and upwelling, B:594
 warm surface water, B:440, 457, 577
- Tsushima Fault
 dextral motion along, B:1317
 and Japan Sea tectonic evolution, B:1315
 offset along, B:1325
- Tsushima Strait, B:423
 age, B:180
 and biosiliceous productivity, B:313
 depth, B:342, 440
 and diatom abundance, B:1219-1220
 diatoms and, B:360
 Japan Sea and, B:362-363, 365
 and Kuroshio Current, A(128):164; B:166-167
 sill depth, B:457
 Site 798, A(128):164-166
 Site 799, A(128):312
 and tectonic subsidence, B:591
 and Tshushima Current, B:69
- tuff
 formation, A(127):190-191
 in logs, A(127):251
- sedimentary structures, A(128):81
 Site 794, A(127):93-94, A(128):81
 Site 795, A(127):189-190, 192
 Site 796, A(127):265-266
 Site 797, A(127):344-345
- tuff, ash-flow
 Onnagawa Formation rhyolitic tuff correlated, B:796
 Site 799, B:793, 803
- tuff, breccia, chemical composition, A(128):287-288
- tuff, hyaloclastite
 Site 794, B:1391
 Site 795, B:1374
- tuff, pumice flow, Site 799, B:1373
- tuff, rhyolitic. *See also* tuff, ash-flow; tuff, pumice flow
 age, B:796
 ash layers correlated, A(128):33
 in geochemical logs, B:1399
 in logs, A(128):339
 Site 799, A(128):35, 243, 265, 292, 354
 and velocity, B:1141
 XRF analyses, A(128):300
- tuff, vitric
 in logs, B:37
 and revised lithostratigraphy, B:1232-1233
 Site 799, B:34, 36, 39, 41-42, 44
- tuff breccia. *See* breccia, tuff
- tuffaceous clayey siltstone. *See* siltstone, tuffaceous clayey
- tungsten, Site 798, B:1370-1371
- turbidite deposits
 seismic expression, A(127):233
 Site 795, A(127):192
 Site 799, A(128):264
- turbidites
 glauconite-bearing, B:68
 sedimentation rate and, B:959
 Site 799, B:104
- turbidity currents, Site 798, B:419, 421
- turbidity deposits, of ash, B:793
- turbidity flows
 of ash layers, A(128):153-154
 Site 799, A(128):259
- tylostyles
 Site 795, B:543
- Tym-Poronaisk Fault, and Japan Sea tectonic evolution, B:1315
- Tyro Basin, paleoenvironment, B:379
- Ulleung Island, ash layers from, A(128):156-157
 unconformity, seismic, and Okushiri Ridge uplift, A(127):312-313
- underconsolidation. *See* consolidation
- United States NW, paleoenvironment, B:337
- uplifting
 in Japan Sea region, B:1343-1344
 Site 799, A(128):312
- uptake rates, fluoride
 Sites 798 and 799, B:66
 vs. sediment depth, B:65
- upwelling
 and dark/light cycles, B:591-594
 enhanced productivity, A(127):267
 evidence from planktonic foraminifers, B:469
 and gyre-margin planktonic foraminiferal assemblages, B:466
 Japan Basin N, A(127):199
 Japan Sea, B:337
 and northern hemisphere glaciation, A(127):271
 radiolarians and, B:299
 and resting-spore laminae, B:551
- and sedimentation rate, A(127):204
 siliceous ooze deposition, A(127):102-103, 191
 Site 795, A(127):199
 Site 796, A(127):274
 Site 797, A(127):349
 and Tsushima Current, B:594
- uranium
 in basement rocks, B:807
 Site 795, B:711, 714
 Site 798, B:1370-1371
 Site 799, B:724
- uranium, logging
 in igneous rocks, A(128):103
 Site 794, A(128):108; B:1418-1419
 Site 796, B:1422
 Site 797, B:1424-1425
 Site 798, B:1400
 Site 799, A(128):366-367; B:1403, 1406-1407
- Uyeda probe. *See* temperature
 vacuainer analyses. *See* hydrocarbons, volatile
 van Krevelen plot
 Site 794, A(127):115-118, 121
 Site 795, A(127):213, 219
 Site 796, A(127):285, 288
 Site 797, A(127):367, 378
 Site 798, A(128):177, 196; B:429
 Site 799, B:429, 629
- vegetation, Japanese archipelago, B:320, 325, 326, 334
- veins
 altered to clays, B:886-887
 and Bagnold effect, B:1180
 in basement, A(128):92-94
 and bedding-parallel extension, B:1176
 chemical composition of alteration clay minerals, B:887
 coarse-grained material in fillings, B:1179
 crystallization stages, A(128):92-93
 and earthquakes, B:1180-1181
 filled tension gash, B:1189
 fillings, B:1178
 and fluid flow/material transport, B:1178-1179
 formation timing, B:1178, 1180
 as healed normal faults, B:1178
 and Japan Sea stress field, B:1187
 lithology correlated, B:1180
 and matrix sieving, B:1179-1180
 and microfaults, B:1180
 order of filling, B:888
 origin, B:1178-1181
 Site 794, B:1182-1183
 Site 798, A(128):143ff; B:1176-1178
 Site 799, A(128):263-264, 269-272; B:1178, 1190-1193
- veins, siliceous
 and phosphate, B:64
 in plagioclase phyric basalt, A(127):217
- velocity
 anisotropy, A(127):229
 basement, A(127):389
 corrected analyses, B:990-1015
 and crustal structure, B:1078-1083
 Japan Sea, B:1279-1280, 1283
 lab and log data compared, B:1279-1280, 1284
 of opal-A to opal-CT transition, B:18, 21
 porosity effect, B:1280
 of rhyolitic tuff, B:1141
 Site 794, A(127):127-134, 139, A(128):102, 105; B:1283
 Site 795, A(127):224-227, 229; B:1283

- Site 796, A(127):290-295; B:1283
 Site 797, A(127):383-389, 391; B:1283
 Site 798, A(128):181, 205; B:1283
 Site 799, A(128):326, 350-351, 359; B:1137, 1140, 1283
 vs. loss on ignition, B:1291
 vs. porosity, A(127):140, 395, B:25, 27, 1005, 1286, 1290
 vs. porosity, logging, B:26
- velocity, logging
 log quality, A(128):103
 Site 794, A(127):134, 139-142, 144-146, A(128):107, 119-120; B:1069
 Site 795, A(127):232, 237-238
 Site 796, A(127):302-305
 Site 797, A(127):393-395, 398, 400-402; B:7
 Site 798, A(128):185-187, 212, 225-228
 Site 799, A(128):332-338, 362-364, 380-382, 393-397
- vertical seismic profile
 depth trace ensemble, A(128):371-372
 energy source, A(128):342
 experimental configuration, A(128):370
 operations, A(128):343
 Site 799, A(128):342-343
- vesicles
 in ash layers, B:791
 in basalt and dolerite, A(128):88-91
 in blue tuff, B:117-119
 chemical composition of alteration clay minerals, B:887
 in dolerite and basalt, B:887-888
 order of filling, B:888
- viscous remanent magnetization (VRM). *See* magnetic properties
- visual microanalysis, methods used, B:57-62
 vitrinite, Sites 798 and 799, B:670
 Vityaz Rise, dredge samples, A(127):177
 Vladivostok, and sea ice, B:167
- void ratio
 Site 794, A(127):128-133
 Site 795, A(127):224-227
 Site 796, A(127):291-293
 Site 797, A(127):383-389
 Site 798, A(128):197-198, 202-205
 Site 799, A(128):345-349
- volcanic activity. *See also* volcanic phases; volcanic pulses
 and ash layer distribution, B:963
 and Eu anomaly, B:731-732
 factors affecting, B:1320-1321
 and Japan Sea, A(128):33; B:1319, 1321
 Site 798, A(128):35, 156, 196-198; B:796-797, 800
 Site 799, A(128):288-291; B:796-797, 800
 Sites 798 and 799 compared, A(128):291-292
 and stress field, B:1319, 1321, 1325
 and thorium/scandium ratio, B:732
 volcanic stages recognized, B:1320-1321
- volcanic eruptions, and nature of ash layers, A(128):152-153
- volcanic front, migration, B:1324, 1330
- volcanic glass. *See* glass, volcanic
- volcanic phases
 Site 798, B:796-797, 800
 Site 799, B:796-797, 800
- volcanic pulses, B:1373-1374. *See* volcanic activity; volcanic phases
 Site 794, B:1374
 Site 795, B:1374
 Site 796, B:1374
 Site 797, B:1373-1374
 Site 798, B:1374
- Site 799, B:1373
 volcanic rock fragments. *See* rock fragments
 volcanoclastic material. *See* ash layers
 volcanism. *See* submarine volcanism; volcanic activity; volcanic pulses
- Vøring Plateau, sediment consolidation, B:1277
- VRM. *See* magnetic properties
- VSP experiment. *See* vertical seismic profile
- Wadati-Benioff zone, Japan Sea, B:1062, 1158
- Wakimoto Formation
 deposition, A(128):18
 foraminifers, B:187
 planktonic foraminifers zonations, B:459
- warming event
 Japan Sea and, B:338
 pollen indicators of, B:479, 484
 Site 798, A(128):162, 164-165
- water content
 of basement, A(127):389
 corrected analyses, B:990-1015
 and electrical resistivity, A(128):111
 Japan Sea, A(127):26
 Site 794, A(127):127-133, A(128):103, 105
 Site 795, A(127):223-228
 Site 796, A(127):290-295
 Site 797, A(127):383-391
 Site 798, A(128):180-181, 197-198, 202-205, 208
 Site 799, A(128):325, 345-349, 354
 vs. density, A(127):233, 395
 vs. porosity, A(127):234
- water-escape structures, Site 799, A(128):268-269
- water ratio
 corrected analyses, B:990-1015
 Site 794, A(127):128-133
 Site 795, A(127):224-227
 Site 796, A(127):291-293
 Site 797, A(127):383-389
- water temperature. *See* temperature, surface-water
- water vapor, and index property data errors, B:986
- Weddell Sea, sediment consolidation, B:1277
- well-logging. *See* logging
- wet-bulk density. *See* density
- wind
 and Japan Sea, B:409, 411, 447
 pollen and, B:333
- winnowing, Site 798, A(128):165
- X-ray diffraction (XRD) analyses
 of chlorite/saponite mixed layer clay, B:886
 CuK- α , lansfordite and magnesite, B:81
 of dark/light cycles, B:579, 584-588
 of diagenetic carbonate, B:79-80, 86-87
 of dolomite, B:79-80
 of lansfordite, B:80
 of magnesite, B:80
 of opal-A to opal-CT transition, B:10, 17-18
 of opal-CT to quartz transition, B:24
 peak height comparisons, B:589
 of rhodochrosite, B:79-80
 of sediments, B:1236-1244
 silica, A(128):292
 Site 795, B:8-10
 Site 797, B:11-14
 Site 798, B:400-401
- X-ray fluorescence (XRF) analyses. *See also* basement; geochemistry, sediment
 of ash layers, A(128):151
 of basalt, B:872
 of basement, B:912-916
 of dark/light cycles, B:569
 geochemical logs compared, B:1025-1029, 1402, 1416-1417, 1420-1421, 1427
 incompatible elements, A(128):69
- Kleiner-Hartigan diagrams, B:1030
 loss on ignition as alteration proxy, B:892-893, 909
 magnetic results correlated, A(127):358-359
 of phosphate, B:64-65
 physical properties of basement compared, B:1283
 of sediments, A(128):151, 292; B:1022-1027, 1031-1035
 ship vs. shore data compared, B:779
 silica data, A(128):281
 Site 794, A(127):123, 126, A(128):68, 96ff; B:872, 892-894, 912-913, 1420-1421
 Site 795, A(127):219, 223; B:913-914
 Site 797, A(127):381-383; B:872, 892-894, 914-916, 1427
 Site 798, B:1022-1027, 1031-1035, 1402
 Site 799, A(128):292
 trace elements, A(128):69
- X-ray radiograph, computer tomogram compared, B:1133
- Yamato ash, tephra marker, B:796
- Yamato Bank
 basement, A(127):333, A(128):247-249
 bathymetry, A(128):245-246
 crustal structure, A(127):330, A(128):248
 and Japan Sea tectonic evolution, B:1315
 Layer 3, A(128):246
 magnetic anomalies, A(128):247
 phosphate source, B:67-68
 sandstone provenance, B:110
 seismic stratigraphy, A(127):399
 subsidence, A(128):295
- Yamato Basin. *See also* Yamato Basin N; Yamato Basin S
 age, A(127):8, 29; B:179, 906-908, 1220
 basement, A(127):29, A(128):23-24, 32-33, 36; B:894, 906-908
 basement chemistry, B:779-789
 basement depth, A(127):6-8
 basement isotopic compositions, B:786-787
 basin floor subsidence, A(127):103
 bathymetry, A(128):126-127
 benthic foraminifers, B:208, 495-496, 500-517, 522-528
 blue tuff, B:127-128
 calcareous nannofossils, B:174-179
 crustal anisotropy, B:1107-1121
 crustal composition, B:928
 crustal extension, B:1345-1346
 crustal structure, A(127):9, 79, 176, A(128):9-11, 24, 70-73, 127-128, 130, 248; B:809, 837, 899, 1083, 1086, 1104, 1107, 1304-1305, 1311-1314, 1318-1319-1341-1346
 crustal structure and tectonic history, B:1318-1319
 depositional history, A(128):81
 diatoms, B:252-258, 264-269, 283-284
 dredge samples, A(127):82
 electrical resistivity, B:1354
 Foram Sharp Line and, B:506, 517
 formation, A(127):10; B:928
 formation age, B:830-831
 free-air gravity anomalies, A(128):127
 geologic history, A(127):72
 geologic setting, B:1147
 heat flow, A(127):27, 73, 330, 390, A(128):127, 246, 250, B:1147-1148, 1150, 1302, 1345
 hotspot activity, B:1345
 interbedded sediments, A(127):25, 29

- Japan Arc basalt correlated, B:903
 Japan Basin age compared, B:1337
 Japan Basin foraminifers compared, B:189
 Lau Basin formation compared, B:928
 Layer 1, A(128):71-72, 127
 Layer 2, A(127):73, A(128):9, 72, 127-128; B:1342
 Layer 3, A(127):73, A(128):71-72, 127-128; B:1342-1343
 and lithospheric stretching models, B:1304-1305, 1307
 lithostratigraphy, B:494, 496, 498-499
 low-Al igneous rocks, B:845
 magma source, B:782-784, 902, 1341
 magmatic history, B:787-788
 magnetic anomalies, A(127):8, A(128):7; B:942, 1314, 1345
 magnetic properties, B:941-942
 magnetostratigraphy, A(127):22
 neodymium isotopes, B:901-903
 oceanic crust, A(127):73
 oceanic lithosphere, B:1354
 oceanographic setting, B:342
 opal-A to opal-CT transition, B:50
 opal-CT to quartz transition, B:50
 oxygenation conditions, B:201-208
 paleodepth, A(127):199; B:189, 201, 1220-1221
 pollen, B:328, 471, 479, 482-483, 485-486
 radiolarians, B:293, 295-296
 radiometric age, B:784-786
 rifting, A(127):29-30
 sedimentation rate, B:287, 297, 1227
 seismic stratigraphy, A(127):399
 stress field, B:1047-1059
 strontium isotopes, B:901-903
 subsidence history, A(127):20, 30, 349; B:209, 1208-1211, 1214
 tectonic evolution, B:787-788, 1187, 1316-1317
 Toyama submarine fan, A(127):11
 volcanic activity, B:796-797
 Yamato Basin N. *See also* Yamato Basin; Yamato Basin S
 age, A(128):67-69
 basement, A(127):77-81, 126-127, A(128):73-75, 117
 bathymetry, A(128):70-71
 biostratigraphy, B:1221-1223
 crustal structure, B:1075-1106
 depositional history, A(127):96
 free-air gravity anomalies, A(127):77, A(128):73
 heat flow, A(128):71-73, 75
 igneous event sequence, A(128):95-97
 magnetic anomalies, A(127):77, A(128):73
 magnetostratigraphy, B:969-970, 1221-1223
 paleoenvironment, A(127):102-103
 rifting history, A(127):143-147
 sedimentation, A(127):83-84, 147, A(128):77-81
 seismic stratigraphy, A(127):143
 Site 794, B:654-656
 surface projections of fault ruptures, A(128):112
 tectonic setting, A(127):81-83, A(128):75-77, 82
 Yamato Basin S. *See also* Yamato Basin; Yamato Basin N
 basement, A(127):323, 331-334, 408-409
 bathymetry, A(127):325, 331
 biostratigraphy, B:1223
 carbonate-compensation depth, A(127):324
 crustal structure, A(127):329-330, 333; B:1083, 1104-1106
 depositional history, A(127):347-351
 dredge samples, A(127):332
 free-air gravity anomalies, A(127):330-331, 335
 geologic history, A(127):323-324
 heat flow, A(127):330, 332
 Layer 1, A(127):330
 Layer 2, A(127):330
 Layer 3, A(127):330
 magnetic anomalies, A(127):330, 334
 magnetostratigraphy, B:973, 1223
 oceanographic conditions, A(127):349
 oceanographic history, A(127):337, 409-410
 paleoenvironment, A(127):356
 rifting, A(127):408-409
 sedimentation, A(127):324, 334-335, 337, 409-410
 seismic stratigraphy, A(127):335, 399
 Site 797, B:560
 subsidence, A(127):323, 409
 tectonic setting, A(127):334, 336
 Yamato Block, B:809
 Yamato Rise, A(127):325, A(128):70
 basement, A(127):9, 333, A(128):73, 247-249
 bathymetry, A(128):245-246; B:110, 1136
 crustal structure, A(127):9, 329-330, 333, A(128):9-11, 245-249, B:1075
 dredge sites, A(128):253
 free-air gravity anomalies, A(128):247
 geologic history, A(128):21-22
 Green Tuff rocks, A(128):249
 heat flow, A(127):330, A(128):246, 250; B:1302-1303
 heavy minerals, B:691
 Layer 2, A(128):246
 Layer 3, A(128):246
 magnetic anomalies, A(128):247; B:1314
 sediment from, A(127):349
 sedimentary rocks dredged, A(128):249
 sedimentation, A(127):11-13
 Site 799, A(128):312
 stress field, A(128):245
 tectonic setting, A(127):8-9, A(128):249-250; B:156
 Yamato Rise E, normal faults, A(127):82-83
 Yamato Seamount Chain, A(128):70
 basement, A(127):333, A(128):73
 crustal structure, A(128):128
 dredge samples, A(127):77-78
 strontium isotopes, B:812
 Yamato Trough, A(128):70
 stress field, A(128):340
 vertical seismic profile, A(128):342-343
 Yansan Fault
 and Japan Sea tectonic evolution, B:1315
 offset along, B:1325
 Yellow River
 fresh water input to Japan Sea, B:458
 origin of dark/light cycle detrital component, B:595
 Young's modulus, Site 794, B:1050
 ytterbium
 in ash layers, B:1386
 Site 798, B:1370-1371
 vs. aluminum, B:737
 yttrium
 alteration and, B:909-911
 basement, A(128):99
 Site 794, B:1363
 Site 795, B:1365
 Site 797, B:1366
 vs. barium, A(128):100
 vs. chromium, A(128):101
 vs. zirconium, A(128):100; B:783
 zeolites. *See also* zeolitization; *names of specific zeolites*
 in basaltic andesite, A(127):217
 in blue tuff, B:119
 formation temperature, B:139
 and opal-A to opal-CT transition, B:10
 in pyroxene plagioclase phyric basalt, A(127):217-219
 SEM photographs, B:30
 Si/Al + Fe ratios, B:139, 141
 Site 797, B:142
 and volcanic glass alteration, B:1375-1377
 zonation of, B:143
 zeolitization. *See also* zeolites
 provenance information destroyed, B:148
 of sandstone, B:137-139
 Zijderveld plots
 of Leg 128 pilot samples, B:971
 Site 794, A(127):104-105, A(128):104-105
 Site 795, A(127):203, 206-207
 Site 796, A(127):277-279
 Site 798, A(128):173
 Site 799, A(128):316, 318-320, 322-323
 zinc
 in dark/light cycles, B:569
 Site 794, B:1363
 Site 795, B:710, 1365
 Site 797, B:1366
 Site 798, B:1370-1371
 in sulfide deposits, A(128):21
 zirconium
 in ash layers, B:1386
 and basalt fractionation models, B:873
 basement, A(128):99
 and fractionation models, B:881
 Site 794, B:1363
 Site 795, B:1365
 Site 797, B:1366
 vs. barium, A(128):100
 vs. niobium, A(128):100; B:783
 vs. titanium, A(128):101
 vs. yttrium, A(128):100; B:783
 vs. zirconium/yttrium ratio, A(128):100
 zirconium/yttrium ratio, vs. zirconium, A(128):100

TAXONOMIC INDEX

- Abies*
Japan Sea, B:328
Neogene climate and, B:337
- abyssalica*, *Silicosigmoilina*, Japan Sea, B:505
- acquiloni*, *Drupptractus*
Japan Sea, B:228
Site 799, B:235
- acquiloni*, *Stylocontarium*
Japan Sea, B:297
Site 794, B:305
- acquiloni*, *Stylocontarium* sp. cf., Japan Sea, B:297
- Actinocyclus ingens*, Pacific Ocean, B:309
- Actinocyclus oculatus*
Japan Sea, B:258
Site 798, B:361
- Aesculus*, Site 796, B:491
- akitaensis*, *Anthocorys*, Site 797, B:307
- akitaensis*, *Anthocorys*?
Japan Sea, B:228
Site 799, B:225, 235
- akitaensis*, *Uvigerina*, Site 798, B:390, 441, 443–444
- Alnus*, Site 795, B:490
- aminensis*, *Bermudezina*, Site 795, B:537
- Ammodiscus macilentus*, Site 795, B:537
- Ammodiscus tenuis*, Site 797, B:537
- Ammodochium rectangulare*
Japan Sea, B:238
Site 798, B:248
- Ammodochium rectangulare* Zone, Japan Sea, B:240
- Ammosphaeroidina* sp., Site 795, B:537
- Ammovertellina* sp., Site 797, B:537
- Amphirhopalum ypsilon*, Site 798, B:235
- Angulogerina kokozuraensis*, Site 798, B:390
- Anthocorys akitaensis*, Site 797, B:307
- Anthocorys?* *akitaensis*
Japan Sea, B:228
Site 799, B:225, 235
- antiqua*, *Thalassiosira*, Japan Sea, B:258
- antiqua antiqua*, *Ebriopsis*
Japan Sea, B:238
Site 799, B:248
- antiqua cornuta*, *Ebriopsis*, Site 798, B:248
- aperta*, *Gephyrocapsa*, Leg 128, B:164
- apertura*, *Pullenia*, Site 798, B:392
- Artemisia*, Site 795, B:491
- asanoi*, *Eucyrtidium*, Site 799, B:236
- asanoi*, *Neogloboquadrina*, Site 797, B:208
- asanoi*, *Planularia*, Site 798, B:389
- asanoi*, *Reticulofenestra*, Site 798, B:164
- Astrononion* sp., Site 797, B:540
- auriculata*, *Globobulimina*, Site 798, B:390
- baccata japonica*, *Karreriella*, Site 798, B:389
- Bathysiphon* sp., Site 797, B:537
- Bermudezina aminensis*, Site 795, B:537
- Betula*
Site 795, B:490
Site 797, B:490
- biconica*, *Fissurina*, Site 798, B:389
- bigelowii*, *Braarudosphaera*, Site 795, A(127):194
- Bolivina*, Site 798, B:557
- Bolivina decussata*, Site 798, B:389
- Bolivina pacifica*
paleoenvironment and, B:379
Site 798, A(128):166; B:373, 389
- Bolivina spissa*, Site 798, B:389
- Bolivinita quadrilatera*, Site 799, B:389
- Botryostrobus bramlettei*, Site 794, A(127):102
- Braarudosphaera bigelowii*, Site 795, A(127):194
- bradyi*, *Cibicidoides*
Japan Sea, B:505
Site 797, B:538
- bradyi*, *eggerella*, Site 795, B:537
- bramlettei*, *Botryostrobus*, Site 794, A(127):102
- Buccella* cf. *kuromatsunaiensis*, Site 797, B:540
- Bulimina rostrata*, Site 795, B:539
- Bulimina striata*
Site 797, B:539
Site 798, B:390
- Buliminella tenuata*, Site 798, B:390
- bulloides*, *Globigerina*
Leg 127, B:193, 200
Leg 128, B:60–61
Site 797, B:591–594
Site 798, B:441–443, 457, 460, 462–463, 465–466, 469, 564
- bulloides*, *Pullenia*, Site 797, B:540
- bulloides*, *Sphaeroidina*, Site 798, B:390
- bulloides* (s.l.), *Globigerina*, Site 798, A(128):165
- bulloides umbilicata*, *Globigerina*, Oki Ridge, B:444
- cabrilloensis*, *Clathrocyclus*
Japan Sea, B:297
Site 794, B:307, 1221
Site 797, B:307
- Calcidiscus leptoporus*, Site 798, B:168
- Calcidiscus macintyreii*
Japan Sea, B:174, 182
Site 798, B:164
- californica*, *Rouxia*, Japan Sea, B:259
- cancellata*, *Cyclammina*
Site 797, B:538
Site 799, B:389
- Carduoideae*, Site 795, B:491
- caribbeanica*, *Gephyrocapsa*
Japan Sea, B:173, 183
Leg 128, B:164
- Carpinus*, Site 797, B:490
- Carya*
Japan Sea, B:484–486
Site 795, B:490
- Cassidulina*, Leg 128, B:62
- Cassidulina japonica*, Site 798, B:389
- Cassidulina norcrossi*, Site 798, B:390
- Cassidulina norvangi*, Site 798, B:390
- Cassidulina* sp. A, Site 798, B:390
- Cassidulinoides tenuis*, Site 799, B:390
- Castanea*, Site 794, B:490
- Catapsydrax parvulus*, Site 797, B:189, 1220, 1223
- Chaetoceros* resting spores, Site 798, B:547–557, 564
- Chilostomella oolina*
Site 797, B:539
Site 799, B:392
- Chilostomella oolina/Cyclammina* sp. C Zone, Site 794, B:517
- Chilostomella ovoidea*, Site 798, B:392
- Chilostomellina fimbriata*, Site 798, B:392
- Chondrites*
Japan Sea, B:1232
Site 798, B:565, 569
Site 799, B:1133
- Cibicidoides bradyi*
Japan Sea, B:505
Site 797, B:538
- Cibicidoides* sp. A, Site 797, B:538
- Cibicidoides wuellerstorfi*
Japan Sea, B:505
Site 797, B:538
- Cibicidoides wuellerstorfi* Zone, Site 797, B:521, 528
- circulus*, *Paramesocena*, Site 798, B:248
- circumtexta*, *Periphyramis*, Site 795, B:304
- Circumtexta pacifica*, Site 795, B:304
- Clathrocyclus cabrilloensis*
Japan Sea, B:297
Site 794, B:307, 1221
Site 797, B:307
- clavatum*, *Elphidium*, Site 798, B:392
- Coccolithus miopelagicus* Subzone, Japan Sea, B:173
- Coccolithus pelagicus*
Site 798, B:167–168
Site 799, B:166
- communis*, *Martinottiella*
Japan Sea, B:1201
Site 795, B:537
Site 797, B:537
Site 798, B:389
- compactus*, *Sphenolithus*, Japan Sea, B:183
- complanata*, *Stainforthia*
Japan Sea, B:388
Site 798, B:390
- compressa*, *Islandiella* cf., Site 797, B:539
- compressa*, *Proteonina*, Site 795, B:537
- compressa*, *Spirosigmoilina*
Site 795, B:537
Site 798, B:379–381, 389
- convexa*, *Thalassiosira*, Japan Sea, B:258
- Corylus*, Site 794, B:490
- Coscinodiscus curvatus*, Site 794, B:1223
- Coscinodiscus marginatus*
Aikawa section, Honshu, B:316
Japan Sea, B:291
Leg 127, B:1219–1220
Leg 128, B:309–314, 1219–1220
North Pacific, B:315–316
Site 433, B:316
Site 581, B:316
Site 794, B:341–357
Site 795, B:341–357
Site 797, B:341–357
Site 798, B:361, 554, 1224
Site 799, B:1224
- crassimargo*, *Cribrostomoides* cf., Site 795, B:537
- Cribrostomoides* cf. *crassimargo*, Site 795, B:537
- crux* var. *stauracanthus*, *Distephanus*
Japan Sea, B:239, 244
Site 798, B:248
- Cryptomeria*
Japan Sea, B:319, 328, 334
Neogene climate and, B:337
- Cunninghamia*, Site 797, B:490
- curvatus*, *Coscinodiscus*, Site 794, B:1223
- curvirostris*, *Rhizosolenia*
Hess Rise, B:360
Japan Sea, B:258
Leg 127, B:581
North Pacific, B:311
Site 798, B:361, 468
- Cycladophora davisiana*, Site 799, B:235
- Cycladophora davisiana cornutoides*
Japan Sea, B:230
Site 799, B:235

- Cycladophora davisiana davisiana*, Japan Sea, B:228
- Cycladophora robusta*
Japan Sea, B:299
Site 794, B:307
- Cyclammina cancellata*
Site 797, B:538
Site 799, B:389
- Cyclammina ezoensis*, Site 795, B:538
- Cyclammina ezoensis* Zone, Site 795, B:521
- Cyclammina orbicularis*, Site 795, B:538
- Cyclammina pusilla*, Site 797, B:538
- Cyclammina* sp. A, Site 797, B:538
- Cyclammina* sp. A/ *Cyclammina*? sp. Zone
age models and, B:531
Site 797, B:521
- Cyclicargolithus floridanus*
Site 795, A(127):195
Site 797, B:179
- Cyrtocapsella tetrapera*
Japan Sea, B:230
Site 794, B:304
Site 797, A(127):356; B:304
Site 799, B:235
- Cyrtocapsella tetrapera* Zone, Site 799, B:226, 1220
- davisiana*, *Cycladophora*, Site 799, B:235
- davisiana*, *Theocalyptra*
Japan Sea, B:297, 299
Site 794, B:306
Site 795, B:303
- davisiana cornutoides*, *Cycladophora*
Japan Sea, B:230
Site 799, B:235
- davisiana davisiana*, *Cycladophora*, Japan Sea, B:228
- decussata*, *Bolivina*, Site 798, B:389
- deflandrei*, *Discoaster*, Japan Sea, B:174
- delmontensis*, *Stichocorys*
Japan Sea, B:228, 230
Site 799, B:225-226, 236
- Dentalina* sp., Site 797, B:539
- Denticulopsis dimorpha*
Japan Sea, B:250
South Atlantic, B:309
- Denticulopsis dimorpha* Zone, Site 795, B:188
- Denticulopsis hustedtii*, Japan Sea, B:250
- Denticulopsis hustedtii* Zone, Site 795, B:189
- Denticulopsis kamtschatica*, Site 795, B:1223
- Denticulopsis katayamae* Zone
Site 794, B:189
Site 797, B:1223
- Denticulopsis praedimorpha*, Japan Sea, B:250
- Denticulopsis praedimorpha* Zone
Japan Sea, B:249-250
Site 795, B:189
- Dictyochoa neopseudofibula*
Japan Sea, B:237-238
Site 798, B:248
- Dictyochoa subarctios*
Japan Sea, B:238-239
Site 798, B:248
- Dictyochoa subarctios* Zone, Japan Sea, B:240-241
- Dictyococcites perplexus*, Japan Sea, B:182
- Didymocyrtes mammifera*, Site 799, B:235
- Didymocyrtes tetrathalamus*
Site 797, B:306
Site 798, B:235
- dimorpha*, *Denticulopsis*
Japan Sea, B:250
South Atlantic, B:309
- Discoaster deflandrei*, Japan Sea, B:174
- Discoaster exilis* Zone, Japan Sea, B:173
- Discoaster hamatus* Zone, Site 797, B:189
- Discoaster kugleri* Subzone, Japan Sea, B:173
- Distephanus crux* var. *stauracanthus*
Japan Sea, B:239, 244
Site 798, B:248
- Distephanus jimlingii*
Japan Sea, B:238
Site 798, B:248
- Distephanus jimlingii* Zone, Japan Sea, B:239
- Distephanus octangulatus*, Site 798, B:248
- Distephanus octangulatus* Zone, Japan Sea, B:241
- Distephanus slavnicii*, Site 798, B:248
- Distephanus speculum*, Site 799, B:248
- doliolus*, *Pseudoemotia*
age models and, B:362
Hess Rise, B:360
Honshu coast, B:360
- Drupptractus acquiloni*
Japan Sea, B:228
Site 799, B:235
- Ebriopsis antiqua antiqua*
Japan Sea, B:238
Site 799, B:248
- Ebriopsis antiqua antiqua* Zone, Japan Sea, B:239-240
- Ebriopsis antiqua cornuta*, Site 798, B:248
- echigoensis*, *Fissurina*, Site 798, B:389
- echigoensis*, *Miliammina*
Leg 127, B:200
Site 798, B:389
Site 799, A(128):311
- Echigoina hataii*, Site 798, B:392
- Eggerella bradyi*, Site 795, B:537
- Eggerella* sp., Site 797, B:537
- Eilohedra rotunda*, Site 798, B:391
- elliptica*, *Mesocena*, Site 799, B:248
- Elphidium* cf. *perforatum*, Site 795, B:540
- Elphidium clavatum*, Site 798, B:392
- Elphidium jenseni*, Site 797, B:540
- Elphidium subarcticum*
Site 797, B:540
Site 798, B:392
- Emiliania huxleyi*
Site 795, A(127):194
Site 798, B:164
- Engelhardia*
Site 796, B:490
Site 797, B:490
- Ephedra*, Site 795, B:490
- Epistominella*, Leg 128, B:62
- Epistominella pulchella*, Site 798, B:391
- Euchitonina furcata*, Site 798, B:235
- Euchitonina* sp., Site 794, B:306
- Eucyrtidium asanoi*, Site 799, B:236
- Eurya*, Site 796, B:491
- Evolvocassidulina kuwanoi*, Site 797, B:539
- ezoensis*, *Cyclammina*, Site 795; B:538
- Fagus*
Japan Sea, B:484-486
Site 795, B:491
Site 797, B:491
- fimbriata*, *Chilostomellina*, Site 798, B:392
- Fissurina biconica*, Site 798, B:389
- Fissurina echigoensis*, Site 798, B:389
- Fissurina orbignyana*, Site 797, B:539
- floriadanus*, *Cyclicargolithus*
Site 795, A(127):195
Site 797, B:179
- Funpingopollenites*, Site 797, B:491
- furcata*, *Euchitonina*, Site 798, B:235
- fusiformis*, *Stainforthia*
paleoenvironment and, B:379
Site 798, B:390
- Gephyrocapsa*, Site 798, A(128):162
- Gephyrocapsa aperta*, Leg 128, B:164
- Gephyrocapsa caribbeanica*
Japan Sea, B:173, 183
Leg 128, B:164
- Gephyrocapsa caribbeanica* Zone, Japan Sea, B:174
- Gephyrocapsa oceanica*
Japan Sea, B:173, 183
Leg 128, B:164
Site 798, B:167-168
- Gephyrocapsa* small, Japan Sea, B:183
- Ginkgo*, Site 795, B:490
- Glabrata subopercularis*, Site 797, B:540
- glacialis*, *Spongotrochus*, Site 795, B:306
- Globigerina*, Site 798, B:569
- Globigerina bulloides*
Leg 127, B:193, 200
Leg 128, B:60-61
Site 797, B:591-594
Site 798, B:441-443, 457, 460, 462-463, 465-466, 469, 564
- Globigerina bulloides* (s.l.), Site 798, A(128):165
- Globigerina bulloides umbilicata*, Oki Ridge, B:444
- Globigerina incompta*, Site 798, B:463
- Globigerina quinqueloba*
Leg 127, B:200
Site 798, B:457, 460, 469
- globigeriniformis*, *Trochammina* cf., Site 795, B:537
- Globigerinita* spp., Site 798, B:457, 460, 469
- Globigerinoides sicanus/Praeorbulina glomerata* Zone, Japan Sea, B:526
- Globobulimina auriculata*, Site 798, B:390
- Globobulimina pacifica*, Site 798, B:390
- Globobulimina perversa*, Site 797, B:539
- Globobulimina pupoides*, Site 797, B:539
- Globobulimina subaffinis*, Site 798, B:390
- Globocassidulina globosa* s.l., Site 797, B:539
- Globocassidulina* sp. A, Site 797, B:539
- Globoquadrina venezuelana*
Leg 127, B:189
Site 797, B:1220
- Globorotalia* cf. *G. fohsi* Zone, Leg 127, B:200
- Globorotalia inflata*, Site 798, B:457, 469
- Globorotalia peripheroronda/Globorotalia quinifalcata* Zone, Japan Sea, B:526
- Globorotalia praescitula*, Site 797, B:189, 1220, 1223
- globosa*, *Nonionella*, Site 798, B:391
- globosa*, *Oolina*
Japan Sea, B:387
Site 797, B:539
- globosa* s.l., *Globocassidulina globosa*, Site 797, B:539
- grande*, *Lychnocanium* sp. cf.
Site 794, A(127):101
Site 797, B:305
- grateloupi*, *Pseudononion*, Site 798, B:392
- Gyroidina orbicularis*, Site 797, B:540
- Gyroidina orbicularis/Gyroidinoides planulatus* Zone
age models and, B:531
Site 797, B:521, 528
- Gyroidinoides planulatus*, Site 797, B:540
- hanzawai*, *Tosaia*, Site 797, B:538

- Hanzawaia tagaensis/Heterolepa praecincta* Zone, Japan Sea, B:526
- Haplophragmoides* sp., Site 795, B:537
- hataii*, *Echigoina*, Site 798, B:392
- helenae*, *Islandiella*, Site 797, B:539
- Helicosphaera ampliapertura* Zone, Site 797, B:1220, 1223
- Helicosphaera perch-nielseniae*, Japan Sea, B:183
- Helicosphaera sellii*
Leg 128, B:164
Site 798, B:164
- heteromorphus*, *Sphenolithus*
Japan Sea, B:173-174, 183
Leg 127, B:189
- hexagona*, *Oolina*, Site 797, B:539
- Hippocrepinella variabilis*, Site 795, B:537
- Hopkinsina morimachiensis* Zone, Japan Sea, B:526
- hustedtii*, *Denticulopsis*, Japan Sea, B:250
- huxleyi*, *Emiliania*
Site 795, A(127):194
Site 798, B:164
- incompta*, *Globigerina*, Site 798, B:463
- inflata*, *Globorotalia*, Site 798, B:457, 469
- ingens*, *Actinocyclus*, Pacific Ocean, B:309
- Islandiella* cf. *compressa*, Site 797, B:539
- Islandiella helenae*, Site 797, B:539
- Islandiella* sp., Site 797, B:539
- jacksonii*, *Thalassiosira*, Japan Sea, B:259
- japonica*, *Cassidulina*, Site 798, B:389
- japonica*, *Sethocyrtis*
Japan Sea, B:297, 301
Site 794, B:305
- japonica*, *Sphaeroidina*, Site 797, B:538
- japonica*, *Thecosphaera*, Site 795, B:305
- jenseni*, *Elphidium*, Site 797, B:540
- jimlingii*, *Distephanus*
Japan Sea, B:238
Site 798, B:248
- jouseae*, *Nitzschia*, B:1219
- Juglans*
Site 794, B:490
Site 795, B:490
- kamtschatica*, *Denticulopsis*, Site 795, B:1223
- kamtschatica*, *Neodenticula*
North Pacific, B:309
Site 795, B:970
Site 798, B:361
Site 799, A(128):302
- Karreriella baccata japonica*, Site 798, B:389
- koizumii*, *Neodenticula*
Japan Sea, B:258
Site 798, B:361
- koizumii*, *Neodenticulopsis*, Site 798, B:1224
- kokazuraensis*, *Angulogerina*, Site 798, B:390
- kuromatsunaiensis*, *Buccella* cf., Site 797, B:540
- kuwanoi*, *Evolvocassidulina*, Site 797, B:539
- labradorica*, *Nonionellina*
Site 797, B:539
Site 798, B:391
- lacunosa*, *Pseudoemiliania*
Japan Sea, B:183
Leg 127, B:581
Leg 128, B:163
Site 798, B:468
Site 799, B:166
- laevis*, *Lagena*, Site 798, B:389
- Lagena laevis*, Site 798, B:389
- Lagena substrata*, Japan Sea, B:387
- Lagena tricaritortuosa*, Site 798, B:389
- langii*, *Sphaeropyle*
Japan Sea, B:297
Site 794, B:305
Site 795, B:303
Site 799, B:236
- Larnacantha polyacantha*, Site 795, B:306
- lentiginosa*, *Thalassiosira*, Southern Ocean, B:309
- lepidula*, *Stilostomella*, Site 798, B:390
- leptoporus*, *Calcidiscus*, Site 798, B:168
- Lipmanella* sp. aff. *L. redondoensis*, Japan Sea, B:297
- Lipmanella* sp. aff. *Theocorys redondoensis*
Japan Sea, B:297
Site 794, B:305
- Liquidambar*
Japan Sea, B:484-486
Site 797, B:491
- Lithartractus tochigiensis*, Site 794, B:305
- Lithopera renzae renzae*, Site 799, B:236
- lobatula*, *Uvigerina*, Site 798, B:390
- Lychnocanium* cf. *L. grande*, Site 794, B:1221
- Lychnocanium nipponicum*
Site 794, A(127):101
Site 797, B:305
- Lychnocanium* sp. cf. *L. grande*
Site 794, A(127):101
Site 797, B:305
- Lychnocanoma* sp.
Japan Sea, B:230
Site 799, B:225-226, 236
- macilentus*, *Ammodiscus*, Site 795, B:537
- macintyreii*, *Calcidiscus*
Japan Sea, B:174, 182
Site 798, B:164
- makiyama*, *Sagarites*, Site 799, B:35, 45
- Makiyama titanni*, Site 799, B:35, 45
- mammifera*, *Didymocyrtis*, Site 799, B:235
- mammilaris*, *Tholospyris*, Site 794, B:305
- marginatus*, *Coscinodiscus*
Aikawa section, Honshu, B:316
Japan Sea, B:291
Leg 127, B:1219-1220
Leg 128, B:309-314, 1219-1220
North Pacific, B:315-316
Site 433, B:316
Site 581, B:316
Site 794, B:341-357
Site 795, B:341-357
Site 797, B:341-357
Site 798, B:361, 554, 1224
Site 799, B:1224
- Marginulina* sp., Site 797, B:539
- Martinottiella communis*
Japan Sea, B:1201
Site 795, B:537
Site 797, B:537
Site 798, B:389
- Martinottiella communis/Cribrostomoides* cf. *crassimargo* Zone, Site 795, B:521
- Martinottiella communis/Globobulimina* *pu-poides* Zone, Site 797, B:524
- Martinottiella communis* Zone, Site 795, B:517
- Martinottiella nodulosa*, Site 798, B:389
- marujamica*, *Thalassiosira*, Japan Sea, B:259
- masudai*, *Valvulineria*, Site 797, B:538
- Melonis nikobarense*
Site 797, B:540
Site 798, B:392
- Melonis pompilioides*
Site 797, B:540
Site 799, B:383, 392
- Melosira*, Site 798, B:554
- Mesocena elliptica*, Site 799, B:248
- Mesocena quadrangula*, Site 798, B:248
- Metasequoia*, Site 796, B:490
- Miliammina echigoensis*
Leg 127, B:200
Site 798, B:389
Site 799, A(128):311
- Miliammina echigoensis* Zone, Leg 127, B:201
- miocenica*, *Nonionella*, Site 797, B:539
- miocenica*, *Thalassiosira*, B:1219
- miocenica*, *Uvigerina*, Site 797, B:539
- Miogypsina-Operculina* Zone, Japan Sea, B:526
- modeloensis*, *Uvigerina* cf., Site 797, B:539
- murrhina*, *Pyrgo*
Site 797, B:538
Site 798, B:389
- naraensis*, *Pseudoparrella*, Site 798, B:391
- Neodenticula kamtschatica*
North Pacific, B:309
Site 795, B:970
Site 798, B:361
Site 799, A(128):302
- Neodenticula kamtschatica* Zone
Leg 127, B:188
Site 796, B:1223
- Neodenticula koizumii*
Japan Sea, B:258
Site 798, B:361
- Neodenticula koizumii* Zone
Leg 127, B:188
Site 795, B:344
Site 796, B:1223
Site 797, B:345
- Neodenticula koizumii-Neodenticula kamtschatica* Zone
Site 794, B:344
Site 795, B:344-345
Site 797, B:345
- Neodenticulopsis koizumii*, Site 798, B:1224
- Neogloboquadrina asanoi*, Site 797, B:208
- Neogloboquadrina pachyderma*
age models and, B:362
Japan Sea, A(128):30
Leg 127, B:188, 193, 200, 218-219, 1220-1221
Leg 128, B:1220-1221
Oki Ridge, B:444
Site 795, A(127):195
Site 797, B:208
Site 798, A(128):165-166; B:457, 459-469
Site 799, A(128):310-311
- Neogloboquadrina pachyderma dextral/Globorotalia orientalis* Zone, Oga Peninsula, B:459
- Neogloboquadrina pachyderma sinistral/Globigerina incompta* Zone, Oga Peninsula, B:459
- Neogloboquadrina pachyderma sinistral/Globigerina quinqueloba* Zone, Oga Peninsula, B:459
- neopseudofibula*, *Dictyochoa*
Japan Sea, B:237-238
Site 798, B:248
- Nephrospyris?* *pervia*
Japan Sea, B:230-231
Site 798, B:235
- nidulus*, *Thalassiosira*, Japan Sea, B:258
- nikobarense*, *Melonis*
Site 797, B:540
Site 798, B:392
- nipponica*, *Plectina*, Site 797, B:537

- nipponicum*, *Lychnocanium*
Site 794, A(127):101
Site 797, B:305
- Nitzschia jouseae*, B:1219
- Nitzschia pliocena*, Japan Sea, B:259
- Nitzschia reinholdii*
Hess Rise, B:360
Site 798, B:361
- Nitzschia rolandii*, Japan Sea, B:259
- nitzschioides*, *Thalassionema*, Site 794, B:1223
- nitzschioides*, *Thalassiothrix*, Site 798,
A(128):160
- nodulosa*, *Martinottiella*, Site 798, B:389
- Nonionella globosa*, Site 798, B:391
- Nonionella miocenica*, Site 797, B:539
- Nonionella stella*
Site 797, B:539
Site 799, B:391
- Nonionellina labradorica*
Site 797, B:539
Site 798, B:391
- norcrossi*, *Cassidulina*, Site 798, B:390
- norvangi*, *Cassidulina*, Site 798, B:390
- oceanica*, *Gephyrocapsa*
Japan Sea, B:173, 183
Leg 128, B:164
Site 798, B:167–168
- octacantha*, *Tetrapyle*
Site 794, B:306
Site 797, B:306
Site 798, B:236
- octangulatus*, *Distephanus*, Site 798, B:248
- oculatus*, *Actinocyclus*
Japan Sea, B:258
Site 798, B:361
- oolina*, *Chilostomella*
Site 797, B:539
Site 799, B:392
- Oolina globosa*
Japan Sea, B:387
Site 797, B:539
- Oolina hexagona*, Site 797, B:539
- orbicularis*, *Cyclammina*, Site 795, B:538
- orbicularis*, *Gyroidina*, Site 797, B:540
- orbignyana*, *Fissurina*, Site 797, B:539
- Oridorsalis tener*, Site 798, B:392
- Oridorsalis umbonatus*
Site 797, B:540
Site 798, B:392
- Oridorsalis umbonatus/Cibicoides wuellerstorfi*
Zone, Site 795, B:517, 528
- ovoidea*, *Chilostomella*, Site 798, B:392
- pachyderma*, *Neogloboquadrina*
age models and, B:362
Japan Sea, A(128):30
Leg 127, B:188, 193, 200, 218–219,
1220–1221
Leg 128, B:1220–1221
Oki Ridge, B:444
Site 795, A(127):195
Site 797, B:208
Site 798, A(128):165–166; B:457, 459–469
Site 799, A(128):310–311
- pacifica*, *Bolivina*
paleoenvironment and, B:379
Site 798, A(128):166; B:373, 389
- pacifica*, *Circumtexta*, Site 795, B:304
- pacifica*, *Globobulimina*, Site 798, B:390
- Parafissurina* sp., Site 797, B:539
- Paramesocena circulus*, Site 798, B:248
- parkeriae*, *Tenuitella*, Leg 127, B:200
- parkeriae*, *Thalmanammina* cf., Site 797, B:537
- parvulus*, *Catapsydrax*, Site 797, B:189, 1220,
1223
- Pasania*
Site 794, B:490
Site 796, B:490
- pelagicus*, *Coccolithus*
Site 798, B:167–168
Site 799, B:166
- perch-nielseniae*, *Helicosphaera*, Japan Sea,
B:183
- peregrina*, *Stichocorys*
Japan Sea, B:297
Site 794, A(127):102
Site 797, A(127):356
Site 799, B:236
- perforatum*, *Elphidium* cf., Site 795, B:540
- Peripyraxis circumtexta*, Site 795, B:304
- Peripyraxis* sp.
Japan Sea, B:301
Site 795, B:306
- perplexus*, *Dictyococcites*, Japan Sea, B:182
- perversa*, *Globobulimina*, Site 797, B:539
- pervia*, *Nephrospyris*?
Japan Sea, B:230–231
Site 798, B:235
- Picea*
Japan Sea, B:328
Neogene climate and, B:337
- Pinus*
Japan Sea, B:319, 328
Neogene climate and, B:337
- Planolites*, Site 798, B:566–567, 569
- Planularia asanoi*, Site 798, B:389
- planulatus*, *Gyroidinoides*, Site 797, B:540
- Plectina nipponica*, Site 797, B:537
- Pleurostomella* sp. A, Site 799, B:390
- pliocena*, *Nitzschia*, Japan Sea, B:259
- polyacantha*, *Larnacantha*, Site 795, B:306
- pompilioides*, *Melonis*
Site 797, B:540
Site 799, B:383, 392
- praebergonii*, *Rhizosolenia*, B:1219
Indian Ocean, B:309
- praedimorpha*, *Denticulopsis*, Japan Sea, B:250
- praescitula*, *Globorotalia*, Site 797, B:189, 1220,
1223
- proboscidea*, *Uvigerina*, Site 797, B:539
- Proteonina compressa*, Site 795, B:537
- Pseudoemiliana lacunosa*
Japan Sea, B:183
Leg 127, B:581
Leg 128, B:163
Site 798, B:468
Site 799, B:166
- Pseudoemotia doliolus*
age models and, B:362
Hess Rise, B:360
Honsu coast, B:360
- Pseudononion grateloupi*, Site 798, B:392
- Pseudoparrella naraensis*, Site 798, B:391
- Pseudoparrella takayanagii*, Site 798, B:391
- pseudoubilica*, *Reticulofenestra*, Japan Sea,
B:183
- Pterocarya*, Site 794, B:490
- pulchella*, *Epistominella*, Site 798, B:391
- Pullenia apertura*, Site 798, B:392
- Pullenia bulloides*, Site 797, B:540
- Pullenia salisburyi*
Site 797, B:540
Site 798, B:392
- Pullenia subcarinata*, Site 797, B:540
- pupoides*, *Globobulimina*, Site 797, B:539
- pusilla*, *Cyclammina*, Site 797, B:538
- Pyrgo murrhina*
Site 797, B:538
Site 798, B:389
- Pyrgo* sp., Site 797, B:538
- quadrangula*, *Mesocena*, Site 798, B:248
- quadrilatera*, *Bolivinita*, Site 799, B:389
- Quercus*
Japan Sea, B:328
Neogene climate and, B:337
- Quercus*, D., Site 794, B:491
- Quercus*, E.
Japan Sea, B:484–486
Site 794, B:491
Site 796, B:491
- quinqueloba*, *Globigerina*
Leg 127, B:200
Site 798, B:457, 460, 469
- Quinqueloculina* sp.
Site 797, B:538
Site 799, B:373
- rectangulare*, *Ammodochium*
Japan Sea, B:238
Site 798, B:248
- redondoensis*, *Lipmanella* sp. aff., Japan Sea,
B:297
- redondoensis*, *Theocorys*
Japan Sea, B:297
Site 794, B:305
- reinholdii*, *Nitzschia*
Hess Rise, B:360
Site 798, B:361
- renzae renzae*, *Lithopera*, Site 799, B:236
- Reophax* sp.
Site 795, B:537
Site 797, B:537
- Reticulofenestra asanoi*, Site 798, B:164
- Reticulofenestra pseudoubilica*, Japan Sea,
B:183
- Reticulofenestra* small
Japan Sea, B:183
Site 797, A(127):352
- Rhizammina* sp., Site 797, B:537
- Rhizosolenia curvirostris*
Hess Rise, B:360
Japan Sea, B:258
Leg 127, B:581
North Pacific, B:311
Site 798, B:361, 468
- Rhizosolenia praebergonii*, B:1219
Indian Ocean, B:309
- Rhus*, Site 794, B:491
- robusta*, *Cycladophora*
Japan Sea, B:299
Site 794, B:307
- robusta*, *Sphaeropyle*
Japan Sea, B:297
Site 794, B:1221
Site 797, B:305
- rolandii*, *Nitzschia*, Japan Sea, B:259
- Rossiella tatsunokuchiensis*, Japan Sea, B:258–
259
- rostrata*, *Bulimina*, Site 795, B:539
- rotunda*, *Eilohedra*, Site 798, B:391
- Rouxia californica*, Japan Sea, B:259
- Rouxia californica* Zone, Leg 127, B:189
- Saccammina sphaerica*, Site 795, B:537
- sadonica*, *Valvulineria*, Site 798, B:390
- Sagarites makiyama*, Site 799, B:35, 45
- salisburyi*, *Pullenia*

- Site 797, B:540
Site 798, B:392
Schizammia? sp.
Site 795, B:537
Site 797, B:537
Sciadopitys, Site 795, B:490
Selaginella selaginoides, Site 794, B:334
selaginoides, *Selaginella*, Site 794, B:334
sellii, *Helicosphaera*
Leg 128, B:164
Site 798, B:164
Sethocyrtis japonica
Japan Sea, B:297, 301
Site 794, B:305
Silicosigmoilina abyssalica, Japan Sea, B:505
slavnicii, *Distephanus*, Site 798, B:248
sp. A, *Cassidulina*, Site 798, B:390
sp. A, *Cibicidoides*, Site 797, B:538
sp. A, *Cyclammia*, Site 797, B:538
sp. A, *Globocassidulina*, Site 797, B:539
sp. A, *Pleurostomella*, Site 799, B:390
speculum, *Distephanus*, Site 799, B:248
sphaerica, *Saccammia*, Site 795, B:537
Sphaeroidina bulloides, Site 798, B:390
Sphaeroidina japonica, Site 797, B:538
Sphaeropyle langii
Japan Sea, B:297
Site 794, B:305
Site 795, B:303
Site 799, B:236
Sphaeropyle robusta
Japan Sea, B:297
Site 794, B:1221
Site 797, B:305
Sphenolithus compactus, Japan Sea, B:183
Sphenolithus heteromorphus
Japan Sea, B:173–174, 183
Leg 127, B:189
Sphenolithus heteromorphus Zone
Japan Sea, B:174
Leg 127, B:201
Site 797, B:1223
Sphenolithus umbrellus, Japan Sea, B:183
Spirosigmoilinella compressa
Site 795, B:537
Site 798, B:379–381, 389
Spirosigmoilinella compressa/Martinottiella communis Zone
Site 794, B:517, 528
Site 798, B:381
Spirosigmoilinella compressa Zone, Leg 127, B:200
spissa, *Bolivina*, Site 798, B:389
Spongaster tetras irregularis
Japan Sea, B:231
Site 798, B:236
Spongastrochus glacialis, Site 795, B:306
Stainforthia complanata
Japan Sea, B:388
Site 798, B:390
Stainforthia fusiformis
paleoenvironment and, B:379
Site 798, B:390
stella, *Nonionella*
Site 797, B:539
Site 799, B:391
Stichocorys delmontensis
Japan Sea, B:228, 230
Site 799, B:225–226, 236
Stichocorys peregrina
Japan Sea, B:297
Site 794, A(127):102
Site 797, A(127):356
Site 799, B:236
Stichocorys wolfii, Site 794, A(127):102
Stilostomella lepidula, Site 798, B:390
striata, *Bulimina*
Site 797, B:539
Site 798, B:390
Styloacontarium acquilonius
Japan Sea, B:297
Site 794, B:305
Styloacontarium sp. cf. *S. acquilonius*, Japan Sea, B:297
Stylochlamyidium venustum, Site 797, B:306
subaffinis, *Globobulimina*, Site 798, B:390
subarcticum, *Elphidium*
Site 797, B:540
Site 798, B:392
subarctios, *Dictyochoa*
Japan Sea, B:238–239
Site 798, B:248
subcarinata, *Pullenia*, Site 797, B:540
subopercularis, *Glabratella*, Site 797, B:540
substriata, *Lagena*, Japan Sea, B:387

takayanagii, *Pseudoparrella*, Site 798, B:391
tatsunokuchiensis, *Rossiella*, Japan Sea, B:258–259
temperei, *Thalassiosira*, Japan Sea, B:259
tener, *Oridorsalis*, Site 798, B:392
tenuata, *Buliminella*, Site 798, B:390
tenuis, *Ammodiscus*, Site 797, B:537
tenuis, *Cassidulinoides*, Site 799, B:390
Tenuitella parkerae, Leg 127, B:200
tetrapera, *Cyrtocapsella*
Japan Sea, B:230
Site 794, B:304
Site 797, A(127):356; B:304
Site 799, B:235
Tetrapyle octacantha
Site 794, B:306
Site 797, B:306
Site 798, B:236
tetras irregularis, *Spongaster*
Japan Sea, B:231
Site 798, B:236
tetrathalamus, *Didymocyrtis*
Site 797, B:306
Site 798, B:235
Thalassinoides, Site 798, B:566–567, 569
Thalassionema nitzschioides, Site 794, B:1223
Thalassionema schraderii Zone, Site 794, B:1221
Thalassiosira antiqua, Japan Sea, B:258
Thalassiosira convexa, Japan Sea, B:258
Thalassiosira jacksonii, Japan Sea, B:259
Thalassiosira lentiginosa, Southern Ocean, B:309
Thalassiosira marujamica, Japan Sea, B:259
Thalassiosira miocenica, B:1219
Thalassiosira nidulus, Japan Sea, B:258
Thalassiosira oestrupii Zone
Leg 127, B:188
Site 794, B:344, 1221
Site 797, B:1223
Thalassiosira temperei, Japan Sea, B:259
Thalassiosira zabelinae, Japan Sea, B:259
Thalassiothrix nitzschioides, Site 798, A(128):160
Thalmammammia cf. *parkerae*, Site 797, B:537
Thecosphaera japonica, Site 795, B:305
Thecalypra davisiana
Japan Sea, B:297, 299
Site 794, B:306
Site 795, B:303
Theocorys redondoensis
Japan Sea, B:297
Site 794, B:305
Tholospyris mammilaris, Site 794, B:305
Tilia, Site 797, B:491
titanni, *Makiyama*, Site 799, B:35, 45
tochiogensis, *Lithatractus*, Site 794, B:305
Tosaia hanzawai, Site 797, B:538
tricarinata, *Triloculina*, Site 798, B:389
tricaritortuosa, *Lagena*, Site 798, B:389
Trifarina sp., Site 797, B:539
Triloculina tricarinata, Site 798, B:389
Trochammia cf. *globigeriniformis*, Site 795, B:537
Tsuga
Japan Sea, B:328
Neogene climate and, B:337

Ulmus, Site 795, B:491
Umbilicosphaera sibogae Zone, Japan Sea, B:174
unbonatus, *Oridorsalis*
Site 797, B:540
Site 798, B:392
umbrellus, *Sphenolithus*, Japan Sea, B:183
Uvigerina akitaensis, Site 798, B:390, 441, 443–444
Uvigerina cf. *modeloensis*, Site 797, B:539
Uvigerina lobatula, Site 798, B:390
Uvigerina miocenica, Site 797, B:539
Uvigerina proboscidea, Site 797, B:539
Uvigerina segundoensis s.l. Zone, Japan Sea B:526
Uvigerina yabei, Site 798, B:390

Valvulineria masudai, Site 797, B:538
Valvulineria sadonica, Site 798, B:390
variabilis, *Hippocrepinella*, Site 795, B:537
venezuelana, *Globoquadra*
Leg 127, B:189
Site 797, B:1220
venustum, *Stylochlamyidium*, Site 797, B:306

wolfii, *Stichocorys*, Site 794, A(127):102
wuellerstorfi, *Cibicidoides*
Japan Sea, B:505
Site 797, B:538

yabei, *Uvigerina*, Site 798, B:390
ypsilon, *Amphirhopalum*, Site 798, B:235

zabelinae, *Thalassiosira*, Japan Sea, B:259
Zelkooa, Site 797, B:491
zones (with letter prefixes)
NP-1, Site 797, B:1221, 1223
NP-2, Site 797, B:1221, 1223
NP-3, Site 797, B:1223
Zoophycos, Japan Sea, B:1232