

## INDEX TO VOLUMES 127 AND 128

This index provides coverage for the *Initial Reports* portions of Volumes 127 and 128 of the *Proceedings of the Ocean Drilling Program* (published as separate leg-specific books) and for the *Scientific Results* portion of those *Proceedings* (published as Volume 127/128, Parts 1 and 2). The reports in Parts 1 and 2 of the *Scientific Results* are not leg-specific but are arranged by subject matter. References to page numbers in the *Initial Reports* are preceded by "A," followed by the appropriate leg reference in parentheses, and then a colon: "A(127):"; those in the *Scientific Results* are preceded by "B" with a colon.

The index was prepared by Wm. J. Richardson Associates, Inc., under subcontract to the Ocean Drilling Program. The index contains two hierarchies of entries: (1) a main entry, defined as a keyword or concept followed by a reference to the page on which that word or concept appears, and (2) a subentry, defined as a further elaboration on the main entry followed by a page reference.

The index is presented in two parts: (1) a Subject Index and (2) a Taxonomic Index. Both parts cover text figures and tables but not core-description forms ("barrel sheets") or core photographs. Also excluded are bibliographic references, names of individuals, and routine front and back matter.

The Subject Index follows a standard format. Geographic, geologic, and other terms are referenced only if they are subjects of discussion. This index also includes taxonomic entries above the generic level, as well as broad fossil groups such as foraminifers and radiolarians. The notation "ff" following a page listing indicates that reference to a topic continues beyond the last page given but is not sequential. This would be the case where one or more figures or tables follow a principal topic of discussion that makes up a major section. A site chapter in the *Initial Reports* is considered the principal reference for that site and is indicated on the first line of the site's listing in the index. A parenthetical reference is made to the appropriate leg. Such a reference to Site 798, for example, is given as "Site 798, A(128):121–236."

The Taxonomic Index is an index relating to significant findings and/or substantive discussions, not of species names *per se*. This index covers three varieties of information: (1) individual genera and species that have been erected or emended formally, (2) biostratigraphic zones, and (3) fossils depicted in illustrations. A taxonomic entry consisting of both genus and species is listed alphabetically by genus and also by species. Biostratigraphic zones are listed alphabetically by genus; zones with letter prefixes are listed under "zones."

For further information, including available electronic formats, contact the Chief Production Editor, Ocean Drilling Program, 1000 Discovery Drive, College Station, Texas 77845-9547, U.S.A.

## SUBJECT INDEX

AACT. *See* aluminum, logging; geochemical logging  
 acanthostyles  
     Site 795, B:542  
 acanthoxeas  
     Site 795, B:542  
 acanthstrongyles  
     Site 795, B:542  
 accelerometer, three-axis feedback-type. *See* downhole seismic experiment  
 acoustic impedance  
     opal-A to opal-CT transition, B:1146  
     Site 799, B:1137–1138, 1140  
 acoustic velocity. *See* velocity  
 ACT. *See* aluminum, logging  
 AF demagnetization. *See* magnetic properties  
 age dating. *See* biostratigraphic zonations; biostratigraphy; radiometric age  
 age spectra  
     Site 794, B:820, 828–832  
     Site 795, B:820–821, 832–833  
     Site 797, B:820, 828–829  
 Aikawa section, diatom abundance, B:316  
 air-gun profiling experiment, B:1109–1114, 1117–1118  
 Aira-Tn, tephra marker, B:796  
 Akita Basin, lithostratigraphy, A(128):17  
 Akita oil field, benthic foraminifers, B:381  
 Akita-Yamagata oil field basalt  
     Site 794 basalt compared, B:918, 1339  
     Site 797 basalt compared, B:921  
 albitization. *See also* feldspar  
     and calcium concentration, B:137  
     occurrence modes, B:136  
     provenance information destroyed, B:148  
     of sandstone, B:136–137  
     silica source, B:148  
     and sodium in interstitial water, B:137  
     and sulfate formation, B:143–144  
     temperature, B:136  
 alginite  
     Site 798, B:429  
     Sites 798–799, B:670  
 alkaline feldspar. *See* feldspar, alkaline  
 alkaline glass, B:1378, 1387  
 alkalinity  
     and carbonate precipitation, A(127):204  
     and diagenetic carbonate, B:94  
     and dolomitization, A(127):362  
     in interstitial water, B:1264  
     at Japan Sea sites, B:1265  
     and organic carbon decomposition, B:1262  
     and organic matter degradation, A(127):362  
     Site 794, A(127):108  
     Site 795, A(127):204  
     Site 796, A(127):279  
     Site 797, A(127):362, 368  
     Site 798, A(128):173, 181  
     Site 799, A(128):317–318, 328  
     and clay minerals, A(127):279–280  
     in interstitial water, B:1264  
     and organic carbon decomposition, B:1262  
     and organic matter degradation, A(127):279–280  
     Site 796, A(127):279–280  
     Site 798, A(128):173, 181  
     Site 799, A(128):317–318, 328  
     amorphous silica. *See* opal-A; silica, biogenic  
     amphibole, in ash layers, B:1379  
     Amur River, and Japan Sea, B:69–70, 442  
     Amurian microplate, A(128):9  
     anaerobic bacterial degradation. *See* methanogenesis; sulfate reduction  
     analcime, Site 797, B:139, 143  
     andesite  
         ash layers, B:793  
         microphotograph, B:859  
         rare earth elements, B:924  
         Site 795, B:850, 854, 1338  
         spidergrams, B:924  
         spinel composition, B:844  
     andesite, basaltic  
         brecciated, A(127):221–222  
         plagioclase pyroxene phric, A(127):174, 217  
         rare earth elements, B:924  
         Site 795, B:918, 1338  
         spidergrams, B:924  
     anhydrite  
         replacing plagioclase, B:148  
         sandstone replacement mineral, B:143  
     Aniai-type flora, Japan Sea, B:249, 486  
     anisochelas  
         Site 795, B:543  
     anisotropy  
         in oceanic crust, B:1110–1115  
         in velocity measurements, A(127):229  
     ankerite  
         and Mossbauer parameters, B:741  
         Site 796, A(127):266  
         Site 799, B:82  
     anomalies. *See* free-air gravity anomalies; magnetic anomalies  
     anorthite  
         Sites 794 and 796, B:126  
         vs. orthoclase, B:122  
     anoxic conditions. *See* oxygenation conditions  
     anoxic-nonsulfidic sediments, Site 799, B:612  
     antimony  
         Site 795, B:711, 714  
         Site 798, B:1370–1371  
         Site 799, B:724–725  
     aperture, vertical, in logs, B:396–397  
     Ar/Ar age. *See* radiometric age  
     arc tholeiitic suite, Site 795, B:1338  
     arenite. *See* feldspathic arenite; lithic arenite; sandstone  
     argon/argon age. *See* radiometric age  
     argon isotopes. *See* isotopes, argon  
     aridity signal, in logs, A(128):188–192  
     arkose. *See* sandstone  
     aromatic species, Site 799, B:627  
     arsenic  
         Site 794, B:1362  
         Site 795, B:711, 714, 1365  
         Site 797, B:1366

augite clinopyroxene, B:885  
 basalt, B:838, 884–888, 899, 901  
 basement, A(127):122–125, A(128):88–91, 321; B:849, 883–889, 891, 908–916, 1265–1267, 1282, 1338  
 basement rock and adjacent pore water elemental variations correlated, B:911  
 and element composition, B:799  
 and element mobility, B:838, 892, 908–916  
 glass shards, A(127):95  
 groundmass, B:885–886  
 hydroclastic shards, A(128):89–90  
 and liquid line of descent calculations, B:911  
 and magnesium enrichment, B:842  
 olivine, B:885  
 plagioclase feldspar, B:884–885  
 and rare earth element mobility, B:911–916  
 secondary minerals, B:838  
 Site 794, B:908–911, 916  
 Site 795, B:918  
 Site 797, B:911  
 temperature, B:839  
 timing, B:838–839  
 and XRF loss on ignition, B:892–893, 909  
 alteration halo, around fractures, A(128):91  
 alternating field demagnetization. *See* magnetic properties  
 Altyn-Tagh Fault, offset along, B:1325  
 aluminum  
     in ash layers, B:1380  
     in augite, B:851–853, 856  
     in dark/light cycles, B:569  
     in Japan Sea sediment, B:1236–1237  
     and parent magma, B:856  
     Site 794, B:654, 656  
     Site 795, B:706  
     Site 796, B:654, 656  
     Site 798, B:654, 656, 722  
     Site 799, B:654, 656, 722  
     vs. chromium, B:733  
     vs. iron, B:732  
     vs. iron/magnesium ratio, B:894  
     vs. lanthanum, B:737  
     vs. ytterbium, B:737  
     XRF and geochemical log analyses compared, B:1028–1029, 1031–1035  
 aluminum, logging  
     clay, A(127):306–307  
     core data correlated, B:1030  
     diatom ooze, A(127):306–307  
     diatom vs. clay content, B:1417  
     opal-A to opal-CT transition, B:22  
     sand, A(127):306  
     Site 794, B:1416, 1420–1421  
     Site 796, B:1416, 1423  
     Site 797, B:7, 1417, 1426–1427  
     Site 798, A(128):212, 215; B:1030, 1401–1402  
     Site 799, A(128):362; B:1404, 1408–1409  
         vs. potassium, A(128):215  
 aluminum activation clay tool (AACT). *See* aluminum, logging; geochemical logging  
 aluminum clay tool (ACT). *See* aluminum, logging  
 aluminum oxide. *See* aluminum  
 aluminum/silica ratio, vs. terrigenous content, B:1035  
 ammonia  
     in clay minerals, A(127):204, 362  
     and organic matter degradation, A(127):204, 362  
     Site 794, A(127):108

Site 795, A(127):204  
 Site 797, A(127):362, 368  
 Site 799, B:610  
 ammonium  
     in clay minerals, A(127):279–280  
     in interstitial water, B:1264  
     and organic carbon decomposition, B:1262  
     and organic matter degradation, A(127):279–280  
     Site 796, A(127):279–280  
     Site 798, A(128):173, 181  
     Site 799, A(128):317–318, 328  
     amorphous silica. *See* opal-A; silica, biogenic  
     amphibole, in ash layers, B:1379  
     Amur River, and Japan Sea, B:69–70, 442  
     Amurian microplate, A(128):9  
     anaerobic bacterial degradation. *See* methanogenesis; sulfate reduction  
     analcime, Site 797, B:139, 143  
     andesite  
         ash layers, B:793  
         microphotograph, B:859  
         rare earth elements, B:924  
         Site 795, B:850, 854, 1338  
         spidergrams, B:924  
         spinel composition, B:844  
     andesite, basaltic  
         brecciated, A(127):221–222  
         plagioclase pyroxene phric, A(127):174, 217  
         rare earth elements, B:924  
         Site 795, B:918, 1338  
         spidergrams, B:924  
     anhydrite  
         replacing plagioclase, B:148  
         sandstone replacement mineral, B:143  
     Aniai-type flora, Japan Sea, B:249, 486  
     anisochelas  
         Site 795, B:543  
     anisotropy  
         in oceanic crust, B:1110–1115  
         in velocity measurements, A(127):229  
     ankerite  
         and Mossbauer parameters, B:741  
         Site 796, A(127):266  
         Site 799, B:82  
     anomalies. *See* free-air gravity anomalies; magnetic anomalies  
     anorthite  
         Sites 794 and 796, B:126  
         vs. orthoclase, B:122  
     anoxic conditions. *See* oxygenation conditions  
     anoxic-nonsulfidic sediments, Site 799, B:612  
     antimony  
         Site 795, B:711, 714  
         Site 798, B:1370–1371  
         Site 799, B:724–725  
     aperture, vertical, in logs, B:396–397  
     Ar/Ar age. *See* radiometric age  
     arc tholeiitic suite, Site 795, B:1338  
     arenite. *See* feldspathic arenite; lithic arenite; sandstone  
     argon/argon age. *See* radiometric age  
     argon isotopes. *See* isotopes, argon  
     aridity signal, in logs, A(128):188–192  
     arkose. *See* sandstone  
     aromatic species, Site 799, B:627  
     arsenic  
         Site 794, B:1362  
         Site 795, B:711, 714, 1365  
         Site 797, B:1366

arsenic (cont.)

## SUBJECT INDEX

Site 798, B:1368-1369  
 Site 799, B:724  
 ash. *See* ash layers  
 ash beds. *See* ash layers  
 ash layers, B:1373-1393. *See also* glass, volcanic; tephrochronology; volcanic activity; volcanic phases; volcanic pulses; specific volcanic rocks  
 acidic, A(128):151, 158, 287, B:793  
 age, A(128):123, 155-156, 288, 300; B:963  
 alteration, A(127):22, 109-110, 363-364, A(128):151, 288; B:701, 1267, 1375-1378  
 alteration effects, B:641, 643, 646, 701  
 aluminum vs. silica oxide composition, B:1380  
 areal distribution, B:791  
 Aso-4 ash, B:581  
 barium anomalies, B:1377-1378  
 basic, A(128):151, 285-286; B:793  
 and bioturbation, A(128):153  
 boron content, B:639-640  
 in Brunhes Chron sediments, B:963  
 calcium, A(127):280; B:1383  
 color, A(128):151  
 core/log correlation, B:398, 401, 1025  
 and core recovery, B:1319-1320  
 crystal fraction, A(128):150; B:791  
 in dark/light cycles, B:581  
 definition, A(128):150  
 erosion, A(128):153  
 feldspar composition, B:1388  
 in FMS logs, A(128):153, 184; B:398, 400  
 frequency, A(127):95-97, 190, 193, 346, 350, A(128):157, 301; B:1313  
 frequency underestimation with depth, B:1375  
 frequency vs. age, B:1319-1320  
 frequency vs. thickness, A(128):159, 300  
 frequency vs. time, B:794, 798  
 glass fraction, A(128):150  
 graded, A(127):186, A(128):151-153, 285; B:793, 802  
 grain size, A(128):150; B:791  
 heterogeneous, A(128):153, 285; B:793, 802  
 homogeneous, A(128):151-152, 285; B:793, 801  
 intermediate composition, B:793  
 and interstitial water chemistry, B:1265  
 iron vs. silica oxide composition, B:1381  
 Japan Sea, A(128):32-33; B:1316, 1319-1321  
 lanthanum vs. ytterbium, B:1386  
 lithology, B:791  
 in logs, A(128):288, 339-340  
 magmatic composition, B:793  
 magnesium content, A(127):280; B:1265, 1382  
 magnetic susceptibility correlated, A(127):105, 199  
 mineral fragment composition, B:1379  
 neodymium vs. zirconium, B:1386  
 origin, A(128):156-157, 291-292  
 and oxygen isotopes, B:52  
 petrography, A(128):150-153, 284-288  
 potassium vs. silica oxide composition, B:1385  
 pyroxene composition, B:1388  
 secondary mineralogy, B:1375-1378  
 sedimentary types, B:793  
 sedimentation processes, A(128):151-153, 284-288  
 seismic stratigraphy, A(128):122, 125, 153-155, 344-351, 379  
 silica diagenetic evolution, B:1375-1377  
 Site 794, A(127):90; B:1374  
 Site 795, A(127):186; B:1374  
 Site 796, A(127):251, 264; B:1374

Site 797, A(127):341; B:1373-1374  
 Site 798, A(128):33, 35, 122-124, 137, 139-140, 150-157, 196-198, B:397-398, 791-793, 963, 1374  
 Site 799, A(128):33, 240, 256, 284-291, 296-297, B:605, 791-793, 963, 1373  
 Sites 798 and 799 compared, A(128):291-292, 301  
 slumps, A(128):285  
 sodium vs. silica oxide composition, B:1384  
 sources, A(128):122, 198; B:791  
 strontium, A(127):280; B:607  
 synsedimentary/post-sedimentary processes, A(128):153  
 on synthetic seismogram, A(128):184  
 tephra markers, A(128):157, 291-292; B:793-796, 798-799  
 thickness, A(128):157, 301; B:791, 1319  
 thickness underestimation with depth, B:1375  
 thickness vs. deposition age, B:963  
 thickness vs. time, B:795, 799  
 thorium/uranium ratio, A(128):339  
 thorium vs. zirconium, B:1386  
 and turbidity flows/slumps, A(128):153-154; B:793  
 volcanic activity, B:800  
 volcanic pulses, B:796-797  
 XRF analyses, A(128):151  
 Asia, sandstone provenance, B:110-111  
 Aso-4, B:581, 796  
 Aso volcano, ash layers from, A(128):157  
 asthenosphere, injection of, B:903  
 Atlantic II Fracture Zone, cation exchange capacity, B:1282  
 Atlantic Ocean N, calcareous nannofossils, B:173  
 Atlantic Ocean NE, paleoenvironment, B:338  
 Atterberg limits, Sites 798 and 799, B:1124, 1126  
 augite  
 altered to clay, B:885  
 anhydrous pressure, B:871  
 augite-liquid relations, B:870-872  
 chemical composition, B:851-853, 867-868  
 and cooling rate, B:852, 870-871  
 crystallization, B:870-872  
 experimental chemical compositions, B:867-868  
 hypersthene-normative, B:864  
 in melting experiments, B:861-862  
 Mg/(Mg+Fe) histograms, B:855  
 minor element content, B:854  
 natural vs. experimental augite compared, B:870-871, 878-879  
 and parent magma, B:854, 856  
 Site 794, B:851-852  
 Site 795, B:852  
 Site 797, B:852-853  
 sodium content, B:858  
 titanium, B:855-856  
 back-arc rifting  
 Japan Basin N, A(127):238  
 Japan Sea, A(127):16, 258, A(128):11  
 mechanisms for, A(127):5  
 Site 794, A(127):84, A(128):82-83  
 Site 795, A(127):179-180  
 Site 796, A(127):258  
 Site 797, A(127):336  
 Yamato Basin S, A(127):336-337  
 back-arc tholeiite. *See* tholeiite, back-arc  
 bacteria. *See also* microbiological studies  
 acetogens, B:766-768  
 aerobic ammonifiers, B:766-768  
 anaerobic, B:767, 770-771  
 anaerobic ammonifiers, B:766-768

anaerobic viable count, B:772  
 and bishomohopanoid acid, B:669  
 chemistry, B:765  
 contamination concerns, B:757-758  
 in dark/light cycles, B:772  
 in deep aquifers, B:755, 759  
 depth distribution of dividing and divided cells, B:766  
 and diagenetic processes, A(128):178-179  
 direct counts, B:756, 765-766  
 distribution, B:769-771  
 dormant, B:759  
 filamentous, B:757-758  
 frequency of dividing and divided cells (FDDC), B:757-758, 765-766, 769, 772  
 growth, B:755-760  
 H-thymidine incorporation, B:757, 759, 767, 769, 771-773  
 heterotrophs, B:766-768  
 importance, B:761  
 incubation, B:756, 764  
 under metabolic stress, B:759  
 and methanogenesis, B:767-769, 772-773  
 methods of study, B:755-756, 761-765  
 nitrate reducers, B:766-768  
 Peru margin, B:755, 757, 759, 761, 769, 773  
 potential activity, B:763, 767-769  
 presence at depth, B:761  
 productivity, B:758-759  
 rates of bacterial activity, B:771-772  
 sampling times, B:756  
 ship-shore transportation, B:762  
 shipboard handling, B:761-762  
 shore lab handling, B:762-765  
 Site 798, A(128):125-126; B:755-776  
 sulfate reducers, B:766-768, 770, 772-773  
 and temperature, B:761  
 total anaerobic, B:767  
 total count, B:758, 766-767, 772  
 viability, B:759, 763-767, 773  
 Bagnold effect, and vein fillings, B:1180  
 Baja California, diagenetic carbonates, B:75  
 Barbados forearc, veins observed, B:1180  
 Barbados Ridge Complex, and chloride, A(128):172  
 barite  
 in ash layer, B:1392  
 authigenic precipitation, B:665  
 crystals, B:664  
 energy dispersive spectra, B:1378  
 estimated ion activity product (IAP)  
 nodules, B:637-638  
 replacement mineral in sandstone, B:143  
 Site 795, B:640  
 and sulfur isotopes, B:637-638  
 barium, B:651-665  
 abundance, B:663  
 accumulation rate, B:660  
 authigenic precipitation, B:654  
 barium front, B:637, 663  
 basement, A(128):99  
 biogenic, B:653, 656, 659  
 burial rates, B:652  
 $C_{org}$  barium ratio, B:653-4554, 661  
 detrital vs. biogenic component, B:657  
 diagenetic redistribution, B:652  
 dissolution, B:659, 661  
 in interstitial water, B:657-659  
 mass accumulation rates, B:665  
 paleoproductivity indicator, B:651-656  
 preservation, B:652  
 and productivity, B:651-656, 661-662

## SUBJECT INDEX

remobilization, B:654, 656-659  
 schematic geochemical diagram, B:664  
 and sediment accumulation rate, B:660-661  
 Site 794, B:653-656, 1362  
 Site 795, B:710, 1365  
 Site 796, B:654, 656  
 Site 797, B:1366  
 Site 798, B:654, 656, 1368-1369  
 Site 799, B:654, 656, 724-725  
 and volcanic glass alteration, B:1377-1378  
 vs. sedimentation rate, B:661  
 vs. yttrium, A(128):100  
 vs. zirconium, A(128):100  
 barium accumulation rate. *See* barium  
 basalt. *See also* basement; specific minerals in basaltic rocks  
 AFM diagram, B:874  
 alteration, B:669-701, 838, 883-889, 891,  
 899, 901, 908-916  
 altered veins, B:886-887  
 augite-liquid relations, B:870-872  
 calc-alkalic evolution trend, B:870, 875  
 chemistry, B:839-842  
 and chromium spinel, B:837-847  
 crystallization history, B:853-854  
 experimental and core data compared, B:870  
 fractionation models, B:872-874  
 grain size, A(127):370  
 lava flows, B:819  
 liquid line of descent, B:869-882  
 major elements, B:869-870  
 neodymium isotopes, B:901-903  
 oxide variations, B:869-870, 873  
 parent magma, B:855-856  
 petrography, B:892  
 phase equilibria, B:870  
 pseudoternary normative projections, B:875  
 sills, B:819  
 Site 794, B:869  
 Site 795, B:1338  
 Site 797, B:869, 899-900, 911  
 strontium isotopes, B:901-903  
 thickness of altered layer, B:699-701  
 tholeiitic evolution trend, B:870, 875  
 vesicles, B:887-888  
 basalt, alkali, Site 797, B:920, 1339-1340  
 basalt, aphanitic, Site 797, A(127):345-346  
 basalt, aphyric  
 alteration, B:885, 887  
 microphotograph, B:859  
 radiometric age, B:822-823, 825  
 Site 794, A(128):68-69, 90-91; B:853, 1339  
 Site 797, A(127):369-370, 375-380; B:850,  
 911  
 basalt, augite phryic, Site 795, B:854  
 basalt, back-arc, isotopes, B:805  
 basalt, calc-alkaline  
 Site 795, B:919  
 Site 797, B:1339-1340  
 basalt, doleritic, Site 794, A(128):68, 86, 88  
 basalt, high-Al, B:893  
 Site 794, B:861  
 Site 797, B:861, 900, 920, 1339-1340  
 basalt, island-arc, isotopes, B:805  
 basalt, mid-ocean ridge (MORB), and helium isotopes, B:747  
 basalt, olivine phryic  
 Site 794, B:849, 853  
 Site 797, B:850, 854-857  
 basalt, olivine plagioclase phryic, Site 794, B:908  
 basalt, olivine-pyroxene microphyric, Site 794,  
 A(128):68, 90

basalt, olivine tholeiitic, liquid line of descent,  
 B:864  
 basalt, phryic pyroxene plagioclase, Site 795,  
 B:918  
 basalt, plagioclase olivine phryic, Site 797,  
 A(127):369  
 basalt, plagioclase phryic  
 crystal accumulation, B:853-854  
 radiometric age, B:823, 826  
 Site 794, B:849, 853  
 Site 795, A(127):174, 217; B:850, 854  
 Site 797, A(127):370-371, 380; B:850, 854  
 basalt, pyroxene phryic, Site 795, B:850  
 basalt, pyroxene plagioclase phryic  
 radiometric age, B:826-827  
 Site 795, A(127):174, 217  
 basalt, tholeiitic  
 Site 794, B:899, 916  
 Site 797, B:900, 1339-1340  
 basement. *See also* fractionation models; mantle source; parent magma; petrology, experimental; radiometric age; X-ray fluorescence (XRF) analyses; specific minerals in basement rocks; specific rock types  
 age, A(127):22, 107, 192-193, 195, 361,  
 A(128):69; B:821, 906-908, 1333-  
 1337  
 age, determination methods, B:1303-1304  
 age, lithospheric stretching models, B:1304-  
 1305  
 alteration, A(127):23, 122-125, 217-219, 280,  
 369-380, A(128):68-69, 321; B:779,  
 849, 883-889, 891, 908-916, 1265-  
 1267, 1282, 1338  
 alteration effects, B:641-643, 646  
 alteration mineralogy, B:884-888  
 arc tholeiite suite, B: 1338  
 baked contacts, A(127):369-380  
 boron as alteration product, B:639  
 brecciation, B:1296  
 calc-alkaline suite, B:1338  
 chemistry, A(127):24-25, 123, 126, 174, 219,  
 223, 325, 381-383, A(128):68-69,  
 96ff, 98; B:779-789, 805-817, 839,  
 872, 892-894, 905-916, 1338-1340  
 chilled margins, A(127):369-380  
 compared to MORBS and BABBs,  
 A(127):126, 219  
 crustal anisotropy, B:1107-1121  
 depth, B:1120  
 drilling rubble, Site 794, A(127):125  
 electrical resistivity, B:1354  
 emplacement, A(127):381  
 eruptive history, A(127):174  
 fractionation models, B:872-874  
 fractures, A(127):217, A(128):69  
 grain size, A(128):90-91  
 groundmass, A(128):88-91, B:849-859, 892  
 igneous event sequence, A(128):95-97  
 interlayered sediments/igneous rocks,  
 A(127):369  
 isotopes, B:779-789, 805-817, 884-888,  
 1338-1341  
 Japan Basin N, A(127):169, 176-177  
 Japan Sea, A(127):24-25, 29, A(128):67, 32-  
 33  
 Kita-Oki Bank, A(128):127-129  
 Kita-Yamato Bank, A(128):247-249  
 Legs 127/128 Site 794 rocks correlated,  
 A(128):93-95  
 logging and logs, A(127):28, A(128):86, 93-  
 95, 103-104; B:1282

Lower Volcanic Complex (LVC), Site 794,  
 B:780  
 magma genesis, B:1340-1341  
 magma suites recognized, B:869-870  
 magmatic affinity, A(128):96ff  
 magnetic intensity, A(127):174  
 magnetic polarity, A(128):86; B:936-938  
 magnetic properties, B:933-945  
 magnetic susceptibility, A(127):174  
 magnetites, A(127):203, 209, 358-359,  
 A(128):101-102  
 major elements, B:869-870, 893  
 Middle Volcanic Unit (MVU), Site 794, B:780  
 mineral composition, B:850-853  
 N-MORB normalized composition, B:1338  
 nature of unit contacts, A(128):68, 88-91  
 neodymium isotopes, B:786-787  
 normalized composition, A(127):25, 126, 223,  
 383  
 Oki Islands, A(128):127-129  
 Oki Ridge, A(128):127-129  
 oxide variation, B:869-870  
 parent magma, B:855-856, 894  
 petrography, A(128):86-91; B:849-850, 891-  
 892  
 phenocrysts, B:849-859, 892  
 physical properties, A(127):26, 135, 139-140,  
 227-228, A(128):102ff; B:1282-  
 1284, 1289-1291  
 primary mineralogy, B:891-892, 908  
 radiogenic isotopes, B:807  
 radiometric age, B:819-836  
 secondary mineralogy, B:891, 908  
 seismic expression, A(127):28-29, 143, 234-  
 236, 312-313  
 sills and flows, A(127):126-127; B:853-855  
 Site 794, A(127):73, 120-127, A(128):32-33,  
 36, 67-69, 86ff, B:849, 883-889,  
 891-894, 906, 927, 933, 1112-1113,  
 1336, 1339, 1354  
 Site 795, A(127):174, 217ff; B:849, 906, 933,  
 1336, 1338-1339  
 Site 797, A(127):325, 369-383; B:849, 891-  
 894, 906-908, 933, 1336, 1339-1340  
 spidergram, B:923-924, 926  
 stratigraphy, B:871, 901  
 strontium isotopes, B:786-787  
 structural features, A(128):91-93; B:1181-  
 1184  
 Takuyo Bank, A(128):248-249  
 tectonic significance, B:1337-1341  
 temperature, A(127):137  
 thermal conductivity, B:1281  
 thickness of altered layer, B:699-701  
 trace elements, B:807  
 unit boundaries correlated, A(128):93-95  
 units, Site 794, A(127):73, 121-125,  
 A(128):68-69, 75, 86-91; B:779-  
 781, 884, 907-908  
 units, Site 795, A(127):174, 217-219, 221;  
 B:909  
 units, Site 797, A(127):369-380; B:910  
 Upper Volcanic Complex' (UVC), Site 794,  
 B:780  
 velocity vs. porosity, A(127):140  
 X-ray fluorescence analyses, B:892-894  
 Yamato Bank, A(128):247-249  
 Yamato Basin, A(127):72, A(128):23-24, 32-  
 33  
 Yamato Basin N, A(127):77-81, A(128):73-  
 75, 117  
 Yamato Basin S, A(127):331-334  
 Yamato Rise, A(128):247-249

basin

basin. *See* pull-apart basin; *particular basins*  
bedding  
  in blue tuff, B:117  
  color-banded, A(127):186, 189  
  dips, Site 799, B:1176, 1178  
Bering Sea, opal flux, B:439  
beryllium isotopes. *See* isotopes, beryllium  
BHTV. *See* borehole televiewer  
bicarbonate, and alkalinity, A(128):173  
biochronology synthesis, B:1219–1228  
bioclastic sand. *See* sand, bioclastic  
biosilica. *See* silica, biogenic  
biosiliceous sediments  
  laminated, B:439–455  
  permeability, B:1128  
biostratigraphic zonations, B:1225–1226. *See also*  
  biostratigraphy; microfossils; *names of*  
  *specific microfossils*  
foraminiferal and diatom zones correlated,  
  B:222–223  
foraminifers, B:200–201  
Japan Sea, A(127):20  
of onshore exposures, A(128):17  
Site 794, A(127):98, 100; B:344–345  
Site 795, A(127):196–197; B:346–347  
Site 796, A(127):271–272  
Site 797, A(127):353; B:348–349  
Site 798, A(128):161  
Site 799, A(128):304–305  
biostratigraphy, B:1219–1228. *See also* biostrati-  
graphic zonations; microfossils; *names of*  
*specific fossil groups*  
age of oldest sediments, B:1335–1337  
datum levels used, B:1222  
high-resolution study, B:193–200  
Japan Sea, A(127):19–22, A(128):28–30  
Leg 127, A(127):19–22  
Leg 128, A(128):28–30  
Legs 127 and 128 correlated, A(128):29  
Neogene foraminifers, B:188–189  
problems with, B:1219  
radiometric ages compared, B:821–822, 1337  
Site 794, A(127):72, 96–103, A(128):99–100;  
  B:187–224, 1221–1223, 1225  
Site 795, A(127):174, 192–199; B:187–224,  
  1223, 1225  
Site 796, A(127):250–251, 269–274; B:187–  
  224, 1223, 1225  
Site 797, A(127):324, 351–357; B:187–224,  
  1223, 1225  
Site 798, A(128):28–30, 124–125, 158–166;  
  B:155–169, 1223–1226  
Site 799, A(128):28–30, 244, 298–312; B:155–  
  169, 1224, 1226  
biotite, in ash layers, B:1379, 1392  
bioturbated sediments. *See* sediments, bioturbated  
bishomohopanoid acid, Sites 798 and 799, B:669,  
  671  
Black Sea  
  oxygenation conditions, B:705  
  pyrite, B:712  
  trace metals, B:1361  
Blake Outer Ridge, carbon isotopes, B:88  
blooms, Site 798, A(128):165  
blue tuff  
  age of Yamato seamount chain samples com-  
    pared, B:126  
  clay minerals, B:119  
  deposition, B:119, 123–126  
  lithology, B:117  
  petrography, B:117–119  
  plagioclase, B:124–126

## SUBJECT INDEX

pyrogenic crystal chemistry, B:116, 121–123,  
  126  
radiometric age, B:116, 123, 126–127  
sedimentology, B:115–120  
Site 794, B:120  
Site 796, B:121  
Sites 794 and 796, B:115–130  
stratification, B:129  
strontium isotopes, B:116, 123, 127  
Yamato Basin and Japan Basin samples com-  
pared, B:127–128  
boninite, spinel composition, B:844  
borehole diameter, and geochemical logging er-  
rors, B:1029–1030  
borehole televiewer  
  fractures, A(127):307, 399  
  hole ellipticity, A(127):307  
  Site 796, A(127):307  
  Site 797, A(127):399, 409  
  stress field determination, B:1047  
boron  
  adsorption/desorption processes, B:640–641  
  and ash layer alteration, B:639–640  
  and basement alteration, B:639, 1266  
  and gas hydrate, B:639  
  in interstitial water, B:638–642  
  and organic matter, B:640  
  and sand transport, B:640  
  Site 794, B:638–641  
  Site 795, B:638–641  
  Site 796, B:638–641  
  Site 797, B:638–641  
  vs. chloride, B:642  
  vs. magnesium, B:642  
boron isotopes. *See* isotopes, boron  
Boso Peninsula, magnetostratigraphy, A(127):199  
bottom simulating reflector (BSR)  
  age, B:1155–1156  
  causes, B:1145–1147  
  and heat flow, B:1147–1148  
  and opal-A to opal-CT transition, B:1149  
  polarity, B:1147  
  and sedimentation rate, B:1147–1148  
  Yamato Basin, B:1155  
Bottom Water, B:577  
bottom water oxygenation conditions. *See oxy-*  
  *genation conditions*  
breccia, tuff. *See also* tuff, rhyolitic  
  in logs, A(128):287  
  Site 799, A(128):243, 287; B:34  
breunnerite, Site 799, B:82  
brittle structures  
  Site 794, B:1181–1183  
  Site 798, B:1181–1182  
  Site 799, B:1182–1183  
broadband seismometer. *See* downhole seismic ex-  
periment  
bromine  
  Site 798, B:1368–1369  
  Site 799, B:724–725  
Brunhes Chron  
  and ash layers, B:963  
  and diatom occurrence, A(128):159  
  and inclination variations, B:960  
  and sediment accumulation rate, B:468  
  and sedimentation rate, B:961  
  silicoflagellate and ebridian zones correlated,  
    A(128):162  
  Site 796, A(127):275  
  Site 797, B:973  
  Site 798, A(128):124  
  warm/cold intervals in, B:459, 464  
Brunhes/Matuyama boundary  
and diatom occurrence, A(128):161  
field behavior during polarity transition,  
  B:970, 972, 974, 978  
and foraminiferal coiling direction, B:208  
and hematite, B:972  
inclination variations, A(128):172  
multiple reversals at, A(127):22  
sedimentation rate, B:961  
silicoflagellate and ebridian zones correlated,  
  A(128):305  
Site 794, B:218, 224, 970, 1221  
Site 795, A(127):199; B:972, 1223  
Site 797, B:218, 224, 973  
Site 798, A(128):30, 156, 165, 167; B:164,  
  563, 973, 1224  
Site 799, A(128):314; B:974, 1224  
BSEM images, faunal/floral morphotaxa recog-  
nized, B:548  
bulk density. *See* density  
Burma earthquake, and downhole seismic experi-  
ment, B:1162, 1168  
burrows  
  in cyclic lithofacies, A(128):140–141  
  in dark/light cycles, B:564–565, 569  
  with organic carbon fillings, B:38  
  and oxygenation conditions, A(128):141;  
    B:1232  
  Site 795, A(127):189  
  Site 796, A(127):265–266  
  Site 797, A(127):344, 346, 348–349  
  Site 798, A(128):143–145  
  Site 799, A(128):262  
iso-butane  
  Site 798, A(128):125, 176, 187  
  Site 799, A(128):244–245, 322  
n-butane  
  Site 798, A(128):125, 176, 187  
  Site 799, A(128):244–245, 322  
C/N ratio  
  in color-banded beds, A(127):365–367  
  in dark/light cycles, A(127):212; B:431  
  Site 794, A(127):113–114, 119  
  Site 795, A(127):211–212, 216; B:708–709  
  Site 796, A(127):283–285, 287  
  Site 797, A(127):365–367, 375  
  Site 798, A(128):189–193; B:425–426, 435–  
    437  
  Site 799, A(128):324, 334–338, 341; B:427,  
    429  
C/S ratio  
  and oxygenation condition, B:1241, 1249  
  Site 794, B:1246–1247, 1255  
  Site 796, A(127):283  
  Site 798, A(128):189–192  
  Site 799, A(128):324, 334–338, 342; B:623–  
    624  
Ca/K atomic ratio, and radiometric dating of blue  
tuff, B:123  
calc-alkalic evolution trend  
  in basalt, B:870, 875  
  and suppression of plagioclase crystallization  
    temperature, B:873  
calc-alkaline glass, B:1378, 1387  
calc-alkaline suite, Site 795, B:1338  
calcareous nannofossils. *See* nannofossils, calcare-  
ous  
calcite  
  abundance in Japan Sea sediments, B:1235–  
    1241  
  in dark/light cycles, B:588  
  isotope data, B:888  
  Site 795, A(127):188  
  Site 799, A(128):289; B:611

## SUBJECT INDEX

cation exchange capacity

in veins, B:887  
 calcite, diagenetic  
   chemical composition, B:81  
   formation depth, B:85  
   isotopic composition, B:83  
   occurrence, B:80  
   replacing orthoclase, B:147  
   in sandstone, B:141–143, 146  
   shallowest occurrence, A(128):273  
 Site 798, A(128):137  
 Site 799, A(128):272; B:75–98

calcium  
   and albitization, B:137  
   alteration and, B:909, 911  
   and ash alteration, A(127):280, 363–364  
   in ash layers, B:1383  
   and basement, A(127):280, B:892  
   in dark/light cycles, B:569  
   and dolomite formation, A(127):363  
   in interstitial water, B:607, 1263, 1268  
 Japan Sea, B:1236, 1265  
   mobility during alteration, B:838  
 Site 794, A(127):109–110  
 Site 795, A(127):174, 205  
 Site 796, A(127):280  
 Site 797, A(127):363–364, 370  
 Site 798, A(128):173–174, 182; B:1368–1369  
 Site 799, A(128):318, 329; B:611  
   sources of, B:143  
   vs. alkalinity, A(128):183  
   vs. magnesium, A(127):115  
   vs. strontium isotopes, B:645  
   XRF and geochemical log analyses compared, B:1028–1029, 1031–1035

calcium, logging  
 Site 794, B:1420–1421  
 Site 796, B:1423  
 Site 797, B:1426–1427  
 Site 798, B:1024  
   Site 799, A(128):364; B:1408–1409

calcium/aluminum ratio, Site 794, B:1236

calcium carbonate ( $\text{CaCO}_3$ ). *See* carbon, inorganic; carbonate content

calcium oxide. *See* calcium

California  
   planktonic foraminifers, A(128):165  
   silicoflagellate, B:237

caliper logging  
   in dolomite, A(128):332–333  
   and lithology, A(127):306  
 Site 794, A(127):142, 147  
 Site 796, A(127):303–306  
 Site 797, A(127):395–396, 403

calthrops  
 Site 795, B:543

carbon, inorganic  
   Japan Sea, A(127):23  
   Site 794, A(127):113  
   Site 795, A(127):209–212  
   Site 796, A(127):283–285  
   Site 797, A(127):365–367  
   Site 798, A(128):125, 176–177, 189–192  
   Site 799, A(128):323–324, 334–338

carbon, organic. *See also* organic matter; Rock-Eval analyses; sediments, organic-rich  
   accumulation rate, B:425, 428, 660  
   and alkalinity, B:1262  
   and ammonium, B:1262  
   in burrows, B:38  
   and carbon isotopes, B:1262–1263  
    $\text{C}_{\text{org}}/\text{barium}$  ratio, B:653–654, 661  
   and cyclic lithofacies, A(128):125, 177  
   in dark-colored beds, A(127):283, 365

in dark/light cycles, B:431–432, 568–569, 579, 586–588, 594, 671  
   decomposition, B:1262–1263  
   and diagenetic carbonate, B:92  
   environment needed for accumulation, B:429–430  
   flux rate, B:423–437  
   glacial/interglacial variation, B:589  
   and gray value, B:589  
   and increased productivity, A(127):283  
 Japan Sea, A(127):24  
   mass accumulation rate, B:425, 428  
   and oceanographic interpretations, A(127):368–369  
   and phosphate, B:1262  
   and productivity, B:430  
   and sedimentation rate, B:628  
   shipboard vs. shorebased analyses compared, B:623, 628  
 Site 794, A(127):113–114, 119; B:656, 1246–1247, 1255  
 Site 795, A(127):209–212, 216; B:706–710, 714  
 Site 796, A(127):251, 283–285, 287; B:656  
 Site 797, A(127):325, 365–367, 375; B:592  
 Site 798, A(128):32, 125, 176–177, 189–194, 196; B:423–426, 435–437, 656, 668–669, 776  
 Site 799, A(128):32, 238–239, 245, 323–324, 334–338, 340; B:38, 423–437, 623–627, 629, 656, 668–669  
   and sulfate, B:1262  
   vs. biogenic barium, B:659  
   vs. free hydrocarbons ( $\text{S}_i$ ), B:631  
   vs. nitrogen, A(128):343–344  
   vs. sulfur, A(128):343–344; B:594, 628, 712–713, 1241, 1254  
 carbon, total organic (TOC). *See* carbon, organic; Rock-Eval analyses  
 carbon isotopic ratios. *See* isotopes, carbon  
 carbon/nitrogen ratio. *See* C/N ratio  
 carbon/sulfur ratio. *See* C/S ratio  
 carbonaceous siltstone. *See* siltstone, carbonaceous  
 carbonate. *See also* carbonate, biogenic; carbonate, diagenetic  
   in dark/light cycles, B:584  
   diagenesis, B:1263–1265, 1267  
   glacial/interglacial variation, B:590  
   in Japan Sea sediments, B:1235–1241  
   logged thickness vs. cored thickness, B:79  
   in logs, B:78–79  
   mass accumulation rate, B:428  
   nodules, B:611–612, 615, 1232  
   occurrence, B:78–79  
 carbonate, authigenic. *See* calcite, diagenetic; carbonate, diagenetic; dolomite; siderite  
 carbonate, biogenic  
   diagenetic phases, A(127):267  
   preservation, A(127):169  
   recrystallization, A(128):318  
   Site 799, B:78  
 carbonate, diagenetic. *See also* carbonate diagenesis  
   abundance and distribution, B:80, 82  
   and calcareous nanofossils, B:166  
   carbon isotopes vs. depth, B:85–90  
   chemical composition, B:81–85  
   compositional zoning, B:83  
   disseminated rhombs, B:78  
   equilibrium oxygen isotope values, B:83–85,  
     89

formation conditions, A(128):277; B:83–87, 89–90, B:888  
   and inorganic geochemical data, A(128):276–277  
   interstitial water chemistry and, B:646  
   isotopic composition, B:83  
   layers, B:78  
   nODULES, B:78  
   occurrence, B:78  
   oxygen isotopes, B:86–88  
   paleoenvironmental implications, B:93–95  
   petrography, B:81  
   provenance, B:148  
   replacing quartz, B:147  
   and revised lithostratigraphy, B:1232–1233  
   in sandstone, B:141–143  
   SEM photographs, B:97–98  
 Site 796, A(127):266–267  
 Site 798, A(128):147–150; B:701–702  
 Site 799, A(128):240, 260, 272–278, 292, 354; B:38, 75–98, 697, 701–702  
   stratigraphically controlled distribution, B:76  
   textural relations, B:81  
   thickness, B:77  
   X-ray diffraction analyses, B:79–80, 86–87  
 carbonate-compensation depth (CCD)  
   and benthic foraminifers, B:504–505  
   and calcareous nanofossils, B:1220  
 Japan Sea, A(127):24, A(128):20; B:155, 208, 440, 1201  
   and microfossil preservation, A(127):19  
 Oki Ridge, B:409  
   and sedimentation rate, B:1228  
 Site 794, A(127):103, 113, 147  
 Site 795, A(127):169, 199, 211  
 Site 796, A(127):247, 283, 315  
 Site 797, A(127):324, 356, 365, 409  
 Site 798, A(128):122, 158; B:168  
 Site 799, A(128):292  
 Yamato Basin, A(127):103  
 carbonate content  
   core analyses, B:404  
   in dark/light cycles, B:431–432, 579, 586–588  
   logging estimates, B:402, 404  
 Site 794, A(127):113–114, 119  
 Site 795, A(127):209–212, 216  
 Site 796, A(127):283–285, 287  
 Site 797, A(127):365–367, 375; B:592  
 Site 798, A(128):32, 189–194; B:401, 435–437, 445–446, 668–669  
 Site 799, A(128):32, 245, 307, 323–324, 334–338, 340; B:668–669  
 carbonate diagenesis. *See also* carbonate, diagenetic and interstitial water oxygen isotopes, B:701–702  
   and magnesium, A(128):174  
   material balance calculation of carbonate precipitation, B:702  
 carbonate minerals  
   formation, A(127):22–23  
   and iron, B:722–723  
 carbonate nodules. *See* nodules, carbonate  
 carbonate preservation  
   Site 794, B:188  
   Site 795, A(127):211  
   Site 797, B:188  
 carboxylic acids  
   Site 798, B:674  
   Sites 798 and 799, B:669–671  
 Cariaco Basin, oxygenation conditions, B:705  
 cation exchange capacity (CEC), and clay alteration minerals, B:1282

CCD. *See* carbonate-compensation depth  
 Ce anomaly  
   and diagenetic fractionation, B:692–693  
   and dolomite, B:92  
   and hydrogenous precipitation, B:731  
   and hydrothermal activity, B:719, 729  
   and lanthanum/ytterbium ratios, B:693–694  
   and light rare earth elements, B:693–694  
   and oxygenation conditions, B:692  
   and sediment contamination of mantle source, B:815  
   Site 794, B:682, 693  
   Site 795, B:683  
   Site 797, B:688–691, 693  
   Site 798, B:726, 735  
   Site 799, B:90–91, 729, 736  
   vs. Eu anomaly, B:737  
   vs. lanthanum/ytterbium ratio, B:737  
   vs. manganese content, B:737  
 Ce/Ce. *See* Ce anomaly  
 celadonite  
   in vesicles, A(127):217  
   and volcanic glass alteration, B:1375  
 cerium. *See also* Ce anomaly  
   distribution, B:692  
   and hydrogenous manganese precipitation, B:726–727  
   Site 794, A(128):101  
   Site 795, B:707  
   Site 798, B:1368–1369  
   Site 799, B:90–91  
 cesium  
   Site 795, B:707  
   Site 798, B:1368–1369  
   Site 799, B:724  
 Chenopodiaceae  
   Site 794, B:491  
 chert  
   cycles in FMS logs, B:1041–1044  
   in FMS logs, A(127):307; B:1039–1040, 1232  
   in logs, A(127):27–28, 302, 394  
   and revised lithostratigraphy, B:1232  
   in sandstone, B:144–148  
   seismic expression, A(127):407–408  
   Site 795, A(127):187–188  
   Site 796, A(127):265  
   Site 797, A(127):343–344, 349  
 China  
   dust storms, B:394  
   loess deposits, B:394  
   subcontinental lithosphere, B:811–812  
 chloride  
   and clay minerals, B:608–610  
   distribution, B:605  
   and opal-A, B:608–610  
   Site 795, A(127):205  
   Site 796, A(127):278–279  
   Site 798, A(128):172, 179  
   vs. boron, B:642  
 chlorine  
   and geochemical logging, B:1023–1024  
   Site 794, A(127):107  
   Site 797, A(127):364, 374  
   Site 799, A(128):317, 328; B:610  
 chlorite  
   in Japan Sea sediments, B:1235–1241  
   in sandstone, B:139  
   as sandstone cement, B:134  
   Site 798, B:411, 416, 418  
 chlorite/saponite mixed-layer clay  
   from altered materials, B:884–886, 888  
   chemical composition, B:885–887  
   and isotopes, B:888

in sandstone, B:139  
 Site 794, B:885  
 in veins, B:886–887  
 X-ray diffraction spectra, B:886  
 chlorophyllinite, Sites 798 and 799, B:669–670  
 chromatography. *See* pyrolysis-gas chromatography  
 chromite, B:840. *See also* spinel; spinel, chromium  
 chromium  
   basement, A(128):99  
   Site 795, B:706  
   Site 797, B:920  
   Site 798, B:1368–1369  
   from terrigenous sources, B:723  
   vs. aluminum, B:733  
   vs. yttrium, A(128):101  
 chromium/barium ratio, basement, A(128):99  
 chromium spinel. *See* spinel, chromium  
 Chron 5, Site 798, A(128):30  
 Chron 6  
   Site 796, A(127):277  
   Site 798, A(128):30  
 Chugoku Backbone Range, uplift, B:526, 528  
 Cichorioideae  
   Site 795, B:491  
 clathrate. *See also* gas hydrate  
   Site 796, A(127):251, 278, 281, 288–289  
 clay  
   aluminum, logging, A(127):306–307  
   and revised lithostratigraphy, B:1229–1230  
   Site 794, A(127):90–92, A(128):77, 81  
   Site 795, A(127):186  
   Site 797, A(127):340–341  
   Site 798, A(128):124, 137  
   Site 799, A(128):256  
 clay, coatings of, B:57  
 clay, diatom  
   and revised lithostratigraphy, B:1230–1232  
   Site 794, A(127):90  
   Site 795, A(127):186  
   Site 796, A(127):261–264  
   Site 797, A(127):341–343  
   Site 798, A(128):124, 137–138  
   Site 799, A(128):256, 260  
 clay, siliceous silty, and revised lithostratigraphy, B:1232  
 clay, silty  
   in dark/light cycles, B:568  
   and revised lithostratigraphy, B:1229–1232  
   Site 794, A(127):90, A(128):77  
   Site 795, A(127):186  
   Site 796, A(127):261–264  
   Site 797, A(127):340–341  
   Site 798, A(128):124, 137–138  
   Site 799, A(128):256  
 clay content  
   and density, B:1278–1279  
   and gamma ray logging, B:6  
 clay minerals. *See also* clay minerals, diagenetic; names of specific clay minerals  
   abundance in Japan Sea sediments, B:1235–1244  
   in altered materials, B:885–888  
   and ammonium, logging, A(127):279  
   and basement alteration, B:908–909, 918  
   in blue tuff, B:119  
   chemical composition, B:885–887  
   compaction structures, B:39  
   crystallization, A(128):320–321  
   D/H ratios, interstitial water, B:642  
   in dark/light cycles, B:569  
   in diagenetic sequence, B:39

fabric, B:35–36, 38  
 and gamma-ray logging, A(127):393  
 glacial/interglacial variation, B:588  
 and low-chloride interstitial fluids, B:608–610  
 major element molar ratios, B:611  
 and rare earth elements, B:691  
 as sandstone cement, B:134–135  
 SEM photographs, B:30  
 Site 794, A(127):94  
 Site 798, B:411, 416, 418  
 Site 799, B:611  
 clay minerals, diagenetic. *See also* clay minerals; names of specific clay minerals  
   in sandstone, B:139–141  
   Site 799, B:35–36  
 clayey diatom ooze. *See* ooze, clayey diatom  
 claystone. *See also* pelite  
   and revised lithostratigraphy, B:1232–1233  
   sediment accumulation rate, A(127):191  
   Site 794, A(127):92–93, A(128):81  
   Site 795, A(127):189  
   Site 796, A(127):264; B:1260  
   Site 797, A(127):343–344  
   Site 798, A(128):124, 138  
   Site 799, A(128):265; B:41–43  
 claystone, dark, deposition, A(127):267  
 claystone, diatom  
   Site 795, A(127):186–187  
   Site 796, A(127):264  
 claystone, pebbly, Site 796, A(127):264–265  
 claystone, siliceous  
   grain shape, B:38  
   grain size, B:38  
   and revised lithostratigraphy, B:1230–1232  
   sediment accumulation rate, A(127):191  
   Site 794, B:1258  
   Site 795, A(127):187–189  
   Site 796, A(127):264–265  
   Site 797, A(127):343–344  
   Site 798, A(128):124, 138  
   Site 799, A(128):260, 264–265; B:39–43  
   in vitric tuff, B:34  
 claystone, silty  
   and revised lithostratigraphy, B:1232–1233  
   Site 794, A(127):93–94  
   Site 797, A(127):343, 345–346  
   Site 799, B:43  
 claystone, silty dolomitic, Site 799, B:41  
 claystone, silty phosphatic, Site 799, B:43  
 claystone, silty siliceous, Site 799, B:40–41  
 claystone, spicular silty, Site 797, A(127):343  
 claystone, tuffaceous, Site 794, A(128):81  
 climate  
   Japan Sea, B:180, 337  
   pollen and, B:320  
   precipitation and, B:326  
   radiolarians and, B:299  
   Site 796, A(127):274  
   Site 797, A(127):356  
   vegetation and, B:325–326  
 clinoptilolite  
   abundance, B:1235–1244  
   in ash layers, B:1390  
   and opal-A to opal-CT transition, B:10  
   SEM photographs, B:30, 1294  
   Site 796, B:139  
   Site 797, A(127):344  
 clinopyroxene  
   in ash layers, B:1379  
   in melting experiment, B:895  
 cluster analysis, Japan Sea, B:525–526, 531, 535  
 CNT-G. *See* porosity, logging  
 cobalt

Site 795, B:710  
 Site 798, B:1368–1369  
 Site 799, B:724  
 coccolithophores  
   Site 799, A(128):310  
 cocoliths  
   Site 799, A(128):307, 309  
 Cochiti Subchron, Site 797, B:1223  
 coiling direction. *See also* coiling ratio; foraminifers, planktonic  
   of planktonic foraminifers, A(128):165, 310–311  
   as surface-water temperature indicator, B:1220  
 coiling dominance (CD) curve, B:459. *See also* coiling direction; coiling ratio; foraminifers, planktonic  
   Site 798 coiling ratios correlated, B:460  
   Site 798 warm/cold events correlated, B:463–465  
 coiling ratio. *See also* coiling direction; coiling dominance (CD) curve; foraminifers, planktonic  
   of N Pacific foraminifers, B:459  
   oxygen isotopes correlated, B:467  
   as paleoclimate indicator, B:463, 467–468  
   of planktonic foraminifers, B:188, 193, 208, 210–215, 218–219  
 collosphaerids  
   Japan Sea, B:296  
 color-banded bedding. *See also* dark/light cycles; lithofacies, cyclic  
   in clay and silty clay, A(127):341  
   in diatom-bearing claystone, A(127):261  
   and organic carbon content, A(127):365  
   origin, A(127):19, 24, 349  
   Site 797, A(127):324, 343  
 color-banding, in dark/light cycles, B:581  
 compaction structures, Site 799, B:38, 39, 48  
 compensated neutron tool (CNT-G). *See* porosity, logging  
 composition, sediment. *See* sediment composition  
 compression  
   along Hokkaido W-Honshu N thrust belt, A(128):76–77  
   and fracture formation, B:1181  
 compressional stress field, Japan Sea and, B:529–531  
 compressional velocity. *See* velocity  
 conductivity model, Japan Arc, A(128):85  
 conglomerate, of porcellanite pebbles, A(127):344–345  
 conifers  
   Japan Sea, B:327–328  
   Japanese archipelago, B:334  
   Site 798, B:318–319  
 connate brackish water, and interstitial water analyses, B:607–608  
 consolidation  
   and Atterberg limits, B:1124  
   in debris flows and slumps, B:1125  
   definitions used, B:1123  
   determined from in situ pore water pressure, B:1124  
   excess pore-water pressure, B:1125–1127, 1129  
   and gassy sediments, B:1129  
   and leakage from artesian water or gas pressure source, B:1129–1130  
   normally consolidated/underconsolidated sediment boundary, B:1124  
   from oedometer tests, B:1125–1127  
   overburden pressure, B:1125–1127, 1129  
   overconsolidation ratio, B:1125–1127, 1129

preconsolidation pressure, B:1125–1127, 1129  
 processes causing, B:1123  
 and sedimentation B:1124–1125, 1129–1130, 1277  
 shear strength/overburden pressure ratio, B:1125–1126  
 Site 798, B:1125–1127  
 Site 799, B:1125, 1127  
 underconsolidation, B:1129–1130  
 void ratio vs. effective pressure, B:1126, 1128  
 convergence  
   along Hokkaido W-Honshu N thrust belt, A(128):76  
   Japan Basin E, A(127):258  
 copper  
   Site 794, B:1362  
   Site 795, B:1365  
   Site 797, B:1366  
   in sulfide deposits, A(128):21  
 core/log correlation  
   and ash layers, B:398, 401  
   and resistivity logging, B:398–400, 402  
   Site 798, B:1024–1025  
 correlation coefficients  
   for core XRF and geochemical log oxides, B:1029  
   in FMS data, B:1040–1041, 1046  
 cristobalite. *See* opal-CT  
 cross correlation, of FMS logs, B:1046  
 crest, lower, structure, B:1342–1343  
 crust, oceanic  
   electrical resistivity structure, B:1351  
   Japan Basin, A(127):73  
   Yamato Basin, A(127):73, B:928  
 crustal anisotropy, and regional stress field, B:1114–1115  
 crustal extension, in Yamato Basin, B:1345–1346  
 crustal structure  
   3.5 km/s layer, meaning of, B:1342  
   4.5 km/s layer thickness variation, B:1120  
   anisotropy, B:1110–1115, 1343  
   electrical resistivity, B:1343  
   and heat flow, B:1302–1303  
   Honshu N shelf, A(128):73  
   Japan Arc, A(128):74  
   Japan Basin, A(127):9, 79, 176, A(128):9–11, 70–72, 74; B:924, 1075, 1083, 1086, 1104, 1107, 1304–1305, 1311–1314, 1341–1346  
   Japan Basin E, A(127):251–252  
   Japan Basin N, A(127):176  
   Japan Sea, A(128):6–7, 9–11; B:1075, 1083, 1104, 1107, 1311–1314, 1333, 1341–1346  
   Japan Sea E, A(128):9–11  
   Japan Trench, A(128):74  
   Kita-Yamato Bank, A(128):248  
   Kita-Yamato Trough, A(128):245–249  
   lateral heterogeneity, B:1110–1112  
   lower crest, B:1342–1343  
   and ocean bottom seismometer data, B:1075  
   Oki Ridge, A(128):127–128  
   Oki Trough, A(128):130  
   Okushiri Ridge, A(127):251–252  
   and paleobathymetry, B:1201  
   and plate reconstruction geometrical constraints, B:1311–1314  
   Takuyo Bank, A(128):249  
   and tectonic history, B:1318–1319  
   and temperature, B:1343  
   Tsushima Basin, B:1311–1314  
   velocity structure, B:1078–1083  
   Yamato Bank, A(127):330, A(128):248

Yamato Basin, A(127):9, 79, 176, A(128):9–11, 24, 70–73, 127–128, 130, 248; B:837, 899, 1083, 1086, 1104, 1107, 1304–1305, 1311–1314, 1341–1346  
 Yamato Basin N, B:1075–1106  
 Yamato Basin S, A(127):329–330, 333; B:1083, 1104–1106  
 Yamato Rise, A(127):9, 329–330, 333, A(128):9–11, 245–249; B:1075  
 Yamato Seamount Chain, A(128):128  
 crystallization age, Site 794, B:821  
 Cupressaceae  
   Site 797, B:490  
 cut fluorescence. *See* fluorescence, cut cycladophorids  
   Site 794, B:307  
 cyclopentane, Site 799, A(128):321–322  
 D-phosphate, Site 799, B:38–39  
 dacite, ash layers, B:793  
 Daijima-type flora, Japan Sea, B:249, 486  
 dark/light cycles, B:1221. *See also* lithofacies, cyclic  
   and bacteria, B:772  
 biogenic component and detrital grain size compared, B:584, 588  
 biogenic opal abundance, B:432  
 burrows, B:564–565, 569  
 C/N ratio, B:431  
 carbon and sulfur analyses, B:579  
 carbonate content, B:431–432, 586–588  
 characterization, B:563  
 clay minerals, B:569  
 color, B:581  
 core continuity and depth corrections, B:578  
 correlations at Site 798, B:562–563  
 dark layers, B:580, 584, 591–594, 597–601  
 density, B:568  
 depositional environment, B:590–594  
 and dropstones, B:581  
 excess silica content, B:1241, 1253  
 first-order rhythms, B:446–447, 559, 563  
 geochemistry of second-order cycles, B:569  
 gray value, B:570–571, 579, 581–584, 586–587  
 hydrogen index, B:431–432  
 and interbedded ash layers, B:581  
 isotopic analyses, B:569–571  
 and Japan Sea circulation, B:574–575  
 layer thickness and organic carbon content correlated, B:594  
 major and minor elements, B:568, 570–571  
 microfossils, B:568  
 and Milankovitch cycles, B:446–447, 571–575  
 and molecular organic fossils, B:670, 675  
 and monsoons, B:574–575  
 nature of basal boundaries, B:569  
 nitrogen content, B:431  
 occurrence, B:562–563  
 opal content, B:446, 569  
 organic carbon, B:430–432, 568, 586–588, 671  
 organic matter, B:446, 564, 668–669  
 origin of cyclicity, B:577–601  
 origin of detrital component, B:594–595  
 oxygen index, B:431–432  
 oxygen isotopes, B:431, 588  
 and oxygenation conditions, B:432, 590–591  
 paleoceanographic significance, B:574–575  
 paleoenvironmental interpretation, B:419–421  
 productivity, B:431–432, 591–594  
 and revised lithostratigraphy, B:1229–1232  
 second-order rhythms, B:446, 559, 563–569  
 sediment composition, B:579, 582–584  
 sedimentological observations, B:569

dark/light cycles (cont.)

## SUBJECT INDEX

Site 795, B:1257  
 Site 797, B:574  
 Site 798, B:393, 421, 446, 458, 559-576  
 Site 799, B:571-574  
 Sites 794, 795, and 797 cycles correlated, B:577-601  
 Sites 797-799 compared, B:574  
 structure/composition, B:564-565  
 sulfur, B:586-588  
 synthetic sequence, Site 798 drillholes, B:566-567  
 thickness, B:581  
 third-order rhythms, B:559, 563  
 trace metals, B:1361  
 X-ray diffraction analyses, B:579, 584-588  
 dark/light sediment cycles. *See* dark/light cycles; lithofacies, cyclic  
 debris flow  
 consolidation, B: 1125  
 seismic expression, B:1140-1141, 1143  
 Site 799, A(128):267  
 declination. *See* magnetic properties  
 deformation, soft-sediment  
 and lithologic boundaries, A(128):142ff  
 Site 798, A(128):141ff  
 Site 799, A(128):241  
 deformation rate analysis (DRA), and stress field, B:1048-1049  
 density. *See also* physical properties  
 acoustic impedance, B:1137-1138  
 basement, A(127):227  
 and clay content, B:1278-1279  
 corrected analyses, B:990-1015  
 in dark/light cycles, B:568  
 data correction methods, B:987  
 data quality, A(127):127, 228-229  
 and diatom content, B:1278-1279  
 errors in wet volume measurement, B:985-986  
 Japan Sea, B:1278-1281  
 and loss on ignition, B:1283  
 and opal-A to opal-CT transition, B:18-19, 21, 1146-1147, 1278-1279, 1288  
 Site 794, A(127):127-134, A(128):102-103, 105; B:1050, 1280-1281  
 Site 795, A(127):222-228, 230, 233; B:1280-1281  
 Site 796, A(127):290-295; B:1280-1281  
 Site 797, A(127):383-390; B:1280-1281  
 Site 798, A(128):180-181, 197-198, 202-205, 208-210; B:1280-1281  
 Site 799, A(128):325-326, 345-349, 354-356; B:1137-1138, 1140, 1280-1281  
 vs. loss on ignition, B:1291  
 vs. opal abundance, B:402  
 vs. porosity, B:25, 27, 986, 989, 1281, 1290  
 vs. water content, A(127):233, 395  
 density, logging  
 Site 794, A(127):139-142, 144-146  
 Site 795, A(127):232, 237  
 Site 796, A(127):302-305, 319-320  
 Site 798, A(128):185-187; 212, 229-232; B:399, 401  
 Site 799, A(128):332-338, 362-364, 383-385, 389-392  
 vs. gamma ray, logging, A(128):215  
 density flow, of blue tuff, B:123-126  
 deposition rate. *See* opal flux; sediment accumulation rate; sedimentation rate; terrigenous flux  
 depositional cycles. *See* lithofacies, cyclic  
 desiccation, as problem for visual microanalysis, B:57  
 detrital accumulation rate. *See* terrigenous flux

detrital minerals  
 eolian origin, B:1238-1239  
 in Japan Sea sediments, B:1241-1249  
 deuterium isotopes. *See* isotopes, deuterium  
 dewatering  
 Site 798, B:1176-1178  
 Site 799, B:1178  
 diachrony, species, of calcareous nannofossils, B:164  
 diactional monoxons  
 Site 795, B:542  
 diatom-bearing clay. *See* clay, diatom-bearing  
 diatom clayey siltstone. *See* siltstone, diatom clayey  
 diatom claystone. *See* claystone, diatom  
 diatom number. *See also* diatoms  
 in dark/light cycles, B:579, 588  
 Site 797, B:593  
 diatom ooze. *See* ooze, diatom  
 diatomaceous clay. *See* clay, diatom  
 diatoms. *See also* biostratigraphic zonations; biostratigraphy; microfossils, siliceous; opal-A; silica, biogenic  
 abundance and opal-A to opal-CT transition, B:1230-1232  
 abundance and preservation, A(127):97, 192-193, 271-272, 351, A(128):28, 159-161, 299-303; B:312, 314, 344-349, 352, 593, 1219  
 abundance and Tsushima Current inflow correlated, B:591  
 age constraints, A(127):271, A(128):69; B:1335  
 BSEM images, B:554  
 in carbonate concretion, A(127):351  
 in dark/light cycles, B:568, 584  
 datum levels, B:251, 258-259  
 and density, B:1278-1279  
 diachronous occurrences, B:309  
 dissolution, B:309-316, 341-342, 353, 1220  
 foraminifers correlated, B:188-189, 222-223  
 freshwater, A(128):249  
 frustule structure, A(127):137  
 internal porosity, B:1277  
 Japan Sea, B:363  
 Kita-Yamato Bank, A(128):249  
 Leg 127, A(127):19-20  
 Leg 128, A(128):28  
 Miocene/Pliocene boundary, A(127):97, 193, 272, 351  
 morphotaxa recognized in BSEM images, B:548  
 and opal-A to opal-CT transition, B:3  
 as paleoenvironmental indicator, B:341-342  
 Pliocene/Pleistocene boundary, A(127):97, 351  
 productivity, A(127):274, A(128):122; B:342  
 Quaternary/Pliocene boundary, A(127):192, 271  
 reworked, A(127):97, 192-193, 273; B:250  
 and sediment velocity, A(127):133  
 SEM photograph, B:28-29, 1292-1293  
 and silica abundance, A(127):362-363  
 Site 794, A(127):97, A(128):99; B:264-269, 284, 344, 1221  
 Site 795, A(127):192-193; B:276-281, 286, 344-345  
 Site 796, A(127):271-273; B:270-275, 285  
 Site 797, A(127):351; B:252-258, 283, 345, 593, 1223  
 Site 798, A(128):124, 159-161; B:361  
 Site 799, A(128):299-303; B:34-35, 361  
 as stratigraphic markers, B:1220  
 and Tsushima Strait closure, B:1219-1220  
 upwelling-specific, A(127):267, 271, 349  
 zonations, A(127):19-20, 97, 192-193, 271-273, 351, A(128):159-161, 299-303; B:249-251, 282, 361-362, 1219, 1225-1226  
 digital downhole seismometer. *See* downhole seismic experiment  
 dikes  
 and crustal heterogeneity, B:1113-1114  
 Site 797, B:942-943  
 discoasters. *See also* nannofossils, calcareous  
 Site 797, A(127):354  
 dissolution  
 of biogenic silica, B:344  
 of calcareous nannofossils, B:166  
 and geologic age, B:310  
 Leg 128, A(128):28, 30  
 and opal abundance, B:310-311  
 of planktonic foraminifers, B:465  
 of radiolarians, A(127):198-199; B:295-297  
 of siliceous microfossils, A(127):169; B:353  
 Site 798, A(128):162, 165  
 Site 799, A(128):310  
 and temperature, B:310  
 dissolved oxygen. *See* oxygen, dissolved  
 DIT. *See* resistivity, logging  
 dolerite  
 alteration, B:883-889  
 isotopic data, B:884-888  
 petrography, B:892  
 sills, A(128):86  
 Site 794, B:849, 869  
 vesicles, B:887-888  
 dolerite, aphyric, A(128):75  
 radiometric age, B:822-826  
 Site 794, A(127):122-125, A(128):68, 88-90; B:908, 1339  
 Site 797, A(127):371-375, 380-381  
 zoned vein alteration, B:886  
 dolerite, olivine, Site 794, A(128):69, 91  
 dolerite, olivine microphyric, Site 794, A(128):68, 89  
 dolerite, plagioclase phryic  
 Site 794, A(127):122-125, A(128):75  
 Site 797, A(127):371  
 dolerite, plagioclase-pyroxene, phryic, leucocratic, Site 794, A(128):68, 88  
 Dolgorae-1 Well  
 paleobathymetry, B:1210  
 subsidence plot, B:1208  
 dolomitic  
 Site 795, A(127):186  
 Site 797, A(127):343  
 dolomite. *See also* ankerite; carbonate, diagenetic; proto-dolomite; siderite  
 beds, A(128):275-276  
 calcium and magnesium abundances, A(127):363-364  
 and Ce anomaly, B:92  
 cementing sandstone, A(128):276  
 and cerium concentration, B:90-91  
 composition, B:82-85  
 in FMS, A(127):307, A(128):184, 368  
 formation, A(127):23; B:85, 93  
 Funakawa Formation, B:75  
 in geochemical logs, B:1399  
 and interstitial water chemistry, B:1265  
 isotopic composition, B:83  
 layers, A(128):274-275  
 in logs, A(127):302, A(128):34, 245, 332-333, 340-341  
 major elements, B:1241

## SUBJECT INDEX

feldspar

Monterey Formation, B:75  
 and Mossbauer parameters, B:740-741  
 nodules, A(127):113, 186-187, 199, 343,  
 A(128):275  
 occurrence, B:80  
 Onnagawa Formation, B:75  
 oxygen isotopes vs. degree of recrystallization, B:83, 89  
 paleoenvironmental implications, B:94  
 patches, A(128):274  
 petrography, B:81  
 precipitation, B:92-93  
 recrystallization categories, B:81, 92-93  
 rhombs, A(128):273-274  
 in sandstone, B:141-143  
 and sedimentation rate, A(128):148  
 seismic stratigraphy, A(128):34, 125; B:1141, 1143  
 SEM photographs, B:98  
 shallowest recurrence, A(128):273  
 Site 795, A(127):187-188  
 Site 796, A(127):266  
 Site 798, A(128):137-138, 147-150, 174  
 Site 799, A(128):240, 260, 272-278, 289, 354; B:38, 40-41, 75-98, 611-612  
 on synthetic seismogram, A(128):184  
 texture, B:81  
 X-ray diffraction analyses, B:79-80  
 dolomite, siliceous, Site 799, B:43  
 dolomite nodules. *See* dolomite  
 dolomitization  
 and alkalinity, A(127):362  
 and interstitial water analyses, B:607  
 Site 799, A(128):320-321  
 downhole seismic experiment. *See also* air-gun profiling experiment; ocean bottom seismometer  
 air-gun profile locations, A(128):112; B:1070, 1077, 1109, 1159  
 air-gun reflection records, B:1076, 1079  
 air-gun shooting, A(128):113-114; B:1162-1163, 1167  
 and anisotropic models, B:1117  
 assignment of cable conductors, A(128):109  
 battery specifications/wiring, B:1067  
 broadband digital telemetry package, B:1064 and crustal structure, B:1342  
 data/power link, B:1064-1065  
 data selection, B:1067  
 digital telemetry of data, B:1063  
 equipment used, A(128):69-70  
 first arrival traveltimes, B:1089  
 frequency response of sensors, B:1158-1159, 1161  
 HIG and NORDA experiments, B:1061  
 horizontal sensor azimuth, B:1159  
 installation, A(128):69-70, 110; B:1067-1068  
 local earthquakes observed, B:1161, 1165-1166  
 logging cable, A(128):106-109; B:1063  
 navigation during, B:1107-1108  
 noise levels, B:1162-1163  
 objectives, A(128):18, 22-23, 83, 106; B:1061, 1157  
 OBS-array locations, B:1070, 1077, 1107, 1109, 1159  
 OBS records, B:1077-1078, 1084, 1160  
 OBSs deployed during real-time experiment, A(128):111, 113; B:1076, 1159  
 off-line experiment, B:1158-1162  
 operations, A(128):68, 85, 113  
 power supply and consumption, B:1066, 1159-1160

pressure housing, B:1063-1064  
 ray tracing results, B:1090-1103  
 real-time experiment, A(128):110-111; B:1068, 1077, 1157-1158  
 recorder unit, A(128):109-110  
 recovery unit, B:1066-1068, 1073  
 seafloor data recorder, B:1065-1066  
 seafloor package deployment, A(128):111; B:1068, 1070-1071  
 seafloor recording/recovery unit specifications, A(128):109  
 seismometer records obtained, B:1077-1078, 1080-1083  
 seismometer specifications, A(128):109  
 seismometer system layout, A(128):84, 114; B:1063, 1161  
 sensors, A(128):106-109; B:1063-1064, 1072, 1158-1160  
 ship positioning during experiment, B:1076-1077  
 Site 794, A(128):36, 68-70, 106ff, 117; B:1061-1073, 1075-1078, 1107-1121, 1157-1171  
 system design, A(128):106-110; B:1061-1063  
 technological difficulties, A(128):106; B:1157  
 teleseismic observations, B:1162, 1167-1168  
 temperature change during, B:1160-1161  
 tilting during, B:1160-1161  
 traveltime, offset distance and basement depth correlated, B:1112-1113  
 velocity power spectrum plot, A(128):115  
 velocity structure, one-dimensional, B:1078-1081, 1085-1086  
 velocity structure, two-dimensional, B:1081-1083, 1086, 1089  
 very long-period data (VLP), B:1159-1161, 1164  
 weather during, A(128): 113; B:1163, 1171  
 downslope transport  
 Site 794, A(127):103  
 Site 795, A(127):199  
 dropstones  
 in dark/light cycles, B:581  
 Japan Sea, B:1229-1230  
 Site 795, B:1256  
 dry-bulk density. *See* density  
 dust, eolian  
 Japan Sea, B:394, 401, 1238-1239  
 in logs, B:403  
 Oki Ridge, B:419  
 origin of dark/light cycle detrital component, B:594-595  
 in Pacific Ocean, B:394, 403  
 dust storms, in China and Mongolia, B:394  
 dysaerobia. *See* oxygenation conditions  
 dysaerobic conditions. *See* oxygenation conditions  
 earthquakes  
 Burma, B:1162  
 during downhole seismic experiment, B:1161-1163, 1165-1166  
 Japan Basin, A(128):74  
 Japan Basin E, A(127):255  
 Japan Sea, A(128):76  
 Japan Sea E, A(128):11  
 Japan Sea quake (1983), B:1047, 1062, 1161  
 and sediment consolidation, B:1130  
 and stress field, B:1054, 1120  
 and vein formation, B:1180-1181  
 East Pacific Rise, sulfide deposits, A(128):21  
 ebridians, B:1220  
 Site 798, A(128):162; B:241-244  
 Site 799, A(128):305; B:241, 246-247  
 zonal data, B:1225-1226  
 zonation, B:237, 239-241  
 eccentricity frequencies, in dark/light cycles, B:446-447, 571, 574-575  
 electrical resistivity, crustal, B:1343, 1351  
 electrical resistivity experiment, B:1351-1359  
 cable used, A(128):114  
 current used, A(128):113, 114, 116; B:1352  
 data collection/operations, B:1352-1354  
 electrodes used, A(128):114, B:1352  
 logging resistivity compared, B:1354  
 methods, A(128):111-115  
 noise source, B:1352  
 objectives, A(128):18, 23, 83, 111  
 operations, A(128):68, 70, 85, 115-116  
 schematic illustration of, A(128):115  
 sensor cable configuration, A(128):116; B:1351-1353  
 shooting ship positions, B:1352  
 signal amplitude, A(128):116; B:1352, 1359  
 signal forms observed, B:1355-1358  
 Site 794, A(128):36, 68, 70, 111-117; B:1343  
 source current time variation, B:1354  
 vertical electrical field ( $E_z$ ) observed, B:1352  
 element mobility, Site 794, B:916  
 ellipticity, of borehole, Site 799, A(128):339-340, 368  
 eolian dust. *See* dust, eolian  
 epidote, in ash layers, B:1379  
 ethane  
 in gas hydrate, A(127):288-289  
 Site 794, A(127):119  
 Site 795, A(127):174, 213-216, 220  
 Site 796, A(127):251, 287  
 Site 797, A(127):368  
 Site 798, A(128):125, 175-176, 187  
 Site 799, A(128):244-245, 321-322, 339  
 ethylene  
 Site 794, A(127):119  
 Site 796, A(127):287  
 Eu anomaly  
 in crest-derived materials, B:731  
 and rhyolitic volcanic activity, B:731-732  
 and siliceous input, B:691  
 Site 794, B:682-683, 916-917  
 Site 795, B:683, 918  
 Site 797, B:688-691, 920  
 Site 798, B:729, 735  
 Site 799, B:729, 736  
 vs. Ce anomaly, B:737  
 Eu/Eu\*. *See* Eu anomaly  
 Eurasian Plate, A(127):5-6, A(128):9  
 europium. *See also* Eu anomaly  
 Site 798, B:1368-1369  
 euxinic conditions. *See* oxygenation condition  
 excess pore-water pressure. *See* consolidation  
 excess silica. *See* silica, excess  
 experimental petrology. *See* petrology, experimental  
 extension, and fracture formation, B:1181  
 F (emplacement mode parameter). *See* magnetic properties  
 F-phosphate, Site 799, B:36-39  
 fabric, of clay minerals, B:35-36, 38  
 n-fatty acids, Sites 798 and 799, B:669-671  
 faults. *See also* microfaults; normal faults; reverse faults; thrust faults  
 Site 794, B:1182-1183  
 with slickensides, A(128):92; B:1185, 1190  
 and slumps, B:1176  
 and stress field analysis, B:1052  
 feldspar. *See also* albitization  
 albitization, B:107, 136-137  
 in ash layers, B:1379, 1388, 1392

feldspar (cont.)

chemical composition, B:106, 136, 139  
and sandstone, B:105–107  
Site 798, B:411–415, 417  
twinning characteristics, B:107  
feldspar, alkali, in ash layers, B:1379  
feldspar, alkaline, in ash layers, B:1392  
feldspar, calcic plagioclase, B:872  
feldspar, orthoclase, B:147  
feldspar, plagioclase  
abundance, B:1235–1244  
albitized, B:105–106, 109, 137–138  
altered, B:141, 144–145, 884–885  
anorthite, B:121–123, 126  
in ash layers, B:1379  
in blue tuff, B:117, 121–126  
in dark/light cycles, B:584  
in dolerite, A(128):88  
effect of water-undersaturated conditions, B:856  
experimental chemical compositions, B:867  
glacial/interglacial variation, B:588  
 $K(0-8)$  vs. mole percent An, B:851  
in melting experiment, B:861–862, 895  
microprobe analyses, B:124–126  
and parent magma, B:856  
and potassium, B:856, 869  
and radiometric age, B:785–786  
replaced by anhydrite, B:148  
in sandstone, B:104  
Site 794, B:850–851  
Site 795, B:850–851  
Site 796, B:107, 136  
Site 797, B:136, 850–851  
Site 799, B:137, 140  
textures, B:136  
vs. illite, B:1245  
vs. quartz, B:1245, 1252  
water-undersaturated conditions, composition effect, B:872  
feldspar, potassium. *See also* feldspar, orthoclase  
altered to kaolinite, B:141  
in ash layers, B:1379, 1390  
in sandstone, B:104  
Site 797, B:105, 137  
Site 799, B:137  
feldspar, sodic, B:1379  
feldspar accumulation rate, Site 798, B:411–420  
feldspathic arenite. *See also* arenite, feldspathic; sandstone  
Site 799, B:105  
ferrobasalt, Site 797, B:861  
fissility  
Site 798, A(128):146ff  
Site 799, A(128):272  
flame structures, Site 799, A(128):268–269; B:1178  
flora, brackish-water, Japan Sea, B:249  
fluid flow  
along thrust faults, A(127):247  
evidence for, A(127):295  
heat transport by, A(127):301  
Site 799, A(128):321  
fluorescence, cut, Site 799, A(128):323; B:623  
fluoride  
and francolite, B:67  
in interstitial water, B:64, 66  
uptake rates, B:65–66  
FMS. *See* formation microscanner  
folds  
along Okushiri Ridge, A(127):255–256  
Site 799, A(128):268  
foliation, clay mineral fabric, B:38  
Foram Sharp Line (FSL)

## SUBJECT INDEX

Japan Sea, B:493, 506  
timing of, B:526, 528–529  
foraminifers  
abundance and preservation, A(127):99–100, 195–198, 273–274, 355–356, A(128):258; B:188  
assemblages, A(127):20, 100–101, 195–198, 273–274, 355–356  
BSEM images, B:548, 557  
and climate fluctuation, A(127):195–197  
diatom zones correlated, B:188–189, 222–223  
morphotaxa recognized in BSEM images, B:548  
paleodepth, A(127):198, 274, 355  
paleoenvironment, A(127):199  
reworked, A(127):195, 273, 355  
Site 794, A(127):99–101, A(128):99; B:187–193  
Site 795, A(127):195–198; B:187–197  
Site 796, A(127):273–274; B:187–200  
Site 797, A(127):355–356; B:187–207, 1223  
Site 799, B:78  
textulariid taxa, A(127):195–198, 356  
and water temperature interpretation, A(127):355  
zonations, B:200–201  
foraminifers, agglutinated  
assemblages, B:188–189  
Site 794, B:500, 528  
Site 795, B:501–502  
Site 797, A(127):355–356; B:503–504  
Site 798, B:373  
foraminifers, benthic, B:1220–1222  
abundance and preservation, A(128):30, 166, 311–312  
assemblages, A(128):166, 311–312; B:188–189  
in dark/light cycles, B:568  
deposition depth, A(128):354  
depth distribution, B:379  
Japan Basin, B:208  
Japan Sea, B:365, 1208  
in laminated sediments, B:551  
Leg 128, A(128):30  
oxygen isotopes, B:443–444  
oxygenation, A(128):166, B:1221  
and paleobathymetric analysis, B:1201  
as paleodepth indicator, A(128):166; B:208, 1220–1221  
preservation, B:366  
Site 794, A(127):101; B:210–215, 495–496, 500, 506–507, 516–517  
Site 795, A(127):197–198; B:500–505, 518–521  
Site 796, A(127):273–274  
Site 797, A(127):355–356; B:210–215, 502–504, 508–515, 522–528  
Site 798, A(128):166; B:367–369, 373–375, 382, 443  
Site 799, A(128):311–312; B:370–373, 376–378  
species distribution, B:208  
Yamato Basin, B:189, 208  
zonation, B:383, 517, 521, 524  
foraminifers, planktonic, B:1220  
abundance and preservation, A(128):30, 165–166, 310–311; B:193, 201, 221, 460  
assemblages, A(128):310–311; B:188–189, 462–463, 466  
and basal sediment age, B:1335  
BSEM images, B:553  
cooling direction, A(128):165, 310–311; B:1220  
cooling ratios, B:188, 193, 208, 210–215, 218–219, 459, 463  
in dark/light cycles, B:568, 584  
dissolution, A(128):165; B:465  
faunal provinces, B:459–462  
gyre-margin fauna, B:459  
Honshu exposures, B:459  
Leg 127, A(127):20  
Leg 128, A(128):30  
monospecific assemblages, A(128):165  
and oxygen isotopes, B:442–443  
paleoclimatic interpretations, B:465–468  
polar-subpolar fauna, B:459, 462  
preservation in diatom-rich sediments, A(128):165  
Q-mode cluster analysis, B:188, 193–200, 208–209, 220, 224  
Site 794, A(127):99–101; B:193–200, 210–215  
Site 795, A(127):195–197  
Site 796, A(127):273–274  
Site 797, A(127):355–356; B:193–200  
Site 798, A(128):124, 165–166; B:442–443, 457–470  
Site 799, A(128):310–311  
subtropic fauna, B:459  
and surface-water temperature, B:187  
transitional fauna, B:459  
upwelling evidence, B:469  
forceps  
Site 795, B:543  
forests  
Japanese archipelago, B:320, 325, 334  
Site 798, B:317  
formation factor  
data quality, A(127):127, 222, 289–290  
Site 794, A(127):128–133, 137  
Site 795, A(127):224–228, 231  
Site 796, A(127):290–295  
Site 797, A(127):383–389, 391  
formation microscanner (FMS)  
ash-flow tuff, B:793  
ash layers, A(128):153, 339–340; B:398, 400, 1025  
chert, A:307; B:1039–1040, 1232  
and core recovery, B:1038–1040  
cross correlations, B:1040–1041, 1045  
cumulate layering in sill, A(128):104  
cyclicity in chert layers, B:1041–1044  
data processing, B:1038–1043  
depth alignments, B:1038–1040  
diagrams, B:1038–1039  
dolomite, A(127):307, A(128):368  
Fourier transform analysis, B:1043, 1046  
hole parameters, A(127):307, 399  
in interbedded sediments, A(127):399  
and intersite lithologic correlation, B:1233  
and Milankovitch cycles, B:1043–1044  
operations, B:1037–1038  
porcellanite, B:1040  
qualitative comparison of Site 794 and 797 logs, B:1039–1040  
signal processing, B:1040–1043  
of siliceous shale, B:1232  
Site 794, A(127):142, A(128):104ff, 109; B: 1037–1046  
Site 796, A(127):307, 310  
Site 797, A(127):399, 408; B:1037–1046  
Site 798, A(128):184–185  
Site 799, A(128):339–341  
Sites 794 and 797 correlated, B:1042–1044  
veins and fractures, A(128):104–106  
Fourier transform analysis, of FMS data, B:1043, 1046

- fractionation models, B:872–874, 881  
fractures  
  ages, A(128):92–93  
  basement, A(127):217, A(128):69, 88–92; B:1181–1184  
  in BHTV data, A(127):307, 399, 409  
  calcite-filled, A(127):188  
  fillings, A(128):91–92; B:1181  
  geometry, B:1181  
  origin, A(128):91–93  
  relative ages, A(128):93  
  Site 794, B:1182–1184  
  Site 798, A(128):143ff  
  and stress regime, B:1181  
fractures, calcite-filled, and normal faults, A(127):188  
frambooids, of pyrite, B:36  
francolite, occurrence, B:65, 67  
free-air gravity anomalies  
  Japan Basin, A(128):74  
  Japan Basin E, A(127):253, 255–256  
  Japan Basin N, A(127):176, 179  
  Japan Sea E, A(128):252  
  Kita-Yamato Trough, A(128):247  
  Oki Ridge, A(128):127, 133  
  Oki Trough, A(128):127  
  Site 794, A(127):81  
  Yamato Basin, A(128):127  
  Yamato Basin N, A(127):77, A(128):73  
  Yamato Basin S, A(127):330–331, 335  
  Yamato Rise, A(128):247  
frictional heating. *See* heating, frictional  
frontal boundary, oceanographic, Japan Sea, B:155–169  
Funakawa Formation, A(127):198, A(128):13  
  deposition, A(128):17  
dolomite, B:75  
foraminifers, B:187  
lithostratigraphy, A(128):354  
fungi spores, Japan Sea, B:485  
gadolinium, logging  
  Site 794, B:1416, 1420–1421  
  Site 796, B:1423  
  Site 797, B:1426–1427  
gamma ray logging  
  of ash-flow tuff, B:793  
  calibrated by strip samples, B:1367  
  and clay content, B:6  
  of clay minerals, A(127):393  
  and core-log depth correlation, B:1024–1025  
  and depth alignments, B:1038–1040  
  lithostratigraphy correlated, B:1416–1417  
  of Olduvai Subchron, B:403  
  of opal-A to opal-CT transition, B:22  
  and oxygen isotopes, B:403  
  paleoclimatic significance, A(128):188–192  
  power spectra, A(128):217, 369; B:407  
  responses, A(128):103–104  
  Site 607, oxygen isotopes correlated, B:405  
  Site 794, A(127):142, 147, 156–159, 164–167, A(128):103–104, 107–108, 119–120; B:1069, 1416, 1418–1419  
  Site 795, A(127):244–245  
  Site 796, A(127):303–306, 319–322; B:1416, 1422  
  Site 797, A(127):393–395, 400–402, 416–421; B:7, 1417, 1424–1425  
  Site 798, A(128):185–187, 212, 215–216, 225–232; B:399, 401, 1398–1400  
  Site 799, A(128):332–338, 362–364, 367, 369, 380–385, 389–397; B:1399, 1403, 1406–1407  
terrigenous content compared, B:1024–1025, 1028  
used for depth shifting of logs, B:1398, 1415  
vs. density logging, A(128):215  
gamma-ray spectroscopy tool (GST). *See* geochemical logging  
gas  
  and magnetic properties, A(127):275  
  sediment consolidation and, B:1129  
gas, thermogenic, Site 799, A(128):322  
gas chromatography. *See* pyrolysis-gas chromatography  
gas hydrate. *See also* clathrate  
  and boron content, B:639  
  bottom simulating reflector (BSR), B:1145, 1147  
  Japan Sea, A(127):24  
  and oxygen isotopes, B:52  
  Site 796, A(127):260, 288–289, 315  
  temperature stability, A(127):288, 299–300  
Gauss Chron  
  and diatom occurrence, A(128):160  
  and N Pacific diatoms, A(128):302  
  silicoflagellate and ebridian zones correlated, A(128):305  
  Site 795, B:970, 1223  
  Site 796, A(127):275  
  Site 797, B:973  
  Site 798, A(128):30, 124–125, 170; B:974  
  Site 799, A(128):303, 314; B:1224  
Gauss/Gilbert boundary  
  Site 794, A(127):109; B:970  
  Site 797, B:973, 1223  
  Site 798, A(128):30; B:1224  
  Site 799, A(128):314; B:975, 1224  
geochemical logging, B:1411–1427. *See also* specific elements analyzed  
  aluminum concentration calculation, B:1398, 1415  
  of ash layers, A(128):339  
  borehole diameter effect, B:1029–1030  
  with boron sleeve, B:1023, 1030  
  calibrated by strip samples, B:1367  
  core comparison, B:1398–1399  
  and core-log depth correlation, B:1024–1025  
  data reduction, B:1395–1399, 1415–1416  
  depth shifting, B:1398, 1415  
  drillpipe iron effect, B:1415  
  elemental weight fraction calculation, B:1398, 1415–1416  
  elements not measured, B:1395, 1415  
  errors in, B:1028–1030  
  industry and ODP data compared, B:1021  
  K, Th, and U calculation, B:1398, 1415  
  Kleiner-Hartigan diagrams, B:1030  
  normalization factors, B: 1029–1030, 1034–1035  
  in opal-A to opal-CT transition, A(127):396–399  
  operations, B:1411–1412  
  and orbital obliquity cycles, B:1021  
  oxide factors for closure normalization, B:1402  
  oxide percentage calculation, B:1398, 1416  
  porosity effect, B:1022–1024, 1029  
  post-cruise processing schematic, B:1397  
  problems in, B:1022–1024  
  relative elemental yield reconstruction, B:1396, 1415  
  Site 794, A(127):142, 150–153, 160–163; B:1416, 1420–1421  
  Site 796, A(127):306–309, 317–318; B:1416, 1423  
Site 797, A(127):396–399, 404–407, 412–415; B:1417, 1426–1427  
Site 798, A(128):187, 233–236; B:1021–1035, 1398–1399, 1401–1402  
Site 799, A(128):366, 386–388, 398–402; B:1399, 1404, 1408–1409  
tool components, B:1022, 1395–1396, 1412–1415  
uses of, B:1021  
X-ray fluorescence analyses compared, B:1025–1029, 1402, 1416–1417, 1420–1421, 1426  
geochemistry, inorganic, B:1261–1274. *See also* geochemical logging; geochemistry, sediment; interstitial water; specific inorganic geochemical measurements  
and ash alteration, A(127):22; B:1265  
and authigenic carbonates, A(128):276–277  
and bacterial degradation of organic matter, A(127):22  
and basement alteration, A(127):23, 205; B:1265–1267  
and biogenic silica diagenesis, B:1263, 1267  
and carbonate chemistry, B:1263–1265, 1267  
and carbonate formation, A(127):22–23, 205  
and clathrates, A(127):278  
of dark/light cycles, B:569  
geochemical intervals, Site 799, A(128):244, 318–321  
Japan Sea, A(127):22–23, A(128):31–32  
new metal-free squeezer, B:635, 648–650  
and organic matter diagenesis, B:1262–1263, 1267  
safety considerations, A(128):322–323  
shorebased analyses of Leg 127 samples, B:635–650  
and silica diagenesis, A(127):23  
Site 794, A(127):72–73, 107ff  
Site 795, A(127):174, 204ff  
Site 796, A(127):251, 278–285  
Site 797, A(127):324–325, 362ff  
Site 798, A(128):31–32, 125, 171ff; B:605–621  
Site 799, A(128):31–32, 244, 316ff, 326–327; B:605–621  
geochemistry, organic. *See also* hydrocarbons, volatile; interstitial gas; molecular organic fossils; Rock-Eval analyses; specific organic geochemical analyses  
of hydrocarbons, A(128):175–176, 321–322  
Japan Sea, A(127):23–24, A(128):32  
Rock-Eval analyses, A(127):114–119, 213, 285, 367, A(128):177  
Site 794, A(127):72–73, 111–119, 121  
Site 795, A(127):174, 209–217  
Site 796, A(127):251, 283–290  
Site 797, A(127):324–325, 365ff  
Site 798, A(128):32, 125, 175ff; B:423–437, 667–675  
Site 799, A(128):32, 244–245, 321ff; B:423–437, 623–633, 667–675  
van Krevelen plot, A(127):121, 219, 288, 367, A(128):196; B:629  
geochemistry, sediment. *See also* geochemical logging; major elements; minor elements; rare earth elements (REE); trace elements of bulk sediment, B:610–611, 613, 615–619  
of carbonates, B:611–612, 615  
of clay-size fraction, B:611, 613, 620–621  
major element molar ratios, B:616–619  
Site 795, B:705–717  
Site 798, B:610–612, 719–737  
Site 799, B:610–612, 719–737

Sites 794-797, B:677-695  
 trace element molar ratios, B:616-619  
 geophysical experiments. *See* downhole seismic experiment; electrical resistivity experiment  
 geosyncline, Japan Sea, A(128):11  
 Gilbert Chron  
 Site 796, A(127):275  
 Site 798, A(128):30  
 Site 799, A(128):303, 314, 316; B:975, 1224  
 glacial/interglacial cycles  
 and biogenic silica variation, B:591  
 carbonate variation, B:590  
 grain size variation, B:589  
 isotopic variations, B:442  
 and Japan Sea circulation, B:457-458, 608  
 and oceanographic front oscillations, B:167-168  
 and opal flux variation, B:446  
 and organic carbon variation, B:589  
 and oxygenation condition, B:590-591  
 and salinity variation, B:591  
 and sediment composition variation, B:588-590  
 sulfur variation, B:589-590  
 and terrigenous flux variation, B:446  
 glacial periods  
 and biogenic opal supply, B:404  
 interglacial periods, B:403-404  
 and loess deposition, B:403  
 glaciation  
 and Japan Sea, B:338, 342-343, 419-421  
 and opal, B:350, 404  
 and oxygen isotopes, B:403  
 and Pacific Ocean, B:528  
 and productivity, B:430  
 Site 796, A(127):274  
 Site 797, A(127):356  
 Site 798, A(128):164  
 Site 799, A(128):312  
 and Tsugaru Strait sill depth, B:351  
 and upwelling, A(127):271, 274  
 vegetation and, B:320  
 glass. *See* alkaline glass; calc-alkaline glass; glass, volcanic; peralkaline glass  
 glass, volcanic  
 andesitic magma series, B:1378  
 chemistries, B:1374-1379  
 dissolution, B:1375, 1389  
 and original magmatic composition, B:1377  
 in sandstone, B:104  
 shoshonite series, B:1378  
 glass shards. *See* shards, glass  
 glauconite  
 allochthonous, B:67, 74  
 autochthonous, B:38, 67, 74  
 in diagenetic sequence, B:39  
 formation, B:71  
 microprobe analyses, B:67, 70  
 and revised lithostratigraphy, B:1232  
 Site 794, B:1259  
 Site 795, A(127):189  
 Site 796, A(127):265  
 Site 797, A(127):344  
 Site 798, B:63-74  
 Site 799, A(128):260, 264; B:38, 63-74  
 glauconitic sand. *See* sand, glauconitic  
 gneiss  
 Oki Islands, A(128):127  
 Goban Spur, inorganic nitrogen, A(127):285  
 gold  
 Site 798, B:1368-1369  
 in sulfide deposits, A(128):21

GPIT. *See* formation microscanner  
 grain density. *See* density  
 grain shape, of siliceous claystone, B:38  
 grain size  
 of ash layers, A(128):150; B:791  
 of blue tuff, B:117  
 of dolerite, A(128):90-91  
 glacial/interglacial variation, B:589  
 of magnetic minerals, B:949-951  
 magnetic properties affected, B:962  
 and permeability, B:1127-1128  
 sandstone, B:100, 104  
 of siliceous claystone, B:38  
 Site 794, A(127):94, 96, 127  
 Site 795, A(127):189, 192  
 Site 797, A(127):346; B:593  
 Site 798, B:1132  
 Site 799, B:1132  
 grainstone, foraminifers, Site 798, B:551  
 Gramineae  
 Site 794, B:491  
 granite  
 Kita-Oki Bank, A(128):127  
 Korea, B:111  
 Oki Islands, A(128):127  
 and sandstone provenance, B:108  
 Yamato Rise, A(128):248-249  
 GRAPE density. *See* density  
 gravity gliding. *See* gravity-induced structures; slumps  
 gravity-induced structures, Sites 798 and 799, B:1175-1176  
 gray analyses. *See* gray value  
 gray value  
 of dark/light cycles, B:570-571, 579, 581-587  
 and organic carbon content, B:589  
 power spectra, B:574  
 Site 797, B:592  
 time-series analyses, B:571  
 Green Tuff, A(128):9, 73  
 and abiogenic origin of methane, B:749  
 deposition, A(128):17  
 Honshu, A(128):127-128  
 Oki Islands, A(128):128  
 Oki Ridge, A(128):127-128  
 and strontium isotopes, B:644  
 Yamato Rise, A(128):249  
 groundmass  
 altered to clay, B:885-886  
 in basalt and dolerite, A(128):88-91  
 in basaltic basement, B:892  
 chemical composition, B:886  
 microphotograph, B:859  
 GST. *See* geochemical logging  
 Guaymas Basin, basalt emplacement, B:1337  
 Gulf of Aden, calcareous nannofossils, B:173  
 Gulf of California, diagenetic carbonates, B:75  
 Gulf of Mexico, microbiological studies, A(128):178  
 hafnium  
 mobility during alteration, B:911  
 Site 795, B:707  
 Site 798, B:1368-1369  
 Site 799, B:724  
 Hannuoba basalt, and subcontinental lithosphere, B:811  
 harmotome  
 in ash layer, B:1393  
 and volcanic glass alteration, B:1377-1378  
 headspace analyses. *See* hydrocarbons, volatile heat flow  
 BSR-derived and probe data compared, B:1148, 1154  
 and crustal radiogenic heat, B:1302-1303  
 estimated from opal-A/opal-CT BSR, B:1156  
 and fluid flux through Toyama Deepsea Fan, B:1148  
 helium isotope ratios correlated, B:747  
 Japan Basin, A(128):71-74, 246; B:1302, 1345  
 Japan Basin E, A(127):252-253  
 Japan Basin N, A(127):174  
 Japan Sea, A(127):26-27, A(128):14; B:1297-1298, 1301-1304, 1375  
 Kita-Oki Bank, A(128):127, 131  
 Kita-Yamato Trough, A(128):246  
 and lithospheric stretching models, B:1304-1305  
 Oki Ridge, A(128):125, 131; B:1302  
 Oki Trough, A(128):131  
 Okushiri Ridge, A(127):30, 300-301; B:1303, 1307  
 and opal-A/opal-CT bottom simulating reflector, B:1147-1148, 1150  
 and sedimentation rate, B:1147-1148, 1302  
 Site 794, A(127):73, 137  
 Site 795, A(127):174, 230-232  
 Site 796, A(127):247, 251, 299-300, 314  
 Site 797, A(127):325, 390  
 Site 798, A(128):125, 213-214  
 Site 799, A(128):338, 365  
 sources, A(127):251  
 topography effect, A(127):300  
 vs. basement age, B: 1305  
 vs. opal-A/opal-CT BSR 2-way traveltimes, B:1152-1153  
 Yamato Basin, A(127):73, 82, 330, A(128):127, 138, 246, 250; B:1147-1148, 1150, 1302, 1345  
 Yamato Basin N, A(128):71-73, 75  
 Yamato Basin S, A(127):332  
 Yamato Rise, A(128):246, 250; B:1302-1303  
 heating, frictional, along thrust faults, A(127):247  
 heavy minerals, and rare earth elements, B:691  
 heavy rare earth elements (HREE), and elemental ratios, B:693, 730-732  
 helium, types of, B:748-749  
 helium isotopes. *See* isotopes, helium  
 hematite  
 at Brunhes/Matuyama boundary, B:972  
 in sandstone, B:144  
 n-heptane, Site 799, A(128):321-322  
 Hess Rise, diatoms, B:360  
 heulandite, Site 797, B:139  
 n-hexane, Site 799, A(128):321-322  
 HI. *See* hydrogen index  
 Hidaka Shear Zone, and Japan Sea tectonic evolution, B:1314-1315  
 high-field-strength elements (HFSE)  
 alteration and, B:911  
 mobility, B:911  
 Site 795, A(127):213  
 Hokkaido, A(127):14, 251  
 basement, A(127):177-178  
 depositional history, A(127):349  
 rock outcrops, A(127):254  
 sandstone provenance, B:110  
 tectonic setting, A(128):75-77  
 terrigenous sediment source, A(127):307  
 Hokkaido W-Honshu N thrust belt, convergence and compression along, A(128):76  
 Honshu, A(127):325, A(128):70  
 basement, A(128):73  
 crustal structure of shelf, A(128):73  
 diatom abundance, B:316  
 Green Tuff, A(128):73, 127-128

Kuroko sulfide deposits, A(128):239, 251  
lithostratigraphy, A(127):1415, A(128):19  
paleobathymetry, B:1200  
planktonic foraminifers, B:459  
rock outcrops, A(127):333–334  
rotation, A(128):9  
tectonic setting, A(128):75–77; B:1316  
uplift, A(128):158  
volcanic events, B:1316

Honshu arc  
compressional stress field and, B:529–531  
diatoms, B:360  
geologic events and, B:536  
movement of, B:528  
paleodepths, B:526

hopanoid acid, Sites 798 and 799, B:669, 671  
hornblende, in ash layers, B:1379  
hotspot activity, in Yamato Basin, B:1345  
HREE. *See* heavy rare earth elements  
Huang Ho River, and freshening of Japan Sea, B:442  
hydrate. *See* gas hydrate  
hydrocarbons, free (S<sub>0</sub>)  
chromatograms, B:633  
Site 799, B:625, 633  
vs. organic carbon, B:631

hydrocarbons, generatable (SZ), Site 799, B:625  
hydrocarbons, heavy, Site 799, B:628–629  
hydrocarbons, high-molecular-weight, Site 799, A(128):324  
hydrocarbons, liquid, Site 799, A(128):323  
hydrocarbons, nonindigenous, Site 799, B:627  
hydrocarbons, total pyrolytic yields (S<sub>0</sub> + S<sub>1</sub>), Site 799, B:624  
hydrocarbons, volatile. *See also names of specific volatile hydrocarbons*  
and opal-A to opal-CT transition, A(128):323  
Site 794, A(127):119  
Site 795, A(127):213–216  
Site 796, A(127):285–289  
Site 797, A(127):368, 378  
Site 798, A(128):32, 125, 175–176, 185–189  
Site 799, A(128):32, 244–245, 321–322, 330–331, 333  
thermogenic origin, A(128):322  
hydroclast, vesicular  
in blue tuff, B:117, 130  
and pressure compensation level, B:119  
hydroclastic shards, in tuffaceous clayey siltstone, A(128):89–90  
hydrogen  
and geochemical logging, B:1023–1024  
organic, Site 799, B:629  
hydrogen index (HI)  
chromatograms compared, B:627  
in dark/light cycles, B:431–432  
and organic matter preservation, B:628  
Site 794, A(127):115–118  
Site 795, A(127):213, 219  
Site 796, A(127):285  
Site 797, A(127):367  
Site 798, A(128):177, 196; B:425–426, 668–669  
Site 799, B:427, 429, 624, 668–669  
vs. marine maceral content, B:670  
hydrogen isotopes. *See* isotopes, hydrogen  
hydrogen sulfide, shallow presence, A(127):259–260  
hydrothermal activity, and Ce anomaly, B:729  
hygromagnaphile elements, spidergram, B:783  
igneous complex. *See* basement  
illite  
glacial/interglacial variation, B:588

in Japan Sea sediments, B:1235–1244  
in sandstone, B:139  
Site 797, B:592  
Site 798, B:411, 416, 418  
vs. kaolinite + chlorite, B:1245  
vs. Plagioclase, B:1245  
vs. smectite, B:139, 1245  
ilmenite, in ash layers, B:1379  
Inazumi Formation, A(128):129  
inclination. *See* magnetic properties  
inclination tool (GPIT). *See* formation microscanner  
incompatible elements. *See* rare earth elements; trace elements; X-ray fluorescence analyses  
index properties. *See* physical properties; *specific index properties*  
Indian Ocean  
benthic foraminifers, B:505  
calcareous nannofossils, B:173  
heat flow anomalies, A(127):301  
induction tool. *See* resistivity, logging  
inertinite, Sites 798 and 799, B:670  
intensity. *See* magnetic properties  
interglacial periods. *See* glacial periods  
interstitial gas  
and helium isotope ratios, B:747–751  
methane, B:749–750  
methane vs. helium content, B:750  
Site 798, B:748  
Site 799, B:748  
interstitial water  
ash layer alteration and, B:646  
barium content, B:657–659  
basement alteration and, B:646  
basement rock and adjacent interstitial water  
elemental variations correlated, B:911  
boron content, B:642  
carbonate formation and, B:646  
D/H ratios, B:642  
and diagenetic reactions, B:699  
and fluid/solid exchange reactions, B:1261  
hydrogen isotopes, B:639, 641–643  
isotopic studies, B:49–56  
organic matter degradation and, B:646  
and oxygen isotope depletion, B:697–703  
oxygen isotopes, B:51–52, 54, 84–85, 639, 641–643, 699–702  
pH, B:94  
silica diagenesis and, B:646  
Site 798, B:607–610  
Site 799, B:607–610  
stable and radiogenic isotopic content, B:635–650  
strontium, B:643–646  
sulfate content, B:639, 657–659  
Inyo dike, and stress field, B:1323  
iron. *See also* Mossbauer spectroscopy  
alteration and, B:911  
in ash layers, B:1381  
in augite, B:851–853  
in basalt, B:870  
in carbonate minerals, B:722–723  
in dark/light cycles, B:569  
depth profiles, B:742  
in experimental augite, B:862–863  
high-spin Fe<sup>2+</sup> in aluminosilicates and carbonates, B:740–741, 744–746  
in Japanese volcanic outcrops, B:874  
in Japan Sea sediment, B:1236  
paramagnetic Fe<sup>3+</sup>, B:739, 741, 745–746  
silicate Fe<sup>2+</sup> content, B:744–745  
Site 795, B:706

Site 798, B:722, 1368–1369  
Site 799, B:90–91, 722  
in veins, B:1178  
vs. aluminum, B:732  
XRF and geochemical log analyses compared, B:1025–1029, 1031–1035  
iron, logging  
Site 794, B:1416, 1420–1421  
Site 796, B:1416, 1423  
Site 797, B:1426–1427  
Site 798, B:1401–1402  
Site 799, B:1404, 1408–1409  
iron/magnesium ratio, in basaltic basement rocks, B:893–894  
iron oxide. *See* iron  
iron sulfide, and magnetic properties, B:947  
island-arc tholeiite. *See* tholeiite, island-arc  
isochelae  
Site 795, B:543  
isochron plots  
Site 794, B:828–832  
Site 795, B:832–833  
Site 797, B:828–829  
isopentanes, Site 799, A(128):321–322  
isoprenoids, Site 799, B:628–629  
isothermal remanent magnetization (IRM). *See* magnetic properties  
isotherms, silica transitions, B:53  
isotopes, argon, in basement rocks, B:822–833  
isotopes, beryllium, and mantle source, B:815  
isotopes, boron  
adsorption/desorption processes, B:640–641  
basement alteration, B:1266  
in interstitial water, B:638–642  
Site 794, B:638–641  
Site 795, B:638–641  
Site 796, B:638–641  
Site 797, B:638–641  
isotopes, carbon  
and carbonate diagenesis, B:85–90  
in chlorite/saponite mixed-layer clays, B:888  
in dark/light cycles, B:569–571  
in interstitial water, B:607  
and methanogenesis, B:88–90  
and organic carbon decomposition, B:1262–1263  
in saponite clays, B:888  
in secondary calcite, B:888  
Site 799, B:614, 1265  
and sulfate reduction, B:88  
vs. oxygen isotopes, B:88  
isotopes, deuterium  
in interstitial water, B:607  
Site 799, B:614  
vs. oxygen isotopes, B:615  
isotopes, helium  
heat flow correlated, B:747  
in interstitial gas, B:747–751  
Japan Sea, B:748  
Japan Trench, B:748–749  
in mid-ocean ridge basalt (MORB), B:747  
Nankai Trough, B:748–749  
Site 799, B:614  
vs. helium to neon ratio, B:749–750  
vs. total carbon to helium ratio, B:750  
isotopes, hydrogen  
in chlorite/saponite mixed-layer clays, B:888  
D/H ratios, B:642  
in interstitial water, B:639, 641–643  
in saponite clays, B:888  
in secondary calcite, B:888  
Site 794, B:641–643  
Site 795, B:641–643

Site 796, B:641-643  
 Site 797, B:641-643  
 vs. isotopes, oxygen, B:643  
 vs. potassium, B:1273  
 vs. rubidium, B:1273  
 isotopes, lead  
   in basement, B:807, 1340-1341  
 Pb-Pb plots, B:808, 813  
 vs. neodymium isotopes, B:810, 813  
 vs. strontium isotopes, B:810  
 isotopes, neodymium  
   in basalt, B:901-903  
   in basement, B:786-787, 807, 893  
   and crustal contamination, B:809  
   and mantle sources, B:903  
   secular variation, B:899  
 Site 794, B:1339-1340  
 Site 795, B:1338-1340  
 Site 797, B:1339-1340  
 vs. lanthanum/samarium ratio, B:811  
 vs. lead isotopes, B:810, 813  
 vs. strontium isotopes, B:809, 812  
 isotopes, oxygen  
   and ash layer alteration, B:52, 641, 701  
   and basalt alteration, B:699-702  
   and basement alteration, B:641-642  
   and benthic foraminifers, B:443-444  
   and carbonate diagenesis, B:701-702  
   in chlorite/saponite mixed-layer clays, B:888  
   coiling ratio correlated, B:467  
   correlated with diatom number in dark/light cycles, B:588  
 D/H ratios, interstitial water, B:642  
 in dark/light cycles, B:431, 569-571  
 in diagenetic carbonate, B:83, 86-88  
 and diagenetic reactions in interstitial water, B:699  
 equilibrium values vs. depth of carbonate formation, B:89  
 gamma-ray logs correlated, B:405  
 and gas hydrate, B:52  
 glacial/interglacial variability, B:442-443  
 gradients with dolomite layers, B:89  
 in interstitial water, B:51-52, 54, 84-85, 607, 639, 641-643, 697-703  
 isotopic stages, B:442-443  
 and Japan Sea, B:442-443, 447  
 loess correlated, B:395  
 and Northern Hemisphere glaciation, B:403  
 oceanographic front oscillations correlated, B:168  
 Oki Ridge stratigraphy, B:444  
 and opal-A to opal-CT transition, B:641-642  
 opal-CT, B:51-54  
 in planktonic foraminifers, B:442-443  
 quartz, B:51-54  
 in saponite clays, B:888  
 in secondary calcite, B:888  
 silica diagenesis and, B:701-702  
 and silica transformation temperature, B:49  
 Site 794, B:641-643  
 Site 795, B:641-643  
 Site 796, B:641-643  
 Site 797, B:593, 641-643  
 Site 798, B:442-443, 698-699  
 Site 799, B:614, 698-700  
 stage boundaries, B:468  
 stratigraphy, B:1221  
 surface-water temperature correlated, B:464-465, 468  
 vs. carbon isotopes, B:88  
 vs. deuterium isotopes, B:615  
 vs. hydrogen isotopes, B:643

vs. potassium, B:1273  
 vs. rubidium, B:1273  
 isotopes, radiogenic  
   in basement rocks, B:805-817  
   in interstitial water, B:635-650  
 isotopes, stable, B:635-650  
 isotopes, strontium  
   ash layer/basement alteration and, B:643  
   in ash layers, B:607  
   in basalt, B:901-903  
   in basement, B:786-787, 807, 893  
   and basement alteration, B:1266  
   in blue tuff, B:123, 127  
   and crustal contamination, B:809  
   in Green Tuff, B:644  
   in interstitial water, B:643-646  
   isotopic ratios vs.  $\text{I}^{\text{Sr}}$ , B:646  
   and seawater alteration of basalt, B:901  
   secular variation, B:899  
 Site 794, B:901-903, 1339-1340  
 Site 795, B:901-903, 1338-1340  
 Site 797, B:901-903, 1339-1340  
 Site 799, B:614  
   and thermal regime, B:644  
 vs. calcium, B:645  
 vs. lead isotopes, B:810  
 vs. magnesium, B:645  
 vs. neodymium isotopes, B:809, 812  
   in Yamato Seamount Chain lavas, B:127, 812  
 isotopes, sulfur  
   in interstitial water, B:637-639  
   and organic carbon decomposition, B:1262  
   in pyrite, B:711-713  
 Site 794, B:637-638  
 Site 795, B:637-638, 640, 708-709, 715  
 Site 796, B:637-638  
 Site 797, B:637-638  
   vs. sulfur content, B:715  
 isotopic stages, Site 798, B:442-443, 468  
 Itoigawa-Shizuoka Tectonic Line, and Japan Sea stress field, B:1185  
 Izu-Bonin Arc  
   basement isotopic composition, B:807  
   pressure compensation level, B:119  
   sediment contamination of mantle source, B:805  
   strontium isotopes vs. neodymium isotopes, B:809  
 Japan. *See also* Japanese archipelago  
   Tertiary system faunal changes, B:493  
 Japan Arc, A(128):5  
   ash, A(128):122, 156-157; B:791  
   conductivity model, A(128):85  
   crust, A(128):74; B:1345  
   geologic history, A(128):1617, 157  
   iron enrichment, B:874  
   paleoposition, B:1331  
   Pb-Pb plots of arc volcanic rocks, B:808  
   rotation, A(127):10, A(128):9, 11-12; B:187, 899, 1054, 1185, 1315, 1317-1318, 1334  
   sandstone provenance, B:110  
   strontium isotopes vs. neodymium isotopes, B:809  
   subcontinental lithosphere, B:811-812  
   tectonic evolution, B:1328-1329  
   volcanic history, B:1311-1331  
   volcanic outcrops, B:874, 882  
   Yamato Basin basalt correlated, B:903  
 Japan archipelago  
   electrical resistivity structure, B:1351  
   radiolarians, B:296  
   separation from Asia, B:531

vegetation, B:320, 325, 334  
 Japan Basin, A(128):70  
   age, A(127):6, 29, A(128):7; B:179, 906  
   ash layers, areal distribution, B:791  
   basement, A(127):6, 29, B:906  
   benthic foraminifers, B:208, 500-505, 518-521  
   biostratigraphy, B:1223  
   blue tuff, B:127-128  
   calcareous nannofossils, B:176, 178  
   crustal structure, A(127):9, 79, 176, A(128):9-11, 71-72, 74; B:924, 1075, 1083, 1086, 1104, 1107, 1304-1305, 1311-1314, 1318-1319, 1341-1346  
   crustal structure and tectonic history, B:1318-1319  
   diatoms, B:276-281, 286  
   earthquakes, A(128):74  
   Foram Sharp Line and, B:506, 517  
   formation age, B:830-831  
   free-air gravity anomalies, A(128):74  
   heat flow, A(127):27, A(128):71-74, 246; B:1302, 1345  
   Layer 1, A(128):71-72  
   Layer 2, A(128):71-72; B:1342  
   Layer 3, A(128):71-72; B:1342-1343  
   and lithospheric stretching models, B:1304-1305, 1307  
   lithostratigraphy, B:496-497  
   magnetic anomalies, A(127):6, 10, A(128):7; B:942, 1314, 1345  
   magnetic properties, B:941-942  
   magnetostratigraphy, A(127):22; B:970-972, 1223  
   oceanic crest, A(127):73  
   oceanographic setting, B:342  
   opal-A to opal-CT transition, B:50  
   opal-CT to quartz transition, B:50  
   oxygenation conditions, B:201-208  
   paleodepth, B:201, 1220-1221  
   pollen, B:328, 471, 480, 484  
   radiolarians, B:295  
   seafloor spreading, B:1345-1346  
   sedimentation rate, B:288, 1227  
   sponge spicules, B:541-543  
   subsidence, A(127):20; B:209, 1208-1211, 1214  
   surface-water temperature, B:209  
   tectonic history, B:1317  
   Yamato Basin, B:189, 1337  
 Japan Basin E  
   crustal structure, A(127):251-252  
   dredge samples, A(127):253  
   earthquakes, A(127):255  
   free-air gravity anomalies, A(127):253, 255-256  
   heat flow, A(127):252-253  
   initiation of convergence, A(127):258  
   magnetic anomalies, A(127):252-254  
   oceanographic history, A(127):259  
   sedimentation, A(127):259  
   stress field, A(127):258  
   tectonic setting, A(127):254-255, 257  
   thrust faults, A(127):255  
 Japan Basin N  
   age, A(127):169, 176  
   back-arc rifting, A(127):238  
   basement, A(127):169  
   bathymetry, A(127):174-176  
   crustal structure, A(127):176  
   depositional history, A(127):190-192  
   dredge samples, A(127):176-178  
   free-air gravity anomalies, A(127):176, 179

geologic history, A(127):169  
heat flow, A(127):174, 176-177  
Layer 1, A(127):176  
Layer 2, A(127):176  
Layer 3, A(127):176  
magnetic anomalies, A(127):176, 178  
oxic conditions, A(127):199  
paleoenvironment, A(127):199  
rifting history, A(127):179  
sedimentation, A(127):179, 181, 238  
seismic stratigraphy, A(127):179, 233-234  
tectonic setting, A(127):178-180  
upwelling, A(127):199  
Japan Sea  
age, A(127):29; B:297, 327, 362, 1208, 1335-1336  
back-arc rifting, A(127):16  
bacteria, B:755-776  
basaltic submarine volcanism, B:874  
basement, A(127):22, 24-25, A(128):6-7; B:749, 1265-1266, 1303-1304  
basin subsidence and volcanic activity, B:796-797  
bathymetry, A(127):6, A(128):6, 10; B:1199, 1276  
benthic environments, B:379  
biosiliceous productivity, B:313  
biostratigraphy, A(127):19; B:187, 1219  
carbonate chemistry, B:1263-1265  
carbonate-compensation depth, B:155, 168, 208, 440, 1201  
climate, B:320  
consolidation/permeability studies, B:1123-1133  
continental crust, B:809  
continental fragments, A(128):5, 10-11  
crustal structure, A(128):6-7, 9-11; B: 1075, 1083, 1104, 1107, 1311-1314, 1333, 1341-1346  
current systems, B:228  
deposition, A(128):15-18; B:1261  
development, B:536  
diatoms, B:309, 363  
dissolved oxygen, A(128):20  
downhole seismic experiment, B:1061-1073, 1157-1171  
earthquakes, A(128):76  
eolian dust, B:394  
faunal change, B:506  
formation, A(127):5, 9-10  
formation age, B:819-836, 1220, 1228  
formation models, A(128):11-13; B:1183-1187  
gateways to, B:362-363  
geochemistry, A(127):22-23  
geographic setting, B:365-366  
geologic history, A(128):16-17  
geologic setting, B:651, 677  
heat flow, A(127):26-27, A(128):14; B:1297-1298, 1301-1303, 1375  
helium isotope ratios, B:748-749  
inorganic geochemistry, B:1261-1274  
lithofacies, B:635  
lithospheric thickness, A(127):255  
lithostratigraphy, A(127):17-19, A(128):13-15; B:679, 1229-1233, 1275, 1315-1316  
logging, A(127):27-28  
magma genesis, B:1340-1341  
magmatic history, B:787-788  
magnetic anomalies, A(127):11; B:1334  
magnetics, A(127):22  
magnetostratigraphy, B:969-982, 1219

mantle, anisotropic velocity model, B:1075  
models for backarc spreading, B:903  
neodymium(strontium isotopes, B:899-904  
oceanographic front, B:155-159  
oceanographic history, A(127):16, 19-22, 24, 30-31, A(128):20-22; B:678-679  
oceanographic setting, A(128):20; B:166-167, 342-343, 394-395, 423, 440, 457-458, 559-560, 577, 605, 651, 705  
onshore exposures, A(128):15-18  
opal-A to opal-CT transition, A(127):31; B:49, 341, 1145-1156  
opal-CT to quartz transition, A(127):31; B:49  
opening of, B:171, 179, 529-531  
and organic matter, A(127):31; B:1262-1263  
origin, B:249  
oxygenation conditions, B:216, 342-343, 423, 430, 678-679, 705, 713-715, 1221  
paleobathymetry, B:1209-1210  
paleoceanography, B:179-180, 298-300, 419-421, 526, 528-529  
paleoclimate, B:403, 459  
paleodepth, B:1228  
paleoenvironment, B:187, 493  
paleogeography, A(128):22  
paleotemperature fluctuations, B:457-470  
phosphorus, B:68-71  
physical properties, A(127):25-26; B:1275-1296  
phytoplankton pigment concentration, B:167  
plate reconstructions, B:1321-1324  
pollen, B:328-336, 395, 471-478  
pre-ODP data available, B:1333  
productivity, A(127):20; B:167, 430, 651  
radiometric ages, B:1336  
rifting, A(127):16; B:891, 899, 1334  
and sea ice formation, A(127):20-22; B:167  
sediment composition, B:1233-1249  
sediment isopachs, A(127):13, A(128):12  
sediment structures, A(128):13-15  
sedimentation, A(127): 10-16, 30-31; B:635, 1361  
sedimentation rate, A(127):22, 24  
seismic reflection profile, A(127):14  
seismic stratigraphy, A(127):28-29  
silica, B:447, 1263  
sill depths, B:299-300  
sites drilled, B:172, 226, 238, 250, 318, 326-327, 479, 494, 543  
slumping, A(128):14  
stress field, A(127):17  
structure, A(127):6; B:780, 792, 1175-1193  
submarine volcanism, A(127):95  
subsidence, A(127):16, 19-20; B:1197-1218, 1343-1344  
surface currents, A(128):21  
surface-water temperatures, B:201, 463-465  
tectonic events and microfossil assemblages, B:300, 526  
tectonic evolution, B:903, 1183-1187, 1311-1331, 1343-1347  
tectonic setting, A(127):5-10, A(128):9; B:1186, 1312  
temperature distribution, A(128):20  
tephra markers, A(128):291-292; B:796  
thermal conductivity, B:1301-1303  
thermal gradient, A(127):26-27; B:1301-1303, 1375  
thermal history, B:49-56, 1297-1309  
thermocline, B:167  
thrust faults, A(127):81-82  
volcanic activity, B:796-797, 829-832, 1319, 1324

Wadati-Benioff zone, B:1062, 1158  
water circulation, A(128):21; B:69-71, 423-424  
water convection, B:394  
Japan Sea Bottom Water, isotope ratios, B:748  
Japan Sea E  
bathymetry, A(127):12  
compression, A(127):16-17, 268, A(128):9  
crustal structure, A(128):9-11  
deformation, B:1314-1315  
earthquakes, A(127):9, A(128):11  
free-air gravity anomalies, A(128):252  
initiation of convergence, A(127):314  
magnetic anomalies, A(128):251  
plate reconstruction, B:1314-1315  
rift propagation, B:1346  
stress field, B:1059  
and strike-slip shear, B:1314-1315  
tectonic history, B:1316-1317  
tectonic map, A(127):83  
thrust faults, A(127):9  
Japan Sea N  
heat flow, A(127):231-232  
stress field, A(127):180-181  
Japan Sea Plate, electrical resistivity structure, B:1351  
Japan Sea Proper Water (JSPW), B:577  
benthic foraminifers and, B:365-366  
Japan Trench, A(127):6, A(128):9  
basement age, B:749  
carbon isotopes, B:88  
crustal structure, A(128):74  
diagenetic carbonates, B:75  
helium isotope ratios, B:748-749  
helium to neon ratios, B:748  
interstitial water oxygen isotopes, B:698  
rare earth elements, B:719  
Jaramillo Event. *See* Jaramillo Subchron  
Jaramillo Subchron  
and diatom occurrence, A(128):159  
inclination variations, A(128):172  
and magnetic intensity variations, B:962  
and sedimentation rate, B:959-960  
silicoflagellate and ebridian zones correlated, A(128):162  
Site 795, B:1223  
Site 797, A(127):359; B:973, 1223  
Site 798, A(128):30, 125, 168; B:164, 564, 973, 1224  
Site 799, A(128):314; B:974, 1224  
joints, Site 794, B:1182-1183  
Juan de Fuca Ridge, basalt emplacement, B:1337  
K/Ar age, Site 794, B:784-786  
Kaena Subchron  
Site 798, B:1224  
Site 799, A(128):314; B:1224  
kaolinite  
abundance, B:1235-1241  
in sandstone, B:139-141  
as sandstone cement, B:134  
Site 798, B:411, 416, 418  
Site 799, B:144  
kaolinite/illite ratio, Site 798, B:411, 416  
Kita-Oki Bank, A(127):325  
basement, A(127):333, A(128):127-129  
bathymetry, A(128):126-127  
heat flow, A(127):330, A(128):127, 131  
seismic stratigraphy, A(127):399  
Kita-Yamato Bank, A(127):325  
basement, A(128):247-249  
bathymetry, A(128):245-246  
crustal structure, A(128):248  
diatoms, A(128):249, 303

freshwater sediments, A(128):249  
 magnetic anomalies, A(128):247  
 phosphate source, B:67–68  
 phosphatic sediments, A(128):249  
 sandstone provenance, B:110–111  
 subsidence, A(128):295  
**Kita-Yamato Trough**, A(127):13, 325. *See also*  
 Yamato Trough  
 n-alkanes, B:669  
 bathymetry, A(128):245–246, 374; B:76  
 benthic foraminifers, B:370–373, 376–378  
 biostratigraphy, B:1224  
 consolidation/permeability studies, B:1123–  
 1133  
 crustal structure, A(128):245–249  
 depositional history, A(128):252–253, 292–  
 298, 353–355  
 diagenesis, B:35–39  
 diatoms, B:361  
 ebridians, B:241, 246–247  
 free-air gravity anomalies, A(128):247  
 geologic history, A(128):21–22  
 heat flow, A(128):246  
 Layer 1, A(128):246  
 Layer 3, A(128):246  
 magnetostratigraphy, B:974–975, 1224  
 normal faults, A(127):334  
 oceanographic conditions, A(128):312, 355–  
 356  
 oceanographic drilling objectives, A(128):19  
 oceanographic history, A(128):252–253  
 organic matter, B:668  
 paleodepth, B:373, 379  
 paleoenvironment, B:381  
 radiolarians, B:225–226, 233–234  
 sediment thickness, A(128):237, 353–354  
 sedimentation, A(128):250; B:1227  
 seismic stratigraphy, A(128):34, 245, 250,  
 344–351; B:1135–1143  
 silicoflagellate, B:241, 246–247  
 Site 799, B:425, 560, 605, 623, 667, 721, 739,  
 1135  
 stress field, A(128):293–294  
 structural cross-section, A(128):303  
 structural setting, B:1135  
 subsidence, A(128):239, 295–298, 355  
 surface-water productivity, A(128):324  
 tectonic evolution, B:1187  
 tectonic setting, A(128):249–250; B:156  
**Kitaura Formation**  
 deposition, A(128):18  
 foraminifers, B:187, 459  
 submarine fan deposition, B:1202  
**Kleiner-Hartigan diagrams**, for core XRF and geochemical log oxides, B:1030  
**Koenigsberger ratio (Q)**  
 in basement, A(127):203  
 Site 795, A(127):199  
**Korea**, granite exposures, B:111  
**Korea Strait**, A(127):355  
**Korean Current**, B:424  
**Korean Peninsula**, A(128):15  
**Korean Plateau**, basement, A(127):9  
**Kuril Basin**, formation, A(127):10  
**Kuroko sulfide deposits**, A(128):21–22, 24, 239,  
 251  
**Kuroshio Current**, A(127):355, A(128):20–21, 164  
 climate and, B:326  
 and diatom assemblages, A(128):159  
 and foraminiferal coiling direction,  
 A(128):165, 310–311  
**Japan Sea and**, A(128):30  
 Site 798, A(128):164–166

Site 799, A(128):312  
 and Tsushima Strait, B:166–167  
**Kyushu**, ash layers from, A(128):156–157  
 laminated sediments. *See* laminations; sediments,  
 laminated  
**laminations**  
 annual or seasonal, B:548, 551  
 in BSEM images, B:549–557  
 calcareous/biosiliceous alternations, B:549  
*Chaetoceros* resting-spore laminae, B:549–  
 551, 555–556  
 in cyclic lithofacies, A(128):140  
 in dark/light cycles, B:581  
 detrital-rich/biosiliceous alternations, B:549  
 in dolomite, B:92  
 foraminifer-rich laminae, B:551  
 mixed-biosiliceous laminae, B:549–551  
 thickness, B:548, 550  
 and upwelling events, B:551  
**La/Yb** values. *See* lanthanum/ytterbium ratio  
**lansfordite**  
 and alkalinity, B:92  
 changed to magnesite, B:80, 91  
 chemical composition, B:82  
 CuK- $\alpha$  vs. depth, B:81  
 formation, B:91–92  
 isotopic composition, B:83  
 occurrence, B:80  
 and organic carbon content, B:92  
 paleoenvironmental implications of, B:94  
 Site 799, B:75–98  
 X-ray diffraction analyses, B:80  
**lanthanum**  
 in ash layers, B:1386  
 Site 795, B:707  
 Site 798, B:1370–1371  
 vs. aluminum, B:737  
**lanthanum/samarium ratio**  
 in basement rocks, B:807  
 vs. neodymium isotopes, B:811  
**lanthanum/ytterbium ratio**  
 and Ce anomaly, B:693–694  
 and manganese accumulation, B:729  
 and migration of heavy rare earth elements  
 (HREE), B:731–732  
 Site 794, B:683  
 Site 795, B:683  
 Site 797, B:688–691  
 Site 798, B:729, 735  
 Site 799, B:729, 736  
 vs. Ce anomaly, B:737  
**large-ion lithophile elements (LILE)**  
 mobility during alteration, B:779, 838  
 Site 795, A(127):219  
 and slab-derived fluids, B:809  
**Last Glacial Maximum**  
 and Japan Sea oceanographic conditions,  
 B:394–395  
 and siliceous microfossil abundance, B:311  
**Lau Basin**  
 formation model, B:928  
 tectonic history, B:1347  
**lava flows**  
 eruptive depth, A(128):95  
 Site 794, B:906, 1339  
 Site 795, B:850, 906, 943  
 Site 797, B:838, 906–908  
 suites observed, B:853–855  
**lava flows, basaltic**, Site 795, B:819  
**Layer 1**  
 Japan Basin N, A(127):176  
 Kita-Yamato Trough, A(128):246  
 Oki Ridge, A(128):127  
**Oki Trough**, A(128):127  
**Yamato Basin**, A(128):127  
**Yamato Basin S**, A(127):330  
**Layer 2**  
 Japan Basin, B:1342  
 Japan Basin N, A(127):176  
 Oki Ridge, A(128):127–128  
 seismic anisotropy, B:1107  
 and thickness of altered basalt, B:701  
 Yamato Basin, A(127):73, A(128):9, 127–  
 128; B:1342, 1345  
 Yamato Basin S, A(127):330  
 Yamato Rise, A(128):246  
**Layer 3**  
 Japan Basin, B:1342–1343, 1345  
 Japan Basin N, A(127):176  
 Kita-Yamato Trough, A(128):246  
 Oki Ridge, A(128):127–128  
 Yamato Bank, A(128):246  
 Yamato Basin, A(127):73, A(128):127–128;  
 B:1342–1343, 1345  
 Yamato Basin S, A(127):330  
 Yamato Rise, A(128):246  
**LDT**. *See* density, logging  
**lead**  
 in basement rocks, B:807  
 Site 794, B:1362  
 Site 795, B:1365  
 Site 797, B:1366  
 in sulfide deposits, A(128):21  
**lead isotopes**. *See* isotopes, lead  
**Leg 13**, trace metal composition, B:1361  
**Leg 19**, silicoflagellate, B:237  
**Leg 31**  
 biostratigraphic zones used, A(128):162  
 consolidation tests, B:1123  
 diatoms, B:250  
 drilling history, A(127):16, A(128):5, 82;  
 B:1197  
 foraminifers, A(127):273  
 Japan Sea formation age, B:1334  
 lithostratigraphy, A(128):13  
 organic carbon, B:430  
 poor carbonate preservation, B:457  
 radiolarians, A(128):162; B:225, 291, 297  
 silicoflagellate, A(128):28; B:237  
 site locations, B:156  
 Toyama Deepsea Fan, B:1148  
**Leg 42B**, trace metal composition, B:1361  
**Leg 57**  
 ash layers, A(127):95–96; B:1319–1320  
 silicoflagellate, B:237  
 volcanic activity, B:796  
**Leg 64**, basalt emplacement, B:1337  
**Leg 65**  
 downhole seismic experiment, B:1061  
 seismometer emplacement, A(128):106  
**Leg 67**  
 downhole seismic experiment, B:1061  
 seismometer emplacement, A(128):106  
**Leg 73**, magnetic properties, B:949  
**Leg 78A**  
 downhole seismic experiment, B:1061  
 seismometer emplacement, A(128):106  
**Leg 78B**, seismometer emplacement, A(128):106  
**Leg 80**, inorganic nitrogen, A(127):285  
**Leg 86**, silicoflagellate, B:237  
**Leg 87**  
 ash layers, B:1319–1320  
 volcanic activity, B:796  
**Leg 88**  
 downhole seismic experiment, B:1061  
 seismometer emplacement, A(128):106

Leg 91  
downhole seismic experiment, B:1061  
seismometer emplacement, A(128):106

Leg 93, sponge spicules, B:541

Leg 112, microbiological studies, A(128):178

Leg 113, sponge spicules, B:541

Leg 120, sponge spicules, B:541

Leg 124, Zijderveld plots, A(127):277

Leg 135  
Lau Basin formation, B:928  
tectonic history, B:1347

Leg 139, basalt emplacement, B:1337

lerite, plagioclase phryic, A(128):75

light rare earth elements (LREE), B:730  
and Ce anomaly, B:693-694

Liman Current, B:342, 423  
cold surface water, B:440, 457  
in Japan Sea, A(128):20-21; B:70

lipids  
in organic matter, A(128):324  
Sites 798 and 799, B:669-671

lipid fraction, in organic matter, A(128):324

lignite, and thermal maturity levels, B:669

liptodetrinitite, Sites 798 and 799, B:670

liquid limit. *See* Atterberg limits

liquid line of descent  
in melting experiments, B:863

mid-ocean ridge basalts and Site 797 sills correlated, B:870

for olivine tholeiitic basalts, B:864

problems with, caused by alteration, B:911

Site 794, B:869-882, 1339

Site 795, B:918, 1338

Site 797, B:869-882, 920

lithium  
and basement alteration, A(127):364; B:1266

and biogenic silica content, A(127):205

and diatom dissolution, A(127):364

in interstitial water, B:1272

and opal-A to opal-CT transition, B:1266

Site 794, A(127):109

Site 795, A(127):205

Site 796, A(127):280-281

Site 797, A(127):364, 371

Site 798, A(128):174-175, 184

Site 799, A(128):318, 332; B:611  
vs. silica, A(128):185

lithodensity tool (LDT). *See* density, logging

lithofacies, cyclic, A(128):27. *See also* color-banded bedding; dark/light cycles  
and bottom water conditions, A(128):141

burrows, A(128):140-141

compositional variability, A(128):140

couplet thickness, A(128):140

cycle boundary characteristics, A(128):140-142

and foraminifers Q-mode cluster analysis, B:200

formation, A(128):283

Japan Sea, A(128):14

laminations, A(128):140

in logs, A(128):34, 125, 185

orbital obliquity, A(128):35

and organic carbon, A(128):125, 177

organic matter, A(128):140

origin, A(128):158

and paleoceanographic changes, A(128):121-122

Site 798, A(128):35, 121-122, 124, 137ff, 195-196; B:156

Site 799, A(128):259, 282-284, 293

lithospheric stretching models, heat flow and basement ages compared, B:1304-1305

lithostratigraphy. *See also* specific sediment types  
basement, B:781, 884, 901  
and biogenic silica dissolution, B:344  
blue tuff, B:120-121  
gamma ray logs correlated, B:1416-1417  
Honshu, A(127):15

intersite lithologic correlation, B:1233

Japan Basin, B:679

Japan Sea, A(127):17-19, A(128):25-28; B:1229-1233, 1275, 1315-1316

Oga Peninsula, B:1203

opal dissolution transition zone (ODTZ) correlated, B:347-349

physical properties correlated, A(127):383-389, A(128):182, 327-329; B:1143

Pohang Basin, B:1205

problems with shipboard identifications, B:1229

revised unit boundaries defined, B:1230-1233

Sakhalin Island, B:1206

and sedimentary structures, B:1177

seismic intervals correlated, B:1140-1143

seismic stratigraphy correlated, A(127):155, 174, 234-238, 243, 251, 312-313, 325, 403-408, 410, A(128):125, 193-194, 224, 351-353, 379

Site 794, A(127):72, 74-77, 91, A(128):25-26, 77-81; B:5, 118, 494, 496, 655, 1049, 1277, 1313, 1336, 1377, 1413

Site 795, A(127):170-174, 186-192, A(128):26; B:5, 118, 655, 706, 1277, 1313, 1336, 1377, 1413

Site 796, A(127):248-250, 261-269, A(128):26; B:5, 105, 118, 133, 655, 1277, 1313, 1336, 1377, 1413

Site 797, A(127):324, 326-330, 340-351, A(128):26; B:5, 105, 118, 133, 496, 498-499, 655, 1277, 1313, 1336, 1376, 1413

Site 798, A(128):25, 27-28, 122-124, 136  
158; B:5, 157, 655, 697, 800, 1277, 1336, 1376

Site 799, A(128):25, 27-28, 35-36, 238-244, 255ff; B:5, 33-48, 76-77, 105, 133, 157, 655, 697, 800, 1137, 1277, 1313, 1336, 1376

Sites 794-796 compared, A(127):269

Sites 794 and 797 correlated, B:1038

units, revised, B:1229-1233, 1250

units, Site 794, A(127):90-94; B:679

units, Site 795, A(127):186-190; B:4, 679, 706

units, Site 796, A(127):261-266

units, Site 797, A(127):340-346; B:4-5, 679

units, Site 798, A(128):124, 137-138; B:156, 697, 973

units, Site 799, A(128):240-244, 256-265; B:76-77, 158-160, 697, 974

and vein abundance, B:1180

vitric tuff, Site 799, B:36

Yamato Basin, B:679

loess  
in China, B:394

clay fraction, B:401

deposition at glacial maxima, B:403

and Northern Hemisphere glaciation, B:421

origin of dark/light cycle detrital component, B:594-595

oxygen isotopes correlated, B:395

and quartz abundance, B:401

source and deposition regions, B:411

logging, B:1395-1409, 1411-1427. *See also* specific logging measurements

aridity signal, A(128):188-192

of ash layers, A(128):288

and Asian dust supply, B:403

of basement, A(128):69, 86

of breccia tuff, A(128):287

of chert and porcellanite layers, A(127):394

correlated with basement recovery, A(128):93-95

of cyclic lithofacies, A(128):125

cyclicity observed, B:398, 405

and intersite correlation, B:1234

Japan Sea, A(127):27-28, A(128):33-34

lithostratigraphy correlated, A(127):393-395

log characteristics, A(128):103-104

log quality and processing, A(127):139, 232, 301-302, 392-393, A(128):103, 183, 329-332

of opal-A to opal-CT transition, B:15-16, 19, 21-22

and opal content variations, B:398-400

of opal-CT to quartz transition, B:23

operations, A(127):27, 139, 142, 183-186, 230, 232, 236, 260, 296, 301-302, 325, 338-339, 389-390, 392, 398, A(128):85, 103, 106, 211, 253-255, 329, 361; B:395-396, 1411-1412

orbital obliquity observed, B:403

paleoclimate, A(128):33-34, 187-192, 341-342; B:393-407

periodicity resolution, B:396-397

physical properties correlated, A(127):139, A(128):338

and sediment composition estimates, B:402-403

seismic stratigraphy correlated, A(128):224, 351-353, 379

of silica diagenetic transitions, B:3

Site 794, A(127):73, 135-142, 144-167, A(128):34, 69; B:1037-1046, 1069, 103ff

Site 795, A(127):174, 230ff

Site 796, A(127):251, 296ff

Site 797, A(127):325, 389ff; B:7, 1037-1046

Site 798, A(128):33-34, 125, 182ff; B:393-407, 1021-1035

Site 799, A(128):34, 245, A:329ff

Sites 798 and 799 compared, A(128):342

stratigraphy, B:1221

temporal resolution, B:396, 404-405

in terrigenous/biogenic opal cycles, B:398

tool string schematic diagrams, B:398

units, Site 799, A(128):332-338

used to determine alteration extent, B:1282

vertical aperture, B:396

of vitric tuff, B:37

logging, geochemical. *See* geochemical logging

long-space sonic tool (LSS). *See* velocity, logging

loss on ignition (LOI)  
as alteration proxy, B:909

and hydrous mineral content, B:1282-1284

and physical properties, B:1283

vs. density, B:1291

vs. elemental data, B:918-921

vs. magnesium, B:839

vs. porosity, B:1290

vs. velocity, B:1291

LREE. *See* light rare earth elements

LSS. *See* velocity, logging

lutetium, Site 798, B:1370-1371

lysocline, and nannofossil preservation, A(128):164

macerals, Sites 798 and 799, B:668-670

magma. *See* mantle composition; mantle source; parent magma

## magma batches

magma batches, Site 794, B:917  
 magma composition, and trace elements, B:1378–1379  
 magma genesis, in Japan Sea, B:1340–1341  
 magma intrusion, and crustal deformation, B:1323  
 magma source  
   depleted asthenospheric mantle, B:1341  
   enriched subcontinental upper mantle, B:1341  
   and neodymium isotopes and titanium correlated, B:902  
   Site 794, B:780–782  
   Yamato Basin, B:782–784, 902  
 magma suites  
   in basement rocks, B:869–870  
   Site 794, B:871, 875  
   Site 797, B:871, 875  
 magmatic affinity, of basement, A(128):96ff  
 magmatic eruption, and blue tuff, B:119, 126  
 magmatic history, Japan Sea, B:787–788  
 magnesiochromite, B:840. *See also* spinel; spinel, chromium  
 magnesite. *See also* breunnerite  
   chemical composition, B:82, 84–85  
   CuK- $\alpha$  vs. depth, B:81  
   derived from lansfordite, B:80, 91  
   formation, B:91–92  
   formation depth, B:85  
   isotopic composition, B:83  
   occurrence, B:80  
   organic carbon content, B:92  
   origin, B:76  
   paleoenvironmental implications, B:94  
   petrography and textural relations, B:81  
   SEM photographs, B:97  
   Site 799, B:75–98  
   X-ray diffraction analyses, B:80  
 magnesite, ferroan, B:745–746  
 magnesium  
   alteration and, B:909, 911  
   and ash alteration, A(127):280, 363–364  
   in ash layers, B:1382  
   and basement alteration, A(127):280; B:892–893, 1265–1266  
   and dolomite formation, A(127):363–364  
   enrichment during alteration, B:842  
   and geochemical logging, B:1395, 1415  
   in interstitial water, B:1263, 1268  
   Japan Sea, B:1236, 1265  
   mobility during alteration, B:838  
   Site 794, A(127):109–110  
   Site 795, A(127):174, 205  
   Site 796, A(127):280  
   Site 797, A(127):363–364, 370  
   Site 798, A(128):173–174, 182  
   Site 799, A(128):318, 329; B:611  
   vs. boron, B:642  
   vs. calcium, A(127):115  
   vs. loss of ignition, B:839  
   vs. smectite, B:1250  
   vs. strontium isotopes, B:645  
 magnesium/aluminum ratio  
   in Japan Sea sediment, B:1236  
   Site 794, B:1249  
   as volcanic input indicator, B:1239  
 magnesium/calcium ratio  
   in interstitial water, B:1263, 1269  
   Japan Sea, B:1265  
   Site 799, A(128):329; B:611  
 magnesium number  
   basement, A(128):98  
   Site 794, B:839  
   Site 797, B:839  
   spinel, B:840–842, 844

## SUBJECT INDEX

magnesium oxide. *See* magnesium  
 magnetic anomalies  
   Japan Basin, A(128):7; B:1314, 1345  
   Japan Basin E, A(127):252–254  
   Japan Basin N, A(127):176, 178  
   Japan Sea, B:1334  
   Japan Sea E, A(128):15, 251  
   Kita-Yamato Bank, A(128):247  
   Oki Ridge, A(128):127, 132  
   Site 794, A(127):80  
   Tsushima Basin, B:1314  
   Yamato and Japan basins compared, B:942  
   Yamato Bank, A(128):247  
   Yamato Basin, A(128):7; B:1314, 1345  
   Yamato Basin N, A(127):77, A(128):73  
   Yamato Basin S, A(127):330, 334  
   Yamato Rise, A(128):247, B:1314  
 magnetic declination. *See* magnetic properties  
 magnetic inclination. *See* magnetic properties  
 magnetic intensity. *See* magnetic properties  
 magnetic minerals. *See also* specific magnetic minerals  
   diagenetic dissolution, B:951  
   formation by thermal demagnetization, B:949, 951  
   grain size, B:949–951  
   and magnetic properties, B:947, 962  
   Site 797, B:949, 951  
 magnetic polarity. *See* magnetostratigraphy  
 magnetic properties. *See also* magnetic minerals; magnetostratigraphy  
   AF demagnetization, B:949, 951  
   in basalt flows vs. dolerite sills, A(128):102  
   in basement, A(127):209, 358–359  
   and basement alteration, A(127):203  
   basement VRM, intersite comparisons, B:941  
   and bioturbation, A(127):275  
   data quality variations, B:975  
   declination, A(127):106–107, 200–202, 276, 358–359, 363, A(128):101–102, 167–170, 313–315; B:960, 977, 979, 1052–1053, 1317  
   demagnetization, A(128):169, 314–315  
   emplacement mode parameter (F), B:934, 936–937, 942  
   equal-area plots, A(127):104–105, 206–207, 278–279, 360–361, A(128):104–105, 173, 316, 318–320, 322–323; B:976  
   frequency dependence susceptibility, B:934, 936–937, 942  
   and gas, A(127):275  
   and geomagnetic field intensity variations, B:962–963, 977  
   inclination, A(127):106–107, 200–202, 276, 358–359, 363, A(128):101–103, 105, 167–170, 173–174, 313–316; B:960, 974, 977–981  
   intensity, A(127):72, 103, 106–107, 174, 200–202, 251, 275–276, 324, 357–359, 363, A(128):101–103, 105, 167–170, 173, 313–315; B:934, 937, 957, 969–973, 975–981  
   intensity decay plots, A(127):104–105, 206–207, 278–279, 360–361, A(128):104–105, 173, 316, 318–320, 322–323  
   isothermal remanent magnetization (IRM), A(127):357, 359; B:972–973, 975  
   Japan Sea, A(127):22, A(128):30–31  
   Koenigsberger ratio (Q), A(127):199, B:934–938  
   magnetic intensity, B:959–967  
   and magnetic minerals, B:951  
   in RCB cores, B:975, 981  
 remagnetization, A(127):203, 359; B:938  
 remanence, A(128):101  
 and rotation of Japan Arc, B:1317–1318  
 and rust contamination, A(127):357; B:973  
 Site 794, A(127):103–105, A(128):30–31, 69, 100–105; B:934–935  
 Site 795, A(127):174, 199ff; B:934, 937  
 Site 796, A(127):251, 275ff  
 Site 797, A(127):324, 357ff; B:934, 937–938, 947–957  
 Site 798, A(128):30–31, 166ff  
 Site 799, A(128):30–31, 312ff  
 soft vs. stable components of remanence, A(128):101  
 and stress field analysis, B:1050  
 susceptibility, A(127):72, 105, 110–111, 174, 199, 203–204, 275, 277, 324, 358, 362, A(128):101–102, 167–171; B:934, 937, 954, 959, 961, 973, 975–977, 979, 981  
 thermal demagnetization, B:938–939, 949, 951, 954  
 vector demagnetization diagrams, B:935, 940, 950, 952–953  
 viscous remanent magnetization (VRM), B:934, 936–937, 939–942, 944–945  
 of volcanic rocks, B:933–945  
 in XCB cores, B:975, 978, 980  
 Yamato and Japan basins compared, B:941–942  
 Zijderveld plots, A(127):104–105, 203, 206–207, 277–279, 360–361, A(128):104–105, 173, 316, 318–320, 322–323; B:971  
 magnetic susceptibility. *See* magnetic properties  
 magnetite, B:853  
   in basaltic andesite, A(127):217  
   magnetostratigraphic analyses and, B:976  
   produced by thermal demagnetization, B:947 and sedimentation rate, B:976  
   Site 797, B:949, 951  
 magnetostratigraphy, B:1219–1228. *See also* magnetic minerals; magnetic properties  
 basement polarity, A(128):86; B:936–938  
 chrons and subchrons, A(127):108  
 core orientation, A(127):103, 199, A(128):167  
 data quality, B:969, 975  
 datums used, B:163, 1222  
 depths to polarity boundaries, B:973  
 diatoms and, B:361–362  
 and drilling disturbance, A(127):103  
 ebridians and, B:240  
 in igneous rocks, A(127):104, 203  
 Japan Sea, A(127):22, A(128):30–31; B:969–982  
 magnetic polarity, B:943  
 problems with, B:1219  
 reference time scale correlated, A(127):22–23, 103–105, 108, 110, 199–203, 208, 275, 280, 359, 364, A(128):170, 174–176, 317; B:970, 972  
 silicoflagellate and ebridian zones correlated, A(128):162; B:240  
 Site 794, A(127):72, 103–105, A(128):69, 100–105; B:969–970, 972, 1221–1223  
 Site 795, A(127):174, 199ff; B:970–972, 1223  
 Site 796, A(127):251, 275ff; B:1223  
 Site 797, A(127):324, 357ff; B:972–973, 1223  
 Site 798, A(128):30–31, 124–125, 166ff; B:399, 417, 972–974, 1223–1224  
 Site 799, A(128):30–31, 244, 312ff, B:972, 974–975, 1224

major element molar ratios  
Site 798, B:616-617  
Site 799, B:611, 618-621  
major elements, B:677-695. *See also specific major elements*  
Al-normalized concentrations, B:687  
alteration and, B:799, 901, 909, 911  
in augite, B:852-854  
in basalt, B:839, 869-870  
basement, A(128):98; B:782, 893  
in dark/light cycles, B:568, 570-571  
and fractionation models, B:881  
in ilmenite, B:856-857  
interelement comparisons, B:690  
of Japan Sea sediment, B:1235-1237, 1246-1248  
in Japanese volcanic outcrops, B:874  
in magnetite, B:856-857  
mobility during alteration, B:838  
in olivine, B:850  
in Plagioclase, B:852-853  
silica, B:682  
Site 794, B:680, 682, 686-687, 782, 839, 858, 869-870, 893, 912-913, 918  
Site 795, B:680-681, 683, 686-687, 913-914  
Site 796, B:681  
Site 797, B:681, 686-688, 839, 858, 869-870, 893, 914-916, 920  
Site 798, B:720-722  
Site 799, B:722-723, 726-727  
in spinel, B:840-842, 856-857  
in volcanic glass, B: 1378  
vs. loss on ignition, B:918, 920  
XRF and geochemical log analyses compared, B:1025-1029, 1031-1035

Mamiya Strait  
depth, B:440  
Japan Sea and, B:362-363  
sill depth, B:457

Mammoth/Gauss boundary, Site 798, A(128):156

Mammoth Subchron, Site 798, B:1224

manganese  
accumulation, B:90-91  
and cerium, B:726-727  
in interstitial water, B:607  
Japan Sea, B:1236  
Site 795, B:710-711, 714  
Site 798, A(128):172-173, 180; B:722, 1263  
Site 799, A(128):317, 328; B:90-91, 610, 722, 727-729  
vs. Ce anomaly, B:737

manganese, hydrogenous, rare earth element source, B:729

manganese nodules, and cerium, B:726-727

manganese oxide. *See manganese mantle*  
anisotropic velocities, B:1075  
composition, B:786-787, 927  
depleted, B:807-815, 903  
heterogeneity, B:927  
processes, B:927-928  
and rare earth mixing model, B:817  
Yamato Basin, B:927

mantle source  
and beryllium isotopes, B:815  
and crustal contamination, B:807, 809-810  
depleted asthenospheric mantle (AM), B:903  
depleted mantle and pelagic sediment mixing plots, B:814  
end-member compositions for mixing calculation, B:815  
enriched mantle II (EMII) component, B:786-787

enriched subcontinental upper mantle (SCUM), B:903  
and Japan Sea basalt, B:903, 1341  
and lithospheric contamination, B:810-812  
N-MORB source depleted mantle (DM) component, B:786-787  
and sediment contamination, B:805  
Site 794, B:917  
and slab-derived fluid contamination, B:809  
and subducted sediment contamination, B:812-815  
Yamato Basin, B:786-787  
for Yamato Seamount Chain, B:812  
marcasite, in vesicle, A(128):90  
marginal basins, Pacific Ocean W, B:1311, 1333-1334  
marginal sea  
formation, A(127):5  
Japan Sea as, B:365-366

Mariana Arc  
Pb-Pb plots of arc volcanic rocks, B:808  
sediment contamination of mantle source, B:805, 812  
strontium isotopes vs. neodymium isotopes, B:809

Mariana Trough  
basement isotopic composition, B:807  
lava flows, B:874  
lava flows, chemical composition, B:882  
Pb-Pb plots of arc volcanic rocks, B:808

Maritime province, and sea ice, B:167

mass accumulation rate. *See also* sediment accumulation rate  
of barium, B:665  
of carbonate, B:428  
of organic carbon, B:425, 428  
Site 798, B:428

matrix sieving, and vein fillings, B:1179-1180

Matuyama/Brunhes boundary, and inclination variations, B:974

Matuyama Chron  
and diatom occurrence, A(128):159  
Site 796, A(127):275  
Site 797, B:973  
Site 798, A(128):124, 170

Matuyama/Gauss boundary  
silicoflagellate and ebridian zones correlated, A(128):162, 305  
Site 794, B:970  
Site 795, B:970  
Site 796, A(127):275  
Site 797, B:973, 1223  
Site 798, A(128):30, 170; B:1224  
Site 799, A(128):314; B:974, 1224

Matuyama/Jaramillo boundary, inclination variations, A(128):172

Matuyama/Olduvai boundary  
Site 798, A(128):164  
Site 799, A(128):307

Mediterranean E, and trace metal composition, B:1361

Meijo-Daisan Seamount, A(128):70

melt distribution, and electrical resistivity, A(128):111

mesostasis  
altered to clay minerals, A(127):217  
Site 795, B:821

methane  
biogenic, A(127):251, 288  
concentrations, A(127):119  
in gas hydrate, A(127):288-290  
Japan Sea, A(127):24  
and methanogenesis, A(127):216

nitrogen correlated, B:750  
origin, B:749-750  
potential production rates, B:771  
shallow presence, A(127):259-260  
Site 794, A(127):119  
Site 795, A(127):174, 213-216, 220  
Site 796, A(127):287-288, 290, 315  
Site 797, A(127):368  
Site 798, A(128):175-176, 180, 187; B:771-772, 776, 1263  
Site 799, A(128):244, 321-322, 339  
and sulfate abundance, A(127):368  
thermogenic, A(127):251, 288  
vs. depth, A(127):220

methane-clathrate. *See* clathrate

methane/ethane ratio ( $C_1/C_2$ )  
in gas hydrate, A(127):288, 290  
Site 795, A(127):220  
Site 796, A(127):287-288, 290  
Site 798, A(128):176, 188  
Site 799, A(128):339

methane/propane ratio ( $C_1/C_3$ )  
Site 798, A(128):176, 188  
Site 799, A(128):339

methanogenesis  
and bacteria, B:767-769, 771  
and carbon isotopes, B:88-90  
and interstitial water analyses, B:607  
and methane content, A(127):216  
and organic carbon decomposition, B:1262-1263  
and organic matter degradation, A(127):207  
Site 796, A(127):267  
Site 797, A(127):368  
Site 798, B:772-773, 776

methylcyclohexane, Site 799, A(128):321-322

methylcyclopentane, Site 799, A(128):321-322

Mg#. *See* magnesium number

Mg/Ca ratio. *See* magnesium/calcium ratio

Mg-calcite, Site 796, A(127):266

microbiological studies  
bacteria, B:755-760, 1262  
methods, B:755-756, 761-765  
objectives, A(128):18, 24-27, 133  
shipboard sampling procedures, A(128):179  
shorebased laboratory studies, A(128):179-180  
Site 798, A(128):24-27, 125-126, 178-180; B:755-776

microfaults  
and Japan Sea stress field, B:1187

Site 798, A(128):143ff

Site 799, A(128):269-272  
and veins, B:1180

microfossils. *See also* biostratigraphic zonations; biostratigraphy; names of individual fossil groups  
abundance and preservation, A(127):19, 102, 192, 194-195, 269-271, 351-352, 354, A(128):28-30, 99, 158-159, 163, 298, 308-309

paleoenvironmental interpretation, A(127):20-22, 102-103, 199, 274, 356, A(128):99-100, 166, 312

phosphatized, B:64

preservation, B:1219

reworked, A(127):270, 356

Site 799, B:34-35

in tuffaceous clayey siltstone, A(128):68, 89

microfossils, calcareous, Leg 128, A(128):28

microfossils, siliceous  
abundance and preservation, B:341-357

microfossils (cont.)

## SUBJECT INDEX

abundance below shallowest opal-CT, A(127):19  
dissolution, A(127):169; B:353  
Leg 127, A(127):19  
Leg 128, A(128):28  
and opal-A to opal-CT transition, B:341  
and opal dissolution transition zone (ODTZ), B:342  
preservation, A(128):137  
as productivity indicators, B:429-430  
Site 794, B:341-357  
Site 795, B:341-357  
Site 797, B:341-357  
micropumice, in ash-flow tuff, B:793  
microquartz, in sandstone, B:144-148  
microtexture, Site 799, B:33-48  
mid-mean ridge basalt. *See* basalt, mid-ocean ridge  
Milankovitch cycles. *See also* eccentricity frequencies; obliquity cycles  
in FMS logs, B:1043-1044  
mineral-liquid-disequilibrium, and spinel zoning, B:844  
minerals, magnetic. *See* magnetic minerals  
minor elements  
  in augite, B:854  
  in basement, B:912-916  
  in dark/light cycles, B:568, 570-571  
  Site 794, B:912-913  
  Site 795, B:913-914  
  Site 797, B:914-916  
  Site 798, B:720-722  
  Site 799, B:722-723, 726-727  
Miocene, early/middle boundary, Site 799, A(128):299; B:1224  
Miocene, middle/late boundary, Site 799, A(128):299; B:796, 1224  
Miocene/Pliocene boundary  
  Site 794, A(127):97  
  Site 795, A(127):193; B:1223  
  Site 796, A(127):272  
  Site 797, A(127):351; B:1223  
  Site 798, B:1224  
  Site 799, A(128):298, 301-302, 305; B:1224  
Mitoku-type flora, Japan Sea, B:486  
Miura Group, veins observed, B:1180  
mixed sediment  
  Site 795, A(127):186-187  
Site 799, A(128):260  
mixed sediment, diatomaceous clayey, Site 799, A(128):256  
Mogami Channel, A(128):70  
Moho  
  Japan Basin, A(128):246  
  Yamato Basin, A(128):246  
  Yamato Rise, A(128):245-246  
molecular organic fossils  
  in dark/light cycles, B:670, 675  
  low-molecular-weight compound concentration, B:673  
  Site 798, B:669-670  
  Site 799, B:669-670  
molybdenum  
  Site 794, B:1362  
  Site 795, B:1365  
  Site 797, B:1366  
  Site 798, B:1370-1371  
monactinal monaxons  
  Site 795, B:542-543  
monaxons  
  Site 795, B:541-543  
Mongolia, dust storms, B:394  
monsoons

and dark/light cycles, B:574-575  
Japan Sea, B:320-321, 326  
Monterey Formation  
  dolomite, B:75  
  lithostratigraphy, A(128):354  
  veins observed, B:1180  
Monzen Formation  
  and age of rifting, B:1202  
  deposition of, A(128):17  
MORB. *See* basalt, mid-ocean ridge  
mordenite, Site 797, A(127):344  
morphotaxa, faunal/flora, recognized in BSEM images, B:548  
Mossbauer spectroscopy, B:739-746  
  absorption areas and sediment chemical compositions compared, B:741-742  
  iron species depth profiles, B:742-746  
  Mossbauer parameters, B:740-742  
  and partial dissolution of sediment samples, B:741  
  typical spectra, B:743  
Mount St. Helens, stress field, B:1323  
mud clasts, Site 799, A(128):267  
mud plumes, Site 799, A(128):268-269; B:1178  
mudstone, Site 799, B:42  
Musasi Basin, dredge sample, A(127):176-177  
Nakanami Formation, A(128):129  
Nankai Trough, A(127):5-6, A(128):9  
  basement age, B:749  
  fissility, A(128):146  
  helium isotope ratios, B:748-749  
  interstitial water oxygen isotopes, B:698  
  veins, A(128):145-146; B:1180  
nannofossils, calcareous  
  abundance and preservation, A(127):98-99, 193-195, 273, 351-355, A(128):28-30, 162-165, 305-310; B:161  
  and authigenic carbonates, B:166  
  and basal sediment age, B:1335  
  and carbonate-compensation depth fluctuations, B:1220  
  datums, Site 798, B:163-166  
  dissolution, A(128):162; B:166  
  diversity, B:175-177  
  events, B:172-174, 178  
  and Japan Sea formation age, B:1220  
  Leg 127, A(127):19-20  
  Leg 128, A(128):28, 30  
  magnetostratigraphy, B:163-164  
  monospecific assemblages, A(128):307  
  and oceanographic front fluctuations, B:167-168, 1220  
  paleoenvironmental interpretation, A(128):164-165  
  preservation determinations, B:161-162  
  reworked, A(127):99, 194  
  and sedimentation rate, B:1043-1044  
  Site 794, A(127):98-99; B:174-176, 1221  
  Site 795, A(127):193-195; B:176, 178  
  Site 796, A(127):273; B:176, 178  
  Site 797, A(127):351-355; B:177-179, 1223  
  Site 798, A(128):162-165; B:155-161, 164-166  
  Site 799, A(128):305-310; B:78, 155-163, 166  
  species diachrony, B:164  
  stratigraphic standards used, B:162-164  
  and surface-water temperature, A(127):273  
  units, Site 798, B:165-166  
  zonations, A(127):20, 98-99, 193-195, 273, 351-355, A(128):162-165, 305-310; B:172-174  
  naphthenic envelope, Site 799, B:627-628

natural gamma-ray tool (NGT). *See* gamma ray, logging  
natural remanent magnetization (NRM). *See* magnetic properties  
navigation, during downhole seismic experiment, B:1107-1108  
neodymium  
  in ash layers, B:1386  
  in basement rocks, B:807  
  Site 798, B:1370-1371  
  vs. strontium, B:902  
  vs. titanium, B:902  
neodymium isotopes. *See* isotopes, neodymium  
neopentane, Site 799, A(128):321-322  
nesquehonite, and alkalinity, B:92  
neutron activation (INAA) analyses  
  of basement, B:912-916  
  Site 794, B:912-913  
  Site 795, B:913-914  
  Site 797, B:914-916  
neutron porosity. *See* porosity, logging  
NGT. *See* gamma-ray logging  
nickel  
  and basalt fractionation models, B:873  
  in dark/light cycles, B:569  
  and fractionation models, B:881  
  Site 794, B:893, 1362  
  Site 795, B:1365  
  Site 797, B:893, 920, 1366  
Niigata Basin, lithostratigraphy, A(128):17  
niobium  
  basement, A(128):99  
  vs. zirconium, A(128):100; B:783  
Nishikurosawa Formation  
  deposition, A(128):17  
  and subsidence history of Oga Peninsula, B:1202  
nitrate, and nitrate-reducing bacteria, B:770  
nitrogen  
  in dark/light cycles, B:431  
methane correlated, B:750  
Site 794, A(127):113-114  
Site 795, A(127):209-212; B:708-709  
Site 796, A(127):283-285  
Site 797, A(127):365-367  
Site 798, A(128):176-177, 189-192  
Site 799, A(128):323-324, 334-338, 341  
  vs. carbon, organic, A(128):343-344  
nodules, carbonate  
  with diatoms, A(127):193  
  Site 795, A(127):186  
  Site 797, A(127):343  
nodules, phosphatic, Site 797, A(127):344, 346  
non-euoxic-anoxic condition. *See* oxygenation condition  
normal faults  
  in basement, A(128):91-93  
  and fractures, A(127):188  
Kita-Yamato Trough, A(127):334  
Oki Ridge, A(128):130  
Oki Trough, A(128):130  
Site 795, A(127):188  
Site 798, A(128):145; B:1181-1182  
Site 799, A(128):263, 267, 269-272, 294-295; B:1182-1183, 1189, 1190  
and stress field, B:1182-1183  
and veins, B:1178  
Yamato Basin, A(128):130  
Yamato Rise, A(128):76  
Yamato Rise E, A(127):82-83  
normalization factors, and geochemical logs, B:1029-1030, 1034-1035  
North American Plate, A(127):5, A(128):9

North Atlantic Deep Water, and biosiliceous sediments, B:439  
 North Korean Current, A(128):21; B:343, 440  
 northern hemisphere glaciation. *See* glaciation  
 Noto Peninsula  
   diatom abundance, B:311  
   and Japan Sea stress field, B:1185  
   rock exposures, A(128):129  
   uplift, A(128):158  
 NRM intensity. *See* magnetic properties  
 Nunivak Subchron  
   Site 797, B:1223  
   Site 799, A(128):314; B:1224  
 nutrient supply, and formation of color-banded bedding, A(127):349-351  
 oblique electrical resistivity experiment. *See* electrical resistivity experiment  
 obliquity cycles  
   in biogenic opal, B:447  
   in dark/light cycles, B:446-447, 571, 574-575  
   in FMS logs, B:1043-1044  
   in geochemical logs, B:1021  
   in logs, B:403, 1221  
   of surface-water temperatures, B:467-468  
 obliquity frequency. *See* obliquity cycles  
 ocean bottom seismometer (OBS). *See also* air-gun profiling experiment; downhole seismic experiment  
 air-gun signals in downhole seismic experiment, B:1162  
 anisotropic models, B:1117  
 array location, B:1070, 1077, 1109, 1159  
 and crustal structure, B:1075, 1342  
 data variance, B:1115  
 downhole seismometers compared, B:1061, 1078, 1084  
 first arrival traveltimes, B:1089  
 frequency response, B:1159, 1162  
 and navigation, B:1107-1108  
 ray tracing results, B:1092-1103  
 record sections obtained, B:1080-1083  
 slowness vectors, B:1118  
 specifications, for downhole seismic experiment, B:1160  
 traveltime, offset distance, and basement depth correlated, B:1112  
 velocity structure, one-dimensional, B:1078-1081, 1085-1086  
 velocity structure, two-dimensional, B:1081-1083, 1086, 1089  
 oceanic crust. *See* crust, oceanic  
 oceanization, Japan Sea, A(128):11  
 oceanographic front  
   and calcareous nannofossils, B:167-168, 1220  
   in Japan Sea, B:155, 167, 458  
   oscillations, B:167-168  
   oxygen isotope curve correlated, B:168  
   and productivity, B:167  
   and sea surface isotherms, B:167  
 oceanographic frontal boundary. *See* oceanographic front  
 ODTZ. *See* opal dissolution transition zone  
 Oga Peninsula, A(128):70  
   age, B:1202  
   basement, B:1202  
   benthic foraminifers, A(127):198; B:379, 381  
   depositional environment, B:1203  
   lithostratigraphy, A(128):17-18; B:1203  
   paleobathymetry, B:1203-1204, 1210  
   planktonic foraminifers exposed, B:459  
   sediment accumulation rate, B:1204  
   stress field, B:1204-1205  
   subsidence/uplift history, B:1202-1205

uplift, A(128):158; B:1202-1205  
 OI. *See* oxygen index  
 Okhotsk microplate, A(128):9  
 Okhotsk Sea  
   diatoms, B:309-310  
   formation, A(128):13  
   and opal flux, B:439  
 Oki-Dogo Island, ash layers from, A(128):156  
 Oki Islands, A(128):126-127  
   basement, A(128):127-129  
   Green Tuff rocks, A(128):128  
   and low-chlorinity pore fluids, A(128):172  
 Oki Ridge  
   ash layers, A(128):122  
   basement, A(128):127-129  
   bathymetry, A(128):126-127  
   benthic foraminifers, B:367-369, 373-375, 382  
   biostratigraphy, B:1223-1224  
   bottom water conditions, A(128):125  
   carbonate-compensation depth, A(128):122, 158; B:409  
   chronostratigraphy, B:317-318  
   consolidation/permeability studies, B:1123-1133  
   crustal structure, A(128):127-128  
   cyclic lithofacies, A(128):195-196  
   depositional history, A(128):157-158, 194-195  
   diatoms, B:361  
   ebridians, B:241-244  
   eolian supply, B:409, 419  
   faulting, A(128):194-195  
   forests, B:317  
   free-air gravity anomalies, A(128):127, 133  
   Green Tuff rocks, A(128):127-128  
   heat flow, A(128):125, 127, 131; B:1302  
   Layer 1, A(128):127  
   Layer 2, A(128):127-128  
   Layer 3, A(128):127-128  
   magnetic anomalies, A(128):127, 132  
   magnetostratigraphy, B:973-974, 1223-1224  
   oceanographic conditions, A(128):166, 195-196  
   oceanographic drilling objectives, A(128):18-19  
   oxygen isotope stratigraphy, B:444  
   paleoclimatic conditions, B:419-421  
   paleodepth, B:373  
   paleoenvironment, B:379-380  
   pollen, B:318-324  
   radiolarians, B:225, 227, 229-230  
   sediment, B:409-422  
   sedimentary rocks dredged, A(128):129  
   sedimentation, A(128):131-132; B: 1227  
   seismic stratigraphy, A(128):34, 125  
   silicoflagellate, B:241-244  
   Site 798, B:393, 409, 423-425, 440, 458-459, 560, 605, 667, 720, 739  
   tectonic setting, A(128):130-131, 134  
   uplift, A(128):35, 122, 157, 194-195; B:155, 373, 419, 1208  
 Oki Ridge Basin, seismic stratigraphy, A(128):122  
 Oki Trough  
   bathymetry, A(128):126-127  
   crustal structure, A(128):130  
   free-air gravity anomalies, A(128):127  
   heat flow, A(128):131  
   Layer 1, A(128):127  
 Okinawa Trough, sulfide deposits, A(128):21  
 Okushiri Island, A(127):251, 254, A(128):76  
 Okushiri Ridge  
   anomalous heat flow, A(127):300-301

basement, A(127):176, 312  
 bathymetry, A(127):251  
 biostratigraphy, B:1223  
 calcareous nannofossils, B:176, 178  
 compression, A(127):255  
 crustal structure, A(127):251-252  
 diatoms, B:270-275, 285  
 dredge samples, A(127):254  
 earthquakes along, A(127):178  
 fluid flow along faults, A(127):275  
 folding, A(127):255-256  
 frictional heating along thrust faults, A(127):247  
 heat flow, A(127):27, 30, 300-301, 314; B:1303, 1307  
 interstitial water chemistry, B:1265  
 magnetostratigraphy, B:1223  
 opal-A to opal-CT transition, B:50  
 opal-CT to quartz transition, B:50  
 physical properties, B:1275  
 pollen, B:471, 481, 484-485  
 sediment thickness, A(127):313-314  
 sedimentation, A(127):256-258; B:288, 1227  
 seismic stratigraphy, A(127):256-258, 307, 312-313  
 slumping, A(127):268  
 tectonic history, A(127):176  
 thermal model, A(127):16  
 uplift, A(127):19, 30, 176, 247, 258, 268, 314; B:1208  
 Olduvai Event. *See* Olduvai Subchron  
 Olduvai Subchron  
   gamma-ray logging, B:403  
   and sedimentation rate, B:975  
   silicoflagellate and ebridian zones correlated, A(128):162  
   Site 794, B:218, 224, 970  
   Site 795, A(127):174, 199; B:970, 1223  
   Site 797, B:224  
   Site 798, A(128):30, 125, 170; B:164, 564, 974, 1224  
   Site 799, A(128):314; B:974, 1224  
 olivine  
   altered to clay, B:885-886  
   experimental chemical compositions, B:866  
   forsterite and, B:850  
   importance in basalt fractionation models, B:873-874  
   in melting experiment, B:861-862, 895  
   and parent magma, B:855-856  
   Site 794, B:850  
   Site 797, B:850  
 Oman margin, and chloride, A(128):172  
 Onnagawa Formation, A(128):13, 265  
   ash-flow tuffs correlated, B:796  
   deposition, A(128):17  
   dolomite, B:75  
   foraminifers, B:187  
   lithostratigraphy, A(128):354  
   opal flux, B:439, 446  
   paleoenvironmental interpretation, B:1202  
 Ontong Java Plateau, magnetic properties, B:951  
 ooze, clayey diatom  
   and revised lithostratigraphy, B:1230-1232  
   Site 794, A(127):90  
   Site 796, A(127):264  
   Site 797, A(127):341-343  
   Site 799, A(128):260  
 ooze, diatom  
   aluminum, logging, A(127):306-307  
   and enhanced productivity, A(127):267  
   and oxygenation conditions, B:350  
   and revised lithostratigraphy, B:1230-1232

sedimentation rate, A(127):204  
 Site 794, A(127):90, A(128):77  
 Site 795, A(127):186  
 Site 797, A(127):341–343  
 Site 798, A(128):124, 137–138  
 Site 799, A(128):256, 260  
**opal**  
 abundance variations seen in logs, B:398–400  
 and geochemical logging, B:1022–1024  
 physical properties correlated, B:400  
 stratigraphy, B:1221  
**opal, biogenic.** *See also* opal-A; opal flux; silica, biogenic  
 abundance, B:443, 446–447, 449–455  
 cycles, B:444–446  
 in dark/light cycles, B:432, 446, 569  
 obliquity cycles, B:447  
 seismic expression, B:1140–1141, 1143  
 Site 798, B:443–446, 449–455  
 variations in supply, B:404  
 vs. age, B:445  
 vs. depth, B:445  
**opal-A.** *See also* opal; opal, biogenic; silica, biogenic  
 abundance, A(127):94; B:1235–1241  
 deepest occurrence, A(127):187  
 and lithium, A(128):174–175  
 and low-chloride interstitial fluids, B:608–610  
 physical properties interrelationships, B:25–27  
 vs. excess silica, B:1253  
**opal-A to opal-CT transition,** A(128):28; B:6–17, 1145–1156. *See also* silica diagenesis  
 and acoustic impedance, B:1146  
 age, A(127): 19; B:251, 1145–1147, 1305–1306  
 as barrier to interstitial water, A(128):244  
 bottom simulating reflector, B:1145–1147, 1149, 1152–1153  
 and chloride concentration, B:609–610  
 and deepest diatom occurrence, A(127):264  
 and density, B:18, 1146–1147, 1288  
 depth, B:1224–1227  
 diatoms, B:3, 249, 313, 361, 1230–1232  
 and dissolution patterns, B:311  
 and dissolved silica maximum, A(127):174  
 factors controlling phase change, B:1145  
 in geochemical logs, B:1417  
 and heat flow, B:1147–1148, 1150, 1156  
 and inorganic geochemical data, A(127):23, 187, 204–209, 362–364, A(128):32, 321; B:607  
 Japan Sea, A(127):31; B:49, 341  
 Leg 127, A(127):19  
 Leg 128, B:359  
 and lithium content, B:1266  
 lithology and, B:17  
 in logs, A(127):139, 251, 302, 393–394, A(128):186–187, 333; B:3, 15–16, 19, 21–22  
 and opal-CT abundance, B:10  
 oxygen isotopes and, B:641–642  
 and physical properties, A(127):26, 127, 133–134, 187, 223, 290–295, 388–389, A(128):33; B:10–18, 1145, 1277–1278, 1284, 1305  
 and porosity, B:3, 6, 16, 18  
 and radiolarian preservation, A(127):198, 355, A(128):303  
 and radiolarians, B:226, 291–292, 295  
 and rate of opal-CT precipitation, B:1306  
 and sedimentation rate, B:1147–1148  
 seismic expression, A(127):28, 143, 169, 179, 234–236, 251, 307–314, 403, 407,

410, A(128):13, 193; B:3, 1141ff, 1305  
 and silica, A(128):125, B:1263  
 and siliceous microfossils, A(128):28; B:341  
 Site 794, A(127):111  
 Site 795, A(127):169, 198; B:4, 50  
 Site 796, A(127):247; B:50  
 Site 797, A(127):343, 351; B:5, 50  
 Site 798, A(128):149, 240  
 Site 799, A(128):238, 260, 278  
 and slumping of slope, A(127):268  
 in synthetic seismograms, B:1146–1148  
 temperature, A(127):19, 137, 232, 347, 392, A(128):150, 280; B:14, 1145–1147, 1152, 1263, 1305–1307, 1375  
 and thermal conductivity, B:1147  
 and thermal gradient, A(127):268, 314  
 and thermal history, Japan Sea, B:1305–1307  
 as thermal marker, B:3  
 thickness, B:16, 1240, 1281–1282  
 time-temperature plot, B:1266, 1288  
 transition temperature and sedimentation rate correlated, B:1306  
 and velocity, B:18  
 and volatile hydrocarbons, A(128):323  
 X-ray diffraction analyses, A(127):347; B:10, 17–18  
 and zeolites, B:10  
**opal abundance**  
 core analyses vs. log estimates, B:404  
 and dissolution patterns, B:310–311  
 logging estimates, B:402  
 and productivity, B:311  
 Site 798, B:401–402  
 vs. density, B:402  
 vs. porosity, B:402  
**opal accumulation rate.** *See* opal flux  
**opal-CT**  
 abundance, A(127):94, A(128):141; B:53, 1235–1241  
 deepest occurrence, A(127):189, A(128):280  
 formation, A(127):109; B:49  
 isotopic temperatures, B:55–56  
 and opal-A to opal-CT transition, B:10  
 oxygen isotopes, B:51–54  
 permeability, B:1263  
 physical properties, B:25–27  
 porosity, B:1263  
 shallowest occurrence, A(127):187, 343, 347, A(128):124, 138, 149, 174, 278  
 Site 795, A(127):190  
 and thermal gradient, B:49  
 time-temperature plot for opal-CT formation, B:1266  
 vs. excess silica, B:1253  
**opal-CT to quartz transition,** B:16–17. *See also* silica diagenesis  
 age, A(127):19  
 and core recovery, B:1038  
 and inorganic geochemical data, A(127):204  
 Japan Sea, A(127):31; B:49  
 in logs, A(127):303, A(128):34, 333; B:3, 23  
 and physical properties, A(127):223–227  
 and porosity, B:3  
 resistivity logging data and, B:1040  
 SEM photographs, B:31  
 and silica content, B:1263  
 silicon content, A(127):307  
 Site 794, B:50  
 Site 795, A(127):169; B:4, 50  
 Site 796, A(127):265; B:50  
 Site 797, A(127):347; B:4, 50  
 Site 799, A(128):238, 242, 260, 280  
 temperature, A(127):19, 137, A(128):280; B:17, 1263, 1375  
 thickness, B:17, 1240, 1281–1282  
 time-temperature plot, B:1288  
 X-ray diffraction analyses, B:24  
**opal dissolution transition zone (ODTZ)**  
 age, B:350  
 Japan Sea, B:342  
 lithostratigraphy correlated, B:347–349  
 and Northern Hemisphere glaciation, B:350  
 photomicrographs, B:355–357  
 and radiolarians, B:345–347  
 Site 794, B:344  
 Site 795, B:345  
 Site 797, B:345  
 thickness, B:347, 350  
**opal flux.** *See also* opal, biogenic; opal-A; silica, biogenic  
 in Bering Sea, B:439  
 in Onnagawa Formation, B:439, 446  
 in Santa Barbara Basin, B:439  
 in Sea of Okhotsk, B:439  
 Site 798, B:439–455  
**operations.** *See particular site*  
**orbital obliquity**, in logs, A(128):34, 342  
**orbital obliquity cycles.** *See* obliquity cycles  
**orbital precession cycles.** *See* precession cycles  
**organic carbon accumulation rate.** *See* carbon, organic  
**organic fossils.** *See* molecular organic fossils  
**organic matter.** *See also* carbon, organic; molecular organic fossils; organic matter decomposition  
 autochthonous, B:627–628  
 bacterial degradation, A(127):22  
 and barium remobilization, B:654  
 and boron content, B:640  
 composition, B:423–429  
 in cyclic lithofacies, A(128):140  
 in dark/light cycles, B:446, 564, 584, 668–669  
 degradation, A(128):172–173, 318; B:646  
 diagenesis, B:1262–1263, 1267  
 and geochemical logs, A(128):187  
 and hydrogen index (HI), B:628  
 and inorganic geochemical data, A(127):204  
 lipid fraction, A(128):324  
 maturity, B:429  
 origin, A(128): 125, 324  
 and paleoenvironmental conditions, B:670–672  
 recycled, B:625  
 and sedimentation rate, B:1265  
 Site 798, A(128):125; B:446, 667–669  
 Site 799, A(128):245, 323–324; B:624, 667–669  
 and thermal maturity, B:671–672  
**organic matter decomposition**  
 and high ammonium values, A(127):279–280  
 and interstitial water analyses, B:607  
 Japan Sea, A(127):31  
 and salinity, A(128):172  
 Site 799, A(128):318  
**organic petrology.** *See* organic matter  
**organic-rich sediments.** *See* sediments, organic-rich  
**orthoclase**, vs. anorthite, B:122  
**orthopyroxene**  
 in ash layers, A(128):151; B:1379  
 in melting experiment, B:895  
**Oshima Island, basement,** A(128):73  
**Oshima Plateau, dredge samples,** A(127):78  
**overburden pressure.** *See* consolidation  
 and stress field analysis, B:1050–1051

overconsolidation. *See* consolidation  
overconsolidation ratio (OCR). *See* consolidation  
oxeas  
Site 795, B:542  
oxic conditions. *See* oxygenation conditions  
oxides. *See specific elements*  
oxides, elemental. *See specific element*  
oxygen  
benthic foraminifers and, B:528  
fugacity, inferred from spinel analyses, B:842-844  
in Japan Sea, A(128):20  
oxygen index (OI)  
in dark/light cycles, B:431-432  
Site 794, A(127):115-118  
Site 795, A(127):213, 219  
Site 796, A(127):285  
Site 797, A(127):367  
Site 798, A(128):177, 196; B:425, 668-669  
Site 799, B:429, 624, 626-627, 668-669  
oxygen isotopes. *See* isotopes, oxygen  
oxygen level, and formation of color-banded bedding, A(127):349  
oxygenation conditions  
and bioturbation, A(127):267, 315  
and burrows, B:1232  
and carbon/sulfur ratio, B:1249  
and Ce anomaly, B:692  
and dark/light cycles, B:432, 574-575, 590-591, 595  
determined by sulfur vs. organic carbon plots, B:708-709  
and diagenetic carbonate, B:93-94  
and dolomite formation, B:92  
glacial/interglacial variation, B:590-591  
Japan Basin, B:201-208  
Japan Basin N, A(127):199  
Japan Sea, B:216, 342-343, 430, 678-679, 705, 713-715, 1221  
Kita-Yamato Trough, A(128):354-355  
and manganese concentration, B:90  
and organic carbon content, A(127):283  
and organic geochemical data, A(127):267  
and oxygen isotopes, B:442-443  
Site 794, A(127):96; B:209  
Site 795, A(127):212; B:708-709, 713-715  
Site 796, A(127):247, 251, 267  
Site 797, A(127):349; B:209, 593  
Site 799, B:76-77, 91-92  
and wind velocity, B:447  
Yamato Basin, B:189, 201-208  
Oyashio Current, B:343  
climate and, B:326  
cold surface water, B:457  
Japan Sea and, A(128):30  
Site 799, A(128):312  
*p*- wave velocity. *See* velocity  
Pacific Bottom Water, helium, B:748  
Pacific Deep Water, A(127):16  
Japan Sea and, B:365  
Pacific Ocean  
benthic foraminifers, B:505  
deep-water inflow, B:528  
eolian dust, B:394  
Japan Sea and, A(128):312; B:180, 299-300  
tectonic setting, A(128):8  
Pacific Ocean, equatorial,  $C_{org}$ /barium ratio, B:653-654  
Pacific Ocean N  
biosiliceous sediments, B:439  
diatoms, A(128):160, 302, 315-316; B:258, 341, 358  
foraminifer coiling ratio, B:459

paleoclimate studies, B:459  
pollen, B:320  
radiolarians, B:228  
silicoflagellate, B:237  
Site 794, A(127):96  
Pacific Ocean NE, planktonic foraminifers, B:465  
Pacific Ocean NW, diatoms, B:309, 360  
Pacific Ocean SW, paleoenvironment, B:338  
Pacific Ocean W  
eolian dust, B:403  
marginal basins, B:1311, 1333-1334  
subduction zones, B:1333-1334  
tectonic setting, A(127):7  
Pacific Plate, A(127):5, A(128):9  
electrical resistivity structure, B:1351  
Pacific Plate oceanic model (PPOM), B:1304  
paleodepth  
benthic foraminifers and, B:504-506  
Japan Basin, B:201  
Japan Sea, B:526, 528-529  
Site 797, A(127):357  
Site 798, B:373  
Site 799, B:373, 379  
Yamato Basin, B:201  
paleoecology, Japanese archipelago, B:320, 334  
paleoenvironment. *See also* specific regions and topics  
pollen and, B:320  
Site 794, A(127):103, A(128):99-100  
Site 795, A(127):199  
Site 796, A(127):274  
Site 797, A(127):357  
Site 798, A(128):166; B:379-380  
Site 799, A(128):312; B:381  
paleomagnetics. *See* magnetic properties; magnetostратigraphy  
paleoproduction. *See* productivity  
palyнологy, Japan Sea, B:317-321, 325-338  
parent magma  
and augite, B:854  
of basalt, B:855-856  
of basement, B:893-894  
estimates, B:894  
Japan Basin, B:927-928  
Japan Sea, B:1341  
origin, B:897  
production depth, B:897  
Site 794, B:844-845, 917  
Yamato Basin, B:927-928  
pebbly claystone. *See* claystone, pebbly  
PEF. *See* photoelectric factor  
pelite, Site 799, B:33  
*n*- pentane, Site 799, A(128):321-322  
peralkaline glass, B:1378  
periodicity, orbital. *See also* obliquity cycles  
in logs, B:396-398  
permeability  
in biosiliceous sediments, B:1128  
and grain size, B:1127-1128  
and opal-CT formation, B:1263  
porosity effect, B:1128-1129  
Site 798, B:1127-1128, 1131-1132  
Site 799, B:39, 1128, 1131-1132  
and temperature measurements, A(127):299  
vs. porosity, B:1132  
Peru/chile Current, diatoms, A(128):16  
Peru margin  
bacteria, B:755, 757, 759, 761, 769, 773  
chloride, A(128):172  
D-phosphate nodules, B:38  
fluoride, B:66  
sediment consolidation, B:1277  
vein fillings, B:1178

petrology, experimental  
AMF diagram for experimental glasses, B:865  
An mole percent, Plagioclase vs. Fo mole percent, B:867  
anhydrous melting-phase relations, B:896  
augite, Cr and Ti vs. Mg/(Mg + Fe), B:867  
augite, Fe/Mg vs. Fe/Mg for coexisting liquid, B:868  
of basaltic basement, B:861-868, 891-898  
chemical compositions of experimental materials, B:863-864, 866-868, 894  
crystallization, B:864-866  
and differential processes of basement basaltic rocks, B:895-897  
equilibrium materials, B:862-863  
exchange coefficients, B:864  
experimental conditions, B:862  
experimental liquid line of descent, B:863  
experimental phase proportions vs. liquid percentage, B:862  
experimental pyroxene-liquid relations and natural augite compared, B:878-879  
experimental results and Site 794 basalt compared, B:870  
Fe/Mg olivine vs. Fe/Mg coexisting liquid, B:866  
melting experiments, B:861, 894-897  
mineral-liquid equilibria, B:862-863  
minerals crystallized, B:861-862, 895-896  
normative projections, B:868  
precision of microprobe analyses, B:862  
and primary magma production, B:897  
procedures used, B:861, 895  
starting materials, B:861-862, 894-895  
pH  
in interstitial water, B:94  
Site 794, A(127):107  
Site 795, A(127):205  
Site 796, A(127):283  
Site 797, A(127):364, 374  
Site 799, A(128):317-318, 328; B:610  
phase equilibria, in basalt, B:870  
phasor dual induction tool (DIT). *See* resistivity, logging  
phenocrysts  
augite, B:851-853  
in basaltic basement, B:892  
ilmenite, B:853  
olivine, B:850  
plagioclase, B:850-851  
Site 794, B:849  
Site 795, B:850  
Site 797, B:850  
spinel, B:853  
Philippine Basin, rare earth elements, B:719  
Philippine Sea  
basement isotope geochemistry, B:805  
Pb-Pb plots of arc volcanic rocks, B:808  
Philippine Sea Plate, A(127):5-6, A(128):9  
electrical resistivity structure, B:1351  
phillipsite, Site 797, B:139  
phonolite, ash layers, B:793  
phosphate. *See also* D-phosphate; F-phosphate; phosphorus  
abundance, B:36  
allochthonous, B:64, 73  
chemistry, B:64-65  
in diagenetic sequence, B:39  
in interstitial water, B:36-38, 64-66, 607, 1264  
microprobe analyses, B:64-67  
and organic carbon decomposition, B:1262  
pelagic, B:36

phosphate (cont.)

## SUBJECT INDEX

peloidal, B:36  
 pristine, B:63-64, 73  
 and sedimentation rate, B:67  
 and siliceous veins, B:64  
 Site 794, A(127):108  
 Site 795, A(127):205  
 Site 796, A(127):279-280  
 Site 797, A(127):362, 368  
 Site 798, A(128):173, 181; B:63-74  
 Site 799, A(128):317-318, 328; B:36-38, 47, 63-74, 610  
 source of allochthonous particles, B:67-68  
 total oxide values, B:65  
 X-ray fluorescence analyses, B:64-65  
 phosphate, Sites 798 and 799, B:64-66  
 phosphatic nodules. *See* nodules, phosphatic  
 phosphatic sediments, Kita-Yamato Brink,  
 A(128):249  
 phosphorus. *See also* phosphate  
 Japan Sea, B:68-69, 1237  
 $P_{org}/P_{inorg}$  ratio, B:71  
 phosphorus oxide. *See* phosphorus  
 photoelectric factor (PEF), and magnesium and sodium content, B:1395, 1415  
 phreatomagmatic eruption, submarine, B:119, 123-126  
 physical properties. *See also* specific physical properties analyses; *specific physical properties logs*  
 basement, A(127):26, 135, 227-228, 389, A(128):102ff; B:1282-1284, 1289-1291  
 basin and rise sites compared, Japan Sea, B:1275-1296  
 consolidation studies, B:1123-1133  
 corrected analyses, B:990-1015  
 data quality, A(127):127, 220-222, 289-290, 382-383; B:987-988  
 determination of wet volumes, B:988-989  
 and diatom abundance, A(127):222  
 error identification, B:985-986  
 index properties, B:985-1015  
 of interbedded basement/sediments, A(127):394; B:1283-1284  
 interrelations at opal-A to opal-CT transition, B:17-18  
 Japan Sea, A(127):25-26, A(128):33  
 lithostratigraphy correlated, A(127):127-135, 222-227, 290-295, 383-389, A(128):182, 327-329; B:1143  
 logs correlated, A(128):338  
 opal-A interrelations, B:25-27  
 and opal-A to opal-CT transition, B:10-16, 1145, 1277-1278, 1281-1282, 1305  
 opal content correlated, B:400  
 opal-CT interrelations, B:25-27  
 permeability studies, B:1123-1133  
 quartz interrelations, B:25-27  
 seismic stratigraphy correlated, A(128):224, 379; B:1143  
 and silica diagenesis, A(128):280; B:3-31, 1281-1282  
 Site 794, A(127):73, 127-136, 138-140, A(128):102ff  
 Site 795, A(127):174, 220ff, 231  
 Site 796, A(127):251, 289-297, B:1275  
 Site 797, A(127):325, 382ff, 392-393, 395  
 Site 798, A(128):33, 180ff  
 Site 799, A(128):33, 324ff; B:1137-1140  
 Sites 794 and 795 compared, A(127):228  
 Sites 794 and 797 compared, A(127):389, 396  
 systematic errors corrected, B:985-1015  
 water loss discrepancy, B:988

X-ray fluorescence basement analyses compared, B:1283  
 phytane, Site 799, A(128):343; B:628-629, 633  
 phytoplankton, Japan Sea, B:167  
 Pisco Formation, veins, B:1180  
 placoliths  
 Site 799, A(128):309  
 plagioclase. *See* feldspar, plagioclase  
 plagioclase feldspar. *See* feldspar, plagioclase  
 plankton blooms. *See* blooms, plankton  
 plastic limit. *See* Atterberg limits  
 plasticity index. *See* Atterberg limits  
 plate reconstructions  
 geometrical constraints, B:1311-1315  
 Japan Sea, B:1321-1324  
 kinematic model used, B:1315  
 Miocene, B:1322-1324  
 Pliocene-Quaternary, B:1321-1322  
 Present time, B:1321  
 reconstruction poles, B:1325  
 tectonic events, B:1316-1317, 1324-1325  
 plateau ages  
 Site 794, B:820  
 Site 795, B:820-821  
 Site 797, B:820, 829  
 Pliocene/Miocene boundary, Site 794, A(127):96  
 Pliocene/Pleistocene boundary  
 and biosiliceous productivity, B:313  
 and diatom assemblages, A(128):160  
 diatoms and, B:250  
 and foraminiferal fauna, B:208  
 and microfossil dissolution, A(128):28  
 Site 794, A(127):97; B:1221  
 Site 795, B:1223  
 Site 796, B:1223  
 Site 797, A(127):351, 355-356; B:1223  
 Site 798, A(128):162, 166; B:1224  
 Site 799, A(128):301; B:1224  
 Pliocene/Quaternary boundary, Site 796, A(127):251  
 Pohang Basin, subsidence/uplift history, B:1205-1206  
 polarity. *See* magnetostratigraphy  
 pollen, B:1221  
 and basal sediment age, B:1335  
 distribution, B:489  
 flora represented, B:479, 484-486  
 and glacial/interglacial periods, B:395  
 Japan Sea, B:328-336, 395  
 marine, B:320-321, 337-338  
 Site 794, B:328, 471, 482, 485  
 Site 795, B:328, 471, 480, 484  
 Site 796, B:471, 481, 484-485  
 Site 797, B:328, 479, 483, 485-486, 1223  
 Site 798, B:318-324  
 Site 799, A(128):302-303  
 sources, B:332-333  
 stratigraphy, B:486, 488  
 zones, B:479, 484  
 polyaxons  
 Site 795, B:543  
 porcellanite  
 conglomerate pebbles, A(127):344-345  
 elastic moduli, A(127):223  
 in logs, A(127):27-28, 394; B:1040  
 and revised lithostratigraphy, B:1230-1232  
 seismic expression, A(127):407-408  
 Site 795, A(127):187  
 Site 799, A(128):260, 264; B:33, 40, 41, 44  
 pore gas. *See* interstitial gas  
 pore space, and clathrate formation, A(127):281  
 pore water. *See* interstitial water  
 porosity. *See also* physical properties

basement, B:1289  
 corrected analyses, B:987, 990-1015  
 data quality, A(127):127, 228-229; B:987  
 and diatom internal porosity, B:1277  
 errors in wet volume measurement, B:985-986  
 and geochemical logging, B:1022-1024, 1029  
 and horizontal stress, B:1053  
 in interbedded sediment/igneous rocks, B:1278  
 Japan Sea, B:1275-1278  
 and loss on ignition, B:1283  
 and normalization factors, B:1035  
 and opal-A to opal-CT transition, B:3, 6, 16, 18, 21, 1277-1278, 1282, 1298  
 and opal-CT formation, B:1263  
 and opal-CT to quartz transition, B:3, 1282  
 permeability and, B:1128-1129  
 Site 794, A(127):127-133, A(128):103, 105; B:6, 1050, 1278-1279  
 Site 795, A(127):223-227, 230; B:6, 1278-1279  
 Site 796, A(127):290-295; B:6, 1278-1279  
 Site 797, A(127):383-389, 391; B:6, 1278-1279, 1300  
 Site 798, A(128):181, 197-198, 202-205, 209; B:6, 1278-1279  
 Site 799, A(128):325, 345-349, 355; B:6, 39, 700, 1278-1279  
 thermal conductivity correlated, B:987  
 velocity and, B:1280  
 vs. density, B:25, 27, 986, 989, 1281, 1290  
 vs. loss on ignition, B:1290  
 vs. opal abundance, B:402  
 vs. permeability, B:1132  
 vs. thermal conductivity, A(127):135, 232, 395; B:25, 27, 989, 1287  
 vs. velocity, A(127):140, 395; B:25, 27, 1005, 1286, 1290  
 vs. water content, A(127):234  
 porosity, logging  
 and geochemical logs compared, B:1023  
 Site 794, A(127):142, 147, 164-167  
 Site 795, A(127):244-245  
 Site 796, A(127):303-306, 319-320  
 Site 797, A(127):395-396, 403, 419-421; B:7  
 Site 798, A(128):185-187, 212  
 Site 799, A(128):383-385, 389-392  
 vs. velocity, B:26  
 potassium  
 alteration and, B:909, 911  
 in ash layers, B:1385  
 and basement alteration, B:892, 1265-1266  
 in clay minerals, A(127):205  
 in igneous rocks, A(128):103  
 in interstitial water, B:1270  
 in Japan Sea sediment, B:1237  
 and magma suite distinction, B:869  
 partitioned into plagioclase, B:856  
 Site 794, A(127):109, A(128):108  
 Site 795, A(127):205  
 Site 796, A(127):280-281  
 Site 797, A(127):364, 371  
 Site 798, A(128):174-175, 184  
 Site 799, A(128):318, 332; B:611, 723  
 vs. hydrogen isotopes, B:1273  
 vs. oxygen isotopes, B:1273  
 vs. sodium, B:732, 858  
 XRF and geochemical log analyses compared, B:1028-1029, 1031-1035

potassium, logging  
 and diatom vs. clay content, B:1417  
 Site 794, B:1416, 1418-1421  
 Site 796, B:1416, 1422-1423  
 Site 797, B:1424-1427

Site 798, B:1400-1402  
 Site 799, B:1403-1404, 1406-1409  
 vs. aluminum, logging, A(128):215  
 potassium/aluminum ratio, Site 794, B:1249  
 potassium/argon age. *See K/Ar age; radiometric age*  
 potassium feldspar. *See feldspar, potassium*  
 potassium oxide. *See potassium*  
 precession cycles  
 and dark/light cycles, B:571, 574-575  
 in FMS logs, B:1043-1044  
 precipitation  
 Japan Sea, B:337, 442-443  
 of silica, A(127):174  
 vegetation and, B:325-326  
 preconsolidation pressure. *See consolidation pressure, hydrostatic, thermal conductivity and, B:1017-1019*  
 pressure compensation level (PCL), and vesicular hydroclasts, B:119  
 primary magma. *See parent magma*  
 primary productivity. *See productivity*  
 pristane, Site 799, A(128):343; B:628-629, 633  
 pristane/phytane ratio, Site 799, B:628  
 productivity  
 and barium content, B:652, 661-662  
 in dark/light cycles, B:431-432, 591-594  
 and diatoms, A(128):122; B:342  
 estimate methods, B:425  
 Japan Sea, A(128):21, B:167, 296, 299, 651  
 Kita-Yamato Trough, A(128):355  
 and opal abundance, B:311  
 and organic carbon, A(127):283; B:430  
 and sea ice, B:311  
 and sedimentation rate, B:1228  
 and silica content, B:694  
 and siliceous microfossil abundance, B:430  
 Site 796, A(127):274  
 and Tsushima Current, B:311  
 propane  
 Site 795, A(127):174, 213-216, 220  
 Site 797, A(127):368  
 Site 798, A(128):125, 176, 187  
 Site 799, A(128):244-245, 321-322, 339  
 proto-dolomite, Site 799, B:611  
 provenance diagrams, and tectonic setting, B:109  
 pull-apart basin, Japan Sea, A(128):12-13  
 pumice, long-tube, B:117, 130  
 pycnometer, and wet sample volume analysis, B:985-986, 988-989  
 pyrite  
 abundance, B:1235-1241  
 in ash layers, B:1392  
 in Black Sea, B:712  
 in dark claystone, A(127):267  
 in dark-colored sediments, A(127):211  
 in dark/light cycles, B:584, 588  
 in diagenetic sequence, B:39  
 in diatom-bearing claystone, A(127):261  
 in dolomite, A(128):148  
 energy dispersive spectra, B:1378  
 frambooidal, B:711  
 and high organic matter, A(127):283  
 and magnetic properties, B:947  
 and Mossbauer parameters, B:740  
 and oxygenation conditions, B:712  
 in sandstone, B:144  
 and sediment demagnetization, A(127):203, 275-276  
 sedimentary, B:711-713  
 Site 795, A(127):186-187, 189; B:717  
 Site 797, A(127):344

Site 799, A(128):243, 265; B:36, 46, 79, 745-746  
 Sites 798 and 799, B:670, 771, 775  
 and sulfur, A(127):212-213, 705-717; B:1249, 1254  
 pyrite, frambooidal, B:706-709  
 pyrite-plus-sulfur (PVS), Site 798, B:771, 775  
 pyroclastic component, sediment gravity flow processes, A(127):267  
 pyroclastic deposits  
 sandstone source rocks, B:107  
 Site 796, A(127):265  
 pyroclastic flows, of blue tuff, B:119  
 pyrogenic crystal chemistry, of blue tuff, B:126  
 pyrolylates, adsorption, A(127):285  
 pyrolysis-gas chromatography, Site 799, B:627-628, 632-633  
 pyroxene. *See also clinopyroxene; orthopyroxene; specific pyroxene minerals*  
 altered to clays, B:885  
 in ash layers, B:1379, 1388  
 chemical composition of alteration clay minerals, B:886  
 in dolerite, A(128):88  
 fractionation models, B:874  
 non-quadrilateral end-member molecules, B:858  
 Site 794, B:851-852  
 Site 795, B:852  
 Site 797, B:852-853  
 pyrrhotite, B:978  
 and magnetic properties, B:947  
 and magnetization, A(128):169  
 Site 797, B:949, 951  
 Q (Koenigsberger ratio). *See magnetic properties*  
 Q-mode cluster analysis, and planktonic foraminififers, B:188, 193-200, 208-209, 220, 224  
 QFR diagram, sandstone classification, B:106  
 quartz  
 abundance, A(127):94, 189; B:53, 1235-1244  
 and Chinese loess deposits, B:401  
 in dark/light cycles, B:584  
 glacial/interglacial variation, B:588  
 isotopic temperatures, B:55-56  
 origin, B:51  
 oxygen isotopes, B:51-54  
 physical properties interrelations, B:25-27  
 polycrystalline, B:107-108  
 replaced by carbonate, B:147  
 rutile inclusions, B:107  
 in sandstone, B:34, 104, 135  
 Site 795, A(127):190  
 Site 797, B:109, 592  
 Site 798, B:401, 411-415, 417  
 tourmaline inclusions, B:108  
 undulatory extinction, B:107  
 vs. excess silica, B:1253  
 vs. plagioclase, B:1245, 1252  
 vs. terrigenous component abundance, B:403  
 quartz, normative component, vs. iron/magnesium ratio, B:894  
 quartz accumulation rate, Site 798, B:411-420  
 quartz/feldspar ratio, Site 798, B:411-415, 417  
 quartz sand. *See sand, quartz*  
 Quaternary/Pliocene boundary  
 Site 795, A(127):192  
 Site 796, A(127):271  
 radiogenic isotopes. *See isotopes, radiogenic*  
 radiolarians. *See also microfossils, siliceous*  
 abundance and preservation, A(127):101-102, 198-199, 274, 356-357, A(128):28, 161-162, 303-305; B:1220 and climate fluctuation, A(127):274, 356

cycladophorid, B:299  
 dilution by detrital components, A(127):198  
 dissolution, A(127):198-199, B:353  
 dissolution-resistant, B:295-297  
 Leg 127, A(127):19-20  
 Leg 128, A(128):28  
 and opal dissolution transition zone, B:345-347  
 Pliocene/Pleistocene boundary, A(127):355  
 preservation, B:292-296, 1220  
 productivity, B:296, 299  
 pyritized, A(127):198  
 refractive index, B:296  
 reworked, A(127):198-199, 274; B:296  
 Site 794, A(127):101-102, A(128):99; B:293, 1221  
 Site 795, A(127):198-199; B:294  
 Site 796, A(127):274  
 Site 797, A(127):356-357; B:295-296  
 Site 798, A(128):161-162; B:225, 227, 229-230  
 Site 799, A(128):303-305; B:225-226, 233-234  
 species diversity, A(127):101-102, A(128):162  
 and water temperature, A(128):162  
 zonations, A(127):20, 101-102, 198-199, 274, 356-357, A(128):161-162, 303-305; B:1220, 1225-1226  
 radiometric age, B:819-836  
 age spectra, B:820-821, 828-836  
 argon isotopes, B:822-827  
 basement, B:1336-1337  
 biostratigraphic age compared, B:1337  
 of blue tuff, B:123, 126-127  
 crystallization age, B:821  
 evaluation of ages obtained, B:821  
 formation age, B:820-821  
 isochron plots, B:828-833  
 of Japan Sea rock samples, B:1334  
 plateau ages, B:820-821  
 radiometric and paleontologic ages compared, B:821-822  
 samples, B:819-820  
 Site 794, A:784-786  
 and timing of volcanic activity in Japan Sea, B:829-832  
 vs. sedimentation rate, B:126  
 radiometric dating. *See radiometric age*  
 rare earth elements (REE), B:677-695, 719-737.  
*See also specific rare earth elements*  
 associated with detrital fraction, B:691  
 basalt, B:900  
 basement, B:780, 912-916  
 and biogenic siliceous input, B:692  
 complexation with organic and inorganic ligands, B:730  
 correlation coefficients, B:691  
 and heavy minerals, B:691  
 LREE enrichment relative to HREE, B:725  
 mixing model of pelagic sediment into depleted mantle, B:817  
 mobility during alteration, B:911-916  
 and parent magma, B:927-928  
 processes controlling distribution, B:692  
 and silica content, B:682  
 Site 794, B:682-684, 688, 912-913, 916-919, 922, 1339  
 Site 795, B:683-685, 688, 913-914, 918, 924, 1338-1339  
 Site 796, B:685  
 Site 797, B:685, 688-691, 914-916, 920-921, 925, 1339-1340

Site 799, B:734  
 sources in sediment column, B:729  
 spidergram, B:783  
 terminology, B:681-682  
 and terrigenous sedimentation, B:691, 732  
 in volcanic glass, B:1387  
 vs. loss on ignition, B:919, 921  
 reconstructions. *See* plate reconstructions  
 recrystallization  
   Site 794, A(128):99  
   Site 799, A(128):303, (128):307  
 Red River Fault, offset along, B:1325  
 redox conditions. *See* oxygenation conditions  
 REE. *See* rare earth elements  
 reflection coefficient, Site 799, B:1138-1141  
 refractory elements. *See also* names of specific refractory elements  
   Site 798, B:706  
 remagnetization, and pyrite, A(127):203  
 resistivity, logging  
   core depths correlated, B:1040  
   core/fog correlation, B:398-400  
   electrical resistivity experiment compared, B:1354  
   of opal-A to opal-CT transition, B:22  
   opal-CT to quartz transition and, B:1040  
   Site 794, A(127):139-142, 144-146, 156-159, A(128):104, 107, 119-120; B:1069  
   Site 795, A(127):232, 237  
   Site 796, A(127):302-305, 321-322  
   Site 797, A(127):393-395, 400-402, 416-418; B:7  
   Site 798, A(128):185-187, 212, 225-228; B:399, 401-402  
   Site 799, A(128):332-338, 362-364, 380-382, 393-397  
   Sites 794 and 797 correlated, B:1039, 1041  
 resolution, in logs, B:396-397, 404-405  
 reverse faults  
   Site 799, A(128):263, 271-272; B:1183, 1189  
   and stress field, B:1182-1183  
   Yamato Basin N, A(128):75-76  
 rhodochrosite  
   and cerium concentration, B:90-91  
   chemical composition, B:81-82, 84-85  
   formation, B:85, 90-91  
   and iron distribution, B:742  
   isotopic composition, B:83  
   and Mossbauer parameters, B:741  
   occurrence, B:80  
   paleoenvironmental implication, B:94  
   petrography and textural relations, B:81  
   SEM photographs, B:97  
   Site 799, B:75-98, 744-745  
   X-ray diffraction analyses, B:79-80  
 ryodacite, volcanic glass, B:1378  
 rhyolite  
   ash layers, B:793  
   tuff, B:731  
   volcanic glass, B:1378  
 rhythmic sedimentation. *See* lithofacies, cyclic  
 rift propagation, in Japan Sea E, B: 1346  
 rifting. *See also* back-arc rifting  
 ripup clasts, in sandstone, B:104  
 rivers, pollen and, B:333  
 Rock-Eval analyses. *See also* hydrogen index (HI); oxygen index (OI); Rock-Eval pyrolysis;  $T_{max}$   
 carbon, organic, A(127):367  
 carbon, total organic (TOC), A(127):115, 213, 285  
 methods, A(127):114

S/S, vs. depth, A(127):121  
 Site 794, A(127):114-120  
 Site 795, A(127):218-219  
 Site 796, A(127):285, 288  
 Site 797, A(127):367, 376-378  
 Site 798, A(128):177, 195; B:425  
 Site 799, B:429  
 source character, A(127):115-118, 205, 285, 367  
 thermal maturity, A(127):118, 285, 367  
 $T_{max}$  vs. depth, A(127):121  
 Rock-Eval pyrolysis. *See also* hydrocarbon; hydrogen index (HI); organic matter; oxygen index (OI); Rock-Eval analyses;  $T_{max}$ ; total pyrolytic yields; transformation ratio (TR); van Krevelen plot  
   Site 798, B:669  
   Site 799, B:627, 669  
 rotaliids  
   Site 794, A(127):101  
 rotation event, Japan Sea, B:179, 528  
 rubidium  
   and ash alteration, B:1266  
   and basement, B:807, 1266  
   in interstitial water, B:1271  
   Site 794, A(127):109, B:1363  
   Site 795, A(127):205, B:1365  
   Site 796, A(127):280-281  
   Site 797, A(127):364, 371; B:1366  
   Site 798, B:1370-1371  
   vs. hydrogen isotopes, B:1273  
   vs. oxygen isotopes, B:1273  
 rutile, as quartz inclusions, B:107  
 S<sub>v</sub>. *See* hydrocarbons, free  
 S<sub>v</sub> + S<sub>t</sub>. *See* hydrocarbons, total pyrolytic yield  
 S<sub>v</sub>/(S<sub>v</sub> + S<sub>t</sub>). *See* transformation ratio  
 S<sub>v</sub>. *See* hydrocarbons, generatable  
 Sado Island  
   and Japan Sea stress field, B:1185  
   uplift, A(128):158  
 Sado Ridge, A(128):70, 76  
   basement, A(128):73  
 Sakhalin Island, A(128):15; B:1206-1207, 1210  
 salinity  
   and clathrate formation, A(127):281  
   glacial/interglacial cycles, B:591, 608  
   and organic matter decomposition, A(128):318  
   and oxygen isotopes, B:442-443, 447  
   Site 794, A(127):107  
   Site 795, A(127):205  
   Site 796, A(127):278-279  
   Site 797, A(127):364, 374  
   Site 798, A(128):172, 179  
   Site 799, A(128):317, 328; B:610  
   and sulfate removal, A(127):205  
 samarium  
   in basement rocks, B:807  
   Site 798, B:1370-1371  
 sand. *See also* sandstone; terrigenous component and clathrate formation, A(127):281  
 composition, A(127):268, B:216  
 differential compaction, B:39  
 and fluid migration, A(128):321  
 and foraminifer abundance, B:193  
 Kita-Yamato Trough, A(128):354  
 in logs, A(127):251  
 microturbidite, B:44  
 provenance, A(127):268  
 sedimentary structures, A(127):268, A(128):243, 265  
 seismic expression, A(127):313  
 shallowest occurrence, A(127):272  
 Site 796, A(127):250, 261-264

Site 799, A(128):258, 265  
 temperature measurements in, A(127):299  
 Toyama Fan, A(128):14  
 sand, bioclastic, Site 797, A(127):343  
 sand, glauconitic  
   origin, A(128):157  
   Site 798, A(128):124, 138, 141  
   Site 799, A(128):241  
 sand, quartz  
   origin, A(128):157  
   Site 798, A(128):124, 138  
   Site 799, A(128):264  
 sandstone. *See also* sand  
   alteration, A(127):345  
   carbonate cementation, B:39, 141  
   cementation, B:132, 134-135  
   classification, B:106  
   compaction, B:132  
   composition, A(127):345  
   deposition, A(127):19  
   diagenesis, B:131-151  
   dolomite-cemented, A(128):276  
   feldspar albitionization, B:131-151  
   feldspar composition, B:105-106, 136, 139  
   framework composition, B:104-105, 113  
   grain size, B:100, 104  
   interbedded with basalt, A(127):345-346  
   laminated, A(127):3347-3349  
   lithic, B:104  
   mineral replacement in, B:131-148  
   normally graded, A(127):264-265, 346, 348  
   oldest deposited, A(127):268-269  
   provenance, B:99-111, 148  
   and revised lithostratigraphy, B:1233  
   sedimentary structures, B:100, 104, 1233  
 Site 796, B:99-113  
 Site 797, A(127):344-346; B:99-113  
 Site 799, A(128):265, 292; B:33-34, 42-43, 99-113  
 source area location, B:110-111  
 source of carbonate, B:141-143  
 source-rock lithology, B:106-109  
 tectonic provenance, B:109-110  
 thickness, B:100, 104  
 zeolitization, B:131-139  
 sandstone, bioclastic, Site 797, A(127):346  
 sandstone, feldspathic, Site 799, B:131  
 sandstone, tuffaceous, Site 796, A(127):265  
 sandstone, volcaniclastic  
   Site 796, B:102, 131  
   Site 797, B:103, 131  
 Santa Barbara Basin, and opal flux, B:439  
 saponite, B:885-888  
 saturation vapor pressure, and index property data errors, B:986, 989  
 scandium  
   Site 795, B:707  
   Site 798, B:1370-1371  
 scanning electron microscope (SEM), and visual microanalysis, B:57-58  
 sea ice  
   in Japan Sea, A(127):20-22; B:167  
   Leg 127, A(127):20  
   and productivity, B:311  
 sea level  
   Japan Sea, B:180  
   Site 798, A(128):166  
 Sea of Okhotsk. *See* Okhotsk Sea  
 sea-surface temperature. *See* temperature, surface-water  
 seafloor spreading, in Japan Basin, B:1345-1346  
 Second Spring Layer, benthic foraminifers and, B:365

sediment accumulation rate. *See also* feldspar accumulation rate; mass accumulation rate; quartz accumulation rate; sedimentation rate  
 and barium distribution, B:660-661  
 of biogenic silica, A(128):122, 158  
 Brunhes Chron, B:468  
 of claystone, A(127):191  
 Japan Sea, A(127):22, 24, A(128):31  
 Oga Peninsula, B:1204  
 of siliceous claystone, A(127):191  
 Site 794, A(127):72, 105-107  
 Site 795, A(127):174, 203-204  
 Site 796, A(127):250-251, 277-278, 281-282  
 Site 797, A(127):324, 359-362  
 Site 798, A(128):31, 124-125, 171ff, 176-177; B:409-420, 446, 563-564  
 Site 799, A(128):31, 244, 259-260, 315-316, 324  
 and sulfate content, B:637, 646  
 sediment composition  
 carbonate components, B:1241  
 of dark/light cycles, B:579, 582-584  
 detrital components, B:1237-1239  
 of fine-grained rocks, B:1367  
 glacial/interglacial variations, B:588-590  
 Japan Sea, B:1233-1249  
 logging estimates, B:402-403  
 major elements, B:1235-1237, 1246-1248  
 of sand, B:216  
 of sandstone, B:104-105, 113  
 siliceous components, B:1239-1241  
 Site 794, A(127):92; B:1235-1237, 1242  
 Site 795, A(127):188; B:8-10, 1237-1240, 1243  
 Site 796, A(127):264  
 Site 797, A(127):345; B:11-14, 1240-1241, 1244  
 Site 798, A(128):138; B:398-400, 409-422  
 Site 799, A(128):258  
 sulfide components, B:1241-1249  
 variations seen in logs, B:398-400  
 sediment geochemistry. *See* geochemistry, sediment  
 sediment grain size. *See* grain size  
 sediment gravity flow deposits  
 and formation of color-banded bedding, A(127):349  
 Site 796, A(127):247, 268, 315; B:100, 110  
 Site 797, A(127):347; B:100  
 sources, A(127):267  
 sediment loading, sediment consolidation and, B:1130  
 sediment texture, Site 799, B:33-48  
 sediment thanatocoenosis, and diatom dissolution, B:309-316  
 sedimentary structures. *See also* specific sedimentary structures  
 in blue tuff, B:117  
 brittle structures, A(128):269-272; B:1181-1183  
 caused by compaction, B:38  
 color-banded bedding, A(127):186  
 dewatering structures, B:1176-1178  
 faults, A(128):143ff  
 gravitational compaction structures, A(128):146ff, 272  
 gravity-induced structures, B:1175-1176  
 Japan Sea, A(128):13-15; B:1183-1187  
 laminations, A(127):95-96, 347-349  
 and lithostratigraphy, B:1177  
 in sandstone, B:100, 104, 1233

in sandstone and siltstone, A(127):324, 345-346  
 in sediment gravity flow deposits, A(127):268 in siltstone, B:1233  
 Site 798, A(128):141ff  
 Site 799, A(128):265-272; B:48  
 soft-sediment deformation, A(127):261-264, A(128):141ff, 241, 266-269  
 thin bedding, A(127):96  
 in tuff, A(128):81  
 veins, A(128):143ff; B:1176-1181  
 sedimentation, rhythmic. *See* dark/light cycles; lithofacies, cyclic  
 sedimentation rate. *See also* sediment accumulation rate  
 age control points, B:351  
 in Brunhes Chron, B:961  
 at Brunhes/Matuyama boundary, B:961  
 and carbonate compensation depth, B:1228  
 and consolidation, B:1277  
 datum levels used, A(127):111, 209, 281, 365 and dolomite, A(128):148  
 error sources, A(128):171  
 and heat flow, B:1302  
 heat flow estimates from opal-A/opal-CT BSR and, B:1147-1148  
 Japan Sea, A(127):22, 24, A(128):31  
 and Jaramillo Subchron, B:959-960  
 and logging resolution, B:396  
 and magnetic intensity, B:966  
 and magnetite degradation, B:976  
 during Olduvai Subchron, B:975  
 and opal-A to opal-CT transition, B:1306  
 and orbital periodicity, B:397-398  
 and organic carbon, B:628  
 and organic matter preservation, B:1265  
 and phosphate formation, B:67  
 Pliocene/Quaternary boundary, A(127):251  
 and productivity, B:1228  
 related to upwelling, A(127):204  
 Site 794, A(127):96, 105-107, 112, A(128):31, 69; B:287, 351, 653, 975, 982, 1043-1044, 1224, 1227  
 Site 795, A(127):174, 203-204, 210; B:288, 351, 653, 975, 982, 1226-1227  
 Site 796, A(127):250-251, 277-278, 281-282; B:288, 653, 1226-1227  
 Site 797, A(127):324, 351, 357, 359-362, 366; B:287, 297, 351, 594, 975, 982, 1043-1044, 1226-1227  
 Site 798, A(128):31, 124-125, 171ff, 176-177; B:407, 443, 446, 548, 563-564, 653, 959, 961, 974-975, 982, 1226-1227  
 Site 799, A(128):31, 244, 259-260, 315-316, 324-325; B:653, 959, 961, 974-975, 982, 1227  
 Sites 794 and 795 compared, A(127):204  
 Sites 798 and 799 correlated, B:961, 965 and sulfate, A(127):362; B:1262  
 and turbidites, B:959  
 vs. barium, B:661  
 vs. depth of phosphate maximum, B:64-66  
 vs. sulfate, B:661  
 Yamato Basin S, A(127):356  
 sediments, biosiliceous, Site 798, B:439-455  
 sediments, bioturbated, in cyclic lithofacies, A(128):124  
 sediments, laminated. *See also* laminations  
 BSEM images, B:553-557  
 in cyclic lithofacies, A(128):124  
 lamination thickness, B:548, 550  
 and low oxygen environment, B:189  
 paleoceanographic significance, B:547-557  
 Site 794, A(127):147  
 Site 799, B:44  
 visual microanalysis methods, B:57-62  
 sediments, organic-rich, formation, A(127):212, 365  
 sediments, subducted, and contamination of mantle source, B:805-817  
 seismic experiment. *See* downhole seismic experiment  
 seismic expression  
 of biogenic opal, B:1140-1141, 1143  
 of debris flows, B:1140-1141, 1143  
 of dolomite, B:1141, 1143  
 of opal-A to opal-CT transition, B:1141, 1143  
 seismic reflection profile  
 data collection and processing, A(127):61-63  
 Japan Sea, A(127):14, A(128):13  
 Kita-Yamato Trough, B:1137-1139  
 reflector origin, B:1140-1142  
 Site 794, A(127):68, 84, 86-87, 153, A(128):83; B:111, 1079  
 Site 795, A(127):181, 183-184, 241-243  
 Site 796, A(127):258, 260-261, 311-312  
 Site 797, A(127):337-338, 340-341, 409  
 Site 798, A(128):135, 218-223; B:395, 410  
 Site 799, A(128):23, 254-255, 260, 375-379; B:1137-1139  
 synthetic seismogram compared, B:1142  
 Yamato Basin, B:1149  
 seismic reflection survey, pre-cruise surveys, A(127):142, 233, 307, 399, A(128):192, 343-344  
 seismic refraction profiles, Yamato Basin N, A(128):112  
 seismic stratigraphy  
 ash layers correlated, A(128):122, 153-155, 344-351, 379  
 of basement, B:779  
 of interbedded sediments/basalts, A(127):408  
 Japan Sea, A(127):28-29, A(128):34  
 Kita-Yamato Trough, A(128):245; B:1135-1143  
 lithostratigraphy correlated, A(127):143, 155, 174, 234-238, 243, 251, 312-313, 325, 403-408, 410, A(128):125, 193-194, 224, 351-353, 379; B:1140-1143  
 logs correlated, A(128):224, 351-353, 379  
 Okushiri Ridge, A(127):256-258, 307  
 physical properties correlated, A(128):224, 379; B:1143  
 seismic intervals, Site 799, B:1136  
 Site 794, A(127):73, 142-143  
 Site 795, A(127):174, 233-238  
 Site 796, A(127):251, 307-314  
 Site 797, A(127):325, 399-410  
 Site 798, A(128):34, 125, 192-194  
 Site 799, A(128):34, 343-353; B:1135-1143  
 Yamato Basin, A(127):83-84, 143  
 seismology. *See* downhole seismic experiment  
 seismometer. *See* downhole seismic experiment  
 SFL. *See* resistivity, logging  
 shards, glass  
 altered, A(127):95  
 in ash layers, B:791  
 in blue tuff, B:117, 119, 130  
 replaced by zeolites, B:139  
 Site 795, A(127):190  
 shards, vitric, in ash-flow tuff, B:793  
 shear joints, Site 794, A(128):92  
 shear strength  
 and consolidation characteristics, B:1124-1125

shear strength (cont.)

## SUBJECT INDEX

overburden pressure ratio, B:1125–1126  
 Site 798, A(128):182, 207, 211  
 Site 799, A(128):327, 353, 360  
 shear zone, Site 799, A(128):270  
 sheet flows, of doleritic basalt, A(128):86  
 Shikibawa Formation, and planktonic foraminifers zonations, B:459  
 Shikoku Basin, rare earth elements, B:719  
 Shikoku–Parce Vela Basin, formation, A(127):10  
 Shiribeshi Basin, A(127):268  
 Shiribeshi Trough  
   bathymetry, A(127):251  
   free-air gravity anomalies, A(127):253  
   sedimentation, A(127):256–257  
   seismic stratigraphy, A(127):307  
 siderite  
   and Mossbauer parameters, B:741  
   shallowest occurrence, A(128):273  
   Site 799, A(128):260, 272, 275–276, 289; B:611–612  
 sigmas  
   Site 795, B:543  
 Sikhote-Alin, basement, A(127):177–178  
 silica  
   in ash layers, B:1380–1385  
   basement, A(128):98  
   and biogenic silica diagenesis, B:1263  
   in dark/light cycles, B:569  
   diagenesis, A(127):190  
   and diatom dissolution, A(127):362–363  
   dissolution, B:350  
   formation, B:52–54  
   in interstitial water, B:1266  
   Japan Sea, B:1235–1237  
   precipitation, A(127):174  
   and rare earth elements, B:682  
   removal rates in Japan Sea, B:447  
   and silicification front, A(127):205  
   Site 794, A(127):108–109; B:682  
   Site 795, A(127):205; B:683  
   Site 796, A(127):280  
   Site 797, A(127):362–363, 369; B:688  
   Site 798, A(128):174–175, 184; B:722  
   Site 799, A(128):318, 320–321, 332; B:611, 722  
   sources, B:148  
   transitions, B:53  
   vs. lithium, A(128):185  
   XRD analyses, A(128):292  
   XRF analyses, A(128):281; B:1025–1029, 1031–1035  
 silica, amorphous. *See* opal-A  
 silica, biogenic. *See also* opal, biogenic; opal-A; opal flux  
   abundance, A(128):150, 259  
   accumulation rate, A(128):122, 158  
   cycles, B:439  
   in dark/light cycles, B:584  
   diagenesis, B:1263, 1267  
   dissolution, A(128):320–321  
   fragmentation degree, B:549  
   glacial/interglacial variation, B:591  
   Japan Sea, B:1239–1241  
   and productivity, B:694  
   and rare earth elements, B:692  
   XRF analyses, A(128):151  
   XRF vs. smear slide analyses, A(128):281  
 silica, diagenetic  
   isotopic studies, B:49–56  
   Japan Sea, B:1239–1241  
   Site 799, B:36, 46  
   XRF vs. smear slide analyses, A(128):281  
 silica, dissolved, in Japan Sea, B:350–351

silica, excess, B:1240–1244, 1253  
 silica, logging  
   and diatom vs. clay content, B:1417  
   Site 794, B:1416, 1420–1421  
   Site 796, B:1416, 1423  
   Site 797, B:1417, 1426–1427  
   Site 798, B:1401–1402  
   Site 799, A(128):364; B:1404, 1408–1409  
 silica/aluminum ratio (Si/Al)  
   Site 795, B:683  
   Site 797, B:688  
 silica diagenesis. *See also* opal-A to opal-CT transition; opal-CT to quartz transition  
   and Ce anomaly, B:682  
   effect of temperature, B:3  
   effects on interstitial water chemistry, B:646  
   and Eu anomaly, B:682  
   and inorganic geochemical data, A(127):23  
   and interstitial water oxygen isotopes, B:85, 701  
 Japan Sea, B:1263, 1267  
   material balance calculation for, B:701  
   and physical properties, A(128):280; B:3–31, 1281–1282  
   and silica content, B:682, 688  
   Site 795, A(127):190  
   Site 797, A(127):347  
   Site 798, A(128):149–150  
   Site 799, A(128):278–281  
   and thermal history, Japan Sea, B:1305–1307  
 silica phase change. *See* opal-A to opal-CT transition; opal-CT to quartz transition  
 silicate, alteration reactions, A(127):110–111  
 siliceous claystone. *See* claystone, siliceous  
 siliceous veins. *See* veins, siliceous  
 silicification front, and silica content, A(127):205  
 silicoflagellate, B:1220  
   Site 798, A(128):162; B:241–244  
   Site 799, A(128):305; B:241, 246–247  
   zonal data, B:1225–1226  
   zonation, B:237, 239–241  
 silicon, at opal-CT to quartz transition A(127):307  
 silicon oxide. *See* silica  
 sill, Japan Sea, B:299–300, 365  
 sills  
   basaltic, B:819  
   bathymetric, A(128):20, 35  
   and crustal heterogeneity, B:1113–1114  
   crystallization history, B:864–865  
   dolerite, A(128):86  
   emplacement, B:894, 927, 938  
   intrusion, A(128):95; B:829–830  
   demagnetization, B:938  
   Site 794, B:837–838, 849, 906, 927, 1339  
   Site 797, B:837–838, 850, 907–908  
   suites, B:853–855  
 silt  
   Site 796, A(127):261–264  
   Site 799, A(128):256, 264; B:43  
 siltstone  
   and revised lithostratigraphy, B:1233  
   sedimentary structures, B: 1233  
 siltstone, carbonaceous, Site 797, A(127):345–346  
 siltstone, diatom clayey, Site 795, A(127):187  
 siltstone, siliceous, Site 799, B:39–40  
 siltstone, tuffaceous clayey, A(128):68, 89–90  
 silty clay. *See* clay, silty  
 silver  
   Site 798, B:1368–1369  
   in sulfide deposits, A(128):21  
 Site 79, physical properties, B:1010–1015  
 Site 116, interstitial water boron content, B:638  
 Site 125, trace metals, B:1361

Site 126, trace metals, B:1361  
 Site 127, trace metals, B:1361  
 Site 128, trace metals, B:1361  
 Site 130, trace metals, B:1361  
 Site 141, paleoenvironment, B:338  
 Site 148, siderite occurrences, B:612  
 Site 173  
   foraminifers, A(128):165  
   paleoclimate, B:459  
   silicoflagellate, B:237  
 Site 290, basement isotope geochemistry, B:805  
 Site 294, rare earth elements, B:719  
 Site 296, rare earth elements, B:719  
 Site 299, A(127):337  
   bathymetry, A(127):6, 78, A(128):72  
   consolidation tests, B:1123  
   oceanic crust, A(127):73  
   planktonic foraminifers, B:459  
   radiolarians, A(128):162, B:225  
   sediment thickness, A(127):84  
   seismic stratigraphy, A(127):179  
   Site 797 sandstone compared, B:111  
   turbidite sands, B:99  
 Site 300  
   bathymetry, A(127):6, A(128):72  
   nanofossils, B:168  
   sand recovered, A(128):14  
 Site 301  
   bathymetry, A(127):6, A(128):72  
   drilling history, A(127):180  
   lithostratigraphy, A(128):13, 249; B:1141  
   nanofossils, B:168, 174  
   sand recovered, A(128):14  
   sediment thickness, A(127):84  
   seismic stratigraphy, A(127):179  
   Site 799 sandstone compared, B:111  
   turbidite sands, B:99  
 Site 302  
   bathymetry, A(127):6, 78, A(128):72  
   consolidation tests, B:1123  
   diatom clay, A(127):84  
   faulting age, A(127):334  
   nanofossils, B:168, 174  
   normal faulting, A(127):83  
   radiolarians, A(128):303, B:226  
   sediment thickness, A(127):84  
   tectonic history, A(128):130  
 Site 303, mantle source contamination, B:815  
 Site 304, mantle source contamination, B:815  
 Site 366, paleoenvironment, B:338  
 Site 395, temperature change during downhole seismic experiment, B:1160  
 Site 397, paleoenvironment, B:338  
 Site 417, alteration and rare earth elements correlated, B:911  
 Site 433, diatom abundance, B:316  
 Site 436  
   ash layers, B:1319–1320  
   basement age, B:749  
   helium isotope ratios, B:749  
   rare earth elements, B:719  
   tectonic setting, B:1312  
 Site 438  
   ash layers, B:1319–1320  
   basement age, B:749  
   diatom abundance, B:311  
   forearc subsidence, B:1213  
   helium isotope ratios, B:749  
   paleobathymetric history, B:1214  
   silicoflagellates, B:237  
   subsidence of Ishikari-Hidaka Basin, B:1199  
   tectonic setting, B:1312  
 Site 439

ash layers, B:1319–1320  
 forearc subsidence, B:1213  
 paleobathymetric history, B:1214  
 subsidence of Ishikari-Hidaka Basin, B:1199  
 tectonic setting, B:1312  
 and volcanic front position, B:1324

Site 440  
 ash layers, B:1319–1320  
 tectonic setting, B:1312

Site 504  
 formation factor and cation exchange capacity correlated, B:1282  
 isotopic analyses, B:888  
 oxygen isotopes, B:699  
 stress field, B:1047

Site 533, nannofossils, B:173

Site 544, paleoenvironment, B:338

Site 578, diatoms, B:350

Site 579  
 biosiliceous productivity, B:311–313  
 diatoms, B:350

Site 580  
 biosiliceous productivity, B:311–313  
 diatoms, B:350  
 radiolarian faunal change, B:447

Site 581  
 diatoms, B:311, 316  
 temperature change during downhole seismic experiment, B:1160

Site 582  
 interstitial water oxygen isotopes, B:698  
 veins observed, B:1180

Site 583  
 basement age, B:749  
 helium isotope ratios, B:748–749  
 interstitial water oxygen isotopes, B:698

Site 584  
 ash layers, B:1319–1320  
 basement age, B:749  
 helium isotope ratios, B:749  
 interstitial water oxygen isotopes, B:698  
 tectonic setting, B:1312

Site 588, paleoenvironment, B:338

Site 590, paleoenvironment, B:338

Site 591, paleoenvironment, B:338

Site 607, oxygen isotopes, B:403

Site 672, veins, B:1180

Site 684, helium isotope ratios, B:748

Site 747, ash layer logging, A(128):339

Site 752, ash layer logging, A(128):339

Site 792, Brunhes/Matuyama boundary, A(127):199

Site 794, A(127):71–167, A(128):67–120  
 aluminum, B:654, 656  
 ash layer chemistry, B:1373–1393  
 augite-liquid relations, B:870–872  
 barium, B:653–657  
 barium as productivity indicator, B:653–656  
 basalt alteration, B:883–889  
 basalt melting experiment, B:891–898  
 basement, B:849–850, 885, 906, 1339  
 basement chemistry, B:779–789, 839, 892–894, 908–911, 916–918  
 basement isotopes, B:786–787, 807–814, 888, 901–902  
 basement magnetic properties, B:934  
 basement mineralogy, B:850–855  
 basement structures, B:1181  
 biostratigraphy, B:1221–1223  
 blue tuff, B:115–130  
 boron, B:638–641  
 crustal anisotropy, B:1107–1121  
 crustal structure, B:1075–1081

diatoms, B:344  
 downhole seismic experiment, B:1061–1073, 1075–1081, 1107–1121, 1157–1171  
 electrical resistivity experiment, B:1351–1359  
 FMS logs, B:1039–1046  
 foraminifers, B:190–193, 210–213  
 geochemical logs, B:1420–1421  
 geothermal analyses, B:1301–1307  
 gray value, B:585  
 hydrogen isotopes, B:641–642  
 inorganic geochemistry, B:637–646, 1261–1274  
 liquid lines of descent, B:870  
 lithostratigraphy, revised, B:1231  
 logs, B:1416, 1418–1419  
 magma suites, B:869  
 magmatic characterization, B:780–784  
 magnetostratigraphy, B:969–970  
 mantle source, B:807–815  
 Olduvai Subchron, B:218  
 operations, A(127):85–90, A(128):83–85  
 organic carbon, B:656  
 oxygen isotopes, B:641–643  
 parent magma, B:855–857, 894  
 physical properties, B:990–993, 1275–1296  
 radiometric age, B:780–784, 820, 824–826  
 sediment composition, B:1236–1237, 1242  
 sediment geochemistry, B:680, 682–684, 691–695, 1246–1249  
 sedimentation rate, B:1224–1225  
 spinel chemistry, B:840–843  
 stress measurements, B:1050–1059  
 strontium isotopes, B:643–646  
 subsidence history, B:1207–1211  
 sulfate, B:637–638, 656  
 sulfur isotopes, B:637–638  
 trace metals, B:1362–1363  
 Yamato Basin tectonics, B:787–788

Site 795, A(127):169–245  
 ash layer chemistry, B:1373–1393  
 augite-liquid relations, B:870–872  
 basement, B:906, 1338–1339  
 basement chemistry, B:913–914, 918–919  
 basement isotopes, B:807–814  
 basement magnetic properties, B:934, 937  
 basement mineralogy, B:850–855  
 basement petrography, B:849–850  
 biostratigraphy, B:1223  
 boron, B:638–641  
 diatoms, B:344–345  
 foraminifers, B:194–197  
 geothermal analyses, B:1301–1307  
 gray value, B:585  
 hydrogen isotopes, B:641–642  
 inorganic geochemistry, B:637–646, 1261–1274  
 lithostratigraphy, B:496–497, 1231  
 magnetostratigraphy, B:970–972  
 mantle source, B:807–815  
 operations, A(127):182–186  
 oxygen isotopes, B:641–643  
 oxygenation conditions, B:711–713  
 parent magma, B:855–857  
 physical properties, B:993–997, 1275–1296  
 physical properties and silica diagenesis, B:3–31  
 pyrite, B:711–713  
 radiometric age, B:820–821, 826–827  
 reworking, A(127):198  
 sediment composition, B:8, 1237–1240, 1243  
 sediment geochemistry, B:680–681, 683–685, 691–695, 706–717  
 sedimentation rate, B:1225  
 strontium, B:643–646  
 subsidence history, B:1207–1211  
 sulfate, B:637–638, 656  
 sulfur, B:637–638  
 zeolitization, B:138–139

Site 797, A(127):323–421  
 ash layer chemistry, B:1373–1393  
 augite-liquid relations, B:870–872  
 basalt melting experiment, B:861–868, 891–898  
 basement, B:906–907, 1339–1340  
 basement chemistry, B:839, 892–894, 911, 914–916, 919–921  
 basement isotopes, B:807–814, 901–902  
 basement magnetic properties, B:934, 937–939  
 basement mineralogy, B:850–855  
 basement petrography, B:850, 891–892  
 biostratigraphy, B:1223  
 boron, B:638–641  
 boron isotopes, B:638–641  
 clay mineral authigenesis, B:139  
 dark/light cycles, B:574, 577–601  
 diatoms, B:345, 348–349  
 feldspar albitization, B:136–137  
 FMS logs, B:1039–1046  
 foraminifers, B:202–207, 213–215  
 fractionation models, B:872–874  
 geochemical logs, B:1426–1427  
 geothermal analyses, B:1301–1307  
 gray value, B:585, 592  
 hydrogen isotopes, B:641–642  
 inorganic geochemistry, B:637–646, 1261–1274  
 liquid lines of descent, B:870  
 lithostratigraphy, B:4, 1231  
 logs, B:7, 1417, 1424–1425  
 magma suites, B:869  
 magnetic minerals, B:949–951  
 magnetostratigraphy, B:973  
 mantle source, B:807–815  
 opal-A to opal-CT transition, B:19  
 operations, A(127):337–340  
 organic carbon, B:586–587, 592

oxygen isotopes, B:641-643  
 parent magma, B:855-857, 894  
 physical properties, B:3-31, 999-1104, 1275-1296  
 radiometric age, B:820, 822-824  
 reworking, A(127):357  
 sandstone provenance, B:99-113  
 sediment composition, B:8-10, 582-584, 586-587, 592, 1240-1241, 1244  
 sediment geochemistry, B:681, 685, 688-695  
 sedimentation rate, B:1225  
 spinel chemistry, B:840-843  
 strontium isotopes, B:643-646  
 subsidence history, B:1207-1211  
 sulfate, B:637-638  
 sulfur, B:586-587, 592  
 sulfur isotopes, B:637-638  
 trace metals, B:1366  
 zeolitization, B:138-139  
 Site 798, A(128):121-236  
 accumulation rates, B:411-419  
 aluminum, B:654, 656  
 ash layers, B:699-701, 791-793, 1373-1393  
 bacteria, B:755-776, 755-769  
 barium, B:654, 656-659  
 basalt alteration, B:699-701  
 biostratigraphy, B:1223-1224  
 C/N ratio, B:435-437  
 carbonate content, B:435-437  
 carbonate diagenesis, B:701-702  
 chronostratigraphy, B:317-318  
 clay mineral composition, B:416, consolidation, B:1124-1127  
 dark/light cycles, B:430-432, 446-447, 559-576  
 feldspar, B:411-419  
 foraminifers, B:460-463  
 geochemical logs, B:1021-1035, 1398, 1400-1402  
 geologic setting, B:425, 440, 560  
 geothermal analyses, B:1301-1307  
 glauconite, B:63-74  
 inorganic geochemistry, B:607-608, 697-703, 1261-1274  
 interstitial gas helium isotopes, B:748-749  
 laminated biosiliceous sediments, B:547-557  
 lithostratigraphy, B:697  
 logs, B:398-400  
 magnetic properties, B:959-967, 975-978  
 magnetostratigraphy, B:973-974  
 methane origin, B:749-750  
 microbiology, B:761-776  
 Mossbauer characterization of sediments, B:739-746  
 opal content, B:443-446, 449-455  
 operations, A(128): 133-136  
 organic carbon, B:425, 435-437, 656  
 organic geochemistry, B:667-675  
 organic matter, B:425, 667-675  
 oxygen isotopes, B:442-443, 698-699  
 permeability, B:1127-1129, 1131-1132  
 phosphate, B:63-74  
 physical properties, B:1006-1010, 1275-1296  
 quartz, B:411-419  
 sediment composition, B:398-400, 411  
 sediment geochemistry, B:610-612, 615-617, 720-737, 1022-1027, 1402  
 sedimentary structures, B:1175-1183  
 sedimentation, B:419-421, 1225  
 seismic stratigraphy, B:1135-1143  
 silica diagenesis, B:701  
 strip sample trace elements, B:1367-1372  
 subsidence history, B:1207-1211

sulfate, B:656  
 surface-water temperature fluctuations, B:463-465  
 synthetic seismogram, B:1142  
 tephrochronology, B:793-797, 800  
 Site 799, A(128):237-402  
 aluminum, B:654, 656  
 ash-flow tuff, B:793  
 ash layers, B:699-701, 791-793, 1373-1393  
 barium, B:654, 654-657  
 basalt alteration, B:699-701  
 biostratigraphy, B:34-35, 1224  
 carbonate diagenesis, B:701-702  
 clay mineral authigenesis, B:139-141  
 consolidation, B:1124-1127  
 dark/light cycles, B:571-574  
 diagenetic carbonates, B:75-98  
 diagenetic mineralogy, B:35-38  
 feldspar albitization, B:137  
 geochemical logs, B:1399, 1403-1408  
 geologic setting, B:425, 560  
 geothermal analyses, B:1301-1307  
 glauconite, B:63-74  
 inorganic geochemistry, B:608-610, 697-703, 1261-1274  
 interstitial gas helium isotopes, B:748-749  
 lithostratigraphy, B:33-34, 76-77, 697  
 magnetic intensity variations, B:959-967  
 magnetic properties, B:975-981  
 magnetostratigraphy, B:974-975  
 methane, B:749-750  
 Mossbauer characterization of sediments, B:739-746  
 operations, A(128):253-255  
 organic carbon, B:425, 427, 623, 626, 656  
 organic geochemistry, B:623-633, 667-675  
 organic matter, B:425, 667-675  
 oxygen isotopes, B:698-699  
 permeability, B:1127-1129, 1131-1132  
 phosphate, B:63-74  
 physical properties, B:1010-1015, 1275-1296  
 pyrolysis-gas chromatography, B:627-628, 632-633  
 Rock-Eval pyrolysis, B:624-627  
 sandstone provenance, B:99-113  
 sediment geochemistry, B:610-612, 618-620, 720-737, 1402  
 sediment structures, B:38-39, 1175-1183  
 sedimentation rate, B:1225-1227  
 silica diagenesis, B:701  
 subsidence history, B:1207-1211  
 sulfate, B:656  
 sulfur, B:623-624  
 tephrochronology, B:793-799  
 thermal conductivity, B:1017-1019  
 slickensides, in tuffaceous clayey siltstone, A(128):90  
 slump folds  
 Site 798, A(128):142  
 Site 799, A(128):261-263, 266-267  
 slumps  
 of ash layers, A(128):153-154, 285  
 and compositional banding, B:1176  
 and consolidation, B:1125  
 extent and magnitude, A(127):268  
 and faults, B:1176  
 Japan Sea, A(128):14  
 on Okushiri Ridge, A(127):268  
 and opal-A to opal-CT transition, A(127):268  
 Site 798, B:1175  
 Site 799, A(128):259, 355; B:1175-1176, 1189  
 smear slide analyses. *See* sediment composition  
 smectite

in blue tuff, B:119  
 in dark/light cycles, B:584-588  
 diagenetic origin, B:1238  
 glacial/interglacial variation, B:588  
 from Japanese Islands, B:595  
 in Japan Sea sediments, B:1235-1244  
 as sandstone cement, B:135  
 Site 797, B:592  
 Site 798, B:411, 416, 418  
 and volcanic glass alteration, B:1375  
 vs. illite, B:1245  
 vs. magnesium, B:1250  
 smectite/illite ratio, Site 798, B:411, 416  
 sodic feldspar. *See* feldspar, sodic sodium  
 in ash layers, B:1384  
 in augite, B:851-853, 858  
 and basement alteration, B:1266  
 and feldspar albitization, B:137  
 and geochemical logging, B:1395, 1415  
 Japan Sea, B:1237  
 and parent magma, B:856  
 Site 794, A(127):107  
 Site 795, A(127):205  
 Site 796, A(127):278-279  
 Site 797, A(127):364, 374  
 Site 798, B:1370-1371  
 Site 799, B:723  
 sources, B:137  
 vs. potassium, B:732, 858  
 sodium/aluminum ratio  
 Site 794, B:1249  
 Site 795, B:683  
 Site 798, B:682  
 sodium/chlorine ratio, in interstitial water, B:1274  
 sodium oxide. *See* sodium  
 sodium/potassium ratio, Site 799, B:723  
 soft-sediment deformation. *See* deformation, soft-sediment  
 soil horizons, formed during interglacials, B:403  
 sonic core monitor (SCM), Site 799, A(128):253, 255  
 sonic velocity. *See* velocity, logging  
 Soya Strait, B:423  
 depth, B:342, 440  
 Japan Sea and, B:70, 362-363, 365  
 sill depth, B:457  
 sphenoliths  
 Site 799, A(128):307, 309  
 sphalerites  
 Site 795, B:543  
 spherically focused log (SFL). *See* resistivity, logging  
 spinel. *See also* spinel, chromium  
 chemistry, B:840-842  
 crystal chemical model, B:840-841  
 magmatic magnesium number inferred from composition, B:840-842  
 magmatic oxygen fugacity from spinel analyses, B:842-844  
 and mineral-liquid disequilibrium, B:844  
 photomicrographs, B:847  
 relict magmatic phase, B:838  
 spinels in other igneous rocks compared, B:840  
 zoned, B:840, 844  
 spinel, chromium, B:853  
 magnesium number vs. Cr (Cr + Al), B:843  
 magnesium number vs. Fe<sup>3+</sup>(Cr + Al + Fe<sup>3+</sup>), B:843  
 as petrogenetic indicator, B:837-847  
 spirasters  
 Site 795, B:543

sponge spicules. *See also* microfossils, siliceous abundance, B:344–349  
 Leg 127, B:292  
 Site 794, B:344  
 Site 795, B:304, 345, 541–543  
 Site 797, A(127):343; B:303, 345  
 Site 799, A(128):260  
 sponges, Site 799, B:35, 45  
 spongodiscids  
 Site 795, B:542  
 spores  
 BSEM images, B:554–556  
 formation and recycling of, B:550–551  
 resting-spore laminae, B:549–551  
 sporinite, Sites 798 and 799, B:670  
 squalene, Site 799, B:627, 632  
 stable isotopes. *See* isotopes, carbon; isotopes, oxygen  
 steranes, in Kita-Yamato Trough, B:669  
 stilbite, and ash alteration, A(128):288  
 strain difference functions, Site 794, B:1054, 1056–1057  
 stratal disruption, Site 798, A(128):142  
 stress field, B:1323–1324  
 azimuthal distribution of, B:1058  
 crack behavior, B:1121  
 crustal anisotropy, B:1114–1115, 1343  
 deformation rate analysis (DRA), B1048–1049  
 determination methods, B:1047  
 from earthquake analysis, B:1054  
 and faults and fractures, B:1181–1183  
 horizontal stress, B:1051–1052  
 and Japan Arc rotation, B:1054  
 Japan Sea, A(127):17, B:1185–1187  
 Japan Sea E, B:1059  
 Kita-Yamato Trough, A(128):293–294  
 and magma ascent, B:1319  
 and Oga Peninsula tectonic history, B:1204–1205  
 porosity effect on horizontal stress, B:1053  
 Site 794, A(127):84; B:1047–1059, 1120  
 Site 795, A(127):180–181  
 Site 796, A(127):258  
 Site 797, A(127):337  
 Site 799, A(128):245  
 strain difference functions, B:1054, 1056–1057  
 stress measurement procedure, B:1049–1050  
 stress-strain curves, B:1051  
 vertical stress, B:1050–1051  
 and volcanic activity, B:796, 1319, 1321, 1325  
 Yamato Rise, A(128):245  
 Yamato Trough, A(128):340  
 strongyles  
 Site 795, B:542  
 strontium  
 and ash layer alteration, A(127):280  
 ash layer/basement alteration and, B:643  
 basement, B:807  
 in interstitial water, B:607, 643–646, 1263  
 isotopic ratios vs.  $\text{Sr}$ , B:646  
 Site 794, A(127):110; B:1363  
 Site 795, A(127):205–207; B:1365  
 Site 796, A(127):280  
 Site 797, A(127):363–364, 370; B:1366  
 Site 798, A(128):173–174, 182  
 Site 799, A(128):318, 329; B:611  
 vs. neodymium, B:902  
 strontium isotopes. *See* isotopes, strontium  
 structures. *See* sedimentary structures  
 styles  
 Site 795, B:542

subduction zones, in Pacific Ocean W, B:1333–1334  
 submarine fans. *See also* Toyama Deep Sea Fan progradation of, A(127):16  
 submarine volcanism  
 Japan Sea, A(127):95  
 Site 795, A(127):219–220  
 subsidence  
 fault-controlled, B:1344  
 in forearcs, B:1213  
 geohistory and backstripping analysis methods, B:1199–1201  
 Japan Sea, B:1209–1214, 1343–1344  
 Oga Peninsula, B:1202–1205  
 onshore sections and offshore wells analyzed, B:1201–1202  
 paleobathymetric analysis methods, B:1200–1201  
 Pohang Basin, B:1205–1206  
 rates, B:1211–1212  
 Sakhalin Island, B:1206–1207  
 and sediment loading effects, B:1211  
 subsidence, B:1200, 1212–1213  
 thermal, B:1344  
 Tsushima Basin, B:1207  
 Sugata Formation, A(128):129  
 sulfate  
 and barite front development, B:637  
 and clathrate formation, A(127):281  
 in interstitial water, B:637–639, 657+59, 1264  
 and methane abundance, A(127):368  
 and organic carbon decomposition, B:1262  
 and organic matter degradation, A(127):362  
 in sandstone, B:143–144  
 and sediment accumulation rate, B:637, 646  
 and sedimentation rate, A(127):362; B:1262  
 Site 794, A(127):108; B:637–638  
 Site 795, A(127):204; B:637–638, 640  
 Site 796, A(127):279; B:637–638  
 Site 797, A(127):325, 362, 368; B:637–638  
 Site 798, A(128):172–173, 180; B:770, 775–776, 1263  
 Site 799, A(128):317, 328; B:610  
 and source of methane, A(127):288  
 and sulfate-reducing bacteria, B:770  
 vs. alkalinity, A(128):183  
 vs. sedimentation rate, B:661  
 sulfate reduction. *See also* sulfate  
 and bacteria, B:767, 771  
 and barium dissolution, B:656, 661  
 and carbon isotopes, B:88  
 and high alkalinity values, A(127):279  
 and interstitial water analyses, B:607  
 and organic carbon decomposition, B:1262–1263  
 and organic matter, A(127):207, B:637  
 potential rates, B:771  
 and sedimentation rate, A(127):207  
 Site 794, A(127):108, 111, 119  
 Site 795, A(127):174, 204  
 Site 796, A(127):251, 267  
 Site 797, A(127):362  
 Site 798, B:770, 772–773, 775–776  
 Site 799, A(128):318  
 and sulfate-reducing bacteria, B:770  
 sulfide, acid-volatile (AVS), Site 798, B:771, 775  
 sulfide deposits  
 in failed rifts, A(128):251–252  
 in Japan Sea sediments, B:1241–1249  
 Kuroko, A(128):24, 239, 251  
 occurrence of, A(128):21–22  
 sulfur

in dark/light cycles, B:579, 586–588  
 as frambooidal pyrite, B:706–709  
 glacial/interglacial variations, B:589–590  
 and organic carbon, B:1241  
 Site 794, A(127):113; B:1246–1247, 1255, 1363  
 Site 795, B:706–710, 714, 209–213  
 Site 796, A(127):283–285  
 Site 797, A(127):365–367; B:592  
 Site 798, A(128):176–177, 189–192  
 Site 799, A(128):323–324, 334–338, 342; B:623–624, 626–627  
 vs. carbon, organic, A(128):343–344  
 vs. organic carbon, B:594, 628, 712–713, 1254  
 vs. pyrite, B:1254  
 vs. sulfur isotopes, B:715  
 sulfur, logging, Site 798, B:1024  
 sulfur/iron ratio  
 and pyrite formation, B:1249  
 Site 794, B:1255  
 sulfur isotopes. *See* isotopes, sulfur  
 Sumisu Rift, Pb-Pb plots of arc volcanic rocks, B:808  
 Sunda Arc, sediment contamination of mantle source, B:805  
 surface-water temperature. *See* temperature, surface-water  
 susceptibility. *See* magnetic properties  
 Sutu Canyon, bathymetry, A(127):251  
 synthetic seismogram  
 and opal-A to opal-CT transition, B:1146–1148  
 seismic field record compared, B:1142  
 Site 794, A(127):143, 154  
 Site 798, A(128):184, 214  
 Site 799, B:1138–1140, 1142  
 Takuyo Bank  
 basement, A(128):248–249  
 bathymetry, A(128):245–246  
 crustal structure, A(128):249  
 tantalum  
 Site 798, B:1370–1371  
 Site 799, B:724  
 Tartar Strait  
 depth, B:342  
 Japan Sea and, B:69–70, 365  
 sea ice, B:167  
 sill depth, B:457  
 tau- $p$  mapping, and one-dimensional velocity structure, B:1078–1080  
 Taxodiaceae  
 Site 795, B:490  
 tectonic provenance. *See* provenance, tectonic  
 tectonic reconstructions. *See* plate reconstructions  
 teleseismic events, and downhole seismic experiment, B:1162, 1167–1168  
 temperature  
 of alteration, B:839  
 and clathrate stability, A(127):281  
 contaminated measurements, A(127):136  
 of diagenetic reactions, B:148–149  
 and electrical resistivity, A(128):111  
 expected decay curve, A(127):298–299  
 and gas hydrate stability, A(127):288, 299–300  
 increase with depth, A(127):135–136  
 Japan Sea, A(127):26–27, A(128):20  
 and lower crust structure, B:1343  
 of opal-A to opal-CT transition, A(128):150, 280; B:14, 1145–1147, 1263, 1305–1306  
 of opal-CT formation, B:49  
 of opal-CT to quartz transition, A(128):280; B:17, 1263

temperature (cont.)

and opal dissolution, B:310  
 in permeable sand, A(127):299  
 Site 794, A(127):73, 140-142  
 Site 795, A(127):174, 230-232  
 Site 796, A(127):251, 296-301  
 Site 797, A(127):390ff, 396-397  
 Site 798, A(128):183-184, 213-214  
 Site 799, A(128):338-339, 365  
 Sites 794-797, B:51, 56  
 and strontium isotopes, B:644  
 typical temperature record, WSTP tool, B:1299  
 vs. opal-A/opal-CT BSR 2-way traveltimes, B:1152  
 vs. thermal resistance, A(127):142, 235, 300, 397; B:1302  
 vs. time, A(127):235  
 WSTP measurement reliability, B:1308-1309

temperature, isotopic  
 opal-CT, B:55-56  
 quartz, B:55-56

temperature, logging  
 Site 798, A(128):125, 183-184, 213  
 Site 799, A(128):338-339

temperature, silica transformation, and oxygen isotopes, B:49

temperature, surface-water  
 calcareous nannofossils and, B:180  
 fluctuations, B:463-465  
 and foraminifers cooling ratio, B:193  
 Japan Basin, B:209  
 Japan Sea, B:201, 337  
 orbital cyclicity of, B:467-468  
 oxygen isotope record correlated, B:464-465, 468  
 and planktonic foraminifers, B:187  
 Site 794, A(127):102-103  
 Site 795, A(127):199  
 Site 796, A(127):273  
 Site 797, A(127):357  
 Site 798, B:457-470  
 Site 799, A(128):310, 312

temperature gradient. *See* thermal gradient

tephra markers  
 Japan Sea, A(128):157, 291-292; B:796

tephrochronology  
 Site 798, A(128):153-157, 196-198; B:793-796  
 Site 799, A(128):284-291; B:793-796, 798-799

terbium, Site 798, B:1370-1371

terrigenous accumulation rate. *See* terrigenous flux

terrigenous component. *See also* sand  
 core analyses, B:404  
 logging estimates, B:402, 404  
 sediment gravity flow deposits, A(127):267  
 Site 796, A(127):250  
 Site 798, B:401  
 vs. quartz abundance, B:403

terrigenous flux  
 glacial/interglacial variations, B:446  
 to Japan Sea, B:409  
 in logs, B:403  
 Site 798, B:419, 446

terrigenous sediment  
 and chromium, B:723  
 influx of, A(127):16  
 and major and minor element chemistry, B:722-723  
 and rare earth elements, B:729, 732  
 X-ray diffraction analyses, B:400-401

tetraxons  
 Site 795, B:543

Th/Sc ratio. *See* thorium/scandium ratio  
 thanatocoenosis, and diatom dissolution, B:309-316  
 thermal conductivity  
 above opal-A to opal-CT boundary, B:1147  
 in basement, B:1281  
 correction factors applied, B:1018  
 data quality, A(127):127  
 hydrostatic pressure and, B:1017-1019  
 Japan Sea, B:1280-1281, 1287, 1298, 1301-1303  
 measuring system schematic, B:1018  
 porosity correlated, B:987  
 in sediments, B:1281  
 Site 794, A(127):127ff, A(128):102, 106; B:1287  
 Site 795, A(127):222ff; B:1287  
 Site 796, A(127):290ff; B:1287  
 Site 797, A(127):383ff, 397; B:1287, 1300  
 Site 798, A(128):181-182, 206-207, 210; B:1287  
 Site 799, A(128):326-327, 338, 352-353, 360, 365; B:1287  
 typical temperature vs. log time plot, B:1018  
 vs. porosity, A(127):135, 232, 395; B:25, 27, 989, 1287

thermal demagnetization. *See* magnetic properties

thermal gradient  
 isotopic, B:56  
 Japan Sea, A(127):26-27; B:1301-1303, 1375  
 and opal-A to opal-CT transition, A(127):268  
 sediment, B:56  
 and silica formation, B:52-54  
 Site 796, A(127):247, 251, 297, 299  
 Site 797, A(127):390  
 Site 798, A(128):125, 183-184, 213-214  
 Site 799, A(128):338-339, 365  
 Sites 794-797, B:51  
 and thickness of opal-CT zone, B:49

thermal history, Japan Sea, B:49-56

thermal maturity  
 and *n*-alkanes, A(128):324  
 of organic matter, B:671-672  
 Site 798, B:669  
 Site 799, B:624, 669-670

thermal resistance  
 Site 796, A(127):299  
 Site 797, A(127):390  
 Site 799, A(128):338  
 vs. temperature, A(127):235, 300, 397; B:1302

thermocline, in Japan Sea N, B:167

thermogenic gas. *See* gas, thermogenic

thermohaline circulation, Japan Sea, B:528-529

tholeiite  
 in basalt, B:870, 875  
 chemistry, B:839  
 rare earth elements, B:922, 925  
 Site 794, A(128):33, 68-69, 96-99, 117; B:779-780  
 Site 795, B:919  
 Site 797, B:920  
 spidergrams, B:923, 926

thorium  
 in ash layers, B:1386  
 in basement rocks, B:807  
 Site 795, B:707  
 Site 798, B:1370-1371  
 Site 799, B:724

thorium, logging  
 Site 794, A(128):103, 108; B:1418-1419  
 Site 796, B:1422  
 Site 797, B:1424-1425  
 Site 798, B:1400

Site 799, A(128):366-367; B:1403, 1406-1407  
 thorium/scandium ratio  
 and volcanic activity, B:723, 732  
 Site 798, B:723, 733  
 Site 799, B:723, 734  
 thorium/uranium ratio, logging  
 in ash layers, A(128):339  
 Site 799, A(128):366

thrust faults  
 fluid flow along, A(127):247, 295, 301  
 frictional heating along, A(127):247, 301  
 Japan Basin E, A(127):255  
 Japan Sea, A(127):81-82  
 Japan Sea E, A(127):9  
 Yamato Basin N, A(128):75-76

Tibetan Plateau, uplift, B:338

titanium  
 in augite, B:851-853, 856, 862-863  
 in Japan Sea sediment, B:1236  
 and parental magma, B:856  
 Site 798, B:722  
 Site 799, B:722  
 vs. Mg/(Mg + Fe), B:858  
 vs. neodymium, B:902  
 vs. zirconium, A(128):101  
 XRF and geochemical log analyses compared, B:1028-1029; 1031-1035

titanium, logging  
 Site 794, B:1420-1421  
 Site 796, B:1423  
 Site 797, B:1426-1427  
 Site 798, B:1401-1402  
 Site 799, B:1404, 1408-1409

titanium/aluminum ratio  
 in augite, B:851-853  
 as indicator of eolian contribution, B:1239  
 Site 794, B:1249

titanium oxide. *See* titanium

titannomagnetite, Site 797, B:949, 951

$T_{max}$   
 Site 798, B:668-669  
 Site 799, B:624-625, 668-669  
 vs. transformation ratio (TR), B:630

tomogram, X-ray radiograph compared, B:1133

Tonga Arc  
 Pb-Pb plots of arc volcanic rocks, B:808  
 and sediment contamination of mantle source, B:812

topography, and heat flow, A(127):300-301

tourmaline, as quartz inclusions, B:108

toxons  
 Site 795, B:542

Toyama Channel, pollen and, B:333

Toyama Deep Sea Fan, A(127):11  
 distribution of deposits, B:1146  
 and heat flow, B:1148  
 sands, A(128):14  
 seismic expression, B:1148

Toyama Trough, A(128):70

TR. *See* transformation yield

trace element molar ratios  
 Site 798, B:616-617  
 Site 799, B:618-621

trace elements, B:677-695, 719-737. *See also* X-ray fluorescence (XRF) analyses; specific trace elements

Al-normalized concentrations, B:689  
 and ash layer magmatic distinctions, B:1378-1379

of basalt, B:839  
 and basaltic basement rocks, B:893

basement, A(128):98; B:781-783, 807, 912-916

- detection limits, INAA analyses, B:1372  
 interelement comparisons, B:690  
 mobility during alteration, B:911–916  
 N-MORB normalized spidergram, B:923–  
     924, 926  
 Site 794, B:682–684, 689, 782–783, 839, 912–  
     913, 918–919, 923  
 Site 795, B:683–685, 689, 705–717, 913–914,  
     924  
 Site 796, B:685  
 Site 797, B:685, 688–691, 839, 914–916, 920–  
     921, 926  
 Site 798, B:722–725, 1367–1372  
 Site 799, B:723–725, 728–731  
 in spinel, B:840–842  
 in strip samples, B:1367–1372  
 vs. loss on ignition, B:918–921
- trace metals, B:1361–1366  
 Site 794, B:1362–1363  
 Site 795, B:1365  
 Site 797, B:1366
- transformation ratio (TR)  
 Site 799, B:624–625  
 vs. T<sub>max</sub>, B:630
- trees, broadleaf, Site 798, B:319
- triaxons  
 Site 795, B:543
- trissocyclids  
 Japan Sea, B:294, 296  
 Site 797, B:303
- Tsugaru Strait, B:423  
 depth, B:342, 440  
 Japan Sea and, B:70, 362–363, 365  
 sill depth, B:351, 457
- Tsushima Basin  
 crustal structure, B:1311–1314, 1318–1319  
 magnetic anomalies, B:1314  
 paleobathymetry, B:1210  
 subsidence history, B:1207–1211, 1214
- Tsushima Current, B:342, 423–424  
 benthic foraminifers and, B:365  
 calcareous nannofossils and, B:180  
 and foraminiferal coiling direction,  
     A(128):165, 310–311  
 inflow and diatom abundance correlated,  
     B:591  
 and Japan Sea, A(128):20–21; B:365  
 and onset of glaciation, B:466  
 and productivity, B:311  
 radiolarians and, B:298–299  
 Site 799, A(128):310, 312  
 and upwelling, B:594  
 warm surface water, B:440, 457, 577
- Tsushima Fault  
 dextral motion along, B:1317  
 and Japan Sea tectonic evolution, B:1315  
 offset along, B:1325
- Tsushima Strait, B:423  
 age, B:180  
 and biosiliceous productivity, B:313  
 depth, B:342, 440  
 and diatom abundance, B:1219–1220  
 diatoms and, B:360  
 Japan Sea and, B:362–363, 365  
 and Kuroshio Current, A(128):164; B:166–167  
 sill depth, B:457  
 Site 798, A(128):164–166  
 Site 799, A(128):312  
 and tectonic subsidence, B:591  
 and Tshushima Current, B:69
- tuff  
 formation, A(127):190–191  
 in logs, A(127):251
- sedimentary structures, A(128):81  
 Site 794, A(127):93–94, A(128):81  
 Site 795, A(127):189–190, 192  
 Site 796, A(127):265–266  
 Site 797, A(127):344–345
- tuff, ash-flow  
 Onnagawa Formation rhyolitic tuff correlated,  
     B:796  
 Site 799, B:793, 803
- tuff, breccia, chemical composition, A(128):287–  
     288
- tuff, hyaloclastite  
 Site 794, B:1391  
 Site 795, B:1374
- tuff, pumice flow, Site 799, B:1373
- tuff, rhyolitic. *See also* tuff, ash-flow; tuff, pumice flow  
 age, B:796  
 ash layers correlated, A(128):33  
 in geochemical logs, B:1399  
 in logs, A(128):339  
 Site 799, A(128):35, 243, 265, 292, 354  
 and velocity, B:1141  
 XRF analyses, A(128):300
- tuff, vitric  
 in logs, B:37  
 and revised lithostratigraphy, B:1232–1233  
 Site 799, B:34, 36, 39, 41–42, 44
- tuff breccia. *See* breccia, tuff
- tuffaceous clayey siltstone. *See* siltstone, tuffaceous clayey
- tungsten, Site 798, B:1370–1371
- turbidite deposits  
 seismic expression, A(127):233  
 Site 795, A(127):192  
 Site 799, A(128):264
- turbidites  
 glauconite-bearing, B:68  
 sedimentation rate and, B:959  
 Site 799, B:104
- turbidity currents, Site 798, B:419, 421
- turbidity deposits, of ash, B:793
- turbidity flows  
 of ash layers, A(128):153–154  
 Site 799, A(128):259
- tylostyles  
 Site 795, B:543
- Tym-Poronaisk Fault, and Japan Sea tectonic evolution, B:1315
- Tyro Basin, paleoenvironment, B:379
- Ulleung Island, ash layers from, A(128):156–157
- unconformity, seismic, and Okushiri Ridge uplift,  
     A(127):312–313
- underconsolidation. *See* consolidation
- United States NW, paleoenvironment, B:337
- uplifting  
 in Japan Sea region, B:1343–1344  
 Site 799, A(128):312
- uptake rates, fluoride  
 Sites 798 and 799, B:66  
 vs. sediment depth, B:65
- upwelling  
 and dark/light cycles, B:591–594  
 enhanced productivity, A(127):267  
 evidence from planktonic foraminifers, B:469  
 and gyre-margin planktonic foraminiferal assemblages, B:466  
 Japan Basin N, A(127):199  
 Japan Sea, B:337  
 and northern hemisphere glaciation,  
     A(127):271
- radiolarians and, B:299  
 and resting-spore laminae, B:551
- and sedimentation rate, A(127):204  
 siliceous ooze deposition, A(127):102–103,  
     191  
 Site 795, A(127):199  
 Site 796, A(127):274  
 Site 797, A(127):349  
 and Tsushima Current, B:594
- uranium  
 in basement rocks, B:807  
 Site 795, B:711, 714  
 Site 798, B:1370–1371  
 Site 799, B:724
- uranium, logging  
 in igneous rocks, A(128):103  
 Site 794, A(128):108; B:1418–1419  
 Site 796, B:1422  
 Site 797, B:1424–1425  
 Site 798, B:1400  
 Site 799, A(128):366–367; B:1403, 1406–1407
- Uyeda probe. *See* temperature
- vacutainer analyses. *See* hydrocarbons, volatile
- van Krevelen plot  
 Site 794, A(127):115–118, 121  
 Site 795, A(127):213, 219  
 Site 796, A(127):285, 288  
 Site 797, A(127):367, 378  
 Site 798, A(128):177, 196; B:429  
 Site 799, B:429, 629
- vegetation, Japanese archipelago, B:320, 325,  
     326, 334
- veins  
 altered to clays, B:886–887  
 and Bagnold effect, B:1180  
 in basement, A(128):92–94  
 and bedding-parallel extension, B:1176  
 chemical composition of alteration clay minerals, B:887  
 coarse-grained material in fillings, B:1179  
 crystallization stages, A(128):92–93  
 and earthquakes, B:1180–1181  
 filled tension gash, B:1189  
 fillings, B:1178  
 and fluid flow/material transport, B:1178–  
     1179  
 formation timing, B:1178, 1180  
 as healed normal faults, B:1178  
 and Japan Sea stress field, B:1187  
 lithology correlated, B:1180  
 and matrix sieving, B:1179–1180  
 and microfaults, B:1180  
 order of filling, B:888  
 origin, B:1178–1181  
 Site 794, B:1182–1183  
 Site 798, A(128):143ff; B:1176–1178  
 Site 799, A(128):263–264, 269–272; B:1178,  
     1190–1193
- veins, siliceous  
 and phosphate, B:64  
 in plagioclase phryic basalt, A(127):217
- velocity  
 anisotropy, A(127):229  
 basement, A(127):389  
 corrected analyses, B:990–1015  
 and crustal structure, B:1078–1083  
 Japan Sea, B:1279–1280, 1283  
 lab and log data compared, B:1279–1280,  
     1284  
 of opal-A to opal-CT transition, B:18, 21  
 porosity effect, B:1280  
 of rhyolitic tuff, B:1141  
 Site 794, A(127):127–134, 139, A(128):102,  
     105; B:1283  
 Site 795, A(127):224–227, 229; B:1283



## SUBJECT INDEX

zirconium/yttrium ratio

- Japan Arc basalt correlated, B:903  
 Japan Basin age compared, B:1337  
 Japan Basin foraminifers compared, B:189  
 Lau Basin formation compared, B:928  
 Layer 1, A(128):71-72, 127  
 Layer 2, A(127):73, A(128):9, 72, 127-128; B:1342  
 Layer 3, A(127):73, A(128):71-72, 127-128; B:1342-1343  
 and lithospheric stretching models, B:1304  
 1305, 1307  
 lithostratigraphy, B:494, 496, 498-499  
 low-Al igneous rocks, B:845  
 magma source, B:782-784, 902, 1341  
 magmatic history, B:787-788  
 magnetic anomalies, A(127):8, A(128):7; B:942, 1314, 1345  
 magnetic properties, B:941-942  
 magnetostratigraphy, A(127):22  
 neodymium isotopes, B:901-903  
 oceanic crust, A(127):73  
 oceanic lithosphere, B:1354  
 oceanographic setting, B:342  
 opal-A to opal-CT transition, B:50  
 opal-CT to quartz transition, B:50  
 oxygenation conditions, B:201-208  
 paleodepth, A(127):199; B:189, 201, 1220-1221  
 pollen, B:328, 471, 479, 482-483, 485-486  
 radiolarians, B:293, 295-296  
 radiometric age, B:784-786  
 rifting, A(127):29-30  
 sedimentation rate, B:287, 297, 1227  
 seismic stratigraphy, A(127):399  
 stress field, B:1047-1059  
 strontium isotopes, B:901-903  
 subsidence history, A(127):20, 30, 349; B:209, 1208-1211, 1214  
 tectonic evolution, B:787-788, 1187, 1316-1317  
 Toyama submarine fan, A(127):11  
 volcanic activity, B:796-797
- Yamato Basin N.** *See also* Yamato Basin; Yamato Basin S  
 age, A(128):67-69  
 basement, A(127):77-81, 126-127, A(128):73-75, 117  
 bathymetry, A(128):70-71  
 biostratigraphy, B:1221-1223  
 crustal structure, B:1075-1106  
 depositional history, A(127):96  
 free-air gravity anomalies, A(127):77, A(128):73  
 heat flow, A(128):71-73, 75  
 igneous event sequence, A(128):95-97  
 magnetic anomalies, A(127):77, A(128):73  
 magnetostratigraphy, B:969-970, 1221-1223  
 paleoenvironment, A(127):102-103  
 rifting history, A(127):143-147  
 sedimentation, A(127):83-84, 147, A(128):77-81  
 seismic stratigraphy, A(127):143  
 Site 794, B:654-656  
 surface projections of fault ruptures, A(128):112
- tectonic setting, A(127):81-83, A(128):75-77, 82  
**Yamato Basin S.** *See also* Yamato Basin; Yamato Basin N  
 basement, A(127):323, 331-334, 408-409  
 bathymetry, A(127):325, 331  
 biostratigraphy, B:1223  
 carbonate-compensation depth, A(127):324  
 crustal structure, A(127):329-330, 333; B:1083, 1104-1106  
 depositional history, A(127):347-351  
 dredge samples, A(127):332  
 free-air gravity anomalies, A(127):330-331, 335  
 geologic history, A(127):323-324  
 heat flow, A(127):330, 332  
 Layer 1, A(127):330  
 Layer 2, A(127):330  
 Layer 3, A(127):330  
 magnetic anomalies, A(127):330, 334  
 magnetostratigraphy, B:973, 1223  
 oceanographic conditions, A(127):349  
 oceanographic history, A(127):337, 409-410  
 paleoenvironment, A(127):356  
 rifting, A(127):408-409  
 sedimentation, A(127):324, 334-335, 337, 409-410  
 seismic stratigraphy, A(127):335, 399  
 Site 797, B:560  
 subsidence, A(127):323, 409  
 tectonic setting, A(127):334, 336
- Yamato Block**, B:809  
**Yamato Rise**, A(127):325, A(128):70  
 basement, A(127):9, 333, A(128):73, 247-249  
 bathymetry, A(128):245-246; B:110, 1136  
 crustal structure, A(127):9, 329-330, 333, A(128):9-11, 245-249, B:1075  
 dredge sites, A(128):253  
 free-air gravity anomalies, A(128):247  
 geologic history, A(128):21-22  
 Green Tuff rocks, A(128):249  
 heat flow, A(127):330, A(128):246, 250; B:1302-1303  
 heavy minerals, B:691  
 Layer 2, A(128):246  
 Layer 3, A(128):246  
 magnetic anomalies, A(128):247; B:1314  
 sediment from, A(127):349  
 sedimentary rocks dredged, A(128):249  
 sedimentation, A(127):11-13  
 Site 799, A(128):312  
 stress field, A(128):245  
 tectonic setting, A(127):8-9, A(128):249-250; B:156
- Yamato Rise E.** normal faults, A(127):82-83  
**Yamato Seamount Chain**, A(128):70  
 basement, A(127):333, A(128):73  
 crustal structure, A(128):128  
 dredge samples, A(127):77-78  
 strontium isotopes, B:812
- Yamato Trough**, A(128):70  
 stress field, A(128):340  
 vertical seismic profile, A(128):342-343
- Yangsan Fault**  
 and Japan Sea tectonic evolution, B:1315
- offset along, B:1325  
**Yellow River**  
 fresh water input to Japan Sea, B:458  
 origin of dark/light cycle detrital component, B:595  
**Young's modulus**, Site 794, B:1050  
**ytterbium**  
 in ash layers, B:1386  
 Site 798, B:1370-1371  
 vs. aluminum, B:737
- yttrium**  
 alteration and, B:909-911  
 basement, A(128):99  
 Site 794, B:1363  
 Site 795, B:1365  
 Site 797, B:1366  
 vs. barium, A(128):100  
 vs. chromium, A(128):101  
 vs. zirconium, A(128):100; B:783
- zeolites.** *See also* zeolitization; names of specific zeolites  
 in basaltic andesite, A(127):217  
 in blue tuff, B:119  
 formation temperature, B:139  
 and opal-A to opal-CT transition, B:10  
 in pyroxene plagioclase phryic basalt, A(127):217-219
- SEM photographs, B:30  
 Si/Al + Fe ratios, B:139, 141  
 Site 797, B:142  
 and volcanic glass alteration, B:1375-1377  
 zonation of, B:143
- zeolitization.** *See also* zeolites  
 provenance information destroyed, B:148  
 of sandstone, B:137-139
- Zijderveld plots**  
 of Leg 128 pilot samples, B:971  
 Site 794, A(127):104-105, A(128):104-105  
 Site 795, A(127):203, 206-207  
 Site 796, A(127):277-279  
 Site 798, A(128):173  
 Site 799, A(128):316, 318-320, 322-323
- zinc**  
 in dark/light cycles, B:569  
 Site 794, B:1363  
 Site 795, B:710, 1365  
 Site 797, B:1366  
 Site 798, B:1370-1371  
 in sulfide deposits, A(128):21
- zirconium**  
 in ash layers, B:1386  
 and basalt fractionation models, B:873  
 basement, A(128):99  
 and fractionation models, B:881  
 Site 794, B:1363  
 Site 795, B:1365  
 Site 797, B:1366  
 vs. barium, A(128):100  
 vs. niobium, A(128):100; B:783  
 vs. titanium, A(128):101  
 vs. yttrium, A(128):100; B:783  
 vs. zirconium/yttrium ratio, A(128):100
- zirconium/yttrium ratio, vs. zirconium, A(128):100

## TAXONOMIC INDEX

- Abies*  
Japan Sea, B:328  
Neogene climate and, B:337
- abyssalica*, *Silicosigmoilina*, Japan Sea, B:505
- acquilonius*, *Druppatractus*  
Japan Sea, B:228  
Site 799, B:235
- acquilonius*, *Stylacontarium*  
Japan Sea, B:297  
Site 794, B:305
- acquilonius*, *Stylacontarium* sp. cf., Japan Sea, B:297
- Actinocyclus ingens*, Pacific Ocean, B:309
- Actinocyclus oculatus*  
Japan Sea, B:258  
Site 798, B:361
- Aesculus*, Site 796, B:491
- akitaensis*, *Anthocorys*, Site 797, B:307
- akitaensis*, *Anthocorys?*  
Japan Sea, B:228  
Site 799, B:225, 235
- akitaensis*, *Uvigerina*, Site 798, B:390, 441, 443–444
- Alnus*, Site 795, B:490
- aminensis*, *Bermudezina*, Site 795, B:537
- Ammodiscus macilentus*, Site 795, B:537
- Ammodiscus tenuis*, Site 797, B:537
- Ammodochium rectangulare*  
Japan Sea, B:238  
Site 798, B:248
- Ammodochium rectangulare* Zone, Japan Sea, B:240
- Ammosphaeroidina* sp., Site 795, B:537
- Amnovertellina* sp., Site 797, B:537
- Amphirhopalum ypsilon*, Site 798, B:235
- Angulogerina kokozuraensis*, Site 798, B:390
- Anthocorys akitaensis*, Site 797, B:307
- Anthocorys?* *akitaensis*  
Japan Sea, B:228  
Site 799, B:225, 235
- antiqua*, *Thalassiosira*, Japan Sea, B:258
- antiqua antiqua*, *Ebriopsis*  
Japan Sea, B:238  
Site 799, B:248
- antiqua cornuta*, *Ebriopsis*, Site 798, B:248
- aperta*, *Gephyrocapsa*, Leg 128, B:164
- apertura*, *Pullenia*, Site 798, B:392
- Artemisia*, Site 795, B:491
- asanoi*, *Eucyrtidium*, Site 799, B:236
- asanoi*, *Neogloboquadrina*, Site 797, B:208
- asanoi*, *Planularia*, Site 798, B:389
- asanoi*, *Reticulofenestra*, Site 798, B:164
- Astrononion* sp., Site 797, B:540
- auriculata*, *Globobulimina*, Site 798, B:390
- baccata japonica*, *Karreriella*, Site 798, B:389
- Bathysiphon* sp., Site 797, B:537
- Bermudezina aminensis*, Site 795, B:537
- Betula*  
Site 795, B:490  
Site 797, B:490
- biconica*, *Fissurina*, Site 798, B:389
- bigelowii*, *Braarudosphaera*, Site 795, A(127):194
- Bolivina*, Site 798, B:557
- Bolivina decussata*, Site 798, B:389
- Bolivina pacifica*  
paleoenvironment and, B:379  
Site 798, A(128):166; B:373, 389
- Bolivina spissa*, Site 798, B:389
- Bolivinita quadrilatera*, Site 799, B:389
- Botryostrobus bramlettei*, Site 794, A(127):102
- Braarudosphaera bigelowii*, Site 795, A(127):194
- bradyi*, *Cibicidooides*  
Japan Sea, B:505  
Site 797, B:538
- bradyi*, *Eggerella*, Site 795, B:537
- bramlettei*, *Botryostrobus*, Site 794, A(127):102
- Buccella* cf. *kuromatsunaiensis*, Site 797, B:540
- Buliminella rostrata*, Site 795, B:539
- Buliminella striata*  
Site 797, B:539  
Site 798, B:390
- Buliminella tenuata*, Site 798, B:390
- bulloides*, *Globigerina*  
Leg 127, B:193, 200  
Leg 128, B:60–61  
Site 797, B:591–594  
Site 798, B:441–443, 457, 460, 462–463, 465–466, 469, 564
- bulloides*, *Pullenia*, Site 797, B:540
- bulloides*, *Sphaeroidina*, Site 798, B:390
- bulloides* (s.l.), *Globigerina*, Site 798, A(128):165
- bulloides umbilicata*, *Globigerina*, Oki Ridge, B:444
- cabrilloensis*, *Clathrocyclas*  
Japan Sea, B:297  
Site 794, B:307, 1221  
Site 797, B:307
- Calcidiscus leptoporus*, Site 798, B:168
- Calcidiscus macytrei*  
Japan Sea, B:174, 182  
Site 798, B:164
- californica*, *Rouxia*, Japan Sea, B:259
- cancellata*, *Cyclammina*  
Site 797, B:538  
Site 799, B:389
- Carduoidea*, Site 795, B:491
- caribeanica*, *Gephyrocapsa*  
Japan Sea, B:173, 183  
Leg 128, B:164
- Carpinus*, Site 797, B:490
- Carya*  
Japan Sea, B:484–486  
Site 795, B:490
- Cassidulina*, Leg 128, B:62
- Cassidulina japonica*, Site 798, B:389
- Cassidulina norcrossi*, Site 798, B:390
- Cassidulina norvangi*, Site 798, B:390
- Cassidulina* sp. A, Site 798, B:390
- Cassidulinoides tenuis*, Site 799, B:390
- Castanea*, Site 794, B:490
- Catapsydrax parvulus*, Site 797, B:189, 1220, 1223
- Chaetoceros* resting spores, Site 798, B:547–557, 564
- Chilostomella oolina*  
Site 797, B:539  
Site 799, B:392
- Chilostomella oolina/Cyclammina* sp. C Zone, Site 794, B:517
- Chilostomella ovoidae*, Site 798, B:392
- Chilostomella fimbriata*, Site 798, B:392
- Chondrites*  
Japan Sea, B:1232  
Site 798, B:565, 569  
Site 799, B:1133
- Cibicidooides bradyi*  
Japan Sea, B:505  
Site 797, B:538
- Cibicidooides* sp. A, Site 797, B:538
- Cibicidooides wuellerstorfi*  
Japan Sea, B:505  
Site 797, B:538
- Cibicidooides wuellerstorfi* Zone, Site 797, B:521, 528
- circulus*, *Paramesocena*, Site 798, B:248
- circumtexta*, *Peripyramis*, Site 795, B:304
- Circumtexta pacifica*, Site 795, B:304
- Clathrocyclas cabrilloensis*  
Japan Sea, B:297  
Site 794, B:307, 1221  
Site 797, B:307
- clavatum*, *Elphidium*, Site 798, B:392
- Coccoilithus miopelagicus* Subzone, Japan Sea, B:173
- Coccoilithus pelagicus*  
Site 798, B:167–168  
Site 799, B:166
- communis*, *Martinottiella*  
Japan Sea, B:1201  
Site 795, B:537  
Site 797, B:537  
Site 798, B:389
- compactus*, *Sphenolithus*, Japan Sea, B:183
- complanata*, *Stainforthia*  
Japan Sea, B:388  
Site 798, B:390
- compressa*, *Islandiella* cf., Site 797, B:539
- compressa*, *Proteonina*, Site 795, B:537
- compressa*, *Spirosigmoilinella*  
Site 795, B:537  
Site 798, B:379–381, 389
- convexa*, *Thalassiosira*, Japan Sea, B:258
- Corylus*, Site 794, B:490
- Coscinodiscus curvatulus*, Site 794, B:1223
- Coscinodiscus marginatus*  
Aikawa section, Honshu, B:316  
Japan Sea, B:291  
Leg 127, B:1219–1220  
Leg 128, B:309–314, 1219–1220  
North Pacific, B:315–316  
Site 433, B:316  
Site 581, B:316  
Site 794, B:341–357  
Site 795, B:341–357  
Site 797, B:341–357  
Site 798, B:361, 554, 1224  
Site 799, B:1224
- crassimargo*, *Cribrostomoides* cf., Site 795, B:537
- Cribrostomoides* cf. *crassimargo*, Site 795, B:537
- crux var. stauracanthus*, *Distephanus*  
Japan Sea, B:239, 244  
Site 798, B:248
- Cryptomeria*  
Japan Sea, B:319, 328, 334  
Neogene climate and, B:337
- Cunninghamia*, Site 797, B:490
- curvatulus*, *Coscinodiscus*, Site 794, B:1223
- curvirostris*, *Rhizosolenia*  
Hess Rise, B:360  
Japan Sea, B:258  
Leg 127, B:581  
North Pacific, B:311  
Site 798, B:361, 468
- Cycladophora davisiana*, Site 799, B:235
- Cycladophora davisiana cornutoides*  
Japan Sea, B:230  
Site 799, B:235

*Cycladophora davisiана davisiана*, Japan Sea, B:228  
*Cycladophora robusta*, Japan Sea, B:299  
 Site 794, B:307  
*Cyclammina cancellata*, Site 797, B:538  
 Site 799, B:389  
*Cyclammina ezoensis*, Site 795, B:538  
*Cyclammina ezoensis* Zone, Site 795, B:521  
*Cyclammina orbicularis*, Site 795, B:538  
*Cyclammina pusilla*, Site 797, B:538  
*Cyclammina* sp. A, Site 797, B:538  
*Cyclammina* sp. A/ *Cyclammina?* sp. Zone age models and, B:531  
 Site 797, B:521  
*Cyclicargolithus floridanus*, Site 795, A(127):195  
 Site 797, B:179  
*Cyrtocapsella tetrapera*, Japan Sea, B:230  
 Site 794, B:304  
 Site 797, A(127):356; B:304  
 Site 799, B:235  
*Cyrtocapsella tetrapera* Zone, Site 799, B:226, 1220  
 davisiana, *Cycladophora*, Site 799, B:235  
 davisiana, *Theocalyptra*, Japan Sea, B:297, 299  
 Site 794, B:306  
 Site 795, B:303  
*davisiана cornutoides*, *Cycladophora*, Japan Sea, B:230  
 Site 799, B:235  
*davisiана davisiана*, *Cycladophora*, Japan Sea, B:228  
*decussata*, *Bolivina*, Site 798, B:389  
*deflandrei*, *Discoaster*, Japan Sea, B:174  
*delmontensis*, *Stichocorys*, Japan Sea, B:228, 230  
 Site 799, B:225-226, 236  
*Dentalina* sp., Site 797, B:539  
*Denticulopsis dimorpha*, Japan Sea, B:250  
 South Atlantic, B:309  
*Denticulopsis dimorpha* Zone, Site 795, B:188  
*Denticulopsis hustedtii*, Japan Sea, B:250  
*Denticulopsis hustedtii* Zone, Site 795, B:189  
*Denticulopsis kamtschatica*, Site 795, B:1223  
*Denticulopsis katayamae* Zone, Site 794, B:189  
 Site 797, B:1223  
*Denticulopsis praedimorpha*, Japan Sea, B:250  
*Denticulopsis praedimorpha* Zone, Japan Sea, B:249-250  
 Site 795, B:189  
*Dictyocha neopseudofibula*, Japan Sea, B:237-238  
 Site 798, B:248  
*Dictyocha subarctios*, Japan Sea, B:238-239  
 Site 798, B:248  
*Dictyocha subarctios* Zone, Japan Sea, B:240-241  
*Dictyococcites perplexus*, Japan Sea, B:182  
*Didymocyrtis mammifera*, Site 799, B:235  
*Didymocyrtis tetrathalamus*, Site 797, B:306  
 Site 798, B:235  
*dimorpha*, *Denticulopsis*, Japan Sea, B:250  
 South Atlantic, B:309  
*Discoaster deflandrei*, Japan Sea, B:174

*Discoaster exilis* Zone, Japan Sea, B:173  
*Discoaster hamatus* Zone, Site 797, B:189  
*Discoaster kugleri* Subzone, Japan Sea, B:173  
*Distephanus crux* var. *stauracanthus*, Japan Sea, B:239, 244  
 Site 798, B:248  
*Distephanus jimlingii*, Japan Sea, B:238  
 Site 798, B:248  
*Distephanus jimlingii* Zone, Japan Sea, B:239  
*Distephanus octangulatus*, Site 798, B:248  
*Distephanus octangulatus* Zone, Japan Sea, B:241  
*Distephanus slavnicii*, Site 798, B:248  
*Distephanus speculum*, Site 799, B:248  
*doliolus*, *Pseudoeunotria* age models and, B:362  
 Hess Rise, B:360  
 Honshu coast, B:360  
*Druppatractus acqilonius*, Japan Sea, B:228  
 Site 799, B:235  
*Ebriopsis antiqua antiqua*, Japan Sea, B:238  
 Site 799, B:248  
*Ebriopsis antiqua antiqua* Zone, Japan Sea, B:239-240  
*Ebriopsis antiqua cornuta*, Site 798, B:248  
*echigoensis*, *Fissurina*, Site 798, B:389  
*echigoensis*, *Miliammina*, Leg 127, B:200  
 Site 798, B:389  
 Site 799, A(128):311  
*Echigoina hatai*, Site 798, B:392  
*Eggerella bradyi*, Site 795, B:537  
*Eggerella* sp., Site 797, B:537  
*Eiloehedra rotunda*, Site 798, B:391  
*elliptica*, *Mesocena*, Site 799, B:248  
*Elphidium cf. perforatum*, Site 795, B:540  
*Elphidium clavatum*, Site 798, B:392  
*Elphidium jenseni*, Site 797, B:540  
*Elphidium subarcticum*, Site 797, B:540  
 Site 798, B:392  
*Emiliania huxleyi*, Site 795, A(127):194  
 Site 798, B:164  
*Engelhardia*, Site 796, B:490  
 Site 797, B:490  
*Ephedra*, Site 795, B:490  
*Epistominella*, Leg 128, B:62  
*Epistominella pulchella*, Site 798, B:391  
*Euchitonita furcata*, Site 798, B:235  
*Euchitonita* sp., Site 794, B:306  
*Eucyrtidium asanoi*, Site 799, B:236  
*Eurya*, Site 796, B:491  
*Evolvocassidulina kuwanoi*, Site 797, B:539  
*ezoensis*, *Cyclammina*, Site 795; B:538  
*Fagus*, Japan Sea, B:484-486  
 Site 795, B:491  
 Site 797, B:491  
*fimbriata*, *Chilostomellina*, Site 798, B:392  
*Fissurina biconica*, Site 798, B:389  
*Fissurina echigoensis*, Site 798, B:389  
*Fissurina orbigniana*, Site 797, B:539  
*floriadanus*, *Cyclicargolithus*, Site 795, A(127):195  
 Site 797, B:179  
*Funpingopollenites*, Site 797, B:491  
*furcata*, *Euchitonita*, Site 798, B:235  
*fusiformis*, *Stainforthia* paleoenvironment and, B:379  
 Site 798, B:390  
*Gephyrocapsa*, Site 798, A(128):162  
*Gephyrocapsa aperta*, Leg 128, B:164  
*Gephyrocapsa caribbeanica*, Japan Sea, B:173, 183  
 Leg 128, B:164  
*Gephyrocapsa caribbeanica* Zone, Japan Sea, B:174  
*Gephyrocapsa oceanica*, Japan Sea, B:173, 183  
 Leg 128, B:164  
 Site 798, B:167-168  
*Gephyrocapsa* small, Japan Sea, B:183  
*Ginkgo*, Site 795, B:490  
*Glabratella subopercularis*, Site 797, B:540  
*glacialis*, *Spongotrochus*, Site 795, B:306  
*Globigerina*, Site 798, B:569  
*Globigerina bulloides*, Leg 127, B:193, 200  
 Leg 128, B:60-61  
 Site 797, B:591-594  
 Site 798, B:441-443, 457, 460, 462-463, 465-466, 469, 564  
*Globigerina bulloides* (s.l.), Site 798, A(128):165  
*Globigerina bulloides umbilicata*, Oki Ridge, B:444  
*Globigerina incompta*, Site 798, B:463  
*Globigerina quinqueloba*, Leg 127, B:200  
 Site 798, B:457, 460, 469  
*globigeriniformis*, *Trochammina* cf., Site 795, B:537  
*Globigerinita* spp., Site 798, B:457, 460, 469  
*Globigerinoides sicanus/Praeorbulina glomerata* Zone, Japan Sea, B:526  
*Globobulimina auriculata*, Site 798, B:390  
*Globobulimina pacifica*, Site 798, B:390  
*Globobulimina perversa*, Site 797, B:539  
*Globobulimina pupoides*, Site 797, B:539  
*Globobulimina subaffinis*, Site 798, B:390  
*Globocassidulina globosa* s.l., Site 797, B:539  
*Globocassidulina* sp. A, Site 797, B:539  
*Globoquadrina venezuelana*, Leg 127, B:189  
 Site 797, B:1220  
*Globorotalia* cf. *G. fohsi* Zone, Leg 127, B:200  
*Globorotalia inflata*, Site 798, B:457, 469  
*Globorotalia peripheronida/Globorotalia quinifalcata* Zone, Japan Sea, B:526  
*Globorotalia praescitula*, Site 797, B:189, 1220, 1223  
*globosa*, *Nonionella*, Site 798, B:391  
*globosa*, *Oolina*, Japan Sea, B:387  
 Site 797, B:539  
*globosa* s.l., *Globocassidulina globosa*, Site 797, B:539  
*grande*, *Lychnocanium* sp. cf., Site 794, A(127):101  
 Site 797, B:305  
*grateloupi*, *Pseudononion*, Site 798, B:392  
*Gyroidina orbicularis*, Site 797, B:540  
*Gyroidina orbicularis/Gyroidinoides planulatus* Zone, age models and, B:531  
 Site 797, B:521, 528  
*Gyroidinoides planulatus*, Site 797, B:540  
*hanzawai*, Tosaia, Site 797, B:538

*Hanzawaia tagensis/Heterolepa praecincta* Zone, Japan Sea, B:526

*Haplophragmoides* sp., Site 795, B:537

*hataii*, *Echigoina*, Site 798, B:392

*helena*, *Islandiella*, Site 797, B:539

*Helicospaera ampliaperta* Zone, Site 797, B:1220, 1223

*Helicospaera perch-nielseniae*, Japan Sea, B:183

*Helicospaera sellii*

- Leg 128, B:164
- Site 798, B:164

*heteromorphus*, *Sphenolithus*

- Japan Sea, B:173-174, 183
- Leg 127, B:189

*hexagona*, *Oolina*, Site 797, B:539

*Hippocrepina variabilis*, Site 795, B:537

*Hopkinsina morimachiensis* Zone, Japan Sea, B:526

*hustedtii*, *Denticulopsis*, Japan Sea, B:250

*huxleyi*, *Emiliania*

- Site 795, A(127):194
- Site 798, B:164

*incompta*, *Globigerina*, Site 798, B:463

*inflata*, *Globorotalia*, Site 798, B:457, 469

*ingens*, *Actinocyclus*, Pacific Ocean, B:309

*Islandiella* cf. *compressa*, Site 797, B:539

*Islandiella helena*, Site 797, B:539

*Islandiella* sp., Site 797, B:539

*jacksonii*, *Thalassiosira*, Japan Sea, B:259

*japonica*, *Cassidulina*, Site 798, B:389

*japonica*, *Sethocystis*

- Japan Sea, B:297, 301
- Site 794, B:305

*japonica*, *Sphaeroidina*, Site 797, B:538

*japonica*, *Thecosphaera*, Site 795, B:305

*jensenii*, *Elphidium*, Site 797, B:540

*jimlingii*, *Distephanus*

- Japan Sea, B:238
- Site 798, B:248

*jouseae*, *Nitzschia*, B:1219

*Juglans*

- Site 794, B:490
- Site 795, B:490

*kamtschatica*, *Denticulopsis*, Site 795, B:1223

*kamtschatica*, *Neodenticula*

- North Pacific, B:309
- Site 795, B:970
- Site 798, B:361
- Site 799, A(128):302

*Karreriella baccata japonica*, Site 798, B:389

*koizumii*, *Neodenticula*

- Japan Sea, B:258
- Site 798, B:361

*koizumii*, *Neodenticulopsis*, Site 798, B:1224

*kokazuraensis*, *Angulogerina*, Site 798, B:390

*kuromatsunaiensis*, *Buccella* cf., Site 797, B:540

*kuwanoi*, *Evolvocassidulina*, Site 797, B:539

*labradorica*, *Nonionellina*

- Site 797, B:539
- Site 798, B:391

*lacunosa*, *Pseudomelania*

- Japan Sea, B:183
- Leg 127, B:581
- Leg 128, B:163
- Site 798, B:468
- Site 799, B:166

*laevis*, *Lagena*, Site 798, B:389

*Lagena laevis*, Site 798, B:389

*Lagena substrata*, Japan Sea, B:387

*Lagena tricaritortuosa*, Site 798, B:389

*langii*, *Sphaeropyle*

- Japan Sea, B:297
- Site 794, B:305
- Site 795, B:303
- Site 799, B:236

*Larnacantha polyacantha*, Site 795, B:306

*lentiginosa*, *Thalassiosira*, Southern Ocean, B:309

*lepidula*, *Stilosomella*, Site 798, B:390

*leptoporoides*, *Calcidiscus*, Site 798, B:168

*Lipmanella* sp. aff. *L. redondoensis*, Japan Sea, B:297

*Lipmanella* sp. aff. *Thecocrys redondoensis*

- Japan Sea, B:297
- Site 794, B:305

*Liquidambar*

- Japan Sea, B:484-486
- Site 797, B:491

*Lithastractus tochiensis*, Site 794, B:305

*Lithopera renzae renzae*, Site 799, B:236

*lobatula*, *Uvigerina*, Site 798, B:390

*Lychnocanium* cf. *L. grande*, Site 794, B:1221

*Lychnocanium nipponicum*

- Site 794, A(127):101
- Site 797, B:305

*Lychnocanium* sp. cf. *L. grande*

- Site 794, A(127):101
- Site 797, B:305

*Lychnocanoma* sp.

- Japan Sea, B:230
- Site 799, B:225-226, 236

*macilentus*, *Ammodiscus*, Site 795, B:537

*macintyrei*, *Calcidiscus*

- Japan Sea, B:174, 182
- Site 798, B:164

*makiyama*, *Sagarites*, Site 799, B:35, 45

*Makiyama titanni*, Site 799, B:35, 45

*mammifera*, *Didymocystis*, Site 799, B:235

*mammilaris*, *Tholosypris*, Site 794, B:305

*marginatus*, *Coscinodiscus*

- Aikawa section, Honshu, B:316
- Japan Sea, B:291
- Leg 127, B:1219-1220
- Leg 128, B:309-314, 1219-1220
- North Pacific, B:315-316
- Site 433, B:316
- Site 581, B:316
- Site 794, B:341-357
- Site 795, B:341-357
- Site 797, B:341-357
- Site 798, B:361, 554, 1224
- Site 799, B:1224

*Marginulina* sp., Site 797, B:539

*Martinottiella communis*

- Japan Sea, B:1201
- Site 795, B:537
- Site 797, B:537
- Site 798, B:389

*Martinottiella communis/Cribrostomoides* cf. *crassimargo* Zone, Site 795, B:521

*Martinottiella communis/Globobulimina* pu-

*poides* Zone, Site 797, B:524

*Martinottiella communis* Zone, Site 795, B:517

*Martinottiella nodulosa*, Site 798, B:389

*marujamica*, *Thalassiosira*, Japan Sea, B:259

*masudai*, *Valvularia*, Site 797, B:538

*Melonis nikobarensis*

- Site 797, B:540
- Site 798, B:392

*Melonis pompilioides*

- Site 797, B:540
- Site 799, B:383, 392

*Melosira*, Site 798, B:554

*Mesocena elliptica*, Site 799, B:248

*Mesocena quadrangula*, Site 798, B:248

*Metasequoia*, Site 796, B:490

*Miliammina echigoensis*

- Leg 127, B:200
- Site 798, B:389
- Site 799, A(128):311

*Miliammina echigoensis* Zone, Leg 127, B:201

*miocenica*, *Nonionella*, Site 797, B:539

*miocenica*, *Thalassiosira*, B:1219

*miocenica*, *Uvigerina*, Site 797, B:539

*Miogypsina-Operculina* Zone, Japan Sea, B:526

*modelensis*, *Uvigerina* cf., Site 797, B:539

*mur rhina*, *Pyrgo*

- Site 797, B:538
- Site 798, B:389

*naraensis*, *Pseudoparella*, Site 798, B:391

*Neodenticula kamtschatica*

- North Pacific, B:309
- Site 795, B:970
- Site 798, B:361
- Site 799, A(128):302

*Neodenticula kamtschatica* Zone

- Leg 127, B:188
- Site 796, B:1223

*Neodenticula koizumii*

- Japan Sea, B:258
- Site 798, B:361

*Neodenticula koizumii* Zone

- Leg 127, B:188
- Site 795, B:344
- Site 796, B:1223
- Site 797, B:345

*Neodenticula koizumii*-*Neodenticula kamtschatica* Zone

- Site 794, B:344
- Site 795, B:344-345
- Site 797, B:345

*Neodenticulopsis koizumii*, Site 798, B:1224

*Neogloboquadrina asanoi*, Site 797, B:208

*Neogloboquadrina pachyderma*

- age models and, B:362
- Japan Sea, A(128):30
- Leg 127, B:188, 193, 200, 218-219, 1220-1221
- Leg 128, B:1220-1221
- Oki Ridge, B:444
- Site 795, A(127):195
- Site 797, B:208
- Site 798, A(128):165-166; B:457, 459-469
- Site 799, A(128):310-311

*Neogloboquadrina pachyderma dextral/ Globorotalia orientalis* Zone, Oga Peninsula, B:459

*Neogloboquadrina pachyderma sinistral/ Globigerina incompta* Zone, Oga Peninsula, B:459

*Neogloboquadrina pachyderma sinistral/ Globigerina quinqueloba* Zone, Oga Peninsula, B:459

*neopseudofibula*, *Dictyocha*

- Japan Sea, B:237-238
- Site 798, B:248

*Nephrosypris? pervia*

- Japan Sea, B:230-231
- Site 798, B:235

*nidulus*, *Thalassiosira*, Japan Sea, B:258

*nikobarensis*, *Melonis*

- Site 797, B:540
- Site 798, B:392

*nipponica*, *Plectina*, Site 797, B:537

*nipponicum*, *Lychnocanium*  
Site 794, A(127):101  
Site 797, B:305

*Nitzschia jouseae*, B:1219

*Nitzschia pliocena*, Japan Sea, B:259

*Nitzschia reinholdii*  
Hess Rise, B:360  
Site 798, B:361

*Nitzschia rolandii*, Japan Sea, B:259

*nitzschioides*, *Thalassionema*, Site 794, B:1223

*nitzschioides*, *Thalassiothrix*, Site 798,  
A(128):160

*nodulosa*, *Martinottiella*, Site 798, B:389

*Nonionella globosa*, Site 798, B:391

*Nonionella miocenica*, Site 797, B:539

*Nonionella stella*  
Site 797, B:539  
Site 799, B:391

*Nonionellina labradorica*  
Site 797, B:539  
Site 798, B:391

*norcrossi*, *Cassidulina*, Site 798, B:390

*norvangi*, *Cassidulina*, Site 798, B:390

*oceania*, *Gephyrocapsa*  
Japan Sea, B:173, 183  
Leg 128, B:164  
Site 798, B:167–168

*octacantha*, *Tetrapyle*  
Site 794, B:306  
Site 797, B:306  
Site 798, B:236

*octangulatus*, *Distephanus*, Site 798, B:248

*oculatus*, *Actinocyclus*  
Japan Sea, B:258  
Site 798, B:361

*oolina*, *Chilostomella*  
Site 797, B:539  
Site 799, B:392

*Oolina globosa*  
Japan Sea, B:387  
Site 797, B:539

*Oolina hexagona*, Site 797, B:539

*orbicularis*, *Cyclammina*, Site 795, B:538

*orbicularis*, *Gyroidina*, Site 797, B:540

*orbigniana*, *Fissurina*, Site 797, B:539

*Oridorsalis tener*, Site 798, B:392

*Oridorsalis umboatus*  
Site 797, B:540  
Site 798, B:392

*Oridorsalis umboatus/Cibicidoides wuellerstorfi*  
Zone, Site 795, B:517, 528

*ovoidea*, *Chilostomella*, Site 798, B:392

*pachyderma*, *Neogloboquadrina*  
age models and, B:362  
Japan Sea, A(128):30  
Leg 127, B:188, 193, 200, 218–219,  
1220–1221  
Leg 128, B:1220–1221  
Oki Ridge, B:444  
Site 795, A(127):195  
Site 797, B:208  
Site 798, A(128):165–166; B:457, 459–469  
Site 799, A(128):310–311

*pacifica*, *Bolivina*  
paleoenvironment and, B:379  
Site 798, A(128):166; B:373, 389

*pacifica*, *Circumtexta*, Site 795, B:304

*pacifica*, *Globobulimina*, Site 798, B:390

*Parafissurina* sp., Site 797, B:539

*Paramesocena circulus*, Site 798, B:248

*parkerae*, *Tenuitella*, Leg 127, B:200

*parkerae*, *Thalmannammina* cf., Site 797, B:537

*parvulus*, *Catapsydrax*, Site 797, B:189, 1220,  
1223

*Pasania*  
Site 794, B:490  
Site 796, B:490

*pelagicus*, *Coccolithus*  
Site 798, B:167–168  
Site 799, B:166

*perch-nielseniae*, *Helicosphaera*, Japan Sea,  
B:183

*peregrina*, *Stichocorys*  
Japan Sea, B:297  
Site 794, A(127):102  
Site 797, A(127):356  
Site 799, B:236

*perforatum*, *Elphidium* cf., Site 795, B:540

*Peripyramis circumtexta*, Site 795, B:304

*Peripyramis* sp.  
Japan Sea, B:301  
Site 795, B:306

*perplexus*, *Dictyococcites*, Japan Sea, B:182

*perversa*, *Globobulimina*, Site 797, B:539

*pervia*, *Nephrosyrpis*?  
Japan Sea, B:230–231  
Site 798, B:235

*Picea*  
Japan Sea, B:328  
Neogene climate and, B:337

*Pinus*  
Japan Sea, B:319, 328  
Neogene climate and, B:337

*Planolites*, Site 798, B:566–567, 569

*Planularia asanoi*, Site 798, B:389

*planulatus*, *Gyroidinoides*, Site 797, B:540

*Plectina nipponica*, Site 797, B:537

*Pleurostomella* sp. A, Site 799, B:390

*pliocena*, *Nitzschia*, Japan Sea, B:259

*polyacantha*, *Larnacantha*, Site 795, B:306

*pompilioides*, *Melonis*  
Site 797, B:540  
Site 799, B:383, 392

*praebergonii*, *Rhizosolenia*, B:1219  
Indian Ocean, B:309

*praedimorpha*, *Denticulopsis*, Japan Sea, B:250

*praescitula*, *Globorotalia*, Site 797, B:189, 1220,  
1223

*proboscidea*, *Uvigerina*, Site 797, B:539

*Proteonina compressa*, Site 795, B:537

*Pseudoeumilia lacunosa*  
Japan Sea, B:183  
Leg 127, B:581  
Leg 128, B:163  
Site 798, B:468  
Site 799, B:166

*Pseudoeunotia doliolus*  
age models and, B:362  
Hess Rise, B:360  
Honshu coast, B:360

*Pseudodonion grateloupi*, Site 798, B:392

*Pseudoparrella naraensis*, Site 798, B:391

*Pseudoparrella takayanagii*, Site 798, B:391

*pseudoumbilica*, *Reticulofenestra*, Japan Sea,  
B:183

*Pterocarya*, Site 794, B:490

*pulchella*, *Epistominella*, Site 798, B:391

*Pullenia apertura*, Site 798, B:392

*Pullenia bulloides*, Site 797, B:540

*Pullenia salisburyi*  
Site 797, B:540  
Site 798, B:392

*Pullenia subcarinata*, Site 797, B:540

*pupoides*, *Globobulimina*, Site 797, B:539

*pusilla*, *Cyclammina*, Site 797, B:538

*Pyrgo murrhina*  
Site 797, B:538  
Site 798, B:389

*Pyrgo* sp., Site 797, B:538

*quadrangula*, *Mesocena*, Site 798, B:248

*quadrilatera*, *Bolivinita*, Site 799, B:389

*Quercus*  
Japan Sea, B:328  
Neogene climate and, B:337

*Quercus*, D., Site 794, B:491

*Quercus*, E.  
Japan Sea, B:484–486  
Site 794, B:491  
Site 796, B:491

*quinqueloba*, *Globigerina*  
Leg 127, B:200  
Site 798, B:457, 460, 469

*Quinqueloculina* sp.  
Site 797, B:538  
Site 799, B:373

*rectangulare*, *Ammodochium*  
Japan Sea, B:238  
Site 798, B:248

*redondoensis*, *Lipmanella* sp. aff., Japan Sea,  
B:297

*redondoensis*, *Thecocorys*  
Japan Sea, B:297  
Site 794, B:305

*reinholdii*, *Nitzschia*  
Hess Rise, B:360  
Site 798, B:361

*renzae renzae*, *Lithopera*, Site 799, B:236

*Reophax* sp.  
Site 795, B:537  
Site 797, B:537

*Reticulofenestra asanoi*, Site 798, B:164

*Reticulofenestra pseudoumbilica*, Japan Sea,  
B:183

*Reticulofenestra* small  
Japan Sea, B:183  
Site 797, A(127):352

*Rhizammina* sp., Site 797, B:537

*Rhizosolenia curvirostris*  
Hess Rise, B:360  
Japan Sea, B:258  
Leg 127, B:581  
North Pacific, B:311  
Site 798, B:361, 468

*Rhizosolenia praebergonii*, B:1219  
Indian Ocean, B:309

*Rhus*, Site 794, B:491

*robusta*, *Cycladophora*  
Japan Sea, B:299  
Site 794, B:307

*robusta*, *Sphaeropyle*  
Japan Sea, B:297  
Site 794, B:1221  
Site 797, B:305

*rolandii*, *Nitzschia*, Japan Sea, B:259

*Rossiella tatsunokuchiensis*, Japan Sea, B:258–  
259

*rostrata*, *Bulimina*, Site 795, B:539

*rotunda*, *Eiloehdra*, Site 798, B:391

*Rouxia californica*, Japan Sea, B:259

*Rouxia californica* Zone, Leg 127, B:189

*Saccammina sphaerica*, Site 795, B:537

*sadonica*, *Valvulineria*, Site 798, B:390

*Sagarites makiyama*, Site 799, B:35, 45

*salisburyi*, *Pullenia*

salisburyi, *Pullenia* (cont.)

## TAXONOMIC INDEX

Site 797, B:540  
 Site 798, B:392  
*Schizammina?* sp.  
 Site 795, B:537  
 Site 797, B:537  
*Sciadopitys*, Site 795, B:490  
*Selaginella selaginoides*, Site 794, B:334  
*selaginoides*, *Selaginella*, Site 794, B:334  
*sellii*, *Helicosphaera*  
 Leg 128, B:164  
 Site 798, B:164  
*Sethocytis japonica*  
 Japan Sea, B:297, 301  
 Site 794, B:305  
*Silicospioolina abyssalica*, Japan Sea, B:505  
*slavnicii*, *Distephanus*, Site 798, B:248  
 sp. A, *Cassidulina*, Site 798, B:390  
 sp. A, *Cibicidoides*, Site 797, B:538  
 sp. A, *Cyclammina*, Site 797, B:538  
 sp. A, *Globocassidulina*, Site 797, B:539  
 sp. A, *Pleurostomella*, Site 799, B:390  
*speculum*, *Distephanus*, Site 799, B:248  
*sphaerica*, *Saccammina*, Site 795, B:537  
*Sphaeroidina bulloides*, Site 798, B:390  
*Sphaeroidina japonica*, Site 797, B:538  
*Sphaeropyle langii*  
 Japan Sea, B:297  
 Site 794, B:305  
 Site 795, B:303  
 Site 799, B:236  
*Sphaeropyle robusta*  
 Japan Sea, B:297  
 Site 794, B:1221  
 Site 797, B:305  
*Sphenolithus compactus*, Japan Sea, B:183  
*Sphenolithus heteromorphus*  
 Japan Sea, B:173–174, 183  
 Leg 127, B:189  
*Sphenolithus heteromorphus* Zone  
 Japan Sea, B:174  
 Leg 127, B:201  
 Site 797, B:1223  
*Sphenolithus umbrellus*, Japan Sea, B:183  
*Spirosigmoilinella compressa*  
 Site 795, B:537  
 Site 798, B:379–381, 389  
*Spirosigmoilinella compressa/Martinottiella communis* Zone  
 Site 794, B:517, 528  
 Site 798, B:381  
*Spirosignoilinella compressa* Zone, Leg 127,  
 B:200  
*spissa*, *Bolivina*, Site 798, B:389  
*Spongaster tetras irregularis*  
 Japan Sea, B:231  
 Site 798, B:236  
*Spongotrochus glacialis*, Site 795, B:306  
*Stainforthia complanata*  
 Japan Sea, B:388  
 Site 798, B:390  
*Stainforthia fusiformis*  
 paleoenvironment and, B:379  
 Site 798, B:390  
*stella*, *Nonionella*  
 Site 797, B:539  
 Site 799, B:391  
*Stichocorys delmontensis*

Japan Sea, B:228, 230  
 Site 799, B:225–226, 236  
*Stichocorys peregrina*  
 Japan Sea, B:297  
 Site 794, A(127):102  
 Site 797, A(127):356  
 Site 799, B:236  
*Stichocorys wolfii*, Site 794, A(127):102  
*Stilostomella lepidula*, Site 798, B:390  
*striata*, *Bulimina*  
 Site 797, B:539  
 Site 798, B:390  
*Stylacontarium acquilonius*  
 Japan Sea, B:297  
 Site 794, B:305  
*Stylacontarium* sp. cf. *S. acquilonius*, Japan Sea,  
 B:297  
*Stylochlamydium venustum*, Site 797, B:306  
*subaffinis*, *Globobulimina*, Site 798, B:390  
*subarcticum*, *Elphidium*  
 Site 797, B:540  
 Site 798, B:392  
*subarcuatus*, *Dictyochea*  
 Japan Sea, B:238–239  
 Site 798, B:248  
*subcarinata*, *Pullenia*, Site 797, B:540  
*subopercularis*, *Glabratella*, Site 797, B:540  
*substriata*, *Lagena*, Japan Sea, B:387  
*takayanagii*, *Pseudoparella*, Site 798, B:391  
*tatsunokuchiensis*, *Rossiella*, Japan Sea, B:258–  
 259  
*temporei*, *Thalassiosira*, Japan Sea, B:259  
*tener*, *Oridorsalis*, Site 798, B:392  
*tenuata*, *Buliminella*, Site 798, B:390  
*tenuis*, *Ammodiscus*, Site 797, B:537  
*tenuis*, *Cassidulinoides*, Site 799, B:390  
*Tenuitella parkerae*, Leg 127, B:200  
*tetrapera*, *Cyrtocapsella*  
 Japan Sea, B:230  
 Site 794, B:304  
 Site 797, A(127):356; B:304  
 Site 799, B:235  
*Tetrapyle octacantha*  
 Site 794, B:306  
 Site 797, B:306  
 Site 798, B:236  
*tetrathalamus*, *Didymocystis*  
 Site 797, B:306  
 Site 798, B:235  
*Thalassinoides*, Site 798, B:566–567, 569  
*Thalassionema nitzschiooides*, Site 794, B:1223  
*Thalassionema schraderii* Zone, Site 794, B:1221  
*Thalassiosira antiqua*, Japan Sea, B:258  
*Thalassiosira convexa*, Japan Sea, B:258  
*Thalassiosira jacksonii*, Japan Sea, B:259  
*Thalassiosira lentiginosa*, Southern Ocean, B:309  
*Thalassiosira marujamica*, Japan Sea, B:259  
*Thalassiosira miocenica*, B:1219  
*Thalassiosira nidulus*, Japan Sea, B:258  
*Thalassiosira oestrupii* Zone  
 Leg 127, B:188  
 Site 794, B:344, 1221  
 Site 797, B:1223

*Thalassiosira temporei*, Japan Sea, B:259  
*Thalassiosira zabelinae*, Japan Sea, B:259  
*Thalassiothrix nitzschiooides*, Site 798, A(128):160  
*Thalmannammina* cf. *parkeriae*, Site 797, B:537  
*Thecosphaera japonica*, Site 795, B:305  
*Theocalyptra davisi*  
 Japan Sea, B:297, 299  
 Site 794, B:306  
 Site 795, B:303  
*Theocorys redondoensis*  
 Japan Sea, B:297  
 Site 794, B:305  
*Tholospyris mammilaris*, Site 794, B:305  
*Tilia*, Site 797, B:491  
*titanni*, *Makiyama*, Site 799, B:35, 45  
*tochigensis*, *Lithatractus*, Site 794, B:305  
*Tosaia hanzawai*, Site 797, B:538  
*tricarinata*, *Triloculina*, Site 798, B:389  
*tricaritortuosa*, *Lagena*, Site 798, B:389  
*Trifarina* sp., Site 797, B:539  
*Triloculina tricarinata*, Site 798, B:389  
*Trochammina* cf. *globigeriniformis*, Site 795,  
 B:537  
*Tsuga*  
 Japan Sea, B:328  
 Neogene climate and, B:337  
*Ulmus*, Site 795, B:491  
*Umbilicosphaera sibogae* Zone, Japan Sea, B:174  
*umbonatus*, *Oridorsalis*  
 Site 797, B:540  
 Site 798, B:392  
*umbrellus*, *Sphenolithus*, Japan Sea, B:183  
*Uvigerina akitensis*, Site 798, B:390, 441, 443–  
 444  
*Uvigerina* cf. *modeloensis*, Site 797, B:539  
*Uvigerina lobatula*, Site 798, B:390  
*Uvigerina miocenica*, Site 797, B:539  
*Uvigerina proboscidea*, Site 797, B:539  
*Uvigerina segundoensis* s.l. Zone, Japan Sea  
 B:526  
*Uvigerina yabei*, Site 798, B:390  
*Valvularia masudai*, Site 797, B:538  
*Valvularia sadonica*, Site 798, B:390  
*variabilis*, *Hippocrepinella*, Site 795, B:537  
*venezuelana*, *Globoquadrina*  
 Leg 127, B:189  
 Site 797, B:1220  
*wolfii*, *Stichocorys*, Site 794, A(127):102  
*wuellerstorfi*, *Cibicidoides*  
 Japan Sea, B:505  
 Site 797, B:538  
*yabei*, *Uvigerina*, Site 798, B:390  
*ypsilony*, *Amphirhopalum*, Site 798, B:235  
*zabelinae*, *Thalassiosira*, Japan Sea, B:259  
*Zelkoova*, Site 797, B:491  
*zones* (with letter prefixes)  
 NP-1, Site 797, B:1221, 1223  
 NP-2, Site 797, B:1221, 1223  
 NP-3, Site 797, B:1223  
*Zoophycos*, Japan Sea, B:1232