

INDEX TO VOLUMES 137 AND 140

This index provides coverage for the *Initial Reports* portions of Volumes 137 and 140 of the *Proceedings of the Ocean Drilling Program* (published separately) and for the *Scientific Results* portion of those *Proceedings* (published as Volume 137/140; this book). The reports in this book (SR Vol. 137/140) are not leg-specific but are arranged by subject matter. References to page numbers in the *Initial Reports* are preceded by "A," followed by the appropriate leg reference in parentheses, and then a colon: "A(137):"; those in the *Scientific Results* are preceded by "B" with a colon.

The index was prepared by tgB, Inc., under subcontract to the Ocean Drilling Program. The index contains two hierarchies of entries: (1) a main entry, defined as a keyword or concept followed by a reference to the page on which that word or concept appears, and (2) a subentry, defined as a further elaboration on the main entry followed by a page reference.

Usually the index is presented in two parts, a subject index and a taxonomic index. As the sheeted-dike section of Hole 504B obviously is nonfossiliferous, the index section is confined to a Subject Index. The index covers text figures and tables but not core-description forms ("barrel sheets") or core photographs. Also excluded are bibliographic references, names of individuals, and routine front and back matter.

The Subject Index follows a standard format. Geographic, geologic, and other terms are referenced only if they are subjects of discussion. A site chapter in the *Initial Reports* is considered the principal reference for that site and is indicated on the first line of the site's listing in the index. A parenthetical reference is made to the appropriate leg; the reference to Site 504, for example, is given as "Site 504, A(137):15-55; A(140):37-200."

For further information, including available electronic formats, contact the Chief Production Editor, Ocean Drilling Program, 1000 Discovery Drive, College Station, Texas 77845-9547, U.S.A.

SUBJECT INDEX

- abrasion loss of weight
hardness, B:347–348
vs. Shore hardness, B:349
- acoustic velocity logging, A(140):113
- actinolite
alteration, A(137):28–29; B:101, 195
composition, B:238
hydrothermal alteration, B:157
replacement silicates, B:210–213
veins, A(140):73–75, 90–91
vs. depth, A(140):66; B:160
See also ferroactinolite
- albite
secondary minerals, B:172, 178–179
veins, A(140):76
vs. depth, A(140):66
- albite percentage, vs. oxygen isotopes, B:103
- alkalinity, vs. depth, A(137):37; B:145
- alteration
basalts, B:43–51
basement, A(137):28–29
borehole fluids, B:141–152
deformation, B:232
dikes, B:275, 278
magnetic properties, B:329
Site 504, A(137):28–29; A(140):64–78, 121–123
strain localization, B:226, 228
veins, B:239–240
vs. depth, A(140):66, 72
vs. silicates, B:67
See also deuteritic alteration; hydrothermal alteration
- alteration halos
hydrothermal alteration, B:158
petrology, A(140):75–76
veins, B:179, 182
- alteration logs, A(140):156–162
- alteration patches
diabase, A(140):70–72
vs. depth, A(140):70
- alteration percentage
vs. barium, B:110
vs. bulk density, B:102
vs. calcium, B:110
vs. chromium, B:109
vs. cobalt, B:109
vs. lanthanum/samarium ratio, B:110
vs. lanthanum/yttrium ratio, B:110
vs. lanthanum, B:110
vs. major-element oxides, B:70–71
vs. nickel, B:109
vs. oxygen isotopes, B:103
vs. samarium, B:110
vs. strontium, B:111
vs. water, B:102
- aluminum, vs. depth, B:173
- aluminum-free minerals, hydrothermal evolution, B:211–213
- aluminum-rich minerals, hydrothermal evolution, B:211–213
- aluminum/titanium ratio, vs. sodium/titanium ratio, B:56
- aluminum logs, vs. depth, B:309
- aluminum oxide
estimate from geochemical logs, B:345–346
percent change from protolith, B:203
pyroxene, B:4
vs. depth, B:91
- vs. magnesium oxide, B:45
vs. temperature, B:135
vs. titanium oxide, B:37
See also calcium oxide/aluminum oxide ratio
- ammonia, vs. magnesium, B:146
- amphibole composition, vs. depth, B:160
- amphibole crystals, veins, A(140):110
- amphibole percentage, vs. oxygen isotopes, B:103
- amphibole veins
deformation, A(140):112
Site 504, B:236
- amphiboles
chemical composition, B:237
geochemistry, B:67–72
geothermometry, B:178
microprobe data, B:210–213
secondary minerals, B:159–160, 169, 172, 174–177
vs. plagioclase, B:179
See also individual minerals
- amygdules, vs. depth, A(140):70
- anatase
deuteritic oxidation, B:332
magnetic minerals, B:257–259
- anhydrite
precipitation, B:144–146, 163–165
saturation vs. depth, B:147
secondary minerals, B:176
veins, A(140):74
- anhydrite crusts, photograph, A(137):39
- anisotropy, magnetic susceptibility, B:249–251
- anorthite, vs. magnesium oxide, B:173
- anorthite percentage, vs. magnesium number, B:9, 39
- anorthosite, crystal clots, A(140):59
- apatite, secondary minerals, B:176
- assimilation, altered oceanic crust, B:43–51
- augite
diabase, B:3–9
formation temperature, B:177–178
phenocrysts, A(140):54–55
- augite, chromian, phenocrysts, A(140):54–55
- backscattered electron imaging, microstructures, B:220–228
- barium
diabase, B:108
lithologic unit 220, B:86
vs. alteration percentage, B:110
vs. chromium, B:111
vs. depth, B:145
vs. magnesium, B:146
- basalts
alteration, B:146–147
geochemistry, B:53–61, 81–97, 117–120, 162
magnetization, B:253–262
mechanical behavior, B:347–351
petrology, A(137):24, 26–27; A(140):52–64, 119–121; B:19–33
physical properties, B:295
- basalts, altered, petrology, B:43–51
- basalts, aphyric, chemical composition, A(140):86–88
- basalts, phyric, cracks, B:224
- basement
seismic studies, A(140):42, 44
stratigraphy, A(137):24, 26
- Bay of Islands massifs, ophiolite, B:283, 287–289
- BHTV. *See* borehole televiewer
- Blow Me Down massif, ophiolite, B:283, 287–289
- borehole fluids, chemistry, A(137):35–37; B:141–152
- borehole televiewer
data analysis, B:293–304
Site 504, A(137):47–50; A(140):113
- boreholes, cross sections, B:297
- boron
reaction zones, B:147–150
vs. depth, B:145
vs. magnesium, B:146
- boundary layers, penetration, A(140):127–128
- breakouts
histograms, B:298
vs. depth, B:299
- breakouts, borehole, stresses, B:296–303
- brecciation
dike contacts, A(140):99
veins, B:234
zoning, A(140):102
- breccia zones, veins, B:239
- brittle deformation
diabase dikes, A(140):123–124
veins, A(140):95; B:239
- bromine, vs. magnesium, B:146
- brunsvigite
microprobe data, B:210–211
molar composition, B:176
- bulk modulus, vs. depth, B:283
- calcium/(calcium + sodium) ratio, melt inclusions
vs. host plagioclase, B:135
- calcium
vs. alteration percentage, B:110
vs. depth, A(137):37; B:145
vs. magnesium, A(137):43; B:146, 151
vs. magnesium number, B:9
vs. sulfate, A(137):43
- calcium oxide
estimate from geochemical logs, B:345–346
vs. alteration percentage, B:70
vs. depth, A(140):88; B:68
vs. magnesium oxide, B:45
vs. water, A(140):90
- calcium oxide/aluminum oxide ratio, vs. magnesium oxide, A(140):92
- calcium oxide/sodium oxide ratio, vs. titanium oxide, A(140):87
- caliper logs, vs. depth, B:316
- carbon dioxide, fluid inclusions, B:194, 197
- chalcopyrite, secondary minerals, A(140):69–70
- CHAOS, computer models, B:54–55
- chilled margins, dikes, 94–99
- chlorine, vs. (sodium + potassium), B:160
- chlorinity
vs. depth, A(137):37; B:145
vs. magnesium, B:146
- chlorite
alteration, A(137):28–29
chemical composition, B:149
geochemistry, B:67–72
geothermometry, B:178–179
hydrothermal alteration, B:157
microprobe data, B:210–211
secondary minerals, B:172–173, 180–181
vein density, A(140):74
vein structure, A(140):90–91
veins, A(140):74
vs. depth, A(140):66

chlorite (cont.)

SUBJECT INDEX

- vs. temperature, B:179
See also brunsvigite; chlorite-smectite;
 diabantite; penninite; picnochlorite;
 talc-chlorite
- chlorite-amphibole veins, deformation,
 A(140):113
- chlorite-smectite
 geothermometry, B:178-179
 hydrothermal alteration, B:157
 microprobe data, B:210-211
 secondary minerals, B:172-173, 180-181
- chlorite/smectite ratio, composition, B:149
- chromium
 diabase, A(140):152; B:108
 percent change from protolith, B:203
 vs. alteration percentage, B:109
 vs. barium, B:111
 vs. lanthanum/samarium ratio, B:115
 vs. lanthanum/ytterbium ratio, B:115
 vs. magnesium number, B:9
 vs. nickel, A(140):92
 vs. strontium, B:111
- chromium oxide
 vs. magnesium number, B:37, 126
 vs. wollastonite, B:169
- clay minerals, mixed-layer, vs. depth, A(140):66
- clays, chemical composition, B:149
- clinopyroxene
 basalts, A(137):24, 26-27
 chemical composition, B:238
 geochemistry, B:67-72, 121-130
 hydrothermal alteration, B:157
 petrography, B:36
 secondary minerals, A(140):69; B:160,
 169-171
 veins, B:236
- cobalt
 diabase, B:108
 percent change from protolith, B:203
 vs. alteration percentage, B:109
- Cocos-Nazca Rift, seismic studies, A(140):44
- Cocos-Nazca Spreading Center, tectonics,
 A(140):41
- coercivity, remanent magnetization, B:266
- coercivity, bulk, vs. depth, B:258
- coercivity/bulk coercivity ratio, vs. depth, B:258
- compressional ultrasonic velocity
 vs. hardness, B:349
 vs. uniaxial compressive strength, B:350
- compressional wave velocity
 cores, B:276-278, 280-283
 Site 504, A(140):108-109, 134
 vs. density, B:280-281
 vs. depth, A(137):32; B:281
 vs. porosity, B:278-279
 vs. shear wave velocity, B:282
- compressional wave velocity/shear wave velocity
 ratio
 Site 504, B:281-283
 vs. depth, B:283
- compressional wave velocity, dry, vs. wet
 compressional wave velocity,
 A(140):136
- compressional wave velocity, wet, vs. dry
 compressional wave velocity,
 A(140):136
- compressive strength, uniaxial, basalt, B:348
- computer models, geochemistry, B:53-61
- convection, heat transfer, B:323
- cooling
 dikes, B:22-23, 25-27, 29-31
 strain localization, B:226, 228
- copper/nickel ratio, vs. magnesium number, B:203
- copper
- depletion, B:204
- diabase, B:200
 percent change from protolith, B:203
 vs. alteration percentage, B:71
 vs. depth, B:68, 92, 164
 vs. sulfur, B:163
 vs. titanium oxide, B:92
 vs. water, A(140):89
 vs. zirconium, B:92
 vs. zirconium/yttrium ratio, B:93
- correlation coefficient patterns, basalts, B:57-59
- Costa Rica Rift
 basalts, B:347-351
 regional geology, A(140):43
 site description, A(137):15-55;
 A(140):37-200
 stratigraphy, B:107-108
 tectonics, A(140):41-44; B:231-232,
 245-247, 298
- cracks
 alteration, A(137):28-29
 augite, B:221
 fluid circulation, B:240
 porosity, B:278
See also en-echelon cracks; microcracks;
 tension gashes; thermal cracks
- cracks, shrinkage, veins, B:213
- crosscutting veins
 deformation, B:234
 distribution, A(140):96-97, 114-116
- crust. *See* oceanic crust
- crustal Layer 2/3, penetration, A(140):127
- crustal Layer 2, fluid evolution, B:191-198
- crustal Layer 2A, fluid flow, B:313-319
- crustal Layer 2C, partial melts, B:47-50
- crystal clots
 clinopyroxene, B:122
 petrology, A(140):55-58
- crystal fracturing, igneous rocks, A(140):84
- crystal growth, clinopyroxene, B:127-129
- crystallization
 diabase, A(140):62-63; B:220, 223
 multiple dikes, B:22-23, 25-27, 29-31
 silicates, B:9-12
 vs. time, B:28-31
- crystallization, post-entrapment, plagioclase,
 B:134-135
- crystallization time, vs. grain size, B:29-31
- Curie temperature
 magnetite, B:257-259
 vs. depth, B:259
- DARWIN, computer models, B:58
- deformation
 alteration, B:232
See also brittle deformation; ductile
 deformation
- demagnetization, thermal, dikes, B:264-266
- density
 dikes, B:275, 278-280
 vs. compressional wave velocity, B:280-281
 vs. depth, B:289
 vs. porosity, B:281
 vs. velocity, A(140):136
- density, bulk
 basalts, B:348
 vs. alteration percentage, B:102
 vs. depth, A(137):31; B:279
 vs. porosity, A(140):126, 128; B:102
 vs. shear wave velocity, A(137):33
- density, dry-bulk
 vs. grain density, A(137):30
 vs. hardness, B:349
 vs. uniaxial compressive strength, B:350
- vs. wet bulk density, A(137):30
- density, grain
 vs. depth, A(137):31
 vs. dry-bulk density, A(137):30
 vs. magnetic susceptibility, A(137):34
- density, wet-bulk, vs. dry-bulk density, A(137):30
- deuteric alteration, titanomagnetite, B:259
- deuterium/hydrogen ratio
 diabase, B:101
 reaction zones, B:147-150
 vs. water content, B:102
- deweylite, microprobe data, B:208-209
- diabantite
 chemical composition, B:149
 molar composition, B:176
- diabase
 chemical composition, A(140):86-87
 chilled dike margins, B:35-42
 clinopyroxene, B:121-130
 comparison to ophiolites, A(140):127
 dikes, B:48, 131-139, 155-166
 fluid inclusions, B:191-194
 geochemistry, B:53-61, 65-97, 107-110,
 117-120
 geothermometry, B:167-189
 hydrothermal alteration, B:199-205, 207-216
 petrology, A(140):52-64, 119-121; B:3-17,
 19-33
 stable magnetic remanence, B:245-252
 strain localization, B:219-229
 veins, B:231-234
- diabase, fresh, vs. altered samples, B:93
- diatoms, borehole fluids, A(137):38-39
- differentiation, magmas, B:59-60
- dike injection
 remanent magnetization, B:267-269
 veins, B:213
- dikelets, igneous rocks, A(140):84-86
- dike margins
 chilled contacts, A(140):94
 petrology, B:35-42
 reorientation, B:247
- dike margins, chilled
 structure, A(140):82
 vs. depth, B:36
- dikes
 alteration, A(137):24, 26-27
 altered basalts, B:43-51
 clinopyroxene, B:121-130
 cores, B:249
 diabase, A(140):52-64, 119-121
 geochemistry, B:107-110, 117-120
 logs, B:308-311
 melt inclusions, B:131-139
 Site 504, B:19-33
 strain localization, B:219-229
 width frequency, B:20
See also dikelets; microdikes
- dikes, sheeted
 boundary layers, A(140):127-128
 cores and logs, B:305-311
 fluid inclusions, B:191-198
 geochemistry, B:65-97, 99-106
 geothermometry, B:167-189
 hydrothermal alteration, B:199-205, 207-216
 magnetic properties, B:327-337
 mineralogy, B:155-166
 natural remanent magnetization, B:263-270
 petrology, A(140):119-121; B:19-33
 rock magnetism, B:253-262
 stable magnetic remanence, B:245-252
 stresses, B:293-304
 veins, B:231-243
- dike width, vs. crystal length, B:23

- diopside
secondary minerals, B:188
veins, A(140):76
- dip
epidote veins, A(140):107
fractures, A(140):99, 102, 116, 143
veins, A(140):86–88, 90–91
- dip angle, vs. number of planes in borehole, A(140):106
- dolerite. *See* diabase
- downhole measurements, A(137):24; A(140):109, 112–116, 125–127
- drilling history, A(137):18–21; A(140):49
- ductile deformation, veins, A(140):92, 94
- dunite, crystal clots, A(140):57–58
- East Pacific Rise, tectonics, A(140):41
- Ecuador Fracture Zone, tectonics, A(140):41
- Ecuador Rift, tectonics, A(140):41
- elastic constants
calculated from velocities and densities, B:284–287
upper crustal rocks, B:273–291
- elasticity, basalt, B:348–349
- electrical images, logging, A(140):112, 116
- element mobility, calculation, B:202–204
- elongate crystals, plagioclase, B:7
- en-echelon cracks, veins, B:233–234
- epidote
dip of veins, A(140):107
secondary minerals, B:160, 173–174, 184–185
veins, A(140):76
vs. depth, A(140):66
- europium
pillow basalts, B:129
vs. strontium, B:134
- fabric, cores, A(140):83–84
- faults. *See* normal faults
- feldspars, secondary minerals, B:160
- ferroactinolite, replacement silicates, B:210–213
- flowmeter studies, A(137):50–51; A(140):114
- flow texture, cores, A(140):83–84
- fluid chemistry, B:141–152
- fluid circulation, veins, B:240
- fluid evolution, alteration, B:195–197
- fluid flow, B:313–319
- fluid inclusions
oceanic crustal Layer 2, B:191–198
See also melt inclusions
- fluid inclusions, liquid-dominated, daughter-mineral-bearing, microthermometry, B:194
- fluid inclusions, liquid-dominated, low-salinity, microthermometry, B:193–194
- fluid inclusions, liquid-dominated, vapor-bubble-absent, microthermometry, B:194
- fluid samplers, design, A(137):39–41
- fluid sampling, A(137):31–42
- foraminifers, borehole fluids, A(137):38–39
- formation microscanner imagery, A(140):112, 116
- formation water, chemistry, B:141–152
- fractional distillation, brines, B:196
- fractionation, basalts, B:59–60
- fracture density
scanning electron microscopy, B:223–226
vs. depth, A(140):118
- fracture porosity, vs. depth, A(140):140–141
- fractures
dikes, B:274–275
formation microscanner imagery, A(140):120
morphology, A(140):117
porosity, B:287
propagation, B:237
reorientation, B:247
Site 504, B:313–319
stereograms, A(140):142
stereographic projections, A(140):119
vs. depth, B:316
fractures, open, sheeted dikes, A(140):97–99, 102
- gabbros
crystal clots, A(140):57–58
residual models, B:60
- gadolinium, estimate from geochemical logs, B:345–346
- Galapagos Gore, tectonics, A(140):41
- Galapagos Hot Spot, tectonics, A(140):41
- Galapagos Rift, tectonics, A(140):41
- Galapagos Spreading Center. *See* Cocos–Nazca Spreading Center
- Galapagos Spreading Segment, tectonics, A(140):41
- Galapagos Triple Junction, tectonics, A(140):41
- geochemical logs, A(140):113–114, 192–197; B:307–308, 339–346
- geochemistry
chilled margins, B:38–42
computer models, B:53–61
oceanic crust, B:327–337
primary and secondary variations, B:65–80
reference sample, B:353–355
sheeted dike complex, B:81–97, 167–189
Site 504, A(137):27–28; A(140):78–82, 121
- geothermal gradient
Site 504, A(137):44–47; A(140):114–116
See also temperature gradient
- geothermal state, oceanic crust, B:323
- geothermometry, sheeted dike complex, B:167–189
- glomerocrysts. *See* crystal clots
- goethite, borehole fluids, A(137):38–39
- gold
diabase, B:117–120
melting calculations, B:204
vs. depth, B:119
- grain boundaries, photograph, B:225
- grain size
diabase, B:220, 223
downhole variations, B:19–33
silicates, B:4
vs. depth, B:22–23
- greenschist facies, alteration, B:279
- Gresen's calculation, mass balance, B:200–202
- groundmass, diabase, A(140):54–55
- Group D composition, B:94
- hafnium, percent change from protolith, B:203
- hardness
basalt, B:347–348
vs. abrasion loss of weight, B:349
vs. compressional ultrasonic velocity, B:349
vs. dry-bulk density, B:349
vs. uniaxial compressive strength, B:350
- heat flow
oceanic crust, B:321–324
Site 504, A(137):46–47
vs. depth, B:323
- heat transfer
convection, B:323
Site 504, A(137):46–47
- heulandite, secondary minerals, B:176, 184–186
- homogenization temperature, vs. depth, B:194–195
- hornblende, actinolitic
composition, B:238
hydrothermal alteration, B:157
vs. depth, B:160
- hornblende, magnesian
replacement silicates, B:210–213
vs. depth, B:160
- hydration, brines, B:196
- hydrofracturing, stresses, B:251–252
- hydrogen isotopes. *See* deuterium/hydrogen ratio
- hydrology, heat flow, B:323–324
- hydrothermal alteration
diabase, B:99–106
geochemistry, B:191–205
geothermometry, B:167–189
magnetic anomalies, B:257–259
sheeted dike complex, B:155–166
- hydrothermal fluids
origin, B:182
stresses, B:251–252
- hydrothermal pulses, fluid inclusions, B:197
- igneous contacts, logs, A(140):150–151
- igneous mineralogy, logs, A(140):146–147
- igneous petrology, A(137):24, 26–27; A(140):52–64, 119–121
- igneous rocks
classification, A(140):52
mineralogy, B:3–17
- ilmenite
deuteric oxidation, B:262, 332
exsolution, B:267
- in-situ reactions, borehole fluids, B:141–152
- inclusions
hydrothermal alteration, B:157–158
See also fluid inclusions; melt inclusions
- index properties
diabase, A(140):133
Site 504, A(137):29; A(140):106, 108
- induced magnetization, basalts, A(140):103–104
- injection, dikes, B:22–23
- injection studies, A(137):50–51
- intrusions, dikes, B:22–23, 25–27
- intrusive contacts, igneous rocks, A(140):84–86
- iridium, melting calculations, B:204
- iron
vs. depth, B:145, 164
vs. magnesium, B:146
- iron/(iron + magnesium) ratio, vs. silicon, B:149, 159, 210
- iron + magnesium/titanium ratio, vs. magnesium/titanium ratio, B:57
- iron logs, vs. depth, B:309
- iron oxide
estimate from geochemical logs, B:345–346
percent change from protolith, B:203
vs. depth, B:91, 176
vs. magnesium oxide, B:45
- isotopes
reference sample, B:353, 355
See also stable isotopes
- Koenigsberger ratio, vs. depth, A(140):124; B:254–256
- lanthanum/samarium ratio
vs. alteration percentage, B:110
vs. chromium, B:115
vs. titanium/zirconium ratio, B:137
vs. ytterbium, B:50
vs. zirconium/yttrium ratio, B:115
- lanthanum/ytterbium ratio
vs. alteration percentage, B:110
vs. chromium, B:115
vs. nickel, B:115
vs. zirconium/ytterbium ratio, B:203
vs. zirconium/yttrium ratio, B:115

lanthanum

SUBJECT INDEX

- lanthanum
 percent change from protolith, B:203
 vs. alteration percentage, B:110
- laterolog
 harmonic analysis, B:310–311
 vs. depth, B:279, 307–310
- laterolog resistivity, B:306–307
- laumontite
 alteration, A(137):29
 secondary minerals, B:176, 184–185
 veins, B:161
- Lawrence Berkeley Laboratory (LBL)
 high-temperature borehole fluid
 sampler, design, A(137):40–41
- least-squares-fit mixing models, basalts, B:57
- lithium
 reaction zones, B:147–150
 vs. depth, B:145
 vs. magnesium, B:146
- lithologic Unit 211, clinopyroxene, B:122–124
- lithologic Unit 227, clinopyroxene, B:123
- lithologic logs, A(140):148–149
- lithophile elements, diabase, B:107–110
- lithostratigraphy, A(137):18, 24, 26–27;
 B:313–319
- Los Alamos National Laboratory (LANL)/Leutert
 borehole samplers, design, A(137):39–40
- magma chambers, petrology, B:49–51
- magmas
 chemistry, B:43–51
 differentiation, B:59–60
 evolution, A(140):42
 rare-earth budget, B:109–110
 segregation, B:136–137
See also primary magmas
- magmatic fluids, brines, B:196
- magnesium
 vs. calcium, A(137):43
 vs. components of borehole fluids, B:146
 vs. depth, A(137):37, 42; B:145
 vs. element mixing lines, B:151
 vs. potassium, A(137):43
 vs. silica, A(137):43
 vs. sodium, A(137):43
 vs. sulfate, A(137):43
See also iron/(iron + magnesium) ratio; (iron +
 magnesium)/titanium ratio
- magnesium/titanium ratio, vs. (iron +
 magnesium)/titanium ratio, B:57
- magnesium number
 basalts, B:54
 vs. anorthite percentage, B:9, 39
 vs. calcium, B:9
 vs. chromium, B:9
 vs. chromium oxide, B:37
 vs. copper/nickel ratio, B:203
 vs. depth, A(140):88
 vs. manganese, B:9
 vs. oxides and ytterbium, B:126
 vs. silicon, B:238
 vs. titanium oxide, A(140):86–87; B:88
 vs. water, A(140):90
- magnesium oxide
 percent change from protolith, B:203
 vs. alteration percentage, B:70
 vs. aluminum oxide, B:45
 vs. anorthite, B:173
 vs. calcium oxide/aluminum oxide ratio,
 A(140):92
 vs. calcium oxide, B:45
 vs. depth, B:91
 vs. iron oxide, B:45
 vs. phosphorus oxide, B:45
- vs. potassium oxide, B:45
 vs. silica, B:45, 54
 vs. sodium oxide, B:45
 vs. strontium, B:48
 vs. titanium oxide, B:45
 vs. trace elements, B:48
 vs. vanadium, B:48
 vs. yttrium, B:48
 vs. zirconium, B:48
- magnetic anomalies, rock magnetism, B:260
- magnetic fabric
 dikes, A(140):105
 structures, B:245–252
- magnetic hardness, remanent magnetization,
 B:269
- magnetic hysteresis, loops, B:331
- magnetic inclination
 vs. depth, A(140):123; B:255, 267
 vs. lithology, B:267–268
- magnetic inclination, stable, frequency, B:260,
 267
- magnetic mineralogy
 oceanic crust, B:327–337
 sheeted dikes, B:253–262
 Site 504, B:263–264
- magnetic properties, minicores, A(140):121
- magnetic reversals, dikes, A(140):105–106
- magnetic susceptibility
 anisotropy, B:249–251
 basalts, A(140):103–104
 cores, A(137):30–31
 vs. depth, A(140):124; B:254–256
 vs. grain density, A(137):34
- magnetite, length of
 average maximum length and equivalent
 diameter, A(140):154–155
 vs. depth, A(140):61–62
 vs. length of plagioclase, A(140):61
- magnetite
 alteration, A(137):28–29
 deuteric oxidation, B:332
 exsolution, B:267
 grain size, A(140):59–61; B:20–22
 hydrothermal alteration, B:158
 thermomagnetic data, B:257–259
- magnetometer logs, vs. depth, A(140):143
- major elements
 basalts, A(137):27–28; B:53–61
 chilled margins, B:38–40, 42
 clinopyroxene, B:123, 125
 diabase, B:114
 igneous rocks, A(140):78–82, 121
 melt inclusions, B:135
 oceanic crust, B:327–337
 partial melts, B:47
 primary and secondary variations, B:65–80
 reference sample, B:353–354
- manganese
 vs. depth, B:145
 vs. magnesium, B:146
 vs. magnesium number, B:9
- manganese oxide, vs. depth, A(140):90; B:92, 164
- mantle, peridotite, B:137–138
- mass balance. *See* Gresen's calculation
- median demagnetizing field
 basalts, A(140):102–103
 vs. depth, A(140):123; B:254–256
- megacrysts, clinopyroxene, B:121–130
- melanite, formation temperature, B:177
- melt inclusions
 plagioclase–olivine–quartz projection, B:135
 plagioclase hosts, B:133
 refractory silicates, B:131–139
See also fluid inclusions
- melting
 altered basalts, B:44, 46–47
 magmas, B:136–138
- metabasalt, fluid inclusions, B:196
- metamorphism
 rare-earth budget, B:109–110
 Site 504, A(140):64–78, 121–123
- metasomatism, calculation, B:202–204
- methane, clathrates, B:194, 197
- microcracks, diabase, B:220–223
- microdikes
 amphibole veins, A(140):105
See also dikes
- microphenocrysts
 plagioclase, B:6
 pyroxene, B:14
- microstructural controls
 fabric, B:245–252
 strain localization, B:219–229
 veins, A(140):91–92, 95–96; B:236–239
- microstructures, diabase, B:220, 223
- microthermometry, high-temperature, melt
 inclusions, B:131–132
- mid-ocean ridges
 melting, B:136–138
 rifting, B:19–33
- Middle American Trench, tectonics, A(140):41
- mineral chemistry, diabase, B:35–38
- mineral clasts, veins, A(140):111
- mineralogy, sheeted dike complex, B:155–166
- minor elements, reference sample, B:353–354
- mixed-layer chlorite. *See* chlorite–smectite
- natural gamma-ray logs, A(140):113–114,
 198–200; B:343–344
- Nazca Plate, tectonics, B:298
- Newfoundland, ophiolite, B:283, 287–289
- nickel
 diabase, B:108, 200
 percent change from protolith, B:203
 vs. alteration percentage, B:109
 vs. chromium, A(140):92
 vs. depth, A(140):89; B:91
 vs. lanthanum/ytterbium ratio, B:115
 vs. zirconium, A(140):92; B:89
See also copper/nickel ratio
- niobium
 vs. depth, B:90
 vs. zirconium, A(137):28; A(140):86; B:90
- normal faults, stresses, B:251
- North Arm massif, ophiolite, B:283, 287–289
- nucleation rates, diabase, B:3–4
- oceanic crust
 alteration, B:182
 drilling, A(137):51–54
 hydrothermal exchange with seawater,
 B:103–104
 sheeted dike complex, B:199–205
 strain localization, B:219–229
- oceanic crust, altered, assimilation, B:43–51
- oceanic crust, upper
 fluid flow, B:313–319
 heat flow, B:321–324
 magnetic properties, B:327–337
 physical properties, B:273–291
- oikocrysts, basalts, A(140):56–58
- oligoclase, secondary minerals, B:172, 178–179
- olivine
 alteration, A(140):67–68, 168–170
 basalts, A(137):24, 26–27
 diabase, B:3–9
 hydrothermal alteration, B:207–216
 petrography, B:35–36

- phenocrysts, A(140):54
 secondary minerals, A(140):64–65, 67–69
- Oman, ophiolite, B:283, 287–289
- ophiolite, physical properties, B:283, 287
- ophitic texture
 pyroxene, B:15
 silicates, B:3–4, 7
- orientation, thin section data, A(140):100–101
- oscillatory zoning, silicates, B:4
- oxidation, magnetic anomalies, B:257–259
- oxygen isotopes
 bulk rocks vs. percent recrystallization, B:163
 diabase, B:102–105
 reaction zones, B:147–150
 secondary minerals, B:160
 vs. albite percentage, B:103
 vs. alteration percentage, B:103
 vs. amphibole percentage, B:103
 vs. depth, B:104–105, 162
 vs. magnesium, B:146, 151
 vs. water/rock ratio, B:105
- Pacific Ocean E
 geochemical logs, B:339–346
 site description, A(137):15–55;
 A(140):37–200
- paleomagnetic corrections, dip, A(140):91
- paleomagnetic reorientation, diabase, B:245–247
- paleomagnetism, A(140):103–106, 125
- palladium
 diabase, B:117–120
 vs. depth, B:119
- Panama Basin, geologic history, A(140):41–44
- Panama Fracture Zone, tectonics, A(140):41
- partial melts, oceanic layers, B:47–50
- particulates, borehole fluids, A(137):38–39
- partitioning, platinum element group, B:203
- Pearce element ratio, basalts, B:54–56
- penninite
 chemical composition, B:149
 microprobe data, B:210–211
- peridotite, mantle, B:137–138
- permeability, packer test, A(137):50–51
- petrography
 alteration, B:168–169
 diabase, B:35–38
 sheeted dike, B:20, 156–158
- petrology
 dike margins, B:35–42
 Site 504, B:3–17
- pH, vs. depth, B:145
- phenocrysts
 basalts, A(137):26–27; A(140):53–55
 clinopyroxene, B:121–130
 geochemistry, B:87
 olivine, B:17
 photograph, B:215–216
 plagioclase, B:5
 pyroxene, B:13
 silicates, B:35–38
- phosphorus/titanium ratio, vs. potassium/titanium ratio, B:55
- phosphorus oxide
 vs. alteration percentage, B:70
 vs. magnesium oxide, B:45
- phyllosilicates
 hydrothermal alteration, B:158, 207–216
 secondary minerals, B:158–159
- physical properties
 Site 504, A(137):29–31; A(140):106,
 108–109, 125
 upper crustal rocks, B:273–291
- picnochlorite
 microprobe data, B:210–211
- molar composition, B:176
- piezoremanence, remanent magnetization, B:269
- pillow lava
 geochemistry, B:38
 petrology, B:19–20
- plagioclase
 basalts, A(137):24, 26–27
 chemistry, B:192
 diabase, B:3–9
 fluid inclusions, B:193–194
 geochemistry, B:67–72
 geothermometry, B:178
 grain boundaries, B:223–225
 grain size, B:20–22
 length distribution, B:21
 petrography, B:36–38
 phenocrysts, A(140):53–54
 post-entrapment crystallization, B:134–135
 secondary minerals, A(140):69
 vs. amphibole, B:179
 zoning patterns, A(140):58–62
- plagioclase, calcic, secondary minerals, B:172,
 178–179
- plagioclase, length of
 average maximum length and equivalent
 diameter, A(140):154–155
 vs. depth, A(140):61–62
 vs. length of magnetite, A(140):61
- plate boundary, tectonics, A(140):41
- platinum element group, sheeted dike complex,
 B:199–205
- poikilitic texture
 pyroxene, B:16
 silicates, B:3–4
- Poisson's ratio, vs. depth, B:282–283
- porosity
 dikes, B:274–275
 fractures, B:315–317
 reaction zones, B:150
 vs. bulk density, A(140):126, 128; B:102
 vs. compressional wave velocity, B:278–279
 vs. density, B:281
 vs. depth, A(137):31; A(140):134–135; B:279
 vs. velocity, A(140):136
 vs. water content, A(137):30
See also fracture porosity
- potassium
 vs. depth, A(137):37; B:145
 vs. magnesium, A(137):43; B:146, 151
See also sodium + potassium
- potassium/titanium ratio, vs. phosphorus/titanium ratio, B:55
- potassium logs, B:343–344
- potassium oxide
 estimate from geochemical logs, B:345–346
 vs. magnesium oxide, B:45
- preferred orientation, plagioclase groundmass,
 A(140):98
- prehnite, secondary minerals, B:160, 174–176,
 184–185
- primary magmas, composition, B:50–51
- pull-apart structures
 clinopyroxene phenocrysts, A(140):110
 photograph, B:235
- pychnochlorite. *See* picnochlorite
- pyrite
 alteration, A(137):28–29
 secondary minerals, A(140):69–70
- pyroxene
 geothermometry, B:177–178
 grain boundaries, B:223–225
 phenocrysts, A(140):54
See also individual minerals
- pyroxenite, crystal clots, A(140):57–58
- quartz
 fluid inclusions, B:194
 veins, A(140):76, 109; B:161
 vs. depth, A(140):66
 quartz-epidote, veins, A(140):73
- rare earths
 chondrite-normalized basalts, B:50
 chondrite-normalized chilled margins, B:41
 chondrite-normalized diabase, B:69, 94,
 112–113, 127–128, 136, 147, 202
 clinopyroxene, B:121–130
 diabase, B:78, 107–116, 200
 hydrothermally altered samples, B:90–93
 lithologic Unit 220, B:86
 magmatic vs. metamorphic origin, B:109–110
 vs. alteration percentage, B:71
 vs. depth, B:68
 vs. zirconium, B:69
- recrystallization
 alteration, B:168–169
 geochemistry, B:163–165
 strain localization, B:226, 228
- reference sample BAS-140, geochemistry, B:79,
 353–355
- reference sample rock BHVO-1, geochemistry,
 B:79
- reference samples, geochemistry, B:79, 353–355
- refractory silicates, melt inclusions, B:131–139
- remanent magnetism, diabase, B:247
- remanent magnetization, anhysteretic
 acquisition, A(140):124; B:332
 anisotropy, A(140):126
 dikes, A(140):104–105
- remanent magnetization, chemical, dikes,
 B:264–269
- remanent magnetization, drilling-induced, dikes,
 B:264–269
- remanent magnetization, isothermal
 acquisition, A(140):125; B:331
 dikes, A(140):104–105; B:264–269
- remanent magnetization, natural
 basalts, A(140):102–103
 dikes, B:254–256
 origin, B:263–270
 vs. depth, A(140):121, 123
- remanent magnetization, thermal, dikes,
 B:264–266, 269
- replacement textures, olivine, B:208–210
- resistivity, vs. depth, A(140):142
- resistivity-natural gamma-ray logs,
 A(140):181–191
- resistivity logs
 Site 504, A(140):113; B:306–307
 vs. depth, B:307–308
- rifting, mid-ocean ridges, B:19–33
- rock magnetism
 oceanic crust, B:327–337
 sheeted dikes, A(140):104–105; B:253–262
- rubidium
 vs. depth, B:145
 vs. magnesium, B:146
- rutile, deuteric oxidation, B:332
- salinity
 vs. depth, B:145, 194
 vs. homogenization temperature, B:194
- samarium
 percent change from protolith, B:203
 vs. alteration percentage, B:110
See also lanthanum/samarium ratio
- sampling tools, borehole fluids, A(137):36, 38–42
- saturation magnetization, vs. depth, B:258
- saturation magnetization/saturation remanence

ratio, vs. depth, B:258
 saturation remanence. *See* saturation magnetization/saturation remanence ratio
 scandium, percent change from protolith, B:203
 seawater/basalt interaction, chemistry, B:146–147
 seawater circulation, fluid inclusions, B:197
 secondary minerals
 alteration, B:72
 hydrothermal alteration, B:158–160
 sheeted dike complex, B:167–189
 thin sections, A(140):164–166
 vs. depth, A(140):65; B:157, 168, 212, 330
 sector zoning, silicates, B:4
 secular variations, paleomagnetism, B:268
 seismic profiles, A(140):95
 Semail massif, ophiolite, B:283, 287–289
 serpentine–stevensite, alteration, B:213
 serpentine
 microprobe data, B:208–209
 secondary minerals, B:173, 183
 vs. depth, A(140):66
 shear deformation, veins, B:238
 shear modulus, vs. depth, B:283
 shear wave velocity
 cores, B:276–278, 280–283
 vs. compressional wave velocity, B:282
 vs. depth, A(137):32
 vs. dry-bulk density, A(137):33
 See also compressional wave velocity/shear wave velocity ratio
 shear zones, chilled margins, A(140):103–104
 sheet silicates. *See* phyllosilicates
 Shore scleroscope method, hardness, B:347
 silica
 estimate from geochemical logs, B:345–346
 percent change from protolith, B:203
 vs. depth, A(137):37; A(140):88; B:91, 145
 vs. magnesium, A(137):43; B:146
 vs. magnesium oxide, B:45, 54
 vs. temperature, B:135
 vs. water, A(140):90
 silicates
 composition, B:3–17
 vs. alteration, B:67
 See also phyllosilicates; refractory silicates
 silicon
 vs. depth, B:160
 vs. iron/(iron + magnesium) ratio, B:149, 159, 210
 vs. magnesium number, B:238
 Site 504, A(137):15–55; A(140):37–200
 alteration and metamorphism, A(140):64–78, 121–123
 altered basalts, B:43–51
 background and objectives, A(137):16–21; A(140):38–44
 basalts, B:347–351
 borehole fluids, B:141–152
 borehole televiewer, A(137):47–50; B:293–304
 chilled dike margins, B:35–42
 cleanout operations, A(137):23–24
 clinopyroxene, B:121–130
 coring, A(137):22–23; A(140):48, 50
 diabase and basalts, B:53–61
 dikes, B:19–33
 downhole measurements, second phase, A(137):24
 downhole measurements, A(140):109, 112–116, 125–127
 downhole measurements prior to drilling, A(137):19
 drilling history, A(137):18, 19
 flowmeter/injection experiment, A(137):50–51

fluid flow, B:313–319
 fluid inclusions, B:191–198
 fluid sampling and chemistry, A(137):31–42
 geochemical logs, B:339–346
 geochemistry, A(140):78–82, 121; B:65–97, 99–110, 117–120, 199–205
 heat flow, B:321–324
 igneous petrology, A(140):52–64, 119–121
 igneous petrology and alteration, A(137):24, 26–29
 magnetic fabric, B:245–252
 melt inclusions, B:131–139
 mineralogy, B:155–166
 natural remanent magnetization, B:263–270
 oceanic crust, B:327–337
 olivine, B:207–216
 operations, A(137):21–24; A(140):44–52
 paleomagnetism, A(140):102–106, 125
 past operational difficulties, A(137):16–19
 petrology, B:3–17
 physical properties, A(137):29–31; A(140):106, 108–109, 125; B:273–291
 reference sample, B:353–355
 remedial and drilling operations, A(137):19–21
 rock magnetism, B:253–262
 sheeted dike complex, B:167–189, 305–311
 site description, A(137):15–55; A(140):37–200
 strain localization, B:219–229
 structure and deformation, A(140):82–88, 90–92, 94–102, 123–124
 summary and conclusions, A(137):51–54
 temperature logging, A(137):42, 44–47
 veins, B:231–243
 skeletal crystals, plagioclase, B:8
 smectite
 vs. depth, A(140):66
 See also chlorite–smectite
 sodium
 vs. depth, A(137):37; B:145
 vs. magnesium, A(137):43; B:146, 151
 See also calcium/(calcium + sodium) ratio
 sodium + potassium, vs. chlorine, B:160
 sodium/titanium ratio, vs. aluminum/titanium ratio, B:56
 sodium oxide
 percent change from protolith, B:203
 vs. alteration percentage, B:70
 vs. magnesium number, B:126
 vs. magnesium oxide, B:45
 vs. wollastonite, B:169
 See also calcium oxide/sodium oxide ratio
 solidification. *See* crystallization
 sphene. *See* titanite
 spinel
 diabase, A(140):152
 petrography, B:36
 phenocrysts, A(140):54
 spinel, chromian
 alteration, A(137):24, 26–27
 hydrothermal alteration, B:157
 stability in basalt and diabase, A(140):62–63
 spreading centers, magmas, B:49
 stable isotopes
 diabase, B:99–106
 sheeted dikes, B:155–166
 See also deuterium/hydrogen ratio; carbon isotopes; oxygen isotopes; strontium isotopes; sulfur isotopes
 stable remanence, dikes, B:266–269
 strain, localization, B:219–229
 stress

borehole televiewer, B:293–304
 cores, B:248
 magnitude, B:298–299, 301–303
 vs. depth, B:302
 See also thermal stress
 stresses, circumferential, vs. maximum horizontal stress, B:302
 stresses, maximum horizontal, orientation, B:301
 strontium
 diabase, B:108
 pillow basalts, B:129
 vs. alteration percentage, B:111
 vs. chromium, B:111
 vs. depth, A(137):37; A(140):89; B:145
 vs. europium, B:134
 vs. magnesium, B:146
 vs. magnesium oxide, B:48
 strontium isotopes
 reaction zones, B:147–150
 vs. depth, B:145
 structural geology, A(140):82–88, 90–92, 94–102, 123–124
 structural logs, A(140):172–180
 sulfate
 vs. calcium, A(137):43
 vs. depth, A(137):37; B:145
 vs. magnesium, A(137):43; B:146, 151
 sulfides
 hydrothermal alteration, B:157–158
 secondary minerals, A(140):69–70
 See also individual minerals
 sulfur
 secondary minerals, B:160–161
 vs. alteration percentage, B:71
 vs. copper, B:163
 vs. depth, B:162–164
 sulfur isotopes
 secondary minerals, B:160–161
 vs. depth, B:162–164
 vs. proportion of sulfate sulfur in lower dike, B:165
 supersaturation, diabase, B:3–4
 talc
 alteration, A(137):28–29
 microprobe data, B:208–209
 secondary minerals, B:173, 183
 vs. depth, A(140):66
 talc–chlorite
 chemical composition, B:149
 microprobe data, B:210–211
 secondary minerals, B:173, 183
 temperature
 fluid entrapment, B:194–195
 vs. depth, A(137):45–47; A(140):46; B:196
 temperature gradient
 vs. depth, A(140):139–141, 144; B:308, 314–316, 322
 See also geothermal gradient
 temperature inversion, Site 504, B:317–318
 temperature logs
 Site 504, A(137):42, 44–47; B:321–324
 vs. depth, B:296, 306, 308, 314, 322
 temperature logs, high-resolution, A(140):112–116
 temperature profile, vs. depth, A(140):144–145
 tension gashes
 amphibole veins, A(140):111
 veins, B:233–234
 textures
 diabase, B:220, 223
 silicates, B:3–4
 thermal conductivity
 cores, A(137):31

- Site 504, A(140):109
 vs. depth, A(140):137–138; B:324
 thermal contraction, stresses, B:251–252
 thermal cracks, backscattered electron imaging, B:222–228
 thermal evolution, diabase, B:195
 thermal stress, dikes, B:295–296
 thorium logs, Site 504, B:343–344
 titanite
 aluminum–titanium–iron diagram, B:177
 deuterium oxidation, B:332
 magnetic minerals, B:257–259
 secondary minerals, B:176, 187
 titanium
 comparison of analyses by electron and ion microprobes, B:134
 hydrothermally altered samples, B:90–93
 immobile elements, B:86–87
 vs. depth, B:160
 vs. titanium/zirconium ratio, A(140):92
 vs. zirconium, B:136
See also aluminum/titanium ratio; (iron + magnesium)/titanium ratio; magnesium/titanium ratio; phosphorus/titanium ratio; potassium/titanium ratio; sodium/titanium ratio; yttrium/titanium ratio; zirconium/titanium ratio
 titanium/zirconium ratio
 vs. depth, A(140):93
 vs. lanthanum/samarium ratio, B:137
 vs. titanium, A(140):92
 vs. ytterbium/zirconium ratio, B:204
 vs. zirconium, B:50
 titanium oxide
 estimate from geochemical logs, B:345–346
 percent change from protolith, B:203
 vs. alteration percentage, B:70
 vs. aluminum oxide, B:37
 vs. calcium oxide/sodium oxide ratio, A(140):87
 vs. copper, B:92
 vs. depth, A(140):88, 93; B:68, 91
 vs. magnesium number, A(140):86–87; B:88, 126
 vs. magnesium oxide, B:45
 vs. water, A(140):89; B:72
 vs. zirconium, B:69
See also zirconium/titanium oxide ratio
 titanohematite, anhysteretic remanent magnetization, A(140):106
 titanomagnetite
 exsolution, B:267
 magnetic anomalies, B:257–259
 secondary minerals, A(140):69
 trace elements
 basalts, A(137):27–28; B:53–61
 chilled margins, B:38–40, 42
 clinopyroxene, B:121–130
 diabase, B:114
 igneous rocks, A(140):78–82, 121
 melt inclusions, B:135
 oceanic crust, B:327–337
 partial melts, B:50
 primary and secondary variations, B:65–80
 reference sample, B:353, 355
 vs. magnesium oxide, B:48
 uranium logs, Site 504, B:343–344
 vanadium, vs. magnesium oxide, B:48
 vein orientations, paleomagnetic corrections, A(140):108
 veins
 alteration halos, B:179, 182
 amphiboles, B:210–213
 chlorite, B:210
 deformation, B:233–234
 density and volume vs. lithology, A(140):75
 dikes, B:274–275
 fill textures, B:233
 formation, A(140):95–96; B:231–243
 geometry and orientation, B:232–233
 hydrothermal alteration, B:158
 petrology, A(140):72–76
 photograph, A(140):71–74; B:227, 242–243
 reorientation, B:248–249
 secondary minerals, B:189
 true dip data, A(140):86–88, 90–91
See also crosscutting veins
 velocity
 vs. density, A(140):136
 vs. depth, A(140):134–135; B:288
 vs. porosity, A(140):136
 velocity, ultrasonic, cores, A(137):29–30
 vugs, alteration, A(137):28–29
 water
 vs. alteration percentage, B:70
 vs. calcium oxide, A(140):90
 vs. copper, A(140):89
 vs. depth, B:68
 vs. magnesium number, A(140):90
 vs. silica, A(140):90
 vs. titanium oxide, A(140):89; B:72
 vs. zinc, A(140):90
 vs. zirconium, A(140):89
 vs. zirconium/yttrium ratio, A(140):89
 water/rock ratio
 mass ratios, B:148–150
 vs. oxygen isotopes, B:105
 water content
 hydrogen isotopes, B:101
 vs. alteration percentage, B:102
 vs. depth, A(137):31
 vs. deuterium/hydrogen ratio, B:102
 vs. porosity, A(137):30
 wehrlite, crystal clots, A(140):57–58
 well logging, A(140):109, 112–116, 125–127; B:313–319, 321–324, 339–346
 well logs, sheeted dike complex, B:305–311
 wollastonite
 vs. chromium oxide, B:169
 vs. sodium oxide, B:169
 xenocrysts, clinopyroxene, B:121–130
 xenoliths, petrology, A(140):56–58
 Young's modulus
 basalt, B:348–349
 correlation diagram between dynamic and static moduli, B:351
 vs. uniaxial compressive strength, B:350
 ytterbium/zirconium ratio, vs. titanium/zirconium ratio, B:204
 ytterbium
 percent change from protolith, B:203
 vs. lanthanum/samarium ratio, B:50
 vs. magnesium number, B:126
See also lanthanum/ytterbium ratio; zirconium/ytterbium ratio
 yttrium
 hydrothermally altered samples, B:90–93
 immobile elements, B:86–87
 vs. alteration percentage, B:71
 vs. depth, B:68
 vs. magnesium oxide, B:48
See also zirconium/yttrium ratio
 yttrium/titanium ratio, vs. zirconium/titanium ratio, B:55
 zeolites
 alteration, A(137):28–29
 chemical composition, B:239
 secondary minerals, B:176, 184–185
 veins, B:236
 zinc
 depletion, B:204
 depletion with depth, B:93–94
 vs. alteration percentage, B:71
 vs. depth, A(140):90, 145; B:68, 92, 164
 vs. water, A(140):90
 zirconium
 hydrothermally altered samples, B:90–93
 immobile elements, B:86–87
 percent change from protolith, B:203
 vs. alteration percentage, B:71
 vs. copper, B:92
 vs. depth, A(140):89, 93; B:68, 90
 vs. magnesium oxide, B:48
 vs. nickel, A(140):92; B:89
 vs. niobium, A(137):28; A(140):86; B:90
 vs. rare earths, B:69
 vs. titanium, B:136
 vs. titanium/zirconium ratio, B:50
 vs. titanium oxide, B:69
 vs. water, A(140):89
 vs. zirconium/yttrium ratio, B:90, 94
See also titanium/zirconium ratio; ytterbium/zirconium ratio
 zirconium/titanium ratio, vs. yttrium/titanium ratio, B:55
 zirconium/titanium oxide ratio, vs. depth, B:68
 zirconium/ytterbium ratio, vs. lanthanum/ytterbium ratio, B:203
 zirconium/yttrium ratio
 vs. copper, B:93
 vs. depth, A(140):89; B:90
 vs. lanthanum/samarium ratio, B:115
 vs. lanthanum/ytterbium ratio, B:115
 vs. water, A(140):89
 vs. zirconium, B:90, 94
 zoning patterns
 plagioclase, A(140):58–62
 secondary minerals, A(140):77–78