

INDEX TO VOLUME 148

This index provides coverage for both the *Initial Reports* and *Scientific Results* portions of Volume 148 of the *Proceedings of the Ocean Drilling Program*. References to page numbers in the *Initial Reports* are preceded by “A” with a colon (A:), and to those in the *Scientific Results* (this book), by “B” with a colon (B:). In addition, reference to material on CD-ROM is shown as “bp:CD-ROM.”

The index was prepared by Earth Systems, under subcontract to the Ocean Drilling Program. The index contains two hierarchies of entries: (1) a main entry, defined as a keyword or concept followed by a reference to the page on which that word or concept appears, and (2) a subentry, defined as an elaboration on the main entry followed by a page reference.

The index covers text, figures, and tables but not core-description forms (“barrel sheets”), core photographs, or thin-section descriptions; these are given in the *Initial Reports*. Also excluded from the index are bibliographic references, names of individuals, and routine front and back matter.

The index follows a standard format. Geographic, geologic, and other terms are referenced only if they are subjects of discussion. This index also includes broad fossil groups such as nannofossils and radiolarians. A site chapter in the *Initial Reports* is considered the principal reference for that site and is indicated on the first line of the site’s listing in the index. Such a reference to Site 896, for example, is given as “Site 896, A:123–192.”

For further information, including available electronic formats, contact the Chief Production Editor, Ocean Drilling Program, 1000 Discovery Drive, College Station, Texas 77845-9547, U.S.A., E-mail: pub_production@ODP.TAMU.EDU

VOLUME 148 SUBJECT INDEX

- acoustic velocity array sonde logging, A:75–76
- acoustic well logs, vs. depth, A:81
- actinolite
- dike complex, B:410–411
 - petrography, A:47
 - photograph, A:51
 - secondary minerals, A:45–53; B:77, 422–423, 427
 - vein dips, A:63
 - vs. depth, A:52
 - See also* chlorite–actinolite veins
- actinolite veins
- dip histograms, B:232
 - petrology, A:49–51
 - secondary minerals, B:85
 - structure, B:231–233
- active faults, image analysis, B:379, 384
- albite
- petrography, A:47
 - vs. depth, A:52
- albitization, chemical effects, B:49
- alteration
- comparison of Sites 504 and 896, A:149–150; B:158
 - composition, B:439–442
 - dikes, B:97–109
 - geochemistry, A:57–60; B:156–157, 192–196
 - gold, B:453–454
 - heat flow, B:293–294
 - heterogeneity, A:45–47
 - image analysis, B:368–369, 379
 - log, A:116–117, 189, bp:CD-ROM
 - magnetic properties, B:225
 - mineralogy, B:421–425, 427–428
 - nitrogen, B:6
 - photograph, B:202–206, 208
 - property changes, B:171–189
 - secondary minerals, B:71–86
 - sheeted dike complex, B:47
 - Site 504, A:45–53
 - Site 896, A:141–150
 - timing, B:137–138
 - vs. depth, B:48, 61, 155, 454
 - vs. rock chemical compositions, B:49
 - zoning, B:419–421
 - See also* hydrothermal alteration; metamorphism; metasomatism; oxidative alteration; secondary minerals; veins
- alteration, bulk-rock, geochemistry, B:12–13
- alteration halos
- color, B:444
 - composition, A:50–53, 141; B:426, 440
 - dikes, B:97–109
 - fluid inclusions, B:94
 - geochemistry, B:52, 156–157
 - isotopes, B:65
 - mineralogy, B:155, 427–428
 - petrology, A:46; B:420
 - secondary minerals, B:86, 168
 - veins, B:261–263
 - volume percent vs. depth, A:49
 - vs. depth, B:442
 - zoning, B:154
- alteration halos, black, alteration, B:180–183
- alteration halos, colored, basalts, B:157–158
- alteration halos, low-temperature, black halos, B:180–183
- alteration patches
- amphibole and chlorite, B:103
 - average element concentration, B:52
 - background alteration, A:48–49
 - composition, B:427–428
 - geochemistry, B:461–463
 - intersection with actinolite vein, A:49
 - isotopes, B:65
 - petrology, A:46
 - photograph, A:48
 - primary depletion, B:50
 - sheeted dike complex, B:47
 - volume percent vs. depth, A:49
- alteration percentage
- vs. depth, B:428
 - vs. trace elements, A:62
- alteration rates, microorganisms, B:199
- alteration space, vs. depth, B:374
- alteration types, trace elements, B:465
- aluminum. *See* iron/(iron + aluminum) ratio
- aluminum, tetrahedral
- secondary phyllosilicates, B:125
 - vs. depth, B:80
 - vs. interlayer potassium, B:125
- aluminum, total, secondary phyllosilicates, B:125
- aluminum oxide
- electron microprobe data along basaltic glass alteration, B:210
 - profiles across microbially processed glass, B:200
 - vs. depth, A:60, 62, 157; B:14, 18, 136, 422, 484
 - vs. loss on ignition, B:139
 - vs. magnesium number, A:59, 151
 - vs. magnesium oxide, B:19, 23, 30–31, 34–35
 - vs. silica, B:197–199
 - vs. sub-basement depth, A:159
 - vs. titanium oxide, B:138
- amphibole composition, vs. depth, B:425
- amphibole veins
- crenulation, B:234
 - fiber fillings, B:234
 - isotopes, B:65
 - photograph, B:231, 233
- amphiboles
- alteration patches, B:103
 - composition plotted on Leake's diagram, B:106
 - electron microprobe data, B:105
 - fibers in veins, A:66
 - modal data, B:66
 - photograph, A:67
 - secondary minerals, B:76–80, 427–428
 - vs. depth, B:60
- analcime
- composition, B:124
 - photograph, A:146
 - strontium isotopes, B:137
- analcime veins, alteration, B:443
- anatase, basalts, B:473, 475–477
- anhydrite
- electron microprobe data, B:107
 - secondary minerals, B:426, 429
- anorthite
- petrography, A:47
 - vs. calcium oxide/sodium oxide ratio, B:26
 - vs. magnesium oxide, B:428
- apatite, secondary minerals, B:77
- aragonite
- composition, B:124
 - fluid inclusions, A:148
 - secondary minerals, A:141; B:153–154
 - secondary minerals in breakout, A:56
 - See also* calcium carbonate; calcite; carbonate content
- aragonite veins
- photograph, A:146
 - stereographic projection, A:165
- Archaea
- DNA products, B:211
 - oligonucleotide probe hybridization, B:209
 - See also* bacteria; chain reactions; DNA; microbial activity; microorganisms
- Archie's formula, porosity, B:320
- augite
- percentage vs. depth, A:138
 - secondary minerals, B:428
 - See also* clinopyroxene
- bacteria
- basaltic glass alteration, B:196–198
 - photograph, B:205
 - See also* Archaea; chain reactions; DNA; microbial activity; microorganisms
- basaltic melts, density, A:140
- basalts
- alteration, B:156–157, 171–189, 419–421, 441
 - alteration mineralogy, B:172–173
 - alteration petrography, B:172–173
 - comparison of geochemistry in Sites 504 and 896, A:150–151; B:13
 - crystallization, B:21–35
 - geochemistry, A:150–155; B:10–13
 - hydrothermal alteration, B:122–123
 - magnetic field, B:335–336
 - major elements, B:483–487
 - neutron absorption cross section, B:389–394
 - thermal conductivity, B:397–400
 - volcanic stratigraphy, A:129–132
- basalts, mid-ocean-ridge
- hydrothermal alteration, B:6
 - isotopic profiles, B:57–69
- basement
- neutron absorption cross section, B:389–394
 - oceanic crust, A:32–34; B:340–341
 - structure, B:384
 - structure geophysical measurements, B:267–268
- bathymetric map, Site 896, A:125
- borehole deviation, vs. depth, A:77–79; B:310
- borehole elongation
- azimuth histogram, A:174
 - fractures, B:314
 - stresses, B:323–325, 384
 - vs. depth, B:329, 386, 406
- borehole fluid geochemistry, A:53–57; B:111–118
- borehole size, vs. depth, A:77
- borehole spike, sodium bromide, A:56–57
- boron, vs. depth, B:422
- boron isotopes, vs. depth, B:423
- branching textures
- photograph, A:135
 - quench textures, A:133
- breccia
- dip histograms, B:255
 - hydrothermal alteration, B:123
 - isotopes, B:145
 - lithostratigraphy, B:437–439
 - matrix, B:443

- permeability, B:446
 petrology, A:146–148
 photograph, A:147–149, 164–165, 252–254
 sedimentary vs. tectonic origin, B:250–251
 structure, B:266
 structure and deformation, A:155–156
 textures, A:131–132
 vs. depth, B:270–271, 438
 breccia, fault, structure, B:251
 breccia, jigsaw-puzzle, vein networks, B:251–256
 breccia, nontectonic, mid-ocean ridges, B:251
 brecciation, pillow lava, B:269
 brines, electrical conductivity, B:302
 bromine, seawater, A:56–57
 bulk rocks, alteration, A:141–144
- calcite
 composition, B:124
 secondary minerals, A:141; B:153–154, 173, 426
See also aragonite; calcium carbonate; carbonate content
- calcium
 vs. depth, A:55, 128
 vs. magnesium, A:55
- calcium carbonate
 alteration geochemistry, B:135–137
 chemical composition, B:124, 166
 isotopes, B:146
 photograph, A:149
 secondary minerals, B:153–154, 173
 vs. depth, B:443
See also aragonite; calcite; carbonate content
- calcium oxide
 electron microprobe data, B:487
 electron microprobe data along basaltic glass alteration, B:210
 profiles across microbially processed glass, B:200
 vs. alteration, B:49
 vs. depth, A:60, 62, 157; B:48, 136, 422, 484
 vs. loss on ignition, B:139
 vs. magnesium number, A:59, 151
 vs. magnesium oxide, B:19, 31, 34–35, 197–199
 vs. strontium isotopes, B:144
 vs. sub-basement depth, A:159
 vs. titanium oxide, B:138
- calcium oxide/sodium oxide ratio
 vs. anorthite content, B:26
 vs. depth, B:22
 vs. magnesium oxide, B:30, 34–35
- caliper logs, borehole elongation azimuth histogram, A:78, 174
- carbon dioxide
 alteration, B:173, 175
 diabase, B:4
 hydrothermal circulation, B:447–448
 vs. depth, B:422
 vs. nitrogen, B:6
 vs. oxygen isotopes, B:144
 vs. strontium isotopes, B:144
- carbon isotopes
 alteration geochemistry, B:132–137
 vs. depth, B:146, 423
 vs. oxygen isotopes, B:148
 vs. strontium isotopes, B:148
- carbonate + clay veins
 dip histograms, B:265
 orientation and distribution, B:264
 poles, B:266
- carbonate content
 X-ray diffraction data, A:143
See also saponite–carbonate–phillipsite veins
- carbonate fibers, photograph, A:161–162
- carbonate veins
 dip histograms, B:265
 history, B:448
 petrology, A:145
 photograph, A:161–163
 Site 896, B:155–156
 vs. depth, B:438
 X-ray diffraction and cathodoluminescence data, B:283
- cataclastic zones
 fractures, B:242
 photograph, A:66
 structure, B:235–236
 thin sections, A:64
- cathodoluminescence, diabase, B:84
- cation exchange capacity
 electrical conductivity, B:298–299
 porosity, B:320–321
- celadonite
 chemical composition, B:125, 160–162, 168–170
 electron microprobe data, B:124–126
 secondary minerals, B:153, 173, 444
See also saponite–celadonite–iron oxides mixtures
- celadonite–nontronite mixtures
 alteration, B:173
 secondary minerals, B:153
- celadonite–nontronite–saponite mixtures,
 secondary minerals, B:153
- celadonite–saponite mixtures, alteration, B:173
- celadonitic minerals
 alteration, B:173
 veins, B:155–156
- cerium, vs. alteration, B:49
- cesium
 vs. loss on ignition, B:141
 vs. potassium oxide, B:141
- chabazite, secondary minerals, A:141; B:208
- chain reactions, polymerase
 alteration, B:209
 microbial activity, B:211
- chalcopyrite
 isotopic profiles, B:63–64
 secondary minerals, A:45–53
- chemical profiles, mixing, B:118
- chemical properties, alteration, B:171–189
- chemostratigraphy, basalts, B:14–16
- chilled margins
 diabase, A:42–43
 microfaults, A:70
 photograph, A:41
 pillow lava, A:132–137
 structure and deformation, A:61, 63, 156–157
 thin sections, A:64–65
- chlorine, seawater, A:56–57
- chlorite
 alteration patches, B:103
 chemical composition, B:107, 124, 163
 iron oxide weight percent, B:83
 modal data, B:66
 petrography, A:47
 secondary minerals, A:45–53, 141; B:77, 82, 86
 sheeted dike complex, B:410–411
 vs. depth, A:52
See also quartz–chlorite–sulfide veins
- chlorite, groundmass, electron microprobe data, B:106
- chlorite veins
 dip histograms, B:232
 dips, A:63
 Site 504, A:50–51
- Site 896, B:269–270
 stereographic projection, A:165
 structure, B:231–233
- chlorite–actinolite veins, structure, B:231–233
- chlorite–smectite mixtures
 secondary minerals, B:77, 82
 veins, B:269–270, 426
- chloritization, chemical effects, B:49
- chromite, electron microprobe data, B:105
- chromium
 vs. alteration percentage, A:62
 vs. depth, A:61–62, 158; B:14, 48, 137
 vs. iron, B:28
 vs. loss on ignition, B:140
 vs. magnesium number, A:59, 156; B:28
 vs. sub-basement depth, A:159
 vs. water content, A:62, 160
 clasts, breccias, B:250–251
- clasts, basaltic, photograph, A:164–165
- clay, vermicular, photograph, A:165
- clay + pyrite veins, petrology, A:145
- clay fibers, photograph, A:162
- clay minerals
 chemical composition, B:164–166
 electron microprobe data, B:176
 identification, A:141
 major oxides, B:163
 photograph, B:214
 secondary minerals, B:152–153
 X-ray diffraction data, A:142
See also iddingsites; phyllosilicates
- clay selvage, photograph, A:145
- clay veins
 dip histograms, B:265
 orientation and distribution, B:264
 petrology, A:145
 photograph, A:146, 161
 stereographic projection, A:165
See also carbonate + clay veins
- clinopyroxene
 alteration, B:173, 428
 background alteration, A:47
 chemical composition, B:153–154
 photograph, A:40
 secondary minerals, B:77, 80–81
 sheeted dike complex, B:410–411
- clinopyroxene phenocrysts, A:40, 132, 137–140; B:24, 28, 104
- Cocos–Nazca spreading center. *See* Galapagos Spreading Center
- coercivity
 basalts, B:471
 vs. depth, B:181
- color, alteration, B:172
- comb textures
 photograph, A:135
 quench textures, A:134
- compressional wave velocity
 basalts, B:367–368
 histogram, A:173
 vs. bulk density, A:171
 vs. confining pressure, B:412–413
 vs. depth, A:20, 35, 74–76, 170, 172–173; B:412, 414, 428
 vs. porosity, B:372
 vs. wet-bulk density, B:371
- convection cycle, volcanic rocks, B:436–437
- copper
 depletion, B:51, 429
 vs. alteration, B:49
 vs. depth, A:61, 158; B:48, 61, 137, 423
 vs. loss on ignition, B:140
 vs. sulfur, B:140
 vs. titanium oxide + iron oxide, B:140

- vs. zinc, B:53
 corrensite, veins, B:269–270
 Costa Rica Rift
 alteration mineralogy, B:151–170
 electrical images, B:375–388
 fractures, B:242, 317–329
 geochemical logging, B:389–394
 geology, B:436–437
 layer 2, B:261–279
 magma chambers, B:17
 site description, A:27–121
 crack-seal mechanism
 fibrous veins, A:158
 veins, B:272, 281–288
 cross-cutting relations
 photograph, A:69
 structure and deformation, A:156
 thin sections, A:66
 crystal growth, crack-seal veins, B:284–285
 crystal size distribution
 basalts, A:139
 thin sections, A:187, bp:CD-ROM
 crystallinity, alteration, B:180
 crystallization
 diabase, B:44–45
 primary melts, B:21–35
See also fractional crystallization;
 fractionation index
 Curie temperature
 alteration, B:178–183
 vs. depth, B:179
 vs. oxidation ratio, B:182
 deformation
 log, A:121, 191, bp:CD-ROM
 seismic reflection, A:60–61, 63–68
 degassing, nitrogen, B:6
 demagnetization, alternating-field
 basalts, A:160, B:474
 diabase, A:70
 orthogonal vector representations, A:166
 dendritic growth, photograph, A:134
 density, vs. depth, B:370
 density, bulk
 vs. compressional wave velocity, A:171
 vs. depth, A:35, 74, 170, 172
 vs. porosity, A:171
 vs. unconfined compressive strength, B:404
 density, grain
 basalts, B:367–368
 vs. depth, A:35
 vs. unconfined compressive strength, B:404
 density, wet-bulk
 basalts, B:367–368
 vs. compressional wave velocity, B:371
 deuterium
 vs. depth, B:423
See also hydrogen isotopes
 diabase
 cathodoluminescence, B:84
 electrical conductivity, B:304
 fractures, B:307–315, 317–329
 geochemistry, A:57–60; B:39–55
 gold, B:453–454
 isotopic profiles, B:57–69
 lithologic units, A:37–38
 metamorphism, B:97–109
 mineral abundance in hand specimens and thin
 section, A:42
 nitrogen, B:3–7
 photograph, A:51
 secondary minerals, B:86
 strengths, B:401–407
 structure, B:229–243
 trace elements, B:455–466
 dike complexes, sheeted
 alteration, A:45–53; B:419–421
 diabase, A:42
 fluid inclusions, B:87–95
 fractures, B:307–315
 geochemistry, B:39–55
 glassy texture, A:130
 hydrothermal alteration, B:417–434
 isotopic profiles, B:57–69
 magnetic field, B:334
 margin structure and deformation, A:156–157
 metamorphism, B:97–109
 nitrogen geochemistry, B:3–7
 petrology, B:410–411
 relation of lower part to hydrothermal vent
 fluids, B:66–67
 secondary minerals, B:71–86
 structure, B:229–243
 trace elements, B:455–466
See also semidikes
 diopside
 secondary minerals, B:85
See also amphiboles
 DNA
 microbial activity, B:207–212
 microbially processed glass, B:200
See also RNA
 dolerite. *See* diabase
 downhole electronic gauge, pressure–time records,
 A:175
 downhole measurements
 nitrogen, B:3–7
 Site 504, A:73–81
 Site 896, A:166–175; B:375–388
 structural data, B:236–237
 drilling and recovery, vs. depth, A:38, 130
 dual laterolog logging, A:75–76
 Ecuador Fracture Zone, geologic setting, A:29;
 B:436–437
 electrical conductivity
 salinity, B:297–305
 sample vs. fluid conductivity, B:300
 surface conductivity, B:301–303
 temperature correction, B:300–303
 vs. strengths, B:405
 vs. temperature, B:302
 electrical images, B:375–388
 electrical logs
 fractures, B:307–315
 vs. depth, B:319
 epidote
 electron microprobe data, B:107
 secondary minerals, B:77, 83, 423, 427–429
 epidote veins, A:50–51, 63
 epidotization, chemical effects, B:49
 epifluorescence microscopy
 microbial activity, B:210–211
See also fluorescence
 europium, anomalies, B:50
 exsolution, carbon dioxide, B:50
 fabrics, amphibole veins, B:233
 fabrics, cataclastic, thin sections, A:64–65
 fabrics, igneous, thin sections, A:64–65
 failure, tensile, diabase, B:406–407
 faulting, fault zones, B:430
 faults
 heat flow, B:294–295
 hydrofractures, B:249
 fibers
 amphiboles, A:66
 photograph, A:67
 fibropalagonite
 glass replacement, B:155
See also palagonite
 fluid circulation
 heat flow, B:294–295
 mixing in boreholes, B:111–118
 pressurized system, B:356–357
 vs. temperature, B:303
 fluid inclusions
 carbonate veins, B:283–284
 chemistry, B:26
 ice melting temperatures, B:284
 petrology, A:148
 sheeted dikes, B:87–95
 fluid pressure
 hydrofractures, B:247–250
 surface conductivity, B:301–303
 fluid salinity
 plagioclase inclusions, B:90–91
See also salinity
 fluorescence. *See* epifluorescence microscopy
 foliation, magnetic susceptibility anisotropy, A:72,
 168
 formation factor
 basalts, B:376
 electrical conductivity, B:299–300, 302–303
 vs. porosity, A:171; B:377
 vs. total porosity, B:378
 Formation MicroScanner imaging
 Costa Rica Rift, B:375–388
 downhole measurements, B:237–242
 fractures, B:310–312, 319–320
 logging, A:76–77
 fractional crystallization
 diabase, B:44–45
 lava, A:139
See also crystallization
 fractionation index
 lavas, B:17
 vs. major elements, A:59
 vs. trace elements, A:59
 fracture
 azimuth and dip, B:384
 depth vs. dip, B:236
 depth vs. strike, B:240
 diabase, B:317–329
 downhole electrical images, B:307–315
 fault zones, B:430
 Formation MicroScanner imaging, B:237,
 239–240, 310–312
 horizontal and vertical porosity with depth,
 B:241
 image analysis, B:378–379
 structure, B:234–235, 265–266
 thin sections, A:64–65
 true dip histograms, A:69
 upper oceanic crust, B:246–250
 variation of orientation with depth, B:241
 veins, A:157–158
 fractures, open, dips, A:66–68
 Galapagos Spreading Center, site description,
 A:27–121
 gallium, vs. loss on ignition, B:140
 geochemical cycles, nitrogen, B:6
 geochemical fluxes, hydrothermal circulation,
 B:447–448
 geochemical logs
 neutron absorption cross section, B:389–394
 Site 896, A:172, 183
 vs. depth, A:22, 173
 geochemistry
 hydrothermal alteration, B:119–150
 igneous rocks, B:439

- lava, B:9–19
 lower sheeted dike complex, B:455–466
 secondary minerals, B:71–86
 sheeted dike complex, B:39–55
 Site 504, A:53–60, 150–151; B:111–118
 geophysical combination logging, A:22, 75–76
 geothermal gradient
 vs. depth, B:292–294
 See also heat flow; temperature
 geothermometry
 fluid inclusions, B:89–94
 secondary minerals, B:77
 See also microthermometry
 glass
 chemical composition, B:167
 chemistry, B:22–23
 major elements, B:483–487
 photograph, B:208
 replacement, B:155
 secondary textures, A:134
 glass, basaltic
 alteration, B:191–206
 composition, B:125
 microbial activity, B:207–214
 photograph, B:214
 glass inclusions
 background alteration, A:48
 photograph, A:137
 glass rims, alteration, B:202–204
 glassy/variolitic zone, quench textures, A:133
 glomerocrysts
 basalts, A:136–137
 classified by mineralogical type, A:115, 188,
 bp:CD-ROM
 diabase and basalt, A:44–45
 photograph, A:137
 goethite, secondary minerals, B:173, 189
 gold
 upper oceanic crust, B:453–454
 vs. depth, B:454
 grain size
 alteration, B:180
 diabase, A:43–44
 median destructive field, B:223–224
 Grant plots, alteration patches vs. diabase, B:53
 granularity, alteration, B:180
 greenschist-facies alteration, fluid inclusions, B:91
 hafnium
 vs. depth, B:464
 vs. loss on ignition, B:141
 See also thorium/hafnium ratio;
 zirconium/hafnium ratio
 heat capacity, vs. temperature, B:399
 heat flow
 downhole measurements, B:291–296
 fracture zones, B:437
 vs. depth, B:292–294
 See also geothermal gradient; temperature
 heat flow maps, A:127
 hematite, secondary minerals, A:45–53
 honeycomb textures, photograph, A:134
 hornblende
 secondary minerals, B:77
 See also amphiboles
 hyaloclastite
 breccia, A:132; B:268
 petrology, A:146–148
 photograph, A:149
 hydration, alteration, B:159
 hydraulic fracturing
 breccia, A:158
 upper oceanic crust, B:246–250
 hydrogen isotopes
 vs. depth, B:423
 See also deuterium
 hydrostatic pressure, effect on thermal
 conductivity, B:397–400
 hydrothermal alteration
 comparison to hydrothermal effects in
 ophiolites, B:431
 fluid introduction, A:148–149
 geochemistry, B:39–55
 isotope profiles, B:57–69
 nitrogen, B:6
 oceanic crust, B:417–434
 plagioclase inclusions, B:91–94
 processes, A:52–53
 secondary minerals, B:71–86
 upper ocean crust, B:119–150
 See also alteration
 hydrothermal alteration, ridge flank, oceanic crust,
 B:435–450
 hydrothermal fluids
 alteration, B:444
 alteration halos, B:158
 basement, B:446
 hydrothermal vents, dikes, B:66–67
 hysteresis
 alteration, B:178–180
 basalts, B:471
 temperature cycle, B:302
 iddingsite
 chemical composition, B:164–166
 electron microprobe data, B:176
 fillings, B:155
 major oxides, B:163
 secondary minerals, B:153, 172–173, 187
 veins, B:155–156
 See also clay minerals; phyllosilicates
 igneous contacts
 deformation, A:61–63
 log, A:110, bp:CD-ROM
 structure, B:235–236, 266–267
 igneous and metamorphic geochemistry
 Site 504, A:57–60
 Site 896, A:150–151
 igneous rocks
 geochemistry, A:57–60
 lithology log, A:112, 185, bp:CD-ROM
 mineralogy log, A:111, 184, bp:CD-ROM
 petrology, A:37–45, 129–141, 439
 ilmenite
 basalts, B:473, 475–477
 electron microprobe data, B:105
 major and trace elements, B:475
 ilmenite–titanite alteration, chemical effects, B:49
 image analysis
 alteration, B:368–370
 fractures, B:378–379
 resistive units, B:379, 380–383
 implosion, breccias, B:254
 inclusions
 igneous contacts, B:235–236
 temperature, B:21–35
 incompatible elements, depletion, B:50
 injection tests, pressure, B:363
 injection tests, constant-rate, permeability,
 B:355–356
 intergranular texture
 diabase, A:41–42
 photograph, A:42
 international reference rocks JB2 and JB3, trace
 elements, B:463
 interstitial materials, background alteration, A:48
 iron
 alteration, B:177–178
 alteration enrichment, B:159
 vs. chromium, B:28
 vs. depth, B:61, 136
 vs. loss on ignition, B:139
 vs. magnesium, B:149
 vs. oxygen isotopes, B:144, 148
 vs. strontium isotopes, B:144
 vs. uranium, B:142
 See also magnesium/(magnesium + iron) ratio
 iron/(iron + aluminum) ratio, prehnites and
 epidotes, B:425
 iron/(iron + magnesium) ratio
 transition zone, B:425
 vs. aluminum total, B:125
 vs. interlayer potassium, B:125
 vs. octahedral occupancy, B:125
 iron logs
 vs. depth, A:173
 vs. silicon logs, A:172
 iron oxide
 chemical composition, B:163
 electron microprobe data, B:487
 electron microprobe data along basaltic glass
 alteration, B:210
 profiles across microbially processed glass,
 B:200
 total weight in chlorites, B:83
 vs. depth, A:60, 157; B:14, 83, 136, 422, 484
 vs. iron oxides, B:170
 vs. loss on ignition, B:139
 vs. magnesium number, A:59, 151
 vs. magnesium oxide, B:31, 34–35
 vs. major oxides, B:170
 vs. manganese oxide, B:212
 vs. silica, B:156, 197–199
 vs. titanium oxide, B:138, 197–199
 See also saponite–celadonite–iron oxides
 mixtures; titanium oxide + iron oxide
 iron oxide/(iron oxide + magnesium oxide) ratio
 vs. depth, B:424
 vs. potassium oxide, B:156
 iron oxyhydroxide
 chemical composition, B:124, 165
 secondary minerals, B:153, 172–173, 444
 iron oxyhydroxide veins
 dip histograms, B:265
 petrology, A:145
 true dip histogram, A:164
 vs. depth, B:438
 isotopes
 hydrothermal alteration, B:119–150
 See also stable isotopes; individual isotope
 types
 Koenigsberger ratio
 basalts, A:160
 diabase, A:70
 vs. depth, A:166
 lanthanum, vs. magnesium oxide, B:53
 laumontite
 composition, B:108
 petrography, A:47
 secondary minerals, B:77, 429
 veins, A:50–51
 lava
 fractional crystallization, A:139
 geochemistry, B:9–19
 textures, A:130–131
 lava flows, veins, B:272
 lava, massive, with complex oxidative alteration
 halos, A:130
 lawsonite, composition, B:108
 layer 2, upper

- seismic structure, B:369–370
 structural evolution, B:245–279
 veins, B:281–288
- layer 2/3 boundary
 seismic layers, B:346
 structure, B:242
 well-logging, B:409–414
- leaching, nitrogen, B:6
- lineation
 magnetic susceptibility anisotropy, A:72, 168
 microfaults, A:70
- lithium
 vs. depth, B:137, 422, 464
 vs. loss on ignition, B:140
 vs. potassium oxide, B:141
- lithologic units, igneous rocks, A:37–38, 129
- lithology
 control on alteration style, B:158
 igneous rocks, A:112
- lithostratigraphy
 comparison of Sites 504 and 896, A:140–141
 secondary mineral distribution, B:59
 Site 504, A:29–30, 32; B:418–419
 Sites 504 and 896, B:437–439
 vs. depth, B:419, 471
- longitudinal splitting, breccias, B:254
- loss on ignition
 vs. depth, B:136
 vs. major oxides, B:139
 vs. oxygen isotopes, B:144
 vs. strontium isotopes, B:144
- maghemitization, black halos, B:180–183
- magma chambers, fractionation, B:17
- magmatic differentiation
 basalts, B:16
 geochemistry, B:39–55
 trends within a solidification zone, B:53
- magmatic minerals, replacement, B:155
- magnesium
 removal, B:114–115
 vs. calcium, A:55
 vs. carbonate trace elements, B:149
 vs. depth, A:55, 128; B:112, 115
 vs. oxygen isotopes, B:148
 vs. silica, A:55
 vs. sodium, B:112
See also iron/(iron + magnesium) ratio
- magnesium/(magnesium + iron) ratio
 amphiboles, B:426
 vs. depth, B:136
 vs. loss on ignition, B:139
 vs. silicon, B:80
 vs. titanium oxide, B:138
- magnesium number
 vs. chromium, B:28
 vs. major elements, A:59, 151
 vs. trace elements, A:59, 156
- magnesium oxide
 electron microprobe data, B:487
 electron microprobe data along basaltic glass
 alteration, B:210
 profiles across microbially processed glass,
 B:200
 vs. alteration, B:49
 vs. anorthite, B:428
 vs. calcium oxide, B:197–199
 vs. calcium oxide and aluminum oxide, B:19
 vs. depth, A:60, 62, 157; B:48, 136, 422, 484
 vs. loss on ignition, B:139
 vs. magnesium number, A:59, 151
 vs. major oxides, B:23, 30–31, 34–35
 vs. number of cations in octahedral sites of
 clay minerals, B:163
- vs. tantalum, B:142
 vs. titanium oxide, B:138
See also iron oxide/(iron oxide + magnesium
 oxide) ratio
- magnetic anomalies
 magnetic logging, B:331–338
 models, A:80
 vs. depth, B:336
- magnetic data
 Site 504, A:71
 Site 896, A:167
- magnetic domains, particle size, B:476–477
- magnetic field
 components vs. depth, B:332–334
 strength vs. depth, A:80
 vs. depth, A:174
- magnetic field logs
 magnetic anomalies, B:331–338
 Site 504, A:78–80, 169–173
 Site 896, A:169–173
- magnetic inclination, vs. depth, A:174; B:222
- magnetic inclination, stable, vs. depth, A:70
- magnetic intensity
 vs. applied demagnetization field, A:72
 vs. depth, A:174
- magnetic oxide mineralogy, upper oceanic crust,
 B:467–482
- magnetic properties
 alteration, B:171–189
 basalts, A:168; B:217–226
 comparison of Sites 504 and 896, B:225
 upper oceanic crust, B:467–482
- magnetic susceptibility
 alteration, B:183
 anisotropy, A:71
 basalts, B:218, 221, 472–473, 477
 vs. depth, B:181, 482
- magnetic susceptibility, bulk
 vs. depth, A:72, 166; B:222, 224
 vs. median destructive field, B:224
 vs. temperature, B:224
- magnetic susceptibility, high-temperature, basalts,
 B:221–223
- magnetic susceptibility anisotropy
 lineation vs. foliation, A:168
 lower hemisphere equal-area projection, A:73,
 170
 Site 896, A:160–161, 169
- magnetite
 background alteration, A:48
 basalts, B:473, 475–477
 electron microprobe data, B:105
 grain size of phenocrysts and intergranular
 material, A:114
 grain-size distribution, A:43–44
 intergranular material, A:42
 isotopic profiles, B:63–64
 major and trace elements, B:475
 photograph, A:41
 secondary minerals, A:45–53
- magnetite, average, vs. depth, A:44
- magnetization, induced
 basalts, A:159–160
 diabase, A:69–70
- major elements
 alteration, B:126–134, 192–196
 basalts, B:22, 390
 diabase, A:57–60; B:39–53
 glass, B:167, 483–487
 magnetic minerals, B:475, 478–479
 magnetite, B:475
 vs. magnesium oxide, B:23
- manganese
 vs. magnesium, B:149
- vs. oxygen isotopes, B:148
- manganese oxide
 vs. alteration, B:49
 vs. depth, A:60, 62, 157; B:48, 137, 423–424
 vs. iron oxide, B:212
 vs. loss on ignition, B:139
 vs. magnesium number, A:59, 151
 vs. sub-basement depth, A:159
- marcasite, secondary minerals, B:173
- massive units, basalts, A:129
- median destructive field
 downhole changes, B:223–224
 vs. depth, B:222
- megacrysts
 diabase and basalt, A:44–45
 percentage vs. depth, A:138
 photograph, A:137
 point-count data, A:187, bp:CD-ROM
- melt inclusions
 chemistry, B:24–26
 experimentally homogenized in olivine, B:35
 experimentally homogenized in plagioclase,
 B:33
- melts, crystallization, B:21–35
- metamorphic rocks, geochemistry, A:57–60
- metamorphism
 diabase dike alteration, B:97–109
 ophiolites, B:431
 Site 504, A:45–53
 Site 896, A:141–150
See also igneous and metamorphic
 geochemistry
- metasomatic zonation, secondary minerals, B:102
- metasomatism, diabase dike alteration, B:97–109
- miarolitic voids
 fillings, B:155
 secondary minerals, B:187
- microbial activity
 basaltic glass, B:207–214
 glass alteration, B:191–206
See also Archaea; bacteria; chain reactions;
 DNA; microorganisms
- microbially processed glass, oxide ratios,
 B:197–200
- microfaults
 dips, A:68
 lower hemisphere projection, A:70
 photograph, A:70
 veins, A:157–158
- microfracture
 diabasic dike rocks, B:235
 Formation MicroScanner imagery, B:240–241
 poles superimposed on a Kamb-contour plot,
 B:235
 structure, B:234–235
- microlites
 petrography, B:172–173
 photograph, B:208, 214
- microorganisms
 basaltic glass alteration, B:196–198
See also Archaea; bacteria; chain reactions;
 DNA; microbial activity
- microstructure, oceanic crust, B:365–374
- microthermometry
 fluid inclusions, B:89–94
See also geothermometry
- mineral chemistry, secondary minerals, B:99–103
- mineral precipitation, hydrofractures, B:250
- mineralogical properties, alteration, B:171–189
- modal composition, vs. depth for sheeted dikes,
 B:414
- MORB. *See* basalts, mid-ocean ridge
- natrolite

- composition, B:124
secondary minerals, A:141
strontium isotopes, B:137
veins, B:443
- natural gamma ray logs
Site 504, A:76–77, 99–109
Site 896, A:180–182
- natural gamma ray–resistivity–velocity logs,
A:178–179
- natural remanent magnetization intensity, vs.
depth, B:222, 224
- neutron absorption cross section, basalts,
B:389–394
- nickel
vs. alteration percentage, A:62
vs. depth, A:61–62, 158; B:15, 19, 48, 137
vs. loss on ignition, B:140
vs. magnesium number, A:59, 156
vs. water content, A:62, 160
- niobium
mobilization, B:47–50
vs. depth, B:464
vs. loss on ignition, B:141
vs. zirconium, B:466
See also zirconium/niobium ratio
- nitrogen
downhole measurements, B:3–7
vs. carbon dioxide, B:6
vs. depth, B:5
- nontronite
chemical composition, B:160–162, 168–170
See also celadonite–nontronite mixtures;
celadonite–nontronite–saponite
mixtures; phyllosilicates
- nucleotide sequences
microbial pits, B:212
See also oligonucleotide probe hybridization
- oceanic crust
alteration, B:423
alteration rates, B:199–200
in situ, B:229–243
magnetic anomalies, B:331–338
structure, B:267–268
well-logging, B:409–414
- oceanic crust, upper
alteration, B:171–189, 435–450
alteration mineralogy, B:151–170
electrical conductivity, B:299–300
gold, B:453–454
hydrothermal alteration, B:57–69, 119–150,
417–434
magnetic oxide mineralogy, B:467–482
permeability, B:353–363
seismic profiles, B:339–347
structural evolution, B:245–259
- oligonucleotide probe hybridization, Archaea,
B:209
- olivine
alteration, A:45–53; B:172–173
background alteration, A:47
chemical composition, B:153–154
percentage vs. depth, A:138
photograph, A:40
See also plagioclase/olivine ratio
- olivine phenocrysts
frequency histogram, A:138
percentage vs. depth, A:38
photograph, A:142, 144
Site 504, A:39–40; B:103
Site 896, A:132, 137–140; B:23, 27, 34–35,
167
- ooze, lithostratigraphy, B:437–439
- ophiolites, alteration, B:431
- oxidation
alteration, A:148–149; B:159
black halos, B:180–183
microbial pits, B:212
pillow lava, A:142
- oxidation ratio
alteration, B:178–180
alteration halos, B:156–157
vs. Curie temperature, B:182
vs. depth, B:182, 423, 440
vs. potassium oxide, B:178
- oxidative alteration
composition, B:108
percentage vs. depth, A:144
- oxygen isotopes
alteration geochemistry, B:132–137
dikes, B:429
hydrothermal alteration, B:57–69
vs. carbon isotopes, B:148
vs. carbonate trace elements, B:148
vs. depth, B:61, 143, 146, 423
vs. iron and volatile content, B:144
vs. strontium isotopes, B:65, 143, 148
whole rock vs. percent alteration, B:64
- packer experiments
permeability, B:357
pressure–time records, B:357
Site 896, A:174–175
- packer inflation, tests, B:359–360
- palagonite
alteration, B:208
chemical composition, B:167
See also fibropalagonite
- paleomagnetism
Site 504, A:68–71
Site 896, A:158–161; B:217–226
- Panama Fracture Zone, geologic setting, A:29;
B:436–437
- parageneses, alteration, B:98
- permeability
heat flow, B:295
in situ vs. calculated bulk, B:355–356
upper oceanic crust, B:353–363
vs. depth, B:354, 361, 363, 420
- permeability, bulk, vs. depth, A:34
- petrography
alteration, B:172–173
diabase, B:42–43
hydrothermal alteration, B:122–123
- petrology, A:32–34
- phenocrysts
chemistry, B:22–24
diabase, A:38–41
mineralogy, A:137–140
percentage vs. depth, A:138
petrology, A:132
point-count data, A:187, bp:CD-ROM
- phillipsite
composition, B:124
photograph, A:149
secondary minerals, A:141; B:154, 173, 186,
208
veins, B:155–156
See also saponite–carbonate–phillipsite veins
- phosphorus oxide
electron microprobe data along basaltic glass
alteration, B:210
microbial pits, B:212
vs. depth, A:60; B:18, 136, 422, 440
vs. loss on ignition, B:139
vs. magnesium number, A:59
vs. uranium, B:142
- phyllosilicates
alteration, B:427–428
characteristics, B:156
veins, B:268–270, 273–274
See also clay minerals; iddingsite
- phyllosilicates, secondary, composition, B:125
- phylogenetic sequences, microbial pits, B:212
- physical properties
basalts, B:376
oceanic crust, B:365–374
relation to strength, B:401–407
Site 504, A:71–73
Site 896, A:161–168
- pillow basalts
alteration, B:419–421
image analysis, B:380–383
lithostratigraphy, B:437–439
photograph, A:145–146
temperature, B:21–35
titanomaghemite, B:479
- pillow lava
carbonate + clay veins, B:264
electron microprobe data, B:485–486
glass alteration, B:191–206
planar and horizontal contact, A:131
rim replacement, B:155
textures, A:130–131
with glassy and chilled rind, A:131
with variolitic zone, A:131
- plagioclase
alteration, B:428
average grain size of groundmass vs. depth,
A:44
average grain size vs. depth, A:49
background alteration, A:47
chemical composition, B:153–154
fluid inclusions, B:91–94
glomerocrysts, A:44–45
grain size of phenocrysts and intergranular
material, A:114
grain-size distribution, A:43–44
intergranular material, A:41
percentage vs. depth, A:138
photograph, A:39; B:208
secondary minerals, B:73–76
sheeted dike complex, B:410–411
thickness of mantle on phenocrysts and mean
grain size of intergranular material,
A:39
zoning, A:39
- plagioclase/olivine ratio
basalts, A:138
percentage vs. depth, A:138
- plagioclase/phenocrysts ratio, vs. phenocrysts,
A:139
- plagioclase, secondary, modal data, B:66
- plagioclase melts, density, A:140
- plagioclase phenocrysts
alteration, B:172–173
composition, A:38–39, 132, 137–140;
B:23–26, 30–31, 33–34, 104
frequency histogram, A:138
percentage vs. depth, A:38
- planar structure
Formation MicroScanner imagery, B:238–240
Rose diagrams of azimuth, B:386
- plumose texture
photograph, A:134
quench textures, A:133
- pore space
alteration minerals, B:372–373
basalts, B:367–368
vs. depth, B:374
- pore-fluid pressure, upper oceanic crust,
B:246–250

- porosity
 alteration, B:422
 basalts, B:376–377
 diabase, B:317–329
 electrical conductivity, B:299–300
 fractures, B:241
 hydrofractures, B:247–250
 image analysis, B:379
 relation to azimuth, B:386
 resistivity, B:309–310
 structure, B:321–322
 thermal parameters, B:398–399
 vs. acoustic velocity, B:377
 vs. bulk density, A:171
 vs. compressional wave velocity, B:372
 vs. depth, A:35, 74, 78, 170, 172; B:310, 321–322, 324–328, 379, 386–388, 400, 420
 vs. formation factor, A:171; B:377
 vs. unconfined compressive strength, B:404
 porosity, bulk, vs. depth, A:34
 porosity, total, vs. porosity derived from acoustic velocity, B:378
 potassium, enrichment, B:159, 200
 potassium, interlayer
 secondary phyllosilicates, B:125
 vs. iron/(iron + magnesium) ratio, B:125
 vs. tetrahedral aluminum, B:125
 potassium oxide
 alteration, B:175
 electron microprobe data along basaltic glass alteration, B:210
 microbial pits, B:212
 profiles across microbially processed glass, B:200
 secondary phyllosilicates, B:125
 vs. depth, A:157; B:12, 136, 422, 440, 484
 vs. iron oxidation ratio, B:178
 vs. iron oxide, B:170
 vs. iron oxide/(iron oxide + magnesium oxide) ratio, B:156
 vs. loss on ignition, B:12, 139
 vs. magnesium number, A:151
 vs. magnesium oxide, B:30–31, 34
 vs. number of cations in octahedral sites of clay minerals, B:163
 vs. octahedral occupancy, B:125
 vs. oxide percentage, B:394
 vs. sodium oxide, B:30, 197–199
 vs. strontium isotopes, B:144
 vs. titanium oxide, B:30, 138
 vs. water content, B:177
 vs. whole-rock trace elements, B:141
 prehnite
 alteration, B:423, 426, 428
 composition, B:108
 pressure
 packer experiments, B:357–360
 vs. time, A:175
 vs. velocity, B:367
 pseudomorphs
 olivine, B:108
 saponite, B:185–186
 pumpellyite, petrography, A:47
 pyrite
 alteration, A:45–53; B:173
 isotopic profiles, B:63–64
 secondary minerals, B:154–155
 vs. depth, B:438
 See also clay + pyrite veins
 pyroxene
 alteration, B:422–423
 intergranular material, A:41–42
 See also augite; clinopyroxene
 pyroxene, quadrilateral component, diabase, B:8
 pyroxene phenocrysts, percentage vs. depth, A:38
 pyrrhotite, isotopic profiles, B:63–64
 quartz
 alteration, A:45–53; B:429
 veins, A:50–51
 quartz–chlorite–sulfide veins, composition, A:63
 quench textures, chilled margins, A:132–134
 radioactivity, natural, vs. depth, B:376, 379
 rare earths
 alteration, B:127–134
 depletion in diabases and patches, B:50
 mobilization, B:47–50
 vs. depth, B:464
 rare earths, chondrite-normalized, in rocks and secondary minerals, B:142
 reaction zones
 composition, B:108
 mixing, B:118
 secondary minerals, B:77, 80–81
 recrystallization
 alteration, B:421
 crack-seal veins, B:284–285
 isotopic profiles, B:63–64
 remanent magnetization, basalts, B:217–226
 remanent magnetization, anhysteretic basalts, A:160; B:471–473, 477
 diabase, A:71
 remanent magnetization, isothermal basalts, B:221, 471–473, 477
 diabase, A:70–72
 intensity, A:72, 166
 remanent magnetization, natural
 alteration, B:178–183
 basalts, A:159; B:472–473, 477
 diabase, A:68–69
 vs. depth, B:482
 See also natural remanent magnetization intensity
 remanent magnetization, saturation
 alteration, B:178–180
 basalts, B:471–472
 downhole changes, B:223
 vs. depth, B:180
 vs. temperature, B:178–180
 resistivity
 basalts, B:377
 fractures, B:308–309, 318–319
 salinity, B:303–304
 vs. depth, A:21, 74, 76, 79, 170, 173; B:309, 319, 376, 424
 See also natural gamma ray–resistivity–velocity logs
 resistivity logs, vs. depth, A:35
 resistivity–velocity logs, A:88–98
 ridge-flank circulation, alteration, B:138–141
 RNA
 ribosomal, microbial activity, B:207–212
 See also DNA
 rock magnetics
 alteration, B:178–180
 basalts, A:160; B:217–226, 471–472
 diabase, A:70–71
 rubidium
 vs. alteration, B:62
 vs. depth, B:137, 464
 vs. loss on ignition, B:140
 vs. magnesium number, A:59, 156
 vs. potassium oxide, B:141
 rutile, basalts, B:473, 475–477
 salinity
 electrical conductivity, B:302–303
 vs. depth, B:92–93
 vs. homogenization temperature, B:91–93
 saponite
 alteration halos, B:445
 chemical composition, B:125, 157–159, 162–163, 165, 168–170
 electron microprobe data, B:124
 fillings, B:155
 glass replacement, B:155
 photograph, A:142, 144, 149
 replacing olivine, B:185
 replacing plagioclase, B:186
 secondary minerals, B:153, 173, 187
 strontium isotopes, B:149
 vs. depth, B:443
 See also celadonite–nontronite–saponite mixtures; celadonite–saponite mixtures; phyllosilicates
 saponite veins
 composition, B:155–156, 442–443
 vs. depth, B:438
 saponite–carbonate–phillipsite veins, petrology, A:145
 saponite–celadonite–iron oxides mixtures, secondary minerals, B:153
 scandium
 vs. depth, A:61, 158; B:464
 vs. magnesium number, A:59, 156
 Schlumberger logs
 Site 504, A:75–77
 Site 896, A:168–169
 seawater, fraction surface, simulation, B:115
 seawater, sodium bromide, A:56–57
 seawater alteration
 basaltic glass, B:211–212
 rates, B:199–200
 secondary minerals
 alteration, B:98, 173
 alteration halos, B:168, 441
 basalts, B:152–155
 geochemistry, B:71–86
 hydrothermal alteration, B:123–124
 identification, A:141
 modal proportions, A:119, bp:CD-ROM; B:66
 variations of mineralogy and style with depth, A:144
 vs. depth, A:144; B:59, 72, 155, 419, 441, 470
 sediments, oceanic crust, B:340
 seismic profiles, pressure–time traces, B:350
 seismic profiles, vertical, upper oceanic crust, B:339–347
 seismic sections, Site 896, A:126
 seismic sources, effect of ship’s hull, B:349–352
 seismic stratigraphy, Site 504, A:33
 seismic structure, layer 2, B:369–370
 seismic waves, amplitude scaling, B:342
 seismograms, oceanic crust, B:340
 semidikes
 petrology, B:103
 See also dike complexes
 sheaf-spherical texture
 photograph, A:135
 quench textures, A:133
 shear wave velocity, vs. confining pressure, B:413
 ship hull, seismic sources, B:349–352
 sigma values, neutron absorption cross section, B:389–394
 silica
 electron microprobe data, B:487
 electron microprobe data along basaltic glass alteration, B:210
 profiles across microbially processed glass, B:200

- vs. alteration, B:49
 vs. depth, A:60, 62, 157; B:48, 136, 422, 484
 vs. iron oxide, B:156
 vs. loss on ignition, B:139
 vs. magnesium, A:55
 vs. magnesium number, A:59, 151
 vs. magnesium oxide, B:31, 34
 vs. major oxides, B:197–199
 vs. titanium oxide, B:138
- silicon
 vs. depth, B:60, 425
 vs. magnesium/(magnesium + iron) ratio, B:80
- silicon logs
 vs. depth, A:173
 vs. iron logs, A:172
- simulation model, borehole fluid mixing,
 B:111–118
- Site 504, A:27–121
 alteration, A:45–53; B:435–450
 background and objectives, A:29–34
 basement, A:32–34
 borehole fluid geochemistry, A:53–57
 coring summary, A:36
 downhole measurements, A:34, 73–81
 drilling history, A:32
 electrical conductivity, B:297–305
 fluid inclusions, B:87–95
 fractures, B:307–315, 317–329
 geochemical comparison with Site 896, B:16
 geochemistry, B:3–7, 39–55, 111–118,
 455–466
 geologic setting, A:29
 gold, B:453–454
 hydrothermal alteration, B:417–434
 igneous and metamorphic geochemistry,
 A:57–60
 igneous petrology, A:37–45
 isotopes, B:57–69
 layer 2/3 boundary, B:409–414
 lithostratigraphy, A:29–30, 32
 magnetic anomalies, B:331–338
 magnetic properties, B:467–482
 metamorphism, A:45–53
 metasomatism and metamorphism, B:97–109
 nitrogen, B:3–7
 operations, A:34–37
 paleomagnetism, A:68–71
 permeability, B:353–363
 physical properties, A:34, 71–73; B:401–407
 regional geology map, A:31
 secondary minerals, B:71–86
 seismic profiles, B:339–347
 seismic stratigraphy, A:33
 site description, A:27–121
 structure and deformation, A:60–61, 63–68
 structure of dike complex, B:229–243
 temperature, B:291–296
 thermal conductivity, B:397–400
- Site 677, concentrations of calcium and
 magnesium vs. depth, A:128
- Site 678, concentrations of calcium and
 magnesium vs. depth, A:128
- Site 896, A:123–192
 alteration, A:141–150; B:171–189, 435–450
 alteration mineralogy, B:151–170
 background and objectives, A:124–128
 bathymetric map, A:125
 coring summary, A:129
 downhole measurements, A:166–175;
 B:375–388
 geochemical logging, B:389–394
 geochemistry, A:150–151; B:9–19
 glass, B:483–487
 heat flow maps, A:127
- hydrothermal alteration, B:119–150
- igneous and metamorphic geochemistry,
 A:150–151
- igneous petrology, A:129–141
 layer 2, B:245–259, 261–279
 magnetic anomalies, B:331–338
 magnetic properties, B:467–482
 microbial activity, B:207–214
 operations, A:128–129
 paleomagnetism, A:158–161; B:217–226
 permeability, B:353–363
 physical properties, A:161–166; B:365–374
 pillow lavas, B:191–206
 site description, A:123–192
 structure and deformation, A:151–158
 veins, B:281–288
- slickenlines, microfaults, A:70
- slickensides, vein faults, B:268
- slug tests, permeability, B:354–355
- smectite
 alteration, A:45–53, B:172–173
 electron microprobe data, B:107
 secondary minerals, A:141
 veins, B:255, 269–270, 273–274
See also chlorite–smectite mixture
- sodium
 vs. depth, B:112
 vs. magnesium, B:112
- sodium bromide, borehole spike, A:56–57
- sodium oxide
 electron microprobe data along basaltic glass
 alteration, B:210
 profiles across microbially processed glass,
 B:200
 vs. alteration, B:49
 vs. depth, A:60, 62, 157; B:14, 48, 136, 422,
 484
 vs. loss on ignition, B:139
 vs. magnesium number, A:59, 151
 vs. magnesium oxide, B:23, 30–31, 34–35
 vs. potassium oxide, B:30
 vs. strontium, B:66
 vs. strontium isotopes, B:144
 vs. titanium oxide, B:30, 138
See also calcium oxide/sodium oxide ratio
- sonic logs, seismic layers, B:346
- spene. *See* titanite
- spinel
 alteration, B:173
 electron microprobe data, B:105
 occurrence log, A:113, 186, bp:CD-ROM
 photograph, A:41
See also ulvospinel
- spinel, chromian
 background alteration, A:48
 chemical composition, B:153–154
- spinel phenocrysts, composition, A:40–41, 132;
 B:23–24, 27–28
- stable isotopes, hydrothermal alteration, B:57–69
- staining, microbial activity, B:210–211
- strength
 physical properties, B:401–407
 vs. electrical conductivity, B:405
- strength, unconfined, compressive
 vs. depth, B:406
 vs. porosity and density, B:404
- stresses
 calibration, B:403
 diabase, B:317–329
 fractures, B:308, 314
 hydrofractures, B:247
 mid-ocean ridges, B:257–258
 physical properties, B:401–407
- strontium
 vs. alteration, B:62
 vs. alteration percentage, A:62
 vs. depth, A:61–62, 158; B:15, 137
 vs. loss on ignition, B:140
 vs. magnesium, B:149
 vs. magnesium number, A:59, 156
 vs. oxygen isotopes, B:148
 vs. sodium oxide, B:66
 vs. strontium isotopes, B:144, 148
 vs. water, A:160
 vs. water content, A:62
- strontium isotopes
 alteration geochemistry, B:132–137
 hydrothermal alteration, B:57–69
 vs. alteration, B:65
 vs. alteration-sensitive geochemical
 parameters, B:144
 vs. carbon isotopes, B:148
 vs. depth, B:61, 143, 146, 150, 423
 vs. oxygen isotopes, B:65, 143, 148
 vs. strontium, B:148
 zeolites, B:149
- structure
 log, A:120, 191, bp:CD-ROM
 sheeted dike complex, B:229–243
- structure and deformation
 Site 504, A:60–61, 63–68
 Site 896, A:151–158
- subophitic textures, photograph, A:42
- sulfide
 alteration, B:173
 background alteration, A:48
See also quartz–chlorite–sulfide veins
- sulfur
 depletion, B:51, 429
 vs. alteration, B:49
 vs. copper, B:140
 vs. depth, A:61, 158; B:48, 61, 137, 423
 vs. loss on ignition, B:140
 vs. recrystallization, B:64
 vs. tantalum, B:142
 vs. uranium, B:142
 vs. zinc, B:140
- sulfur isotopes, vs. depth, B:61, 423
- surface conductivity, porosity, B:320
- talc
 alteration, A:45–53
 electron microprobe data, B:107
 petrography, A:47
 secondary minerals, B:77, 82
 vs. depth, A:52
- tantalum
 vs. depth, B:464
 vs. loss on ignition, B:141
 vs. magnesium oxide, B:142
 vs. niobium, B:466
 vs. potassium oxide, B:141
 vs. sulfur, B:142
 vs. zirconium, B:466
- temperature
 calculated vs. run in olivine phenocrysts, B:34
 calculated vs. run in plagioclase phenocrysts,
 B:31
 crystallization, B:21–35
 downhole measurements, B:291–296
 stresses, B:323
 vs. depth, A:36, 75, 172; B:92–93, 117, 292,
 294–295
 vs. heat capacity, B:399
 vs. time and pressure, B:400
 temperature, borehole, mixing effects, B:115–117
 temperature logs
 Site 504, A:74–75

- Site 896, A:168
textures, alteration, B:172
thermal conductivity
 basalts, B:397–400
 heat flow, B:295
 vs. depth, A:74–75, 170, 172; B:400
 vs. temperature, B:399
thermal diffusivity, vs. temperature, B:399
thermal evolution, plagioclase inclusions, B:91–94
thermal parameters, basalts, B:398–399
thermomagnetic curves, alteration, B:179
Thermus, microbial pits, B:212
tholeiites, phyric olivine, petrology, A:141
thorium
 vs. depth, B:464
 vs. loss on ignition, B:141
 vs. potassium oxide, B:141
 vs. zirconium, B:466
thorium/hafnium ratio, vs. depth, B:464
thorium/uranium ratio, vs. uranium, B:142
titanite
 alteration, B:426
 basalts, B:473, 475–477
 electron microprobe data, B:107
 major and trace elements, B:475
 secondary minerals, B:77
 See also ilmenite–titanite alteration
titanium
 mobilization, B:47–50
 vs. depth, B:60, 425
titanium oxide
 electron microprobe data along basaltic glass
 alteration, B:210
 profiles across microbially processed glass,
 B:200
 vs. alteration, B:49
 vs. alteration percentage, A:62
 vs. depth, A:60, 62, 157; B:15, 18, 48, 136,
 422, 440, 484
 vs. iron oxide, B:197–199
 vs. loss on ignition, B:139
 vs. magnesium number, A:59, 159
 vs. magnesium oxide, B:23, 30–31, 34–35, 53
 vs. major elements, B:138
 vs. oxide percentage, B:394
 vs. potassium oxide, B:30
 vs. silica, B:197–199
 vs. sodium oxide, B:30
 vs. sub-basement depth, A:159
 vs. water content, A:62, 160
titanium oxide + iron oxide
 vs. copper, B:140
 vs. sulfur, B:140
titanomaghemite
 alteration, B:178–183
 backscattered electron images, B:479–480
 basalts, A:160; B:473, 475–477
 magnetic susceptibility, B:221–223
 major and trace elements, B:478–479
titanomagnetite
 background alteration, A:48
 basalts, A:160; B:473, 475–477
 chemical composition, B:153–154
 magnetic susceptibility, B:221–223
 secondary minerals, B:173, 180–183, 188–189
tortuosity, porosity, B:320
trace elements
 alteration, B:127–134
 basalts, B:15–16, 390, 393
 diabase, A:57–60; B:39–53
 lower sheeted dike complex, B:455–466
 magnetic minerals, B:475, 478–479
 magnetite, B:475
 titanite, B:475
tracer experiments, borehole spike, A:56–57
transmissivity, packer experiments, B:356
traveltime
 oceanic crust, B:341–343
 vs. depth, A:81, 85–87; B:342–343
tritium
 mixing, B:117–118
 predicted vs. measured in borehole fluids,
 B:112
ulvospinel
 titanomaghemite, B:479
 See also spinel
uralitization, chemical effects, B:49
uranium
 alteration geochemistry, B:131
 vs. iron, B:142
 vs. loss on ignition, B:141
 vs. phosphorus oxide, B:142
 vs. potassium oxide, B:141
 vs. sulfur, B:142
 vs. thorium/uranium ratio, B:142
 See also thorium/uranium ratio
vanadium
 vs. alteration percentage, A:62
 vs. depth, A:61–62, 158; B:15, 19, 137
 vs. loss on ignition, B:140
 vs. magnesium number, A:59, 156
 vs. sub-basement depth, A:159
 vs. water content, A:62, 160
variolitic texture
 photograph, A:133–134
 quench textures, A:133
 See also glassy/variolitic zone
vein faults
 equal-area stereonet, B:268
 structure, B:265–266
vein networks, structure, B:250–251
vein orientations, structure and deformation,
 A:157
veinlets, photograph, A:147
veins
 alteration, B:426
 comparison of Sites 504 and 896, B:271–272
 composition, A:50–53
 composition and abundance vs. depth, A:147
 crosscutting relations, B:263
 deformation, A:63–66
 dikes, B:97–109
 dip angles vs. depth, B:271
 distribution of true dips for a randomly
 oriented set of planes, A:165
 equal-area stereographic projections, B:265
 evolution, A:163
 faults and fractures, A:157–158
 fluid inclusions, B:94
 formation mechanism, A:66
 Formation MicroScanner imagery, B:237
 histogram of veins greater than 22 mm wide,
 A:147
 hydrothermal alteration, B:123
 isotopes, B:65, 145
 Kamb-contour stereographic equal-area
 projection, A:164
 log, A:118, 190, bp:CD-ROM
 mechanism of formation, A:154
 microstructural characteristics, A:63–64
 mineral composition, B:155–156, 421–425,
 427–428
 morphology, A:153–154
 number logged per meter of core recovery,
 A:52
 occurrence vs. depth, A:65
 orientation and distribution, B:263–265
 orientation with stable paleomagnetic
 declination, A:63
 petrology, A:46, 144–146
 photograph, A:51, 164–165; B:102, 253–254
 poles superimposed on a Kamb-contour plot,
 B:233
 relations in thin section, A:52
 structure, B:231–233, 250–257, 261–266
 structure and deformation, A:151–155
 thin sections, A:64–65
 true dip histogram, A:164
 variations with depth, A:146; B:270–271
 vs. depth, B:438–443
 vs. proportion of massive units, B:442
 veins, crack-seal, structure, B:256–257, 281–288
 veins, fibrous
 photograph, B:287–288
 stereographic projections, A:165; B:282
 stereographic projections and Kamb-contour
 diagrams, B:267
 structure, B:256–257, 261–262
 structure and deformation, A:152–153
 true dip histogram, A:164
 types, B:282–284
 veins, nonfibrous
 structure and deformation, A:152
 true dip histogram, A:164
 veins, volume percent, vs. depth, A:49
velocity
 basalts, B:366–368
 oceanic crust, B:409–410
 vs. confining pressure, B:367
 vs. depth, A:81; B:321–346, 370
 See also natural gamma ray–resistivity–
 velocity logs; resistivity–velocity logs
velocity, acoustic
 basalts, B:376
 vs. depth, B:376
 vs. porosity, B:377
vertical seismic profile logging, Site 504, A:80–81
vesicles, fillings, B:155
volcanic rocks
 alteration, B:419–421, 435–450
 nitrogen geochemistry, B:3–7
volcanic stratigraphy
 comparison of Sites 504 and 896, A:141
 Site 896, A:129–132
 vugs, fillings, B:155
wairakite, composition, B:108
water content
 alteration, B:175, 177
 diabase, B:4
 glasses, B:30
 vs. alteration, B:49
 vs. depth, B:5, 48, 423
 vs. iron oxides, B:170
 vs. oxygen isotopes, B:144
 vs. potassium oxide, B:177
 vs. strontium isotopes, B:144
 vs. trace elements, A:62, 160
water gun shots, pressure–time traces, B:350
wavefield imaging, basement, B:343–344
weathering, glass, B:191–206
well-logging
 Formation MicroScanner imagery, B:375–388
 oceanic crust, B:409–414
 Site 504, A:73–81
 Site 896, A:166–175
 See also different kinds of logs; geophysical
 combination logging
Young's modulus, vs. unconfined compressive

- strength, B:404
- yttrium
- vs. alteration percentage, A:62
 - vs. depth, A:61–62, 158; B:16, 19, 137
 - vs. loss on ignition, B:140
 - vs. magnesium number, A:59, 156
 - vs. sub-basement depth, A:159
 - vs. water content, A:62, 160
- zeolite veins, vs. depth, B:438
- zeolites
- chemical composition, B:108, 124, 166
 - secondary minerals, B:154, 426
 - strontium isotopes, B:149
 - veins, B:442–443
 - See also* phillipsite; individual minerals
- zeolites, fibrous, photograph, A:146
- zinc
- depletion, B:51, 429
 - diabase, B:4
 - relation to gold, B:453–454
 - vs. alteration, B:49
 - vs. copper, B:53
 - vs. depth, A:61–62, 158; B:5, 48, 61, 137, 423, 454
 - vs. loss on ignition, B:140
 - vs. magnesium number, A:59, 156
 - vs. sub-basement depth, A:159
 - vs. sulfur, B:140
 - vs. titanium oxide + iron oxide, B:140
 - vs. water content, A:62, 160
- zirconium
- mobilization, B:47–50
 - vs. alteration, B:49
 - vs. alteration percentage, A:62
 - vs. depth, A:61–62, 158; B:16, 18, 48, 137
 - vs. loss on ignition, B:1400
 - vs. magnesium number, A:59, 156
 - vs. magnesium oxide, B:53
 - vs. niobium, B:466
 - vs. sub-basement depth, A:159
 - vs. tantalum, B:466
 - vs. thorium, B:466
 - vs. water content, A:62, 160
- zirconium/hafnium ratio, vs. depth, B:464
- zirconium/niobium ratio, vs. depth, B:464
- zoning
- fractures, B:312–314
 - plagioclase, A:39, 134–136
 - zoning, complexed, photograph, A:136
 - zoning, oscillatory, photograph, A:135
 - zoning, sector, photograph, A:136