

OCEAN DRILLING PROGRAM

LEG 194 SCIENTIFIC PROSPECTUS

MARION PLATEAU

Dr. Alexandra Isern
Co-Chief Scientist
Ocean Studies Board
National Research Council
2101 Constitution Avenue, NW
Washington, DC 20418
USA

Dr. Flavio Anselmetti
Co-Chief Scientist
Geologie
Dept. Erdwissenschaften (D-ERDW)
NO H 50
Sonneggstrasse 5
ETH Zentrum
CH-8092 Zurich
Switzerland

Dr. Jack Baldauf
Deputy Director of Science Operations
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
USA

Dr. Peter Blum
Leg Project Manager and Staff Scientist
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
USA

July 2000

Material in this publication may be copied without restraint for library, abstract service, educational, or personal research purposes; however, this source should be appropriately acknowledged.

Scientific Prospectus No. 94

First Printing 2000

Distribution

Electronic copies of this publication may be obtained from the ODP Publications homepage on the World Wide Web at: <http://www-odp.tamu.edu/publications>

This publication was prepared by the Ocean Drilling Program, Texas A&M University, as an account of work performed under the international Ocean Drilling Program, which is managed by Joint Oceanographic Institutions, Inc., under contract with the National Science Foundation. Funding for the program is provided by the following agencies:

Australia/Canada/Chinese Taipei/Korea Consortium for Ocean Drilling
Deutsche Forschungsgemeinschaft (Federal Republic of Germany)
Institut National des Sciences de l'Univers-Centre National de la Recherche Scientifique (INSU-CNRS; France)
Ocean Research Institute of the University of Tokyo (Japan)
National Science Foundation (United States)
Natural Environment Research Council (United Kingdom)
European Science Foundation Consortium for the Ocean Drilling Program (Belgium, Denmark, Finland, Iceland, Ireland, Italy, The Netherlands, Norway, Portugal, Spain, Sweden, and Switzerland)
People's Republic of China

D I S C L A I M E R

Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the views of the National Science Foundation, the participating agencies, Joint Oceanographic Institutions, Inc., Texas A&M University, or Texas A&M Research Foundation.

This Scientific Prospectus is based on precruise JOIDES panel discussions and scientific input from the designated Co-chief Scientists on behalf of the drilling proponents. The operational plans within reflect JOIDES Planning Committee and thematic panel priorities. During the course of the cruise, actual site operations may indicate to the Co-chief Scientists and the Operations Manager that it would be scientifically or operationally advantageous to amend the plan detailed in this prospectus. It should be understood that any proposed changes to the plan presented here are contingent upon approval of the Director of the Ocean Drilling Program in consultation with the Science and Operations Committees (successors to the Planning Committee) and the Pollution Prevention and Safety Panel.

Technical Editors: Karen K. Graber and Lori J. Cagle

ABSTRACT

Cretaceous rifting in the western Coral Sea (offshore northeast Australia) formed continental fragments that are now capped by carbonate platforms. The Marion Plateau carbonate platform, which has grown on one of these fragments, provides a natural laboratory to study the causes, magnitudes, and effects of sea-level change on continental margin sediments. One of the fundamental controls on the nature and geometry of continental margin sediment deposition is sea level; however, much of the information on the relationship between sea-level and depositional facies is qualitative. Leg 194 coring will provide a superb and unique opportunity to determine the absolute magnitude of the major Cenozoic sea-level falls.

The drilling strategy outlined for Leg 194 utilizes the stratigraphic relationship between a lower to middle Miocene second-order highstand carbonate platform complex and an upper Miocene second-order lowstand platform complex to establish the magnitude of the middle Miocene N12-N14 sea-level fall. An important characteristic of this platform relationship is that the proposed sites are essentially located along a single strike line without intervening structural elements. Thus, subsidence of the platform will have affected all sites equally, enabling determination of the true amplitude of the sea-level fall that caused a shift in the locus of carbonate platform deposition.

The carbonate platforms and adjacent slopes of the Marion Plateau also preserve a superb record of third-order sea-level variations within a mixed carbonate-siliciclastic depositional environment. High-resolution seismic data collected for the Leg 194 site surveys provide quasi-three-dimensional images of Oligocene-Pliocene depositional geometries. The correlation of these seismic images with drill core and logging data will provide a synoptic view of depositional processes in a mixed carbonate-siliciclastic carbonate platform setting.

INTRODUCTION

Importance of Determining the Magnitude of Eustatic Sea-Level Variations

Measuring the amplitude and timing of eustatic sea-level fluctuations is essential both for the establishment of an accurate eustatic sea-level curve for the Phanerozoic and for the accurate interpretation of sediment sequences on continental margins. Several attempts have been made to determine the amplitude of glacioeustatic fluctuations, including passive-margin sequence stratigraphy (Vail et al., 1977; Vail and Hardenbol, 1979; Haq et al., 1987); modeling of sedimentary depositional regimes (Watts and Thorne, 1984); calibration of the oxygen isotope curve (Majors and Mathews, 1983; Miller et al., 1987; Williams, 1988); and analysis of the depositional history of carbonate sediments on atolls (Schlanger and Premoli-Silva, 1986; Halley and Ludwig, 1987; Moore et al., 1987; Lincoln and Schlanger, 1987, 1991). These analyses yield a wide range of results, and although the different independent data sets often agree with regard to the timing of sea-level events, significant differences between estimates for the magnitude of sea-level fluctuations remain. The establishment of a eustatic sea-level curve has major implications for global stratigraphic correlation and basin analysis, and defining the amplitude of such a curve remains one of the major challenges in sea-level research (COSOD II, 1987; Sahagian and Watts, 1991; JOIDES Planning Committee, 1996). The excellent record of Miocene sea-level fluctuations preserved in the carbonate platforms of the Marion Plateau, southern Coral Sea (Figs. 1, 2), provides an ideal opportunity to test sea-level models and quantify the magnitude of eustatic variations.

Growth Phases of the Marion Plateau and their Record of Sea-Level Variations

Carbonate platforms and their slopes are sensitive indicators of sea-level variations, as they predominantly record growth during sea-level highstands and shutdown during sea-level lowstands. Sampling through carbonate platforms records sea-level effects in a "dipstick" fashion. On the other hand, sediments on platform margins and slopes record sea-level variations as alternations of shallowing and deepening sequences. The geometric relationships between the carbonate platforms and adjacent slope sediments of the Marion Plateau have been clearly imaged by seismic data, enabling the investigation, correlation, and dating of sediment sequences. The information recovered from Leg 194 sites (Fig. 2) will provide an independent basis for development and assessment of the global sea-level curve.

The lower-middle Miocene MP2 platform appears to have formed as a series of transgressive and highstand system tracts (Figs. 3, 4). Five highstand events are recorded by MP2 (MP2a - MP2e),

providing a record of third-order sea-level variations. Only MP2e was sampled during Leg 133, and thus the age at which the other events occurred is not known (Davies, McKenzie, and Palmer-Julson, 1991). Newly acquired site survey seismic data indicates that MP2 prograded over its former slope sediments. These pulses of progradation are strongly controlled by sea-level fluctuations, as was shown for the progradation of the Great Bahama Bank (Eberli and Ginsburg, 1989; Eberli, Swart, and Malone, et al., 1997).

The upper Miocene MP3 platform began to form during a lowstand on the outer slope sediments of MP2. The MP3 phase subsequently evolved into a series of highstand systems tracts but remained structurally lower than the top of MP2 for most of its history (Fig. 3). The upper Miocene MP3 platform records four sea-level cycles (MP3a-MP3d; Fig. 4). The sea-level rise during MP3d corresponds to the last phase of platform growth. This rapid sea-level rise, in conjunction with other environmental factors, resulted in the drowning of most of the MP3 platform (Pigram et al., 1992).

At present, it is difficult to compare the growth phases seismically imaged within the MP2 and MP3 platforms to global events, as the exact timing of their development will not be known until the sequences are cored. In addition, the internal structure of the MP3 platform, whose top is exposed with a nondepositional hardground at the seafloor, shows no internal seismic structure because of the nonpenetrating seismic signal and the probably well-cemented and homogeneous lithology. However, the recovery and dating of sediments from these sequences will provide important information on Miocene sea-level events and their influence on continental margin sedimentation. Data from these sediments may also be used in conjunction with other "sea-level" legs cored as part of the Ocean Drilling Program (ODP) global sea-level strategy.

To determine the sea-level event stratigraphy on the Marion Plateau through drilling, it will be necessary to establish

1. The depositional history of the Miocene carbonate platforms (Fig. 3) of the Marion Plateau by
 - establishing a detailed chronostratigraphy for each platform phase;
 - determining the depositional environment of each platform phase;
 - determining the age and duration of each unconformity;
 - inferring the paleowater depth of each phase; and
 - establishing the total thickness of each platform.

2. The amplitude of the middle Miocene (N14-N12) sea-level fall by determining
 - the age, depth, and paleowater depth of the older (MP2) platform; and
 - the age, depth, and paleowater depth of the initial phase of the younger (MP3) platform.

To establish the magnitude of the sea-level fall that led to the formation of the lowstand MP3 platform, it is first necessary to determine the paleowater depth of the top of the lower-middle Miocene MP2 platform (Fig. 3). Leg 133 coring (Sites 816 and 826) showed that the top of this platform consisted of a tropical reefal assemblage deposited in water depths no greater than 20 m (Davies, McKenzie, Palmer-Julson, et al., 1991). This depth defines the approximate point from which sea level began to fall (Pigram et al., 1992). Sampling evidence indicates that the top of MP2 has been subjected to subaerial exposure. The dissolution and erosion that is likely to have resulted from exposure would have made the present-day top of MP2 lower than it was originally, introducing an error in determining the highest position of sea level immediately before the fall. The extent of erosion is difficult to quantify, but it can be expected that the loss would be small because the high diagenetic potential of these tropical carbonates would tend to create a carbonate pavement that would be difficult to erode. Any sediment loss from the top of MP2 will result in the underestimation of the true amplitude of sea-level fall (Pigram et al., 1993).

The low-point of the sea-level fall is defined by the paleowater depth at the time the first sediments of the upper Miocene lowstand MP3 platform were deposited (Pigram et al., 1992; Fig. 2). The MP3 platform was not sampled during Leg 133 drilling, and therefore the biofacies that compose this platform can only be inferred seismically. The seismic characteristics of MP3 are poorly imaged and difficult to assess, but appear to have both "tropical" (vertically accreted) and "temperate" (mound like) signatures. The presence of cooler water fauna would indicate that the depth of platform initiation was deeper than that for purely tropical carbonate. Without sampling the MP3 platform, we can only speculate on the paleowater depth of the MP3 formation. Three possible scenarios are

- If MP3 is entirely tropical, its formation depth was likely to be ~20-25 m, resulting in a N12-N14 sea-level fall of 185-190 m. This eustatic change is greater than other estimates for this time interval (30-90 m, Miller et al., 1987; >100 m, Vail and Hardenbol, 1979).
- If MP3 is subtropical in composition, the depth of initiation could have been ~50-70 m. Thus, the

N12-N14 sea-level fall would be ~135-155 m.

- It is also possible that sea level fell below the level on which MP3 was established, thus also affecting the estimate of sea-level change, but no seismic evidence supports this conclusion (Pigram et al., 1992).

The Influence of Subsidence on Sea-Level Magnitudes

The inability to remove tectonic subsidence effects from relative sea-level signatures has hindered the quantification of eustatic sea-level variations in many areas. However, a sea-level shift that occurs between two sites of equal tectonic subsidence will provide an accurate record of the magnitude of eustatic change.

For the difference between the top of MP2 and the initiation of MP3 to be an accurate measure of the N12-N14 eustatic fall, it is necessary to demonstrate that there is no differential subsidence along the drilling transect. There are two lines of evidence to support this. First, the Marion Plateau is not structurally compartmentalized and therefore behaves as a single structural entity (Symonds et al., 1988). Seismic lines between the proposed sites show that there are no structural elements, such as faults, between the sites that could cause them to have relative differential subsidence. Second, because the Marion Plateau basement surface is planated with minimal dip to the northeast, depths to basement surface contours can be considered isosubsidence lines (Fig. 5). The eight proposed sites are all near the 1-s basement contour, indicating a minimal basement gradient between the sites and thus negligible differential subsidence.

Calibration of eustatic sea-level variations can only be realistically estimated on slowly subsiding, structurally well-understood margins where an accurate tectonic subsidence history can be established and where sites of equal tectonic subsidence, which have both the highstand and the lowstand history preserved, can be located. The advantage of such areas is that although falling sea level follows the slow tectonic subsidence of the platform, the relative depth change recorded between two sites is self-correcting because they both subside by the same amount. For the predicted middle Miocene Marion Plateau subsidence rates, the increase in water depth at both sites as a result of tectonic subsidence (<10 m) is an order of magnitude less than the eustatic sea-level change over the same interval. The tectonic component of sea-level change is therefore within the error of paleowater depth estimates achievable in these sediments.

BACKGROUND

Geologic Setting of the Marion Plateau

The Marion Plateau is located between 18°S and 23°S, seaward of the south central Great Barrier Reef on the northeastern Australian continental margin. This plateau is the most southerly of the northeast Australian marginal plateaus, forming a deeper extension of the Queensland continental shelf. The plateau is bounded along its northern margin by the Townsville Trough; by the Cato Trough along the eastern margin, and by the south central Great Barrier Reef to the west (Fig. 6). The Marion Plateau is part of a slowly subsiding margin. It is believed that the plateau top remained exposed throughout much of the Paleogene and became planated to form a gently northward-dipping, relatively smooth plateau surface (Pigram et al., 1993).

Tectonics of the Marion Plateau

The eastern Coral Sea has been affected by two distinct tectonic events. The earlier event, late Jurassic-Early Cretaceous in age, was responsible for the formation of the Queensland and Townsville Basins, which underlie the present-day bathymetric features of the Queensland and Townsville Troughs (Fig. 6). These basins formed because of oblique extension along pre-existing Paleozoic structural trends (Struckmeyer and Symonds, 1997). The Queensland and Townsville Basins do not appear to have been affected by the later tectonism responsible for seafloor spreading in the Tasman and Coral Sea Basins (Struckmeyer and Symonds, 1997).

In the Late Cretaceous, rifting in the Coral Sea Basin created numerous continental fragments, which are now capped by carbonate platforms, such as the Marion and Queensland Plateaus (Fig. 6). Rifting in the Coral Sea was an extension of Late Cretaceous (80 Ma) seafloor spreading in the Tasman Basin, which extended to the north to form the Cato Trough and the Coral Sea Basin by 65 Ma (Fig. 6; Weissel and Hayes, 1971; Hayes and Ringis, 1973; Shaw, 1978). Spreading is believed to have ceased along the length of this system by the earliest Eocene (52 Ma; Gaina et al., 1999). Thus, the main physical elements of the western Coral Sea were in place by the early Tertiary (Davies et al., 1989). Although the exact structural style and development history of the rift system is still not completely understood, it is clear that the late Jurassic-Early Cretaceous rifting event controlled the gross architecture of the margin in addition to the form of the high-standing structural elements on which the carbonate platforms in the area are located.

The Marion Plateau is a largely undeformed basement block with faults occurring only on its margins. Basement along the northern margin consists of gently dipping ramps that gradually deepen toward the Townsville Trough until a fault is encountered. Normal extensional faults along this northern margin are restricted to the edge of the plateau and include both down-to-basin faults with dips to the north-northwest and normal faults of opposite polarity that dip beneath the plateau (Symonds, et al., 1988). The eastern margin of the plateau is free of major structural offsets (Mutter and Karner, 1980), and the slope is apparently simple and continuous. Faults along this margin are steeply dipping to vertical and the margin of the plateau downfaults into the Cato Trough. The southern part of the plateau is formed by a southeasterly plunging, gently arched basement high. The top of the arch is unstructured, and faults are confined to the flanks of the arch and appear on conventional seismic data to be high-angle, down-to-basin normal faults (Pigram et al., 1993).

The basement of the Marion Plateau is likely to be similar to that of the Queensland Plateau to the north. This basement was cored during Leg 133 (Sites 824 and 825; Fig. 1) and consists of fine-grained, dark gray, poorly foliated, well-lithified quartz-feldspar-mafic metasediment or metavolcanic rocks. These rocks are similar to those found in the onshore Queensland Hodgkinson Province (Ordovician-Devonian), which outcrops as part of the northern Tasman Fold Belt (Feary et al., 1993). Planation of the surface occurred during subaerial exposure in the Mesozoic and Paleogene prior to the deposition of Megasequence A.

As stated previously, no direct sampling of the basement under the Marion Plateau has occurred, but seismic data and a recently developed plate model indicate that basement crustal blocks of the Queensland and Marion Plateaus had roughly similar tectonic histories in regard to rifting and extension (Gaina et al., 1999; Struckmeyer and Symonds, 1997). The presence of shallow-water (~20 m) carbonate sediments directly overlying basement at Sites 824 and 825 on the Queensland Plateau indicates that the planated basement surface of the Queensland Plateau was at or near sea level immediately prior to the onset of sedimentation. Using this information, we can estimate the thickness of the crust under the Queensland Plateau. Assuming average crustal density, the upper surface of a 30-km-thick crust would exist at sea level. Tectonic subsidence models show almost no change in the depth of the Queensland Plateau surface between 25-10 Ma (Fig. 7). Thus, we can conclude that the early Tertiary opening of the Coral Sea resulted in little crustal thinning on the Queensland Plateau. Otherwise, we would not observe thermal subsidence rates equal or close to zero in the early Tertiary.

These results also show that thermal subsidence from the earlier Late Jurassic-Early Cretaceous event could no longer be detected on the Queensland Plateau during the Neogene.

Subsidence History of the Marion Plateau

The tectonic histories of the Marion and Queensland Plateaus are well constrained by Leg 133 sites and extensive multichannel seismic data. Subsidence curves for these plateaus have been produced using both benthic foraminifera (Fig. 8; Katz and Miller, 1993) and geohistory modeling (Fig. 7; Müller et al., 2000). Geohistory models were calculated using integrated geophysical logs, biostratigraphic/lithologic information, and seismic reflection data (Müller et al., 2000). These models predict post-9-Ma subsidence of 1300 ± 200 m in the Queensland Trough and 650 ± 200 m on the western margin of the Queensland Plateau and post-5-Ma subsidence of 500 ± 30 m on the southern margin of the Queensland Plateau and 660 ± 50 m on the northern margin of the Marion Plateau (Fig. 7; Müller et al., 2000). Although the Marion and Queensland Plateaus are located on a passive margin ~1000 km south of the Pacific-Australian plate boundary, geohistory models predict a greater amount of post-9-Ma subsidence than simple elastic models do. This subsidence occurred in pulses between 9 and 5 Ma on both plateaus. It is difficult to account for this observed subsidence, either by means of thrust loading in Papua New Guinea or by a combination of such thrust loading and in-plane stresses originating from collision along the Australian-Pacific plate boundary (Müller et al., 2000).

Müller et al. (2000) suggest that the observed post-9-Ma tectonic subsidence of the Queensland and Marion Plateaus and Queensland Trough is largely caused by dynamic surface topography resulting from Australia's northeastern margin overriding a slab burial ground and modulated by flexural deformation resulting from collision tectonics north of Australia. This conclusion is supported by shear-wave tomography data (Zhang and Tanimoto, 1993) that shows a north-northwest to south-southeast trending band of anomalously high velocities in the upper mantle at depths between 300 and 650 km. Although unproven, this explanation appears to be the most reasonable for all available data.

Post-9-Ma subsidence rates on the Marion Plateau are much lower than those of third-order sea-level changes and can thus be differentiated from glacial eustasy. In addition, any unaccounted subsidence will be the same for all sites along the transect as they were selected along lines of equal subsidence. Although we will attempt to quantify in detail the additional water depth added to all sites as a result

of tectonic subsidence, this increase is likely to be less than 10 m between N12-N14 and thus will not greatly affect our attempts to quantify eustatic sea-level variations.

Stratigraphy of the Marion Plateau: Evidence from Prior Drilling

Stratigraphies for the Marion Plateau were obtained during ODP Leg 133 (Fig. 9), and these data supplement previously acquired extensive seismic surveys over the plateau (Fig. 10). Both of these data sets have enabled a description of the Marion Plateau depositional history. Initiation of shallow marine carbonate sedimentation on the Marion Plateau began during the latest Paleogene as the sea transgressed across the metasedimentary basement of the plateau (Davies, McKenzie, Palmer-Julson, et al., 1991). We have no direct age controls on the sediments that form Megasequence A as they have not yet been sampled. Their age is inferred by their stratigraphic position under Megasequence B (Pigram et al., 1993; Figs. 3 and 10). These first sediments over basement are believed to be primarily siliciclastics, with temperate water carbonates occurring in the eastern part of the sequence.

Sedimentary facies recovered during Leg 133 and their correlation to seismic profiles indicate that tropical reef development was initiated on the Marion Plateau in the early Miocene, and by the middle Miocene there was extensive reef growth on the plateau (Davies, McKenzie, Palmer-Julson et al., 1991). These reefal sediments are part of Megasequence B and include the aggrading and prograding MP2 carbonate platform (Figs. 3, 10).

In the late middle Miocene, carbonate bank productivity rapidly diminished on the Marion Plateau, as shown by a reduced fine-grained bank-derived component in slope sediments. This decline was primarily the result of subaerial exposure resulting from a sea-level regression that caused the demise of the MP2 platform. During the low sea-level interval between 11 and 7 Ma, the MP3 platform was initiated on the eastern side of the Marion Plateau. Despite the fact that MP3 developed during a lowstand, the platform continued its development during subsequent highstand intervals. MP2, on the other hand, did not reinitiate even after being reflooded during the subsequent sea-level increases. During the development of MP3 the western two-thirds of the Marion Plateau was exposed, forming a broad, low-relief karstic surface. Unlike MP2, MP3 has not completely drowned but is now restricted to the area of Saumarez Reef. The limited sampling of MP3 inhibits speculation on the cause for partial drowning, although some likely factors are reduced sea-surface temperatures (Isern et al., 1996) and increased terrigenous inputs causing increased water-column turbidity (Pigram et al., 1993).

Carbonate production from the Pliocene to Holocene never again achieved the areal extent that existed in the Neogene. Instead, hemipelagic drift sediments dominated sedimentation on the Marion Plateau.

SCIENTIFIC OBJECTIVES

The Marion Plateau provides ideal targets to address the causes, magnitudes, and effects of sea-level change on continental margin sediments. Although one of the fundamental controls on the nature and geometry of continental margin deposition is sea-level fluctuations, much of the information on the relationship between sea-level and depositional facies is qualitative.

Extensive seismic surveys over the Marion Plateau have enabled the description of the various depositional sequences found on the plateau (Figs. 2, 10). These data, along with sedimentological information from Leg 133 cores and dredge samples, have been used to develop a growth model for carbonate platform phases on the Marion Plateau and the relationship of these platforms to sea-level change (Fig. 3). Coring and downhole measurements during Leg 194 will address the following objectives:

- *Magnitude of the N12-N14 second-order eustatic sea-level fall*

Although difficult, determining the magnitude of eustatic sea-level fluctuations is significant not only for the establishment of a Phanerozoic sea-level curve but also for understanding the stratigraphic response of continental margin sediments to sea-level forcing. Leg 194 coring provides an opportunity to determine the absolute magnitude of one of the major Cenozoic sea-level falls.

- *Oligocene-Pliocene, third-order sea-level fluctuations*

Coring will recover an Oligocene-Pliocene sea-level record preserved in the carbonate platform growth phases of the Marion Plateau. Seismically, this record appears to include a complete third-order event stratigraphy from 30 to 4 Ma within the second-order sea-level falls that dominate the sequence stratigraphic framework of the Marion Plateau. Analyses of these variations, and the higher order fluctuations contained within them, will provide

information on the timing and influence of sea level on the carbonate growth phases of the Marion Plateau.

- *Sedimentary facies change and development of sequence stratigraphic units controlled by sea-level changes*

Leg 194 coring will recover a detailed record of carbonate and siliciclastic sediment facies variations resulting from mixing of sediment from carbonate banks on the Marion Plateau and the continental margin of Australia. In general, carbonate sediment export increases during sea-level highstands as carbonate banks have additional accommodation space for increased growth of carbonate-producing organisms. On the other hand, terrigenous sedimentation generally decreases during highstands because of the elevated erosional base level. Leg 194 coring will enable detailed analysis of the cumulative result of these different sedimentological responses to sea-level forcing.

- *Impact of potential sequence of temperate to subtropical carbonates*

It is likely that the MP3 platform is at least partially composed of temperate to subtropical carbonates. Thus, the results of this coring will rely on and extend the results of Leg 182 (cool-water carbonates of the Great Australian Bight). Stable isotopic data from Leg 133 Site 811 showed that during the late Miocene, sea-surface temperatures (SST) were cool (~20°-22°C), as were global sea-surface temperatures (Isern et al., 1996). Given the more southerly location of the MP3 platform with respect to Site 811 (Queensland Plateau), temperatures were probably similar to, if not cooler than those over the Queensland Plateau. Sea-surface temperatures at or below 20°C would not prevent tropical coral growth but would make it more likely that the MP3 platform was constructed of a “cooler,” more sub-tropical bioassemblage. If MP3 is indeed composed of cooler water fauna, the depth of platform initiation probably would be deeper than for purely tropical carbonate. Should the early growth phases of MP3 be temperate in nature, documenting the transition of these cooler water forms into the tropical forms existing for most of the MP3 growth phase will be an important outcome of Leg 194.

- *Fluid flow and diagenesis within pure carbonate and mixed siliciclastic/carbonate depositional environments*

The mechanisms, rates, and distributions of fluid transport through carbonate platforms and reef structures are critical to the understanding of diagenetic processes (Buddemeier and Oberdorfer, 1986) and for the geochemical cycling of many elements. Fluid movements have the ability to chemically alter the mineralogic composition of the sediment by converting metastable minerals such as high-Mg calcite and aragonite to more stable calcite and dolomite (Mullins et al., 1984; Simms, 1984). Alteration of carbonate sediments to dolomite has been significant in both the Bahamas (Varenkamp et al., 1991) and the carbonate platforms of northeast Australia (McKenzie et al., 1993; Davies, McKenzie, Palmer-Julson, et al., 1991). Recent studies have shown that carbonate sediments off northeast Australia were dolomitized by multigenerational fluids flowing through the platforms (McKenzie et al., 1993). The movements of fluids through the Queensland Plateau were demonstrated by using $^{87}\text{Sr}/^{86}\text{Sr}$ isotopic ratios and the Sr composition of interstitial waters (Elderfield et al., 1993).

Fluid flow can also alter the sedimentary structure, permeability, and porosity of the deposit, thus having important effects on flow pathways and reservoir potential. The existence of fluid flow has been described in tropical carbonate platforms such as the Great Bahama Bank and the Queensland Plateau (Eberli et al., 1997; Elderfield et al., 1993) and also in temperate water carbonates (Feary, Hine, Malone, et al., 2000). However, the mechanisms causing this flow are neither well documented nor understood.

- *Role of climatic and paleoceanographic change in the tropical South Pacific and its influence on carbonate platform development.*

Leg 194 coring will enable the investigation of paleoceanographic variations in the western Coral Sea and will provide the opportunity to correlate these variations with changes in sea level. Changes in paleocirculation in the study area have been modified both by the movement of continental fragments resulting from local rifting events and by the northward movement of the Indo-Australian Plate and its collision with the Asian Plate. Northward movement of the Indo-Australian Plate caused significant variations in climate because of movement across climatic boundaries. These changes, in addition to global climatic variations, had dramatic influences on the depositional environments in the Coral Sea, which today are dominated by tropical carbonates. An important paleoceanographic objective is to determine the effects of

changes in paleocirculation and climate on the development of carbonate platforms and reefs off northeast Australia.

DRILLING STRATEGY

The establishment of a sea-level curve for the Miocene in the Coral Sea region is critically dependent on determining the facies and age of the MP2 and MP3 platforms of the Marion Plateau (Figs. 2, 10). Typically, precise dating of warm shallow-water carbonate platforms is difficult because of the broad stratigraphic range of larger foraminifers and diagenetic alteration of the sediments. Therefore, the drilling strategy described here involves paired holes chosen so that one is located within predicted shallow-water facies and a second is located downslope to obtain correlative facies in which planktonic forms are preserved for high resolution dating. Five primary and 10 contingency sites are proposed according to this strategy and defined on multichannel seismic lines collected in 1999 (Tables 1, 2; Figs. 11-34 and Site Summary section).

Primary Sites

The operational time estimates indicate that the five highest priority sites can be cored and logged in the time allocated for Leg 194 (Table 2). These include two platform sites (Sites CS-01 and 06), two sites to sample the adjacent paleoslope sediments (Sites CS-02 and 05), and a distal site (Site CS-10) to acquire a high-quality biostratigraphy that can be correlated to the other sites using high-resolution seismic data. The platform sites will first be penetrated to basement using the advanced hydraulic piston corer and the extended core barrel (APC/XCB) (e.g., Site CS-01) and/or the rotary core barrel (RCB) where the formation is too indurated (e.g., Sites CS-01 and 06). In an attempt to achieve better recovery in reefal carbonate intervals of interest (maximum 300 m per site), we plan to deploy the developmental advanced diamond core barrel (ADCB; see below). All sites are to be drilled to basement (including up to one barrel of basement rock) to form a facies transect from a position within the shallow facies of MP2, across the platform edge, and downslope to platform MP3. Sites between the two shallow phases of platform facies are designed to establish whether lowstand signals can be detected in slope sediments (Sites CS-03A and CS-05A). If such signals can be seen, it may be possible to establish rates as well as amplitudes of sea-level fluctuations.

The order of operations will greatly depend on weather conditions. Should conditions be favorable after transit from Townsville (1.2 days) and the Hydrate Autoclave Coring Equipment (HYACE) tests (2 days, see below), sites would be drilled in the following order: CS-01A, CS-02A, CS-06, CS-05, and CS-10A. After completing operations, including potential contingency plans (see below), there will be an 8-day transit to Guam.

Contingency Sites/Holes

Should time be available after completing operations at the five primary sites, the following contingency plans will be evaluated and ranked depending on available Leg 194 results. Three types of contingency plans have been developed:

- Additional sites: (in order of priority) CS-08A, CS-03A, CS-12A, CS-09A, CS-11A, CS-13A, CS-14A, CS-15A, CS-16A, and CS-17A.
- Third holes to improve stratigraphic coverage: CS-10A, CS-02A, CS-05A.
- Additional ADCB coring: Site CS-06A to recover the upper reefal carbonate interval and also Site CS-12A, if it is drilled.

All primary and some contingency sites have been approved by the Pollution Prevention and Safety Panel (PPSP). Sites CS-03A, CS-05A, CS-06A, and CS-09A are located within the Great Barrier Reef Marine Park. As such, it is imperative that care be taken during coring to prevent impacts on the reef ecosystems. All Leg 194 shipboard operations will follow the guidelines established by the Great Barrier Reef Marine Park Authority (GBRMPA) for scientific research within park boundaries.

As the site survey seismic data was collected using differential Global Positioning System (GPS) navigation, it is not expected that we will have to use the *JOIDES Resolution* to collect survey data prior to drilling Leg 194 sites.

Advanced Diamond Core Barrel (ADCB) Coring System

The ADCB coring system was designed to improve core recovery in indurated sediments and rocks by using a diamond-impregnated drill bit that cuts a core by grinding instead of chipping as does the conventional RCB system. The outer diameter and cutting area of the ADCB are smaller than those of the conventional RCB system (outer diameter = 7.5 in vs. 9.875 in), which

potentially increases the penetration rates. In addition, circulation around the freshly cut core is better constrained, and flushing away recovered core material is minimized. The recovered ADCB core has a larger diameter than conventional RCB cores (3.345 in vs. 2.312 in), which results in greater recovery in friable sediments while also increasing available material for analysis.

Because this tool is still in development, there are a few operational limitations. First, only 300 m of smaller diameter drill pipe needed for the ADCB has been purchased by ODP to date. This means to core an interval greater than 300 m, the drill string needs to be tripped to the surface, a RCB bit must be deployed to ream the hole to the bottom of the first 300-m interval, and the pipe must be tripped again to deploy the next ADCB bit to continue coring. Second, the logging tools cannot be deployed through the ADCB bit. Thus, if downhole logging is desired, an additional pipe trip is needed to deploy the logging bottom-hole assembly (BHA). Third, as the diamond coring bit is more susceptible to wear, a pipe trip may also be required to change the bit before 300 m are penetrated. In addition, core splitting, sampling, and archiving issues have not been addressed for the larger core diameter and would have to be optimized during the cruise.

The ADCB system will be deployed for the first time on Leg 193. Since the recovery of reefal carbonates has traditionally been difficult with the ODP arsenal of tools, and because the recovery of these lithologies is a primary Leg 194 objective, the use of this system will be important for Leg 194. Deployment of this developmental tool at one or two sites is therefore an essential use of Leg 194 operational time. Unless some fatal flaws are discovered during Leg 193, the ADCB will be used at the first site to evaluate the usefulness of this tool in these sediments. Subsequent deployments will only occur if operational time does not exceed the time estimates and if coring results are truly superior to those obtained with the RCB system.

HYACE Testing

In July 2000, two days were added to the original Leg 194 operations schedule to accommodate feasibility tests of the developmental HYACE tool, which is a gas hydrate sampling and monitoring system. The HYACE was modified from the ODP Pressure Core Sampler (PCS) at the Technische Universität Berlin with support from the European Commission and from ODP. The tests will be carried out at the very beginning of the leg at one of the designated contingency sites. At least five dedicated personnel will sail for these tests. Since a full technical and scientific

staff is needed during the scientific operations of Leg 194, the HYACE test personnel will be replaced by ODP technical staff arriving with a boat from shore two days into the cruise.

LOGGING PLAN

Downhole logging will be an essential component of Leg 194 scientific operations, particularly at those sites where low core recovery is likely (Sites CS-01A and CS-06A). All sites will be logged with standard logging tool strings (triple combo and Formation MicroScanner [FMS]/sonic), together with deployment of the well seismic tool (WST) and the geologic high-resolution magnetic Tool (GHMT; if available). The characteristics of these logging tools are as follows (additional information on these tool strings can be found at <http://www.ldeo.columbia.edu/BRG>; all logging tools Schlumberger):

- The triple combo toolstring includes the dual induction tool (DITE) that measures resistivity from deep and shallow induction, the accelerator porosity sonde (APS) that measures porosity from epithermal neutron measurements, and the hostile environment litho-density sonde (HLDS) that measures bulk density from Compton scattering and provides an indication of general lithology from the photoelectric effect. Commonly, the hostile environment natural gamma-ray sonde (HNGS) is added to this tool string. Depending on the success of its deployment during Leg 191, the multichannel gamma-ray logging tool (MGT) will be added to the triple combo, providing a vertical resolution three times higher than the standard gamma-ray sonde.
- The FMS/sonic tool string includes the Formation MicroScanner, which measures resistivity at centimeter resolution on four pads moving along the borehole, the general purpose inclinometry tool (GPIT), and the dipole sonic imager (DSI), which measures compressional and shear wave velocity, as well as cross-dipole and Stoneley waveforms.
- The WST is a single-axis check-shot tool used for zero-offset vertical seismic profiles (VSPs). It consists of a single geophone that is used to record acoustic waves generated by an air gun located near the sea surface.

- The GHMT includes the nuclear magnetic remanence sonde (NMRS), which measures total magnetic field, and the susceptibility measurement sonde (SUMS), which measures the magnetic susceptibility from induction.

The Leg 194 logging plan has been designed to achieve the following objectives:

- Core-log-seismic correlation.** A detailed correlation between cores, logs, and the extensive suite of high-quality seismic reflection data (including the closely spaced site survey grids and regional two-dimensional lines) will be critical for understanding the three-dimensional architecture of the Marion Plateau sequences and for compiling a detailed sequence stratigraphy. Accordingly, check-shot (WST) surveys, at 30 to 50 m spacing, should be run at all sites.
- Detailed description of cored sequences.** Integrated interpretation of the FMS and geophysical logs (triple-combo) will provide an essential complement to drill cores for describing the lithostratigraphy, particularly in intervals where core recovery is incomplete. Geophysical logs will also be important for characterizing the lithostratigraphic response of the carbonate sediments to sea-level and climatic fluctuations.
- Characterization of sedimentary diagenesis.** Sedimentary diagenesis in carbonates is expressed through changes in porosity and chemical precipitation of soluble elements (such as uranium) that lead to large changes in log properties seen on sonic, resistivity, density, and neutron gamma-ray logs.
- Detection of terrigenous sediment input to the sedimentary section.** The Marion Plateau is a mixed carbonate-siliciclastic depositional environment; thus, the GHMT tool will be useful for detecting variations in terrigenous sediment content intermixed with the platform and pelagic carbonate sediments, particularly in those intervals characterized by low core recovery. These data will be also be useful for intersite sediment correlation.
- Assessment of fracture networks and basic fluid properties.** Fine-scale characteristics of sedimentary bedding, including pore spaces, bioturbation, and fractures, will be imaged using the FMS. The assessment of fracture networks and fluids will be accomplished using sonic and resistivity logs that, together with FMS images, should help achieve leg fluid-flow objectives.

SAMPLING PLAN

All sampling of recovered cores will be subject to the rules described in the ODP Sample Distribution Policy (<http://www-odp.tamu.edu/curation/sdp.htm>). Prior to Leg 194, the sampling program for the leg will be developed and approved by the Sampling Allocation Committee (SAC), consisting of the Co-Chief Scientists (Anselmetti and Isern), Staff Scientist (Blum), and Curator. The initial sampling plan is preliminary and can be modified depending upon actual material recovered and collaborations that may evolve between scientists during the leg. In the final shipboard sampling plan, sample requests will be closely linked to proposed postcruise research.

During the leg, samples for measurements of ephemeral properties will be collected. These measurements include those for organic geochemistry for safety monitoring (free gas and 20-cm³ sediment samples), interstitial water chemistry (whole rounds 5-15 cm in length), and biostratigraphy (core catcher samples). After splitting of the cores, samples for moisture content (20 cm³) will also be routinely collected. The study of fluid movement within the sedimentary section is an important goal of Leg 194. Thus, to characterize these fluid movements and sediment diagenesis, the collection of whole-round samples for pore-water geochemical analyses is anticipated to be at a higher resolution than typical for other ODP cruises.

Except for Sites CS-01A and CS-06A, all Leg 194 sites will be double cored in an attempt to recover a complete stratigraphic section. No archive halves (permanent or temporary) will be sampled aboard ship, and the permanent archive will be designated postcruise. Shipboard sampling for Leg 194 postcruise studies will be kept to a minimum. Only low-resolution sampling to support sediment facies description/characterization and facilitate pilot studies will be carried out aboard ship, and shipboard participants will only collect enough material for 6-months of postcruise research. The remaining samples required by participants and all high-resolution sampling will be deferred until after the cruise to enable the development of a composite sedimentary section from the multiple holes. All shipboard scientists will participate in shipboard sampling according to the shift schedule developed at the start of the leg.

REFERENCES

- Buddemeier, R.W., and Oberdorfer, J.A., 1986. Internal hydrology and geochemistry of coral reefs and atoll islands: key to diagenetic variations. *In* Schroeder, J.H., and Purser, B.H. (Eds.), *Reef Diagenesis*: Berlin (Springer-Verlag), 9-11.
- COSOD II, 1987. Rep. 2nd Conf. Scientific Ocean Drilling. Washington/Strasbourg (JOIDES/European Sci. Found.).
- Davies, P.J., Symonds, P.A., Feary, D.A., and Pigram, C. J., 1989. The evolution of the carbonate platforms of northeast Australia. *In* Crevello, P.D., Wilson, J.L., Sarg, J.F., and Read, J.F. (Eds.), *Controls on Carbonate Platform and Basin Development*, Spec. Publ.-Soc. Econ. Paleontol. Mineral., 44:233-258.
- Davies, P.J., McKenzie, J.A., Palmer-Julson, A., et al., 1991. *Proc. ODP, Init. Repts.*, 133 (Pts. 1,2): College Station, TX (Ocean Drilling Program).
- Eberli, G.P., and Ginsburg, R.N., 1989. Cenozoic progradation of the northwestern Great Bahama Bank: a record of lateral platform growth and sea level fluctuations. *In* Crevello, P.D., Wilson, J.L., Sarg, J.F., and Read, J.F. (Eds.), *Controls on Carbonate Platform and Basin Development*, Spec. Publ.-Soc. Econ. Paleontol. Mineral., 44:339-351.
- Eberli, G.P., Swart, P.K., Malone, M., et al., 1997. *Proc. ODP, Init. Repts.*, 166: College Station, TX (Ocean Drilling Program).
- Elderfield, H., Swart, P.K., McKenzie, J.A., and Williams, A., 1993. The strontium isotopic composition of pore waters from Leg 133: northeast Australian margin. *In* McKenzie, J.A., Davies, P.J., and Palmer-Julson, A., et al., *Proc. ODP, Sci. Results*, 133: College Station, Texas (Ocean Drilling Program), 473-480.
- Feary, D.A., Hine, A., Malone, M., et al., 2000. *Proc. ODP, Init. Repts.*, 182 [CD-ROM] Available from: Ocean Drilling Program, Texas A&M University, College Station TX (77845-9547, USA).
- Feary, D.A., Champion, D.C., Bultitude, R.J., and Davies, P.J., 1993. Igneous and metasedimentary basement lithofacies of the Queensland Plateau (Sites 824 and 825). *In* McKenzie, J.A., Davies, P.J., and Palmer-Julson, A., et al., *Proc. ODP, Sci. Results*, 133: College Station, Texas (Ocean Drilling Program), 535-540.
- Gaina, C., Müller, R.D., Royer, J.-Y., and Symonds, P., 1999. The evolution of the Louisiade Triple Junction. *J. Geophys. Res.*, 104:12,927-12,939.
- Halley, R.B., and Ludwig, K.R., 1987. Disconformities and Sr-isotope stratigraphy reveal a Neogene sea-level history from Enewetak Atoll, Marshall Islands, Central Pacific. *Geol. Soc. Amer. Abstr.*, 19:691.

- Haq, B.U., Hardenbol, J., and Vail, P.R., 1987. Chronology of fluctuating sea levels since the Triassic. *Science*, 235:1156-1167.
- Hayes, D.E., and Ringis, J., 1973. Seafloor spreading in the Tasman Sea. *Nature*, 243:454-458.
- Isern, A.R., McKenzie, J.A., and Feary, D.A., 1996. The role of sea-surface temperature as a control on carbonate platform development in the western Coral Sea. *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, 124:247-272
- JOIDES Long Range Plan, 1996. Understanding our dynamic Earth through ocean drilling.
- Katz, M.E., and Miller, K.G., 1993. Neogene subsidence along the northeast Australian margin: benthic foraminiferal evidence. In McKenzie, J.A., Davies, P.J., and Palmer-Julson, A., et al., *Proc. ODP, Sci. Results*, 133: College Station, Texas (Ocean Drilling Program), 75-92.
- Lincoln, J.M., and Schlanger, S.O., 1987. Miocene sea level falls related to the geologic history of Midway Atoll. *Geology*, 15:454-457.
- Lincoln, J.M., and Schlanger, S.O., 1991. Atoll stratigraphy as a record of sea level change: problems and prospects. *J. Geophys. Res.*, 96:6727-6752.
- Majors, R.P., and Mathews, R.K., 1983. Isotopic composition of bank margin carbonates on Midway Atoll: amplitude constraints on post-early Miocene eustasy. *Geology*, 11:335-338.
- McKenzie, J.A., Isern, A.R., Elderfield, H., Williams, A., and Swart, P., 1993. Strontium isotope dating of paleoceanographic, lithologic, and dolomitization events on the northeast Australian margin, Leg 133. In McKenzie, J.A., Davies, P.J., and Palmer-Julson, A., et al., *Proc. ODP, Sci. Results*, 133: College Station, Texas (Ocean Drilling Program), 489-498.
- Miller, K.G., Fairbanks, R.G., and Mountain, G.S., 1987. Tertiary oxygen isotope synthesis, sea-level history, and continental margin erosion. *Paleoceanography*, 2:1-19.
- Moore, T.C., Loutit, T.S., and Greenlee, S.M., 1987. Estimating short-term changes in eustatic sea-level. *Paleoceanography*, 2:625-637.
- Müller, R.D., Lim, V.S.L., and Isern, A.R., 2000. Late tertiary tectonic subsidence on the northeast Australian passive margin: response to dynamic topography? *Mar. Geol.*, 162:337-352.
- Mullins, H.T., Heath, K.C., Van Buren, M., and Newton, K., 1984. Anatomy of a modern open-ocean carbonate slope: northern Little Bahama Bank. *Sedimentology*, 31:141-168.
- Mutter, J.C. and Karner, G.D., 1980. The continental margin off northeast Australia. In Henderson, R.A. and Stephenson, P.J. (Eds.), *The Geology and Geophysics of Northeast Australia*, Geol. Soc. Aust. Queensl. Div., 47-49.
- Pigram, C.J., Davies, P.J., Feary, D.A., and Symonds, P.A., 1992. Absolute magnitude of the second-order middle to late Miocene sea-level fall, Marion Plateau, northeast Australia. *Geology*, 20:858-862.

- Pigram, C.J., Davies, P.J., and Chaproniere, G.C.H., 1993. Cement stratigraphy and the demise of the early-middle Miocene carbonate platform on the Marion Plateau. *In* McKenzie, J.A., Davies, P.J., and Palmer-Julson, A., et al., *Proc. ODP, Sci. Results*, 133: College Station, Texas (Ocean Drilling Program), 499-512.
- Sahagian, D.L., and Watts, A.B., 1991. Introduction to the Special Section on measurement, causes, and consequences of long-term sea-level changes. *J. Geophys. Res.*, 96:6585-6589.
- Schlanger, S.O. and Premoli-Silva, I., 1986. Oligocene sea-level falls recorded in the mid-Pacific atoll and archipelagic apron settings. *Geology*, 14:392-395.
- Shaw, R.D., 1978. Seafloor spreading in the Tasman Sea: a Lord Howe Rise - eastern Australian reconstruction. *Bull. Aust. Soc. Explor. Geophys.*, 9:75-81.
- Simms, M., 1984. Dolomitization by groundwater flow systems in carbonate platforms. *Trans. Gulf Coast Assoc. Geol. Soc.*, 34:411-420.
- Struckmeyer, H.I.M., and Symonds, P.A., 1997. Tectonostratigraphic evolution of the Townsville Basin, Townsville Trough, offshore northeastern Australia. *Aust. J. Earth Sci.*, 44:799-817.
- Symonds, P.A., Davies, P.J., Feary, D.A., and Pigram, C.J., 1988. Geology of the northeast Australian margin basins. *In* Queensland 88 - Exploration and Development, PESA (QLD) Petroleum Symposium, 61-77.
- Vail, P.R., and Hardenbol, J., 1979. Sea-level change during the Tertiary. *Oceanus*, 22:71-79.
- Vail, P.R., Mitchum, R.M., Jr., Todd, R.G., Widmeir, J.M., Thompson, S., III, Sangree, J.B., Bubb, J.N., and Hatelid, W.G., 1977. Seismic stratigraphy and global changes in sea level. *In* Payton, C.E. (Ed.), *Seismic Stratigraphy: Applications to Hydrocarbon Exploration*. AAPG Memoir 26:49-212.
- Varenkamp, V.C., 1991. Episodic dolomitization of late Cenozoic carbonates in the Bahamas. *J. Sed. Petrol.* 61:1002-1014.
- Watts, A.B., and Thorne, J., 1984. Tectonics, global changes in sea level and their relationship to stratigraphical sequences at the U.S. Atlantic continental margin. *Mar. Pet. Geol.*, 1:319-339.
- Weissel, J.K. and Hayes, D.E., 1971. Evolution of the Tasman Sea reappraised. *Earth Planet. Sci. Lett.*, 36:77-84.
- Williams, D.F., 1988. Evidence for and against sea-level changes from the stable isotope record of the Cenozoic. *In* Wilgus, C.K., Hastings, B.S., Kendall, C.G.St.C., Posamentier, H.W., Ross, C.A., and Van Wagoner, J.C. (Eds.), *Sea-Level Changes: an Integrated Approach*. Spec. Publ.-Soc. Econ. Paleontol. Mineral. 42:31-36.
- Zhang, Y.-S., and Tanimoto, T., 1993. High-resolution global upper mantle structure and plate tectonics. *J. Geophys. Res.*, 98:9793-9823.

Table 1. Leg 194 proposed sites.

Site	Priority	Line	CDP	Fig.	Latitude (S)	Longitude (E)	Water Depth (m)	Sediment Thickness (m)	Target Depth (m)	Requested Penetration (m)	PPSP Approval
CS-01A	1	MAR-29	4486	15	20°14.53'	151°47.54'	342	474	483	600	Yes
		MAR-29	2315	16							
CS-02A	1	MAR-23	2180	17	20°14.10'	151°59.03'	368	379	388	500	Yes
		MAR-23	720	18							
CS-03A	2	MAR-37	1268	19	20°47.57'	152°16.51'	320	445	454	550	Yes
		MAR-37	1782	20							
CS-05A	1	MAR-51	1290	21	20°57.75'	152°44.31'	331	469	478	575	Yes
		MAR-51	1750	22							
CS-06A	1	MAR-70	348	23	21°00.38'	152°51.40'	314	561	570	670	Yes
		MAR-70	5610	24							
CS-08A	2	MAR-63	1296	25	21°04.60'	153°03.98'	358	513	522	650	Yes
		MAR-63	1814	26							
CS-09A	2	MAR-45	1300	27	20°54.66'	152°35.05'	342	477	486	600	Yes
		MAR-45	1814	28							
CS-10A	1	MAR-04	7050	29	20°24.18'	152°40.23'	431	484	493	600	Yes
		MAR-04	7898	--							
CS-11A	3	MAR-20	15348	30	20°03.40'	151°45.77'	360	491	500	650	Pending
CS-12A	2	MAR-20	15110	30	20°04.45'	151°47.08'	360	491	245	650	Pending
CS-13A	3	MAR-20	8260	31	20°34.53'	152°24.55'	369	526	535	650	Pending
CS-14A	3	MAR-20	4980	32	20°48.89'	152°42.58'	370	508	517	650	Pending
CS-15A	3	MAR-20	3780	33	20°54.14'	152°49.19'	365	469	478	650	Pending
CS-16A	3	MAR-20	2690	34	20°58.90'	152°55.21'	312	561	571	670	Pending
CS-17A	3	MAR-20	1270	35	21°05.09'	153°03.05'	336	519	528	650	Pending




Figure 1. Map showing the location of DSDP Sites 209 and 287 and ODP Leg 133 sites off northeastern Australia.




Figure 2. Map showing location of Leg 194 proposed primary sites (black circles), contingency sites (black squares), and Sites 815 and 816 from Leg 133 (black-rimmed white circles). Solid lines show the location of multichannel seismic lines from the Australian Geological Survey Organization (thinner lines = Survey 75) and from the Leg 194 site survey (heavier lines = MAR data). Gray shaded areas show the estimated extent of the MP2 (early to middle Miocene) and MP3 (late Miocene to Pliocene) platforms. Dashed line shows the boundary of the Great Barrier Reef Marine Park.




Figure 4. Marion Plateau sea-level events vs. age compared to those from Haq et al. (1987).




Figure 5. Structure contour map for basement of the Marion Plateau (after Pigram et al., 1992). Contours are effectively isosubsidence lines for total subsidence of basement. Locations of proposed sites are shown. Note that the sites generally strike parallel to the flexural axis and have undergone similar total subsidence.




Figure 6. Map showing the major structural features of the Coral Sea.




Figure 7. Water-loaded tectonic subsidence (i.e., with the isostatic sediment load removed) for ODP Leg 133 Sites 812, 814, 811/825, 815, and 823, assuming constant eustatic sea level (a) and using eustatic sea-level variations of Haq et al. 1987 (b). The latter is not shown for Site 823, as the errors in water depth (vertical error bars) are much larger than eustatic sea-level variations. Shading around error bars indicates the area in which the true subsidence curve should occur. Comparisons between (a) and (b) allows evaluation of the potential effect of eustatic variations on tectonic subsidence models. For instance, the first model (a) for Site 814 shows a gently subsiding platform until about 5 Ma, whereas the second model (b), including eustacy, shows a tectonic subsidence pulse between 14 and 12 Ma. Therefore, the latter may be entirely due to the input of an ill-constrained eustatic sea-level curve.




Figure 8. Paleobathymetric histories for the Queensland and Marion Plateaus as inferred from benthic foraminiferal data plotted vs. age from Katz and Miller (1993). Present-day water depths (m) are shown for all sites.




Figure 9. Stratigraphic summary of previously cored sites located near Leg 194 sites. All sites were cored during Leg 133 except for Site 209, which was cored during DSDP Leg 21.




Figure 10. Interpreted AGSO Line 75-64 with proposed sites and major sequence stratigraphic units. The facies geometry used to calibrate the amplitude of the Miocene N12-N14 sea-level fall is shown by the relationship between platforms MP2 and MP3. Dashed lines indicate sites that have been chosen using other seismic lines but that still can be identified within a similar facies on this line. More detail on site placement can be seen on subsequent detailed sections. C2 is a carbonate growth phase occurring during Megasequence C.




Figure 11. High-resolution site survey seismic (two-way traveltime) from Line MAR-13 (Fig. 2) with locations of Leg 194 Sites CS-01A and CS-02A indicated.




Figure 12. High-resolution site survey seismic (two-way traveltime) from Line MAR-7 (Fig. 2) with locations of Leg 194 Sites CS-03A, CS-05A, CS-06A, CS-08A, and CS-09A indicated.




Figure 13. High-resolution site survey seismic (two-way traveltime) from Line MAR-7 (Fig. 2) with locations of Leg 194 Sites CS-10A indicated.

Table 2. Leg 194 Marion Plateau (Proposal 510)

Operations Plan and Time Estimate:							
Site No.	Location (Lat/Long)	Water Depth (mbrf)	Operations Description	Transit (days)	Drilling (days)	Logging (days)	Total On-site (days)
Sea Voyage from Townsville to CS-01A (195 nmi @ 10.5 kt)				0.8			
CS-01A	20°14.53'S 151°47.54'E	342	Hole A: APC to ref. (~50 mbsf), XCB 70 mbsf Hole B: RCB 483 mbsf, Log. Hole C: RCB 40 mbsf, FFF, ADCB 340 mbsf, Log		0.7 2.7 6.9	0.0 0.8 0.8	0.7 3.5 7.7
Transit from CS-01A to CS-02A (11 nmi @ 10.5 kts)				0.1			
CS-02A	20°14.10'S 151°59.03'E	368	Hole A: APC to refusal (~119 mbsf), XCB to 388 mbsf, Log Hole B: APC to refusal (~119 mbsf), XCB to 388 mbsf		2.2 1.6	0.8 0.0	3.0 1.6
Transit from CS-02A to CS-06A (97 nmi @ 10.5 kts)				0.4			
CS-06A	21°00.38'S 152°51.40'E	314	Hole A: RCB to 570 mbsf, Log Hole B: RCB to 280 mbsf, FFF, ADCB to 570 mbsf, Log		3.6 7.7	0.8 0.8	4.4 8.5
Transit from CS-06A to CS-05A (118 nmi @ 10.5 kts)				0.5			
CS-05A	20°57.75'S 152°44.31'E	331	Hole A: APC to refusal (~170 mbsf), XCB to 478 mbsf Hole B: APC to refusal (~170 mbsf), XCB to 283 mbsf Hole C: Drill to 170 mbsf, RCB to 478 mbsf, Log		2.0 1.1 2.0	0.0 0.0 0.8	2.0 1.1 2.8
Transit from CS-05A to CS-10A (34 nmi @ 10.5 kts)				0.1			
CS-10A	20°24.18'S 152°40.23'E	431	Hole A: APC to refusal (~234 mbsf), XCB to 493 mbsf Hole C: APC to refusal (~234 mbsf), XCB to 493 mbsf, Log		2.0 2.4	0.0 0.8	2.0 3.2
Sea Voyage from CS-10A to Guam (2014 nmi @ 10.5 kt)				8.0			
SUBTOTAL:				9.9	34.9	5.6	40.5
TOTAL OPERATING DAYS:				50.4			
CONTINGENCY SITES:							
CS-03A	20°47.57'S 152°16.51'E	320	Hole A: APC to refusal (~220 mbsf), XCB to 454 mbsf, Log Hole B: APC to refusal (~220 mbsf), XCB to 454 mbsf, Log		2.1 2.0	0.8 0.8	2.9 2.8
CS-06A	21°00.38'S 152°51.40'E	314	Hole C: RCB to 30 mbsf, FFF, reentry, ADCB to 330 mbsf		6.3	0.0	6.3
CS-08A	21°04.60'S 153°03.98'E	358	Hole A: APC to refusal (~150 mbsf), XCB to 522 mbsf Hole B: APC to refusal (~150 mbsf), XCB to 280 mbsf Hole C: Drill to 270 mbsf, RCB to 522 mbsf, Log		2.2 1.1 2.0	0.0 0.0 0.8	2.2 1.1 2.8
CS-09A	20°54.66'S 152°35.05'E	342	Hole A: APC to refusal (~200 m), XCB to 486 m Hole B: APC to refusal (~200 m), XCB to 486 m, Log		2.0 2.2	0.0 0.8	2.0 3.0

Table 2 (cont.). Leg 194 Marion Plateau (Proposal 510)

Contingency Sites:							2.0
Site No.	Location (Lat/Long)	Water Depth (mbrf)	Operations Description	Drilling (days)	63.0 63.0 (days)	Total On-site (days)	
CS-11A	20°03.40'S	360	Hole A: APC to refusal (~50 mbsf), XCB to 70 mbsf	0.7	0.0	0.7	
(01A)	151°45.77'E		Hole B: RCB to 500 mbsf, Log.	2.8	0.8	3.6	
			Hole C: RCB to 40 mbsf, FFF, ADCB to 340 mbsf, Log	6.8	0.8	7.6	
CS-12A	20°04.45'S	360	Hole A: APC to refusal (~50 mbsf), XCB to 70 mbsf	0.7	0.0	0.7	
	151°47.08'E		Hole B: RCB to 245 mbsf	1.3	0.0	1.3	
CS-13A	20°34.53'S	369	Hole A: APC to refusal (~220 mbsf), XCB to 535 mbsf, Log	2.4	0.8	3.2	
(03A)	152°24.55'E		Hole B: APC to refusal (~220 mbsf), XCB to 535 mbsf, Log	2.3	0.8	3.1	
CS-14A	20°48.89'S	370	Hole A: APC to refusal (~200 mbsf), XCB to 517 mbsf	2.1	0.0	2.1	
(09A)	152°42.58'E		Hole B: APC to refusal (~200 mbsf), XCB to 517 mbsf, Log	2.4	0.8	3.2	
CS-15A	20°54.14'S	365	Hole A: APC to refusal (~170 mbsf), XCB to 478 mbsf	2.0	0.0	2.0	
(05A)	152°49.19'E		Hole B: APC to refusal (~170 mbsf), XCB to 283 mbsf	1.1	0.0	1.1	
			Hole C: Drill to 170 mbsf, RCB to 478 mbsf, Log	2.1	0.8	2.9	
CS-16A	20°58.90'S	312	Hole A: RCB to 571 mbsf, Log	3.6	0.8	4.4	
(06A)	152°55.21'E		Hole B: RCB 280 mbsf, FFF, ADCB to 571 mbsf, Log	7.6	0.8	8.4	
			Hole C: RCB to 30 mbsf, FFF, reentry, ADCB to 330 mbsf	5.9	0.0	5.9	
CS-17A	21°05.09'S	336	Hole A: APC to refusal (~150 mbsf), XCB to 528 mbsf	2.2	0.0	2.2	
(08A)	153°03.05'E		Hole B: APC to refusal (~150 mbsf), XCB to 280 mbsf	1.1	0.0	1.1	
			Hole C: Drill to 270 mbsf, RCB to 528 mbsf, Log	2.0	0.8	2.8	

S209 SEISMIC LINES




Figure 14. Track line for Leg 194 seismic data

Site CS-01A
Line MAR-29 CDP 4486
Water Depth = 342 m
Estimated Penetration = 483 mbsf
Approved Penetration = 600 mbsf




Figure 15. Detailed high-resolution north-south seismic section (two-way traveltimes) used to locate Site CS-01A.

Site CS-01A
Line MAR-29 CDP 2315
Water Depth = 342 m
Estimated Penetration = 483 mbsf
Approved Penetration = 600 mbsf




Figure 16. Detailed high-resolution east-west seismic section (two-way traveltimes) used to locate Site CS-01A.

SITE SUMMARIES

Site: CS-01A

Priority: 1

Position: 20°14.53'S, 151°47.54'E

Water Depth: 342 m

Sediment Thickness: 474 m

Target Depth: 483 mbsf

Approved Maximum Penetration: 600 mbsf

Seismic Coverage: Regional Line MAR-13, shotpoint 4751; crossing Line MAR-29, CDPs 2315 and 4486

Objectives: The objectives of Site CS-01A are the following.

1. Determine the age of the MP2 platform drowning phase.
2. Determine the age and duration of regional unconformities.
3. Determine the total thickness of MP2.
4. Determine the age of the initial marine transgression over basement.
5. Determine the age and nature of the basement.
6. Measure and describe fluid flow within the MP2 platform.
7. Describe the MP2 platform carbonates.
8. Calibrate the seismic sequence stratigraphy.

Drilling Program: Double APC/XCB to refusal (3-4 cores), RCB (~1 core into basement); ADCB in selected intervals

Logging and Downhole Operations: Triple-combo, sonic-FMS, WST, GHMT (if available)

Nature of Rock Anticipated: 33 m of hemipelagic ooze overlying dolomitized reefal carbonates; underlying basement composed of Paleozoic quartz-feldspar mafic metasediment

Site CS-02A
Line MAR-23 CDP 2180
Water Depth = 368 m
Estimated Penetration = 388 mbsf
Approved Penetration = 500 mbsf




Figure 17. Detailed high-resolution north-south seismic section (two-way traveltimes) used to locate Site CS-02A.

Site CS-02A
Line MAR-23 CDP 720
Water Depth = 368 m
Estimated Penetration = 388 mbsf
Approved Penetration = 500 mbsf




Figure 18. Detailed high-resolution east-west seismic section (two-way traveltimes) used to locate Site CS-02A.

Site: CS-02A

Priority: 1

Position: 20°14.10'S, 151°59.03'E

Water Depth: 368 m

Sediment Thickness: 379 m

Target Depth: 388 mbsf

Approved Maximum Penetration: 500 mbsf

Seismic Coverage: Regional Line MAR-13; crossing Line MAR-23, CDPs 720 and 2180

Objectives: The objectives of Site CS-02A are the following.

1. Determine the age and describe the facies of Megasequences B-D, particularly the prograding, proximal slope sediments from MP2.
2. Determine the age and duration of unconformities that can be carried into the platform.
3. Determine the age of initial marine transgression over basement.
5. Determine the age and nature of the basement.
6. Measure and describe fluid flow and diagenetic processes within the MP platform and adjacent sediments.
7. Calibrate the seismic sequence stratigraphy.

Drilling Program: Double APC/XCB to refusal, XCB or RCB (~1 core into basement)

Logging and Downhole Operations: Triple-combo, sonic-FMS, WST, GHMT (if available)

Nature of Rock Anticipated: Approximately 119 m of hemipelagic ooze overlying ~260 m of periplatform ooze, wackestones with some mudstones, siltstones, and turbidites; underlying basement composed of Paleozoic quartz-feldspar mafic metasediment

Site CS-03A
Line MAR-37 CDP 1268
Water Depth = 320 m
Estimated Penetration = 454 mbsf
Approved Penetration = 550 mbsf




Figure 19. Detailed high-resolution northwest-southeast seismic section (two-way traveltimes) used to locate Site CS-03A.




Figure 20. Detailed high-resolution northeast-southwest seismic section (two-way traveltimes) used to locate Site CS-03A.

Site: CS-03A

Priority: 2

Position: 20°47.57'S, 152°16.51'E

Water Depth: 320 m

Sediment Thickness: 445 m

Target Depth: 454 mbsf

Approved Maximum Penetration: 550 mbsf

Seismic Coverage: Intersection of regional Line MAR-07 (shotpoint 223) with Line MAR-44 (shotpoint 3584); crossing Line: MAR-37 at shotpoints 887 and 630, CDPs 1782 and 1268

Objectives: The objectives of Site CS-03A are the following.

1. Determine the age and describe the facies of Megasequences B-D.
2. Determine the age of the initial marine transgression over basement.
3. Determine the age and describe the facies of lowstand deposits.
4. Determine the age and nature of the basement.
5. Measure fluid-flow processes within the MP2 platform and adjacent sediments.
6. Determine the lithologic signature of basinward unconformities.
7. Calibrate the seismic sequence stratigraphies.

Drilling Program: Double APC/XCB to refusal, XCB or RCB (~1 core) into basement

Logging and Downhole Operations: Triple-combo, sonic-FMS, WST, GHMT (if available)

Nature of Rock Anticipated: Approximately 180 m of hemipelagic ooze overlying ~265 m of periplatform ooze, wackestones with some siltstones and mudstones; underlying basement composed of Paleozoic quartz-feldspar mafic metasediment

Site CS-05A
Line MAR-51 CDP 1290
Water Depth = 331 m
Estimated Penetration = 478 mbsf
Approved Penetration = 575 mbsf




Figure 21. Detailed high-resolution north-south seismic section (two-way traveltimes) used to locate Site CS-05A.

Site CS-05A
Line MAR-51 CDP 1750
Water Depth = 331 m
Estimated Penetration = 478 mbsf
Approved Penetration = 575 mbsf




Figure 22. Detailed high-resolution east-west seismic section (two-way traveltimes) used to locate Site CS-05A

Site: CS-05A

Priority: 1

Position: 20°57.75'S, 152°44.31'E

Water Depth: 331 m

Sediment Thickness: 469 m

Target Depth: 478 mbsf

Approved Maximum Penetration: 575 mbsf

Seismic Coverage: Regional Line MAR-07, shotpoint 2261; crossing Line Mar-51, shotpoint 871, CDPs 1290 and 1750

Objectives: The objectives of Site CS-05A are the following.

1. Determine the age and facies description of Megasequences B-D, particularly the initiation of the MP3 platform.
2. Determine the age and duration of the unconformities that can be carried into the MP3 platform and those separating each sequence in the proximal slope adjacent to MP3.
3. Determine the paleowater depth of the initial growth phase of MP2.
4. Determine the age and nature of the condensed section equivalent to MP2.
5. Determine the age and nature of the basement.
6. Measure fluid flow processes within the MP3 platform and adjacent sediments.
7. Determine Pliocene-Holocene paleoceanography from the Megasequence D drift deposit.
8. Calibrate the seismic sequence stratigraphy.

Drilling Program: Double APC/XCB to refusal, XCB or RCB (~1 core into basement)

Logging and Downhole Operations: Triple-combo, sonic-FMS, WST, GHMT (if available)

Nature of Rock Anticipated: Approximately 196 m of hemipelagic ooze overlying ~273 m of periplatform ooze, wackestones with some siltstones, mudstones, and turbidites; underlying basement composed of Paleozoic quartz-feldspar mafic metasediment

Site CS-06A
Line MAR-70 CDP 348
Water Depth = 314 m
Estimated Penetration = 570 mbsf
Approved Penetration = 670 mbsf




Figure 23. Detailed high-resolution east-west seismic section (two-way traveltimes) used to locate Site CS-06A.




Figure 24. Detailed high-resolution northwest-southeast seismic section (two-way traveltime) used to locate Site CS-06A.

Site: CS-06A

Priority: 1

Position: 21°00.38'S, 152°51.40'E

Water Depth: 314 m

Sediment Thickness: ~561 m

Target Depth: ~570 mbsf

Approved Maximum Penetration: 670 mbsf

Seismic Coverage: Intersection of Lines MAR-07, shotpoint 2801, CDP 5610, and MAR-70, shotpoint 170, CDP 348

Objectives: The objectives of Site CS-06A are the following.

1. Determine the Initiation and facies development of the MP3 platform.
2. Determine the age and paleowater depth of the initial growth phase of MP3.
3. Describe the MP3 platform carbonates
4. Determine the age and duration of unconformities separating each platform phase.
5. Determine the age and nature of the condensed section equivalent to MP2.
6. Measure the fluid flow processes within the MP3 platform.
7. Calibrate the seismic stratigraphy.

Drilling Program: RCB (~1 core into basement), ADCB an interval of ~300 m.

Logging and Downhole Operations: Triple-combo, sonic-FMS, WST, GHMT (if available)

Nature of Rock Anticipated: Approximately 501 m of dolomitized framestone, packstone, and wackestone with a thin cover of periplatform ooze; underlying basement composed of Paleozoic quartz-feldspar mafic metasediment




Figure 25. Detailed high-resolution northwest-southeast seismic section (two-way traveltime) used to locate Site CS-08A.

Site CS-08A
Line MAR-63 CDP 1814
Water Depth = 358 m
Estimated Penetration = 522 mbsf
Approved Penetration = 650 mbsf




Figure 26. Detailed high-resolution northeast-southwest seismic section (two-way traveltimes) used to locate Site CS-08A.

Site: CS-08A

Priority: 2

Position: 21°04.60'S, 153°03.98'E

Water Depth: 358 m

Sediment Thickness: 513 m

Target Depth: 522 mbsf

Approved Maximum Penetration: 650 mbsf

Seismic Coverage: Regional Line MAR-07 shotpoint 3721; crossing Line MAR-63 at shotpoints 903 and 644, CDPs 1814 and 1296

Objectives: The objectives of Site CS-08A are the following.

1. Determine the age and describe the facies of Megasequences A-D.
2. Determine the age and paleowater depth of the initial phase of MP3.
3. Determine the duration of the unconformities separating each platform phase.
4. Determine the nature of the basement.
5. Measure the fluid flow processes within the MP3 platform and adjacent sediments.
6. Calibrate the seismic stratigraphy.

Drilling Program: Double APC/XCB to refusal, XCB or RCB (~1 core into basement)

Logging and Downhole Operations: Triple-combo, sonic-FMS, WST, GHMT (if available)

Nature of Rock Anticipated: Approximately 69 m of hemipelagic ooze overlying ~444 m of periplatform ooze, wackestones with some siltstones, mudstones, and turbidites; underlying basement composed of Paleozoic quartz-feldspar mafic metasediment

Site CS-09A
Line MAR-45 CDP 1300
Water Depth = 342 m
Estimated Penetration = 486 mbsf
Approved Penetration = 600 mbsf




Figure 27. Detailed high-resolution northwest-southeast seismic section (two-way traveltimes) used to locate Site CS-09A.




Figure 28. Detailed high-resolution northeast-southwest seismic section (two-way traveltimes) used to locate Site CS-09A.

Site: CS-09A

Priority: 2

Position: 20°54.66'S, 152°35.05'E

Water Depth: 342 m

Sediment Thickness: 477 m

Target Depth: 486 mbsf

Approved Maximum Penetration: 600 mbsf

Seismic Coverage: Regional Line MAR-07 shotpoint 1601; crossing Line MAR-45 at shotpoints 903 and 646, CDPs 1814 and 1300

Objectives: The objectives of Site CS-09A are the following.

1. Determine the age and facies description of Megasequences B-D, particularly the initiation of the MP3 platform.
2. Determine the age and duration of the unconformities separating each platform phase.
3. Determine the paleowater depth of the initial growth phase of MP3.
4. Determine the age and nature of the condensed section equivalent to MP2 and the basement.
5. Measure fluid flow processes within the Marion Plateau.
6. Determine the nature of the mounds imaged in the seismic data.
7. Calibrate the seismic sequence stratigraphy.

Drilling Program: Double APC/XCB to refusal, XCB or RCB (~1 core into basement)

Logging and Downhole Operations: Triple-combo, sonic-FMS, WST, GHMT (if available)

Nature of Rock Anticipated: Approximately 181 m of hemipelagic ooze overlying ~296 m of periplatform ooze, wackestones with some siltstones, mudstones and turbidites; underlying basement composed of Paleozoic quartz-feldspar mafic metasediment

Site CS-10A
Line MAR-15 CDP 7050
Water Depth = 431 m
Estimated Penetration = 493 mbsf
Approved Penetration = 600 mbsf



Figure 29. Detailed high-resolution east-west seismic section (two-way traveltimes) used to locate Site CS-10A.

Site: CS-10A

Priority: 1

Position: 20°24.18'S, 152°40.23'E

Water Depth: 431 m

Sediment Thickness: 484 m

Target Depth: 493 mbsf

Approved Maximum Penetration: 600 mbsf

Seismic Coverage: Intersection of regional Lines MAR-04 (shotpoint 3945, CDP 7898) and MAR-15 (shotpoint 3558 and CDP 7050)

Objectives: The objectives of Site CS-10A are the following.

1. Determine the chronostratigraphy for Megasequences A-D.
2. Determine the age of the initial marine transgression over basement.
3. Determine the age and facies of lowstand deposits.
4. Determine the age and nature of basement.
5. Calibrate the seismic stratigraphy.

Drilling Program: Triple APC/XCB (~1 core into basement)

Logging and Downhole Operations: Triple-combo, sonic-FMS, WST, GHMT (if available)

Nature of Rock Anticipated: Approximately 122 m of hemipelagic ooze overlying ~312 m of periplatform ooze, wackestones with some siltstones, mudstones, and turbidites; underlying basement composed of Paleozoic quartz-feldspar mafic metasediment




Figure 30. Regional seismic section (two-way traveltimes) used to locate Sites CS-11A and CS-12A.

Site: CS-11A

Priority: 2

Position: 20°03.40'S, 151°45.77'E

Water Depth: 360 m

Sediment Thickness: 500 m

Target Depth: ~500 mbsf

Approved Maximum Penetration: 650 mbsf

Seismic Coverage: Regional Line MAR-20, shotpoint 7670, CDP 15348

Objectives: The objectives of Site CS-11A are the following.

1. Determine the age of the MP2 platform drowning phase.
2. Determine the age and duration of regional unconformities.
3. Determine the total thickness of MP2.
4. Determine the age of the initial marine transgression over basement.
5. Determine the age and nature of the basement.
6. Measure and describe fluid flow within the MP2 platform.
7. Describe the MP2 platform carbonates.
8. Calibrate the seismic sequence stratigraphy.

Drilling Program: Single APC/XCB to refusal, XCB or RCB (~1 core into basement)

Logging and Downhole Operations: Triple-combo, sonic-FMS, WST, GHMT (if available)

Nature of Rock Anticipated: Approximately 49 m of hemipelagic ooze overlying ~442 m of periplatform ooze and dolomitized reefal carbonates; underlying basement composed of Paleozoic quartz-feldspar mafic metasediment

Leg 194
Scientific Prospectus
Page 66

blank

Site: CS-12A

Priority: 2

Position: 20°04.45'S, 151°47.08'E

Water Depth: 360 m

Sediment Thickness: 500 m

Target Depth: 245 mbsf

Approved Maximum Penetration: 650 mbsf

Seismic Coverage: Regional Line MAR-20, shotpoint 7551, CDP 15110

Objectives: The objectives of Site CS-12A are the following.

1. Determine the water depth of the shallowest part of the MP2 platform
2. Determine the age of the MP2 platform drowning
3. Determine the age and duration of unconformities that can be carried into the platform.
4. Calibrate the seismic sequence stratigraphy.

Drilling Program: Single APC/XCB or RCB to 245 mbsf; possibly ADCB of reefal carbonates

Logging and Downhole Operations: None

Nature of Rock Anticipated: Approximately 32 m of hemipelagic ooze overlying 213 m of periplatform ooze and dolomitized reefal carbonates




Figure 31. Regional seismic section (two-way traveltme) used to locate Site CS-13A.

Site: CS-13A

Priority: 2

Position: 20°34.53'S, 152°24.55'E

Water Depth: 369 m

Sediment Thickness: 526 m

Target Depth: 535 mbsf

Approved Maximum Penetration: 650 mbsf

Seismic Coverage: Regional Line MAR-20 shotpoint 4126, CDP 8260

Objectives: The objectives of Site CS-13A are the following.

1. Determine the age and describe the facies of Megasequences B-D.
2. Determine the age of the initial marine transgression over basement.
3. Determine the age and describe the facies of lowstand deposits.
4. Determine the age and nature of the basement.
5. Measure fluid-flow processes within the MP2 platform and adjacent sediments.
6. Determine the lithologic signature of basinward unconformities.
7. Calibrate the seismic sequence stratigraphies.

Drilling Program: Double APC/XCB (~1 core into basement)

Logging and Downhole Operations: Triple-combo, sonic-FMS, WST, GHMT (if available)

Nature of Rock Anticipated: Approximately 124 m of hemipelagic ooze overlying ~326 m of periplatform ooze, wackestones with some siltstones, mudstones, and turbidites; underlying basement composed of Paleozoic quartz-feldspar mafic metasediment




Figure 32. Regional seismic section (two-way traveltimes) used to locate Site CS-14A.

Site: CS-14A

Priority: 2

Position: 20°48.89'S, 152°42.58'E

Water Depth: 370 m

Sediment Thickness: 508 m

Target Depth: 517 mbsf

Approved Maximum Penetration: 650 mbsf

Seismic Coverage: Regional Line MAR-20, CDP 4980

Objectives: The objectives of Site CS-14A are the following.

1. Determine the age and facies description of Megasequences B-D, particularly the initiation of the MP3 platform.
2. Determine the age and duration of the unconformities separating each platform phase.
3. Determine the paleowater depth of the initial growth phase of MP3.
4. Determine the age and nature of the condensed section equivalent to MP2 and the basement.
5. Measure fluid flow processes within the Marion Plateau.
6. Determine the nature of the mounds imaged in the seismic data.
7. Calibrate the seismic sequence stratigraphy.

Drilling Program: Double APC/XCB (~1 core into basement)

Logging and Downhole Operations: Triple-combo, sonic-FMS, WST, GHMT (if available)

Nature of Rock Anticipated: Approximately 184 m of hemipelagic ooze overlying ~324 m of periplatform ooze, wackestones with some siltstones, mudstones, and turbidites; underlying basement composed of Paleozoic quartz-feldspar mafic metasediment




Figure 33. Regional seismic section (two-way traveltimes) used to locate Site CS-15A.

Site: CS-15A

Priority: 2

Position: 20°54.14'S, 152°49.19'E

Water Depth: 365 m

Sediment Thickness: 469 m

Target Depth: 478 mbsf

Approved Maximum Penetration: 650 mbsf

Seismic Coverage: Regional Line MAR-20, CDP 3780

Objectives: The objectives of Site CS-15A are the following.

1. Determine the age and facies description of Megasequences B-D, particularly the initiation of the MP3 platform.
2. Determine the age and duration of the unconformities that can be carried into the MP3 platform and those separating each sequence in the proximal slope adjacent to MP3.
3. Determine the paleowater depth of the initial growth phase of MP2.
4. Determine the age and nature of the condensed section equivalent to MP2.
5. Determine the age and nature of the basement.
6. Measure fluid flow processes within the MP3 platform and adjacent sediments.
7. Determine Pliocene-Holocene paleoceanography from the Megasequence D drift deposit.
8. Calibrate the seismic sequence stratigraphy.

Drilling Program: Double APC/XCB (~1 core into basement)

Logging and Downhole Operations: Triple-combo, sonic-FMS, WST, GHMT (if available)

Nature of Rock Anticipated: Approximately 203 m of hemipelagic ooze overlying ~266 m of periplatform ooze, wackestones with some siltstones, mudstones, and turbidites; underlying basement composed of Paleozoic quartz-feldspar mafic metasediment




Figure 34. Regional seismic section (two-way traveltimes) used to locate Site CS-16A.

Site: CS-16A

Priority: 2

Position: 20°58.90'S, 152°55.21'E

Water Depth: 312 m

Sediment Thickness: 561 m

Target Depth: 571 mbsf

Approved Maximum Penetration: 600 mbsf

Seismic Coverage: Intersection of regional Lines MAR-04 (shotpoint 3945) and MAR-15 (shotpoint 3558); Regional Line MAR-20, CDP 2690

Objectives: The objectives of Site CS-16A are the following.

1. Determine the Initiation and facies development of the MP3 platform.
2. Determine the age and paleowater depth of the initial growth phase of MP3.
3. Describe the MP3 platform carbonates
4. Determine the age and duration of unconformities separating each platform phase.
5. Determine the age and nature of the condensed section equivalent to MP2.
6. Measure the fluid flow processes within the MP3 platform.
7. Calibrate the seismic stratigraphy.

Drilling Program: Single XCB to refusal, RCB (~1 core into basement), ADCB may be used for selected intervals

Logging and Downhole Operations: Triple-combo, sonic-FMS, WST, GHMT (if available)

Nature of Rock Anticipated: Approximately 561 m of hemipelagic ooze, dolomitized framestone, packstone, wackestone with a thin cover of periplatform ooze; underlying basement composed of Paleozoic quartz-feldspar mafic metasediment




Figure 35. Regional seismic section (two-way traveltimes) used to locate Site CS-17A.

Site: CS-17A

Priority: 2

Position: 21°05.09'S, 153°03.05'E

Water Depth: 336 m

Sediment Thickness: 519 m

Target Depth: 528 mbsf

Approved Maximum Penetration: 650 mbsf

Seismic Coverage: Regional Line MAR-20, CDP 1270; Line MAR-63

Objectives: The objectives of Site CS-17A are the following.

1. Determine the age and describe the facies of Megasequences A-D.
2. Determine the age and paleowater depth of the initial phase of MP3.
3. Determine the duration of the unconformities separating each platform phase.
4. Determine the nature of the basement.
5. Measure the fluid flow processes within the MP3 platform and adjacent sediments.
6. Calibrate the seismic stratigraphy.

Drilling Program: Double APC and XCB (~1 core into basement)

Logging and Downhole Operations: Triple-combo, sonic-FMS, WST, GHMT (if available)

Nature of Rock Anticipated: 50 m of hemipelagic ooze overlying 469 m of periplatform ooze, wackestones with some siltstones, mudstones, and turbidites; underlying basement composed of Paleozoic quartz-feldspar mafic metasediment

SCIENTIFIC PARTICIPANTS*

Co-Chief
Flavio S. Anselmetti
Geologisches Institut
Eidgenössische Technische Hochschule - Zentrum
Sonneggstraße 5
Zürich 8092
Switzerland
Internet: flavio@erdw.ethz.ch
Work: (41) 1-632-65-69
Fax: (41) 1-632-10-80

Co-Chief
Alexandra R. Isern
Ocean Studies Board (HA-470)
National Academy of Sciences/National Research Council
2101 Constitution Ave. NW
Washington, D. C. 20418
USA
Internet: aiser@nas.edu
Work: (202) 334-2742
Fax: (202) 334-2885

Staff Scientist
Peter Blum
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
USA
Internet: blum@odpemail.tamu.edu
Work: (979) 845-9299
Fax: (979) 845-0876

Inorganic Geochemist
Tsutomu Yamada
Institute of Geology and Paleontology
Tohoku University
Graduate School of Science
Aobayama
Sendai, Miyagi 980-8578
Japan
Internet: yamada@dges.tohoku.ac.jp
Work: (81) 22-217-6618
Fax: (81) 22-217-6634

*As of 8/00

Organic Geochemist
Timothy S. White
Department of Geosciences
Pennsylvania State University
801 Deike Building
University Park, PA 16802
USA
Internet: dcb@essc.psu.edu
Work: (814) 865-2213
Fax: (814) 865-3191

JOIDES Logging Scientist
Gregor Eberli
Rosenstiel School of Marine and Atmospheric Science
University of Miami
4600 Rickenbacker Causeway
Miami, FL 33149
USA
Internet: geberli@rsmas.miami.edu
Work: (305) 361-4678
Fax: (305) 361-4632

Paleomagnetist
Tesfaye K. Birke
Hawaii Institute of Geophysics and Planetology
University of Hawaii at Manoa
Geology and Geophysics
2525 Correa Road
Honolulu, HI 96822
USA
Internet: tesfaye@soest.hawaii.edu
Work: 808-956-4879
Fax: 808-956-3188

Paleontologist (Foraminifer)
Pamela Hallock Muller
Department of Marine Science
University of South Florida, St. Petersburg
140 Seventh Ave. S
St. Petersburg, FL 33701-5016
USA
Internet: pmuller@marine.usf.edu
Work: (727) 553-1567
Fax: (727) 553-1189

Paleontologist (Foraminifer)
Michael W. Howell
Department of Geological Sciences
University of South Carolina

Leg 194
Scientific Prospectus
Page 80

Earth and Water Sciences Bldg.
Columbia, SC 29208
USA
Internet: howell@geol.sc.edu
Work: (803) 777-6943
Fax: (803) 777-6610

Paleontologist (Nannofossil)
Wuchang Wei
Scripps Institution of Oceanography
University of California, San Diego
Geoscience Research Division
La Jolla, CA 92093-0215
USA
Internet: wwei@ucsd.edu
Work: (858) 534-5844
Fax: (858) 558-2081

Physical Properties Specialist
Brandon E. Dugan
Department of Geosciences
Pennsylvania State University
410 Deike Building
University Park, PA 16802
USA
Internet: dugan@geosc.psu.edu
Work: (814) 863-9663
Fax: (814) 863-8724

Physical Properties Specialist
Garry D. Karner
Lamont-Doherty Earth Observatory
Columbia University
Marine Geology & Geophysics
Route 9W
Palisades, NY 10964
USA
Internet: garry@ldeo.columbia.edu
Work: (914) 365-8355
Fax: (914) 365-8156

Sedimentologist
Nils Andresen
GEOMAR
Christian-Albrechts-Universität zu Kiel
Wischhofstrasse 1-3
Kiel 24148
FRG

Internet: nandresen@geomar.de
Work: (49) 431-600-2835
Fax: (49) 431-600-2941

Sedimentologist
Guido L. Bracco Gartner
Rosenstiel School of Marine and Atmospheric Science
University of Miami
4600 Rickenbacher Causeway
Miami, FL 33149
USA
Internet: brag@geo.vu.nl
Work: (305) 361-4683
Fax: (305) 361-4632

Sedimentologist
Gilles Conesa
Centre de Sédimentologie-Paléontologie
Université de Provence
3 Place Victor Hugo
Marseille, Cedex 03 13331
France
Internet: gilles.conesa@newsup.univ-mrs.fr
Work: (33) 4 91 10 62 64
Fax: (33) 4 91 64 99 64

Sedimentologist
Albert C. Hine
Department of Marine Science
University of South Florida, St. Petersburg
140 7th Avenue South
St. Petersburg, FL 33701
USA
Internet: hine@seas.marine.usf.edu
Work: (727) 553-1161
Fax: (727) 553-1189

Sedimentologist
Cedric Michael John
Institut für Geologie und Paläontologie
Universität Stuttgart
Herdweg 51
Stuttgart 70174
Germany
Internet: cedric.john@geologie.uni-stuttgart.de
Work: (49) 711 1211330
Fax: (49) 711 1211341

Leg 194
Scientific Prospectus
Page 82

Sedimentologist
Pascal F. Kindler
Département de Géologie et de Paléontologie
Université de Genève
Section des Sciences de la Terre
13 rue des Maraichers
Genève 4 1211
Switzerland
Internet: pascal.kindler@terre.unige.ch
Work: (41) 22-702-6649
Fax: (41) 22-205-732

Sedimentologist
Keiichi Sasaki
Low Level Radioactivity Laboratory (LLRL)
Kanazawa University
Wake, Tatsunokuchi-machi
Nomi-gum, Ishikawa 923-1224
Japan
Internet: sasak1@llrl.ku-unet.ocn.ne.jp
Work: (81) 761-51-4440
Fax: (81) 761-51-5528

LDEO Logging Staff Scientist
Heike Delius
Department of Geology
University of Leicester
University Road
Leicester LE1 7RH
United Kingdom
Internet: hd21@le.ac.uk
Work: (44) 116-252-3634
Fax: (44) 116-252-3918

Operations Manager
Ron Grout
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
USA
Internet: grout@odpemail.tamu.edu
Work: (979) 845-2144
Fax: (979) 845-2308

ODP Drilling Engineer
Eddie Wright
Ocean Drilling Program
Texas A&M University

1000 Discovery Drive
College Station, TX 77845-9547
USA
Internet: wright@odpemail.tamu.edu
Work: (979) 845-3207
Fax: (979) 845-2308

Laboratory Officer
William G. Mills
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
USA
Internet: mills@odpemail.tamu.edu
Work: (979) 845-2478
Fax: (979) 845-0876

Marine Lab Specialist: Yeoperson
Jo Ribbens
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
USA
Internet: ribbens@odpemail.tamu.edu
Work: (979) 845-8482
Fax: (979) 845-0876

Marine Lab Specialist: Chemistry
Tim Bronk
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
USA
Internet: bronk@odpemail.tamu.edu
Work: (979) 845-9186
Fax: (979) 845-0876

Marine Lab Specialist: Chemistry
Anne Pimmel
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
USA

Leg 194
Scientific Prospectus
Page 84

Internet: pimmel@odpemail.tamu.edu
Work: (979) 845-8482
Fax: (979) 845-0876

Marine Lab Specialist: Downhole Tools, Thin Sections
Sandy Dillard
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
USA
Internet: dillard@odpemail.tamu.edu
Work: (979) 845-2481
Fax: (979) 845-0876

Marine Lab Specialist: Physical Properties
Cyndi J. Prince
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845
USA
Internet: prince@odpemail.tamu.edu
Work: (979) 845-2483
Fax: (979) 845-0876

Marine Lab Specialist: Underway Geophysics
Steve Prinz
Scripps Institution of Oceanography
University of California, San Diego
Ocean Drilling Program
West Coast Repository
La Jolla, CA 92093-0231
USA
Internet: prinz@odpemail.tamu.edu
Work: (858) 534-1657
Fax: (858) 534-4555

Marine Lab Specialist: X-Ray
Jaquelyn Ledbetter
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
USA
Internet: ledbetter@odpemail.tamu.edu
Work: (979) 845-8482
Fax: (979) 845-0876

Marine Lab Specialist: X-Ray
Johanna M. Suhonen
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
USA
Internet: suhonen@odpemail.tamu.edu
Work: (979) 845-9186
Fax: (979) 845-0876

Marine Electronics Specialist
Pieter Pretorius
Ocean Drilling Program
Texas A&M University
1000 Discovery drive
College Station, TX 77845-9547
USA
Internet: pretorius@odpemail.tamu.edu
Work: (979) 845-8482
Fax: (979) 845-0876

Marine Computer Specialist
Margaret Hastedt
Ocean Drilling Program
Texas A&M University
1000 Discovery Dr.
College Station, TX 77845
USA
Internet: hastedt@odpemail.tamu.edu
Work: (979) 862-2315
Fax: (979) 458-1617

Marine Computer Specialist
Erik Moortgat
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
USA
Internet: moortgat@odpemail.tamu.edu
Work: (979) 845-7716
Fax: (979) 845-0876

ODP/TAMU Application Developer
David Fackler
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive

Leg 194
Scientific Prospectus
Page 86

College Station, TX 77845
USA
Internet: fackler@odpemail.tamu.edu
Work: (979) 845-3031
Fax: (979) 458-1617