

OCEAN DRILLING PROGRAM
LEG 175 SCIENTIFIC PROSPECTUS
BENGUELA CURRENT

Dr. Wolfgang Berger
Co-Chief Scientist, Leg 175
Mail Code A-015
SIO-Geological Research Division
University of California/San Diego
La Jolla, CA 92093-0215
U.S.A.

Dr. Gerold Wefer
Co-Chief Scientist, Leg 175
Fachbereich Geowissenschaften
Universität Bremen
Postfach 33 04 40
D-28334 Bremen
Germany

Dr. Carl Richter
Staff Scientist, Leg 175
Ocean Drilling Program
Texas A&M University Research Park
1000 Discovery Drive
College Station, Texas 77845-9547
U.S.A.

Paul J. Fox
Director
of Science Operations
ODP/TAMU

James F. Allan
Interim Manager
Science Services
ODP/TAMU

Timothy J.G. Francis
Deputy Director
of Science Operations
ODP/TAMU

April 1997

Material in this publication may be copied without restraint for library, abstract service, educational, or personal research purposes; however, republication of any portion requires the written consent of the Director, Ocean Drilling Program, Texas A&M University Research Park, 1000 Discovery Drive, College Station, Texas 77845-9547, U.S.A., as well as appropriate acknowledgment of this source.

Scientific Prospectus No. 75

First Printing 1997

Distribution

Electronic copies of this publication may be obtained from the ODP Publications Home Page on the World Wide Web at <http://www-odp.tamu.edu/publications>.

D I S C L A I M E R

This publication was prepared by the Ocean Drilling Program, Texas A&M University, as an account of work performed under the international Ocean Drilling Program, which is managed by Joint Oceanographic Institutions, Inc., under contract with the National Science Foundation. Funding for the program is provided by the following agencies:

Australia/Canada/Chinese Taipei/Korea Consortium for Ocean Drilling
Deutsche Forschungsgemeinschaft (Federal Republic of Germany)
Institut Français de Recherche pour l'Exploitation de la Mer (France)
Ocean Research Institute of the University of Tokyo (Japan)
National Science Foundation (United States)
Natural Environment Research Council (United Kingdom)
European Science Foundation Consortium for the Ocean Drilling Program (Belgium, Denmark, Finland, Iceland, Italy, The Netherlands, Norway, Spain, Sweden, Switzerland, and Turkey)

Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the views of the National Science Foundation, the participating agencies, Joint Oceanographic Institutions, Inc., Texas A&M University, or Texas A&M Research Foundation.

This Scientific Prospectus is based on pre-cruise JOIDES panel discussions and scientific input from the designated Co-Chief Scientists on behalf of the drilling proponents. The operational plans within reflect JOIDES Planning Committee and thematic panel priorities. During the course of the cruise, actual site operations may indicate to the Co-Chief Scientists and the Operations Manager that it would be scientifically or operationally advantageous to amend the plan detailed in this prospectus. It should be understood that any proposed changes to the plan presented here are contingent upon approval of the Director of the Ocean Drilling Program in consultation with the Science and Operations Committees (successors to the Planning Committee) and the Pollution Prevention and Safety Panel.

Technical Editor: Karen K. Graber

ABSTRACT

During Leg 175, we propose to drill between eight and ten advanced hydraulic piston and extended core barrel core sites off the western coast of Africa (Angola and Namibia) to reconstruct the late Neogene history of the Benguela Current and the associated upwelling regime between 5°S and 32°S. The Angola/Namibia system contains one of the great upwelling regions of the world. Like the systems off Peru and California, which were studied during recent Ocean Drilling Program legs, it is characterized by organic-rich sediments that contain an excellent record of productivity history, which can be read on a very fine scale. In addition, this environment provides an excellent setting for natural experiments in diagenesis.

The individual transects reflect a compromise between geographic coverage, accessibility, and time constraints. We aim to study specific "end-member" environments, which collectively comprise one of the most active areas of ocean productivity. One of the major goals is to reconstruct the evolution of the Benguela Current system and its relationship with the onset of glacial cycles in the northern hemisphere. Most of the proposed sites are expected to have high sedimentation rates, offering an opportunity to develop detailed paleoceanographic records, and all proposed sites will extend and refine the partial record of the paleoceanographic and paleoclimatic changes for the late Neogene that was provided by Deep Sea Drilling Project Sites 362 and 532. Sediments will be largely diatomaceous and carbonate-rich clays with variable, and occasionally very high organic carbon contents.

INTRODUCTION

The ocean's role in climatic change through heat transport and control of carbon dioxide is increasingly being recognized. This new awareness, and the urgency that must be accorded the attempt to understand the mechanisms of climatic change, have led to the initiation of large integrated efforts in physical and chemical oceanography. Likewise, the potential of the oceanic record for understanding climatic change has received increased attention in recent years (CLIMAP, 1976; COSOD II, 1987). The Angola/Namibia high-productivity system needs to be studied because of its importance in the global ocean-carbon cycle, and to provide for comparison with the Peru system and the California system. Only by comparing these systems with each other shall we be able to learn which elements of a system are peculiar, and which have general validity through time and on a global scale. To further these goals, Leg 175 will drill at least eight APC/XCB sites off the southwestern coast of Africa (Fig. 1) to study the paleoenvironment of the Benguela Current and Angola/Namibia upwelling system, with emphasis on the late Neogene.

Eastern boundary upwelling is strongly involved in modulation of the carbon cycle and therefore, in control of the partial pressure of carbon dioxide ($p\text{CO}_2$) ("biological pumping", Berger and Keir, 1984; Sundquist and Broecker, 1985; Boyle and Keigwin, 1987; Sarnthein et al., 1988, Berger et al., 1989). It is now generally thought that such pumping is a crucial factor for the explanation of short-term fluctuations in atmospheric CO_2 , of the type seen in ice cores (Barnola et al., 1987). Along these lines of argument, there is a good correlation between productivity indices in the eastern equatorial Pacific and the ice-core record of $p\text{CO}_2$ (Fig. 2). Likewise, there is good correlation between the ice-core record and estimates of CO_2 pressure in surface water (Fig. 3).

On a longer time scale, Vincent and Berger (1985) have postulated that depositional pumping by coastal upwelling is responsible for changing the general level of atmospheric $p\text{CO}_2$. They propose a climatic preconditioning by upwelling-induced carbon extraction from the ocean-atmosphere system for the beginning of the modern ice-cap dominated world. Their argument is based on the

observation that carbon isotopes in deep-sea benthics become ^{13}C enriched just when organic-rich phosphatic sediments begin to accumulate around the Pacific margins (Fig. 4). In this view, eastern boundary upwelling, and therefore upwelling off Angola and Namibia, has global implications for the long-term history of the carbon cycle and climate and for the evolution of life and biogeography on land and in the sea.

To be able to predict the effects of changes in productivity on the CO_2 content of the atmosphere, the interrelationships between ocean circulation, nutrient transport, and the sedimentation of organic compounds and carbonate must be established for each of the important productivity regions. Until now, there is no information on Neogene upwelling fluctuations off Angola and Namibia, a region that is probably of considerable importance for the global carbon cycle.

The most important period for understanding the workings of the present system is the time since the Miocene. Within this period, we see the evolution of the present planetary orography, the buildup of ice-caps on both poles, the development of modern wind and upwelling regimes, and the stepwise increase in North Atlantic Deep Water (NADW) production, which dominates the style of deep circulation in the ocean. The present system is characterized by a strong 100,000-yr climatic cycle, beginning 700,000 years ago (Berger et al., 1996). High-amplitude fluctuations associated with buildup and decay of northern ice sheets began around 2.8 million years ago (Shackleton et al., 1984; Hodell and Venz, 1992) (Fig. 5).

BACKGROUND

The Angola/Namibia system is one of five or six great upwelling regions in the world. It extends over a considerable portion of the western margin of South Africa, with productivity values of $180 \text{ g}^\circ\text{C}/\text{m}^2\text{yr}$, and greater (black areas in Fig. 6). It is characterized by organic-rich sediments, containing an excellent record of productivity history, which, in turn, is closely tied to the regional dynamics of circulation, mixing, and upwelling, as seen in the oxygenation of thermocline waters (Fig. 7). In addition, this environment provides an excellent setting for "natural experiments" in

diagenesis, especially concerning the genesis of economically important resources (e.g., petroleum and phosphate).

Upwelling off southwest Africa is at present centered on the inner shelf and at the shelf edge. The Benguela Current flows roughly parallel to the coast and within ~180 km of it south of 25°S, and then turns to the west over the Walvis Ridge between 23° and 20°S (Fig. 8). At about 20°S, warm, tropical-water masses from the north meet the cold Benguela Current water. Eddies of cold, upwelled water contain radiolarian and diatom skeletons, which are transported from the upwelling area to the northern part of the Walvis Ridge, where they have been sampled at Deep Sea Drilling Project (DSDP) Sites 532 (Hay, Sibuet, et al., 1984) and 362 (Bolli, Ryan, et al., 1978).

According to previous studies, during the last glacial maximum (LGM) eddies formed farther north and the Benguela Current flowed parallel to the coast and over the Walvis Ridge to reach the Angola Basin, finally bearing to the west at about 17°S. Sediments deposited at Site 532 during the LGM apparently confirm the absence of upwelling eddies by containing zero to very few opal skeletons (Hay, Sibuet, et al., 1984; Diester-Haass, 1985). Upwelling may have continued to occur on the African shelf, but the Benguela Current then did not transport that upwelling signal to the Walvis Ridge. However, from the distribution of foraminiferal assemblages at Site 532, it appears that the northeastern Walvis Ridge was in fact characterized by intensified upwelling and a westward expansion of coastal upwelling cells at glacial periods during the last 500,000 yr (Oberhänsli, 1991). The issue of contrasting models of glacial/interglacial upwelling dynamics in this region is unresolved. It hinges on the question of why opaline fossils show contrary abundance variations, with respect to the productivity record from other proxy indicators.

The results from Sites 362 and 532 can be used to reconstruct, tentatively, the evolution of the Benguela Current during the past 10 m.y. This evolution is characterized, on the whole, by increasing rates of accumulation of organic carbon. In addition, there are indications from changing correlations between percent carbonate, percent C_{org} , and diatom abundance that the dynamics of the system undergo stepwise modification. In this connection, as well, a distinct opal maximum in the early Quaternary is of great interest (Fig. 9). The nature of this transition is not clear; perhaps it

is a response to the migration of the polar front to its modern position.

The evolution of the climate of the northern hemisphere, and particularly that of northern Europe, is linked to the exchange of heat between the South Atlantic and the North Atlantic Oceans (Fig. 10). This energy transport, operating over large distances, is involved in the formation and magnitude of polar ice caps. In today's world, a net heat transfer from the South Atlantic to the North Atlantic exists in currents above the thermocline (Fig. 11). A part of the heat contribution from the South Atlantic is believed to originate from the Indian Ocean via the Agulhas Current. The Benguela Current is a connection between the waters north of the polar front in the South Atlantic and the Equatorial Currents of the Atlantic. Northward and southward shifts of the Southern Ocean polar front constrict or expand, respectively, the interchange of heat from the Indian Ocean to the South Atlantic (McIntyre et al., 1989). This interchange presumably has a drastic impact on the heat budget of the Benguela Current and, consequently, that of the entire Atlantic Ocean. Such variations in heat transfers should appear as changes in the course and intensity of currents and productivity regimes and should be recorded in the sediments accumulating along the southwest African margin.

An important element of the heat transfer dynamics is the deep-circulation pattern. Traditionally, the focus in reconstructing this pattern has been on the properties and boundaries of NADW-related water masses, as seen in the $\delta^{13}\text{C}$ of benthic foraminifers. The emphasis has been on glacial-to-interglacial contrast (Fig. 12). This contrast shows that NADW production was greatly reduced during glacial periods (as also reflected in the pattern of carbonate preservation). More recent studies have added much detail to this story (summarized in Bickert and Wefer, 1996) (Fig. 13). It appears that the strength of the NADW is reflected in the differences between eastern and western basins and in gradients within the eastern basin. Information on associated changes at depths above the NADW has been sparse. It must be assumed that the strength of the nutrient maximum underlying the Benguela upwelling regions (Fig. 14) is somehow coupled to the evolution of NADW, which in turn influences dynamics of intermediate water-mass formation to the south. At this point, we do not know how the different cycles are related, so little or nothing can be said about causal relationships.

Paleoceanographic interpretations regarding the history of the Benguela Current are derived mainly from a single location off southwest Africa (Site 532) and must be considered preliminary. Given the indications that the axis and the intensity of the Benguela Current have changed over the past 15 m.y. and that productivity has fluctuated with glacial/interglacial cycles, confirmation and refinement of these ideas is needed. Although DSDP Legs 40, 74, and 75 occupied sites in the Cape and Angola Basins and on the Walvis Ridge, these sites are situated too far offshore to provide the needed information. The Benguela Current and its associated upwelling are not recorded well in the sediments at these sites. Even Sites 362 and 532 on the Walvis Ridge are too far offshore to contain a direct record of upwelling. They receive an indirect record of near-coastal upwelling from material transported to their location by the Benguela Current. Furthermore, modern coring technology (advanced hydraulic piston corer [APC], extended core barrel [XCB]) allows for high-resolution studies by avoiding much of the drilling disturbance present in the Leg 40 cores. Such high-resolution work is crucial if the dynamics of upwelling are to be captured back to the Miocene on a scale of glacial/interglacial cycles. Information from an array of sites situated in the southern and central Cape Basin, on the Walvis Ridge, and in the southern Angola Basin would allow the construction of a coherent picture.

SCIENTIFIC OBJECTIVES

The results from DSDP Sites 362 and 532 suggest that there has been a general northward migration of the Benguela Current upwelling system during the last 14 m.y. Because the shape of the South Atlantic has not changed appreciably during this time, the changes in the upwelling system must reflect large-scale, perhaps global, changes in ocean circulation. Leg 175 will focus primarily on the paleoceanographic and paleoclimatic aspects of the area. However, there is interest in investigating samples from the upwelling area off Angola and Namibia with regard to early diagenetic processes taking place in this unique environment. Possible work includes study of the formation of dolomite (Baker and Kastner, 1981; Kulm et al., 1984), phosphorite (Calvert and Price, 1983), and chert (see articles in Garrison et al., 1984). We also hope to examine the organic-

matter type and distribution as a function of time and climatic cycles. Important questions that are being addressed during Leg 175 include the following:

- Determine the history of the Benguela Current for the late Neogene. Of special interest is the changing response to orbital forcing, as seen in spectral amplitudes and phase relationships (e.g., McIntyre et al., 1989; Schneider, 1991; Berger and Wefer, 1996; Jansen et al., 1996; Schneider et al., 1996; Wefer et al., 1996).
- Study the history of productivity of the upwelling off Angola and Namibia and the influence of the Zaire River, extending available information about the late Quaternary (Bremner, 1983; Jansen et al., 1996) to earlier periods. The history of opal deposition off the Zaire River is of interest (Schneider, 1991), as well as the origin of cycles of carbonate, organic matter-deposition, and diatoms, in each region.
- Determine what kind of oceanographic changes occur simultaneously in the Atlantic Ocean (Agulhas Current, polar front position, equatorial current, Argentine Current) with the shifting of the Benguela Current. Results from Ocean Drilling Program (ODP) Legs 108 and 114 can help define the past equatorial and polar boundaries of the Benguela Current. The final aim is to reconstruct the late Neogene paleocirculation pattern of the South Atlantic Ocean to evaluate implications for the glacial/interglacial heat balance through time between the South and North Atlantic. Of special interest is the identification of changes in modes of circulation, as seen in changes in correlations between proxy variables, as a function of time.
- Determine if changes in the surface-current and upwelling patterns of the Benguela Current cause, or are related to, changes in climates of western South Africa. For example, is the origin of the Namib desert related to the initiation of upwelling off southwest Africa? Sites close to the continent probably contain enough information (clay minerals, grain size of terrigenous material, pollen, phytoliths, and fresh water diatoms) to allow reconstruction of continental climatic changes and to determine whether these changes are synchronous with oceanographic changes (i.e., the establishment of upwelling off southwest Africa).

- Examine the effect of sea-level changes, if any, on sedimentation below the Benguela Current. Published eustatic sea-level curves (Haq et al., 1987) will be useful for this purpose.
- Study early diagenetic processes in environments with very high organic carbon and opal contents, which will offer an interesting contrast to the studies undertaken during Leg 112, off Peru (Suess, von Huene, et al., 1990). The upwelling sediments off the Peruvian active margin are deposited in forearc basins in a disturbed tectonic setting, whereas off Angola/Namibia sedimentation occurs on a steadily sinking passive margin with quite stable conditions. Therefore, we expect a more continuous and longer record in comparison to the sites drilled off Peru, although the sedimentation rate might not be quite as high.

DRILLING STRATEGY

Leg 175 will drill at least eight sites as part of a latitudinal transect between 5°S and 32°S. Time permitting, ten or eleven sites will be attempted. Proposed sites are located in the Lower Congo Basin (LCB), Mid-Angola Basin (MAB), Southern Angola Basin (SAB), Walvis Basin (WB), Northern Cape Basin (NCB), Mid-Cape Basin (MCB), and Southern Cape Basin (SCB). To recover a complete stratigraphic sequence, we anticipate coring two or three holes with the APC at each site.

PROPOSED SITES

1. Lower Congo Basin (LCB)

Two LCB sites will sample a complex environment dominated by riverine input, seasonal coastal upwelling, and incursions from the southern equatorial counter current (Fig. 8). Whereas these two sites represent the same depositional environment, they are located at varying distances from the shelf break, in different water depths, and at different positions with respect to the river plume and

Congo Canyon. Seismic lines and proposed drill sites are shown in Figure 15. Maximum penetration is 200 m, for Sites LCB-1 and LCB-4. Time permitting, Site LCB-3A will be attempted to 200 m, with double APC coring.

2. Mid-Angola Basin (MAB)

The MAB sites, off the bight of Angola near 12°S, were chosen to provide information on "most nearly normal" margin sedimentation, being influenced neither by riverine input, nor by sustained year-round upwelling. Upwelling is greatly influenced by variations in the Angola Thermal Dome. It is seasonal, and productivity is relatively weak compared to that of adjacent regions (Schneider, 1991). This setting allows maximum expression of a pelagic signal in the regional high-productivity record. Proposed drill sites are located on seismic Line GeoB 93-015 (Fig. 16), because the southerly profile Line GeoB 93-017 is significantly influenced by slumping of shelf sediments and turbidity currents. Of the various sites proposed originally, the two shallow sites were determined to be feasible for drilling: Site MAB-1 for a maximum of 200 m, and Site MAB-2 for 120 m.

3. Southern Angola Basin (SAB)

The Southern Angola Basin sites are positioned to sample the northern end of the Angola/Namibia upwelling region. The transect should nicely complement previous results obtained from Walvis Ridge. This transect is important not only for the history of the Benguela Current and coastal upwelling migration, but also for its contribution to the climatic history of southern Africa. The Kunene River, reaching the coast at ~17°S, is at the climatological barrier between an illite zone in arid areas to the south and a kaolinite zone from tropical weathering areas to the north (Bornhold, 1973). The proposed sites are situated on a climatic boundary, and should sensitively reflect changes in the position of continental climatic zones. Suitable drill sites were identified from seismic lines in water depths between 2200 and 3000 m. The bathymetric survey confirmed the complex nature of the depositional environment. Although the survey was not sufficient to analyze all structures in detail, it is clear from the combined HYDROSWEEP and PARASOUND echosounder data set that few potential drill sites may be found in the area. Seismic Line GeoB 93-030 lies across proposed Site SAB-2 (Fig. 17). Stratigraphic data from two gravity cores (GeoB

1023-5, 17°09.4'5, 11°00.7'E, water depth 1918 m; GeoB 1024-2, 17°09.8'E, water depth 2799 m) show high Pleistocene sedimentation rates (10-50 cm/1000 yr) (Wefer et al., 1988; Schneider et al., 1992). We will attempt to drill to 600 m at Site SAB-2. Time permitting, Site SAB-1 will be drilled for double APC coring.

4. Walvis Basin (WB)

Sites WB-B and WB-C, together with DSDP Sites 532 and 362 at water depths of 1331 m and 1325 m, form a transect that is central to the reconstruction of the history of the Benguela Current (Fig. 18). The DSDP sites are seaward of the upwelling center, but contain an upwelling signal that was transported to this location by the Benguela Current and its filaments. At the other end of the transect, proposed Site WB-B will give a better record of the upwelling itself. Cores spanning the late Quaternary from nearby areas show a sedimentation rate of 4-7 cm/1000 yr (Schulz et al., 1992). We will attempt to drill to 600 m at Site WB-B. Time permitting, Sites WB-C and/or WB-A will be drilled for double APC coring.

5. Northern Cape Basin (NCB)

The NCB site will help document the northward migration of the Benguela Current system from the Miocene to the Quaternary, as well as the shoreward/seaward migration of the upwelling center. This site will also provide a record of maximum productivity in the system. Previous work in this area (for a summary see Dingle et al., 1987) has documented anaerobic, in part varved, sedimentation in the upper margin regions. Phosphatic deposits also are abundant (Calvert and Price, 1983). Seismic lines and the proposed drill site are shown in Figure 19.

The results of Emery et al. (1975) and Austin and Uchupi (1982) show a thick hemipelagic wedge sitting on "rifted continental crust." Slumps would not seem to pose a major problem, although hiatuses are anticipated. Noteworthy is the confirmation of a thick sequence below the shelf region. A close tie-in between pelagic and terrigenous sedimentation is expected to be present within the slope record. During the SONNE Cruise SO-86, vertical profiles were shot over multichannel seismic (MCS) Line AM-1, which were collected by the University of Texas to obtain detailed data for the planned site. A first stratigraphy on an 11-m-long core taken in the high-production

upwelling area off Namibia (GeoB 1711-4, 23°18.9'S, 12°22.6'E) from a depth of 1967 m indicated a sedimentation rate of 11 cm/1000 yr (Schulz et al., 1992). We will attempt to drill to 600 m at Site NCB-2B.

6. Mid- and Southern Cape Basin (MCB, SCB)

These sites are located in the southernmost area of the Cape Basin (Figs. 1, 20, 21). The sedimentary records will help in exploring the early history of the Benguela Current in the southern Cape Basin and in detecting possible Agulhas Current influences. The sites are located close to the continent to detect upwelling signals and signals from continental climates (pollen, clay minerals, and coarser terrigenous matter), as well as sea-level changes. South of the proposed transect, the margin becomes too steeply sloped to support undisturbed sediments. Site MCB-A is located along GeoB/AWI Line 96-009 (Fig. 20), Site SCB-1 is located along MSC Line AM-54, collected by the University of Texas and along profile GeoB/AWI 96-003 (Fig. 21). A sedimentation rate of about 5 cm/1000 yr was determined (Schulz et al. 1992) for an 11-m-long core collected from near the same water depth (GeoB 1719-7, 28°55.6'S, 14°10.7'E, water depth 1010 m), but ~150 miles to the north. Time permitting, Site MCB-A will be drilled for double APC coring, on the way to SCB. Site SCB-1 will be attempted to a depth of 600 m.

Site SCB-1, as the last site, could be deepened if time is available at the end of the leg. If it appears that time is indeed available, permission to deepen this hole beyond the 400 m now allowed should be sought promptly, to take account of recommendations by the Ocean History Panel (OHP). Alternatively, second priority sites (SCB-A, SCB-B, SCB-C) will be drilled for APC coring.

SAMPLING STRATEGY

General

Most of the core material to be recovered during Leg 175 will be retrieved by APC, generally by triple coring for the first priority sites and by double coring for any secondary priority sites drilled. One half of the first hole at each site will be the permanent archive half. Micropaleontology and

sedimentology sampling will be done after a composite sampling splice is constructed from the two or more holes drilled at that site. High-resolution sampling is anticipated for most sites (5 cm interval), with 10 to 20 cm³ needed for each sample, depending on the abundance of fossils (especially benthic foraminifers). Sampling schedules will be worked out between the parties involved to optimize stratigraphic coverage and to minimize overlap. Geochemical sampling, which calls for larger volumes, will be done on material from the third hole if it interferes with micropaleontology and sedimentology sampling. Also, whole-round samples will be available for pore-water studies from the third copy, as long as a sample's position is not crucial to filling gaps in the continuous stratigraphic record. If there is a need for a rapid decision on the location of a whole-round sample, but information is insufficient to place critical intervals for continuous stratigraphy, whole-round sections that would result in gaps greater than 15 cm will be avoided. Such sections should be separated by at least 1 m. Sampling for microbiological studies also will follow this strategy. Sampling for physical properties should be done so as not to interfere with stratigraphically sensitive sampling sequences, and to take advantage of available continuous nondestructive measurements. U-channel sampling for high-resolution paleomagnetic and rock-magnetic studies will be carried out along the composite sampling splice (temporary archive), where appropriate.

Ultra-high Resolution Sites

There is a possibility that varves will be encountered in some sites, notably in Walvis Basin. Detailed sampling will be necessary to achieve objectives in such sections. The sampling allocation committee ([SAC], consisting of the Co-chiefs, Staff Scientist, and Curator's representative) will determine details of the sampling pattern in these cases.

Sampling Time Table

Detailed sampling of cores from a given site will proceed after a composite stratigraphy is constructed from cores from the two or more holes drilled at the site. The splice will be constructed, and the stratigraphic information will be distributed to the scientific party in advance of post-cruise sampling to facilitate planning and scientific collaboration. Requests to sample on board, for pilot studies or for projects requiring lower stratigraphic resolution, will be considered

by the SAC.

General Sampling Procedure

Investigators should avoid sampling the center of core halves. Sample plugs should be taken as close to the edges of a core half as is feasible, given the purpose of sampling. Samples may also be taken with the "scoop" tool, which inherently takes samples from the edges of the core half. Large samples taken with the "cookie-cutter" tool, for example for lamina-scale studies, should be shared among interested scientific party members.

Archives

The permanent archive will be the ODP-defined "minimum permanent archive." Once the working half of a section is depleted, the temporary archives for that section will be accessible for sampling. Wherever possible, one quarter of such temporary archives should be preserved by sampling off-center to one side.

Special Core Handling

Large numbers of samples for organic geochemistry analysis may be taken and may need to be frozen.

LOGGING STRATEGY

A total of four sites will be logged during Leg 175 (Sites SAB-2, WB-B, NCB-2B, and SCB-1). Possibly one other site will be logged at the end of the leg, if time is available (e.g., SCB-A). Variations in biogenic carbonate, opal, and detrital deposition associated with climatic, oceanographic, and eustatic changes will be reflected in terms of corresponding changes in physical and geochemical properties. Coring may be discontinuous over deeper intervals because of gas expansion or XCB-coring disturbance. Consequently, downhole log data present an excellent resource for developing a quantitative paleoclimatic and paleoceanographic time series.

Special software for core-core and core-log data integration (CLIP) will be used during Leg 175 to establish composite sedimentary sections vs. depth.

Only holes deeper than 250 m will be logged with a combination of geophysical sensors: Triple-combo tool, the Formation MicroScanner (FMS) associated with the sonic tools, and the geological high-sensitivity magnetic tool (GHMT). The Triple-combo provides measurements of gamma ray, porosity, density, and electrical resistivity, which will be used to describe the lithology, sedimentary fabric, degree of lithification, and fluid composition. The FMS tool string produces high-resolution electrical resistivity images of the borehole wall that can be used to study the structure of bedding, diagenetic features, hiatus, and cyclicity recorded by sediments. The sonic tool coupled to the FMS can be useful to establish synthetic seismograms. By combining acoustic velocity with density evaluations and then convolving them with appropriate wavelet techniques, we can accurately calibrate the seismic lines. The GHMT provides continuous measurements of magnetic susceptibility and the vertical component of the total magnetic field. This latter measurement provides a magnetic reversal stratigraphy, if the magnetization of the sediments is sufficiently strong.

REFERENCES

- Austin, Jr., J.A., and Uchupi, E., 1982. Continental-Oceanic Crustal Transition off Southwest Africa. *AAPG Bull.*, 66:1328-1347.
- Baker, P.A., and Kastner, M., 1981. Constraints on the formation of sedimentary dolomite. *Science*, 213:214-216.
- Barnola, J.M., Raynaud, D., Korotkevich, Y.S., and Lorius, C., 1987. Vostok ice core provides 160,000-year record of atmospheric CO₂. *Nature*, 329:408-414.
- Berger, W.H., 1989. Global Maps of Ocean Productivity. In Berger, W.H., Smetacek, V., and Wefer, G. (Eds.), *Productivity of the Ocean: Present and Past: Dahlem Workshop Reports*, J. Wiley & Sons, Chichester, 429-455.
- Berger, W.H., and Keir, R.S., 1984. Glacial-Holocene changes in atmospheric CO₂ and the deep-sea record. In Hansen, J.E., and Takahashi, T. (Eds.), *Climate processes and climate sensitivity: Geophysical Monograph 29*, American Geophysical Union, Washington, D.C., 337-351.
- Berger, W.H., and Wefer, G., 1996. Central themes of South Atlantic circulation. In Wefer, G., Berger, W.H., Siedler, G., and Webb, D. (Eds.), *The South Atlantic: Present and Past Circulation*, Springer-Verlag, 1-11.
- Berger, W.H., Smetacek, V., Wefer, G., 1989. Ocean Productivity and Paleoproductivity - An Overview. In Berger, W.H., Smetacek, V., and Wefer, G., (Eds.), *Productivity of the Ocean: Present and Past: Dahlem Workshop Reports*, J. Wiley & Sons, Chichester, 1-34.
- Berger, W.H., Bickert, T., Yasuda, M., and Wefer, G., 1996. Reconstruction of atmospheric CO₂ from the deep-sea record of Ontong Java Plateau: The Milankovitch Chron. *Geologische Rundschau*, 85:466-495.
- Bickert, T., and Wefer, G., 1996. Late Quaternary deep water circulation in the South Atlantic: Reconstruction from carbonate dissolution and benthic stable isotopes. In Wefer, G., Berger, W.H., Siedler, G., and Webb, D., (Eds.), *The South Atlantic: Present and Past Circulation*, Springer-Verlag, 599-620.
- Bolli, H.M., Ryan, W.B.F., et al., 1978. *Init. Repts. DSDP*, 40: Washington (U.S. Gov. Printing Office).

- Bornhold, B.D., 1973. Late Quaternary sedimentation in the Eastern Angola Basin. Techn. Rep. Woods Hole, WHOI 73-8, 164 pp.
- Boyle, E.A., and Keigwin, L., 1987. North Atlantic thermohaline circulation during the past 20,000 years linked to high-latitude surface temperature. *Nature*, 330:35-40.
- Bremner, J.M., 1983. Biogenic sediments on the South West African (Namibian) continental margin. In Thiede, J., and Suess, E., (Eds.), *Coastal Upwelling, Its Sediment Record*, Part B, Plenum Press, 73-103.
- Calvert, S.E., and Price, N.B., 1983. Geochemistry of Namibian Shelf Sediments. In Suess, E., and Thiede, J. (Eds.), *Coastal Upwelling, Its Sediment Record*, Part A: Responses of the Sedimentary Regime to Present Coastal Upwelling, NATO Conference Series, Series IV: Marine Sciences, Vol. 10a: 337-375.
- Chapman, P., and Shannon, L.V., 1987. Seasonality in the oxygen minimum layer at the extremities of the Benguela system. In Payne, A.I.L., Gulland, J.A., Brink, K.H. (Eds). *The Benguela and Comparable Retroflection*. Deep-Sea Res. 34:1399-1416.
- Ciesielski, P.F., Kristoffersen, Y., et al., 1988. *Proc. ODP, Init. Repts.*, 114: College Station, TX (Ocean Drilling Program).
- CLIMAP Project Members, 1976. The surface of the ice-age earth. *Science*, 191:1131-1137.
- COSOD II, 1987. Report of the Second Conference on Scientific Ocean Drilling, Joint Oceanographic Institutions for Deep Earth Sampling, European Science Foundation. Strasbourg, 142 pp.
- Dean, W., and Gardner, J., 1985. Cyclic variations in calcium carbonate and organic carbon in Miocene to Holocene sediments, Walvis Ridge, South Atlantic Ocean. In. Hsü, K.J, and Weissert, H. (Eds.), *South Atlantic paleoceanography*, Cambridge University Press, 61-78.
- Diester-Haass, L., 1985. Late Quaternary sedimentation on the eastern Walvis Ridge, SE Atlantic (HPC 532 IPOD Leg 75) and neighbored piston cores. *Mar. Geol.*, 65:145-189.
- Dingle et al., 1987. Deep-sea sedimentary environments around Southern Africa (South-East Atlantic and South-West Indian Oceans). *Annals of the South African Museum*, 98:1-27.
- Duplessy, J.-C., Shackleton, N.J., Fairbanks, R.G., Labeyrie, L., Oppo, D., and Kallel, N., 1988. Deep-water source variations during the last climatic cycle and their impact on the global deep-water circulation. *Paleoceanography*, 3:343-360.

- Emery, K.O., Uchupi, E., Bowin, C.O., Phillips, J., and Simpson, E.S.W., 1975. Continental Margin Off Western Africa: Cape St. Francis (South Africa) to Walvis Ridge (South-West Africa). *AAPG Bull.*, 59:3-59.
- Garrison, R.E., Kastner M., and Zenger, D.H. (Eds.), 1984. *Dolomites of the Monterey Formation and other organic-rich units*. Soc. Econ. Paleont. Mineral., Pacific Section, Spec. Publ. 41, 215 pp.
- Haq, B.U., Hardenbol, J., and Vail, P.R., 1987. Chronology of fluctuating sea-levels since the Triassic. *Science*, 235:1156-1167.
- Hay, W.W., and Brock, J.C., 1992. Temporal variation in intensity of upwelling off southwest Africa. In Summerhayes, C.P., Prell, W.L., Emeis, K.C., (Eds.), *Upwelling Systems: Evolution Since the Early Miocene*. Geol. Soc. Special Publication No 63, pp 463-497.
- Hay, W.W., Sibuet, J.C., et al., 1984. *Init. Repts. DSDP, 75* (2 vols.): Washington (U.S. Govt. Printing Office).
- Hodell, D.A., and Venz, K., 1992. Toward a high-resolution stable isotopic record of the Southern Ocean during the Pliocene-Pleistocene (4.8 to 0.8 Ma). In Kennett, J.P., Warnke, D.A., (Eds.), *The Antarctic Paleoenvironment: A Perspective on Global Change Part One Vol 56* (Antarctic Research Series) American Geophysical Union, Washington D.C., 265-310.
- Jansen, J.H.F., 1985. Middle and Late Quaternary carbonate production and dissolution, and paleoceanography of the eastern Angola Basin, South Atlantic Ocean. In Hsü, K.J., and Weissert, H.J. (Eds.), *South Atlantic paleoceanography*, Cambridge University Press, 25-46.
- Jansen, J.H.F., Ufkes, E., and Schneider, R.R., 1996. Late Quaternary Movements of the Angola-Benguela Front, SE Atlantic, and Implications for Advection in the Equatorial Ocean. In Wefer, G., Berger, W.H., Siedler, G., and Webb, D. (Eds.), *The South Atlantic: Present and Past Circulation*, Springer-Verlag, 553-575.
- Jouzel, J., Barkov, N.I., Barnola, J.M., Bender, M., Chappellaz, J., Genthon, C., Kotlyakov, V.M., Lipenkov, V., Lorius, C., Petit, J.R., Raynaud, D., Raisbeck, G., Ritz, C., Sowers, T., Stievenard, M., Yiou, F., and Yiou, P., 1993. Extending the Vostok ice-core record of paleoclimate to the penultimate glacial period. *Nature*, 364:407-412.
- Kulm, L.D., Suess, E., and Thornburg, T.M., 1984. Dolomites in organic-rich muds of the Peru forearc basin: analogue to the Monterey Formation. In Garrison, R.E., Kastner, M., and

- Zenger, D.H. (Eds.), *Dolomites of the Monterey Formation and other organic-rich units: Pacific Section*, Soc. Econ. Paleont. Mineral., 41:29-47.
- Kroopnick, P., 1980. The distribution of ^{13}C in the Atlantic Ocean. *Earth Planet. Sci. Lett.*, 49:469-484.
- McIntyre, A., Ruddiman, W.F.K., Karlin, K., and Mix, A.C., 1989. Surface water response of the equatorial Atlantic Ocean to orbital forcing. *Paleoceanography*, 4:19-55.
- Miller, R.J., and Russell, G.L., 1989. Ocean heat transport during the last glacial maximum. *Paleoceanography*, 4:141-155.
- Müller, P.J., Schneider, R., Ruhland, G., 1994. Late Quaternary pCO_2 variations in the Angola current: evidence from organic carbon $\delta^{13}\text{C}$ and alkenone temperatures. In Zahn, R., et al. (Eds.), *Carbon cycling in the glacial ocean: constraints on the ocean's role in global change*, Springer, Berlin, Heidelberg, New York, 343-366.
- Oberhänsli, H., 1991. Upwelling signals at the northeastern Walvis Ridge during the past 500,000 years. *Paleoceanography*, 6:53-71.
- Peterson, R.G., and Stramma, L., 1991. Upper-level circulation in the South Atlantic. *Progress in Oceanography* 26:1-73.
- Popp, B.N., Takigiku, R., Hayes, J.M., Louda, J.W., Baker, E.W., 1989. The post-Paleozoic chronology and mechanism of ^{13}C depletion in primary marine organic matter. *Amer. Jour. Sci.*, 89:436-454.
- Rau, G.H., Froelich, P.N., Takahashi, T. Des-Marais, D.J., 1991. Does sedimentary organic delta ^{13}C record variations in Quaternary ocean $[\text{CO}_2(\text{aq})]$?. *Paleoceanography*, 6:335-347.
- Ruddiman, W., Sarnthein, M., Baldauf, J., 1988. *Proc. ODP, Init. Repts.*, 108 (Sections 1 and 2): College Station, TX (Ocean Drilling Program).
- Sarnthein, M., Winn, K., Duplessy J.C., and Fontugne, M.R., 1988. Global variations of surface ocean productivity in low and middle latitudes: influence on the CO_2 reservoirs of the deep ocean and the atmosphere during the last 21,000 years. *Paleoceanography*, 3:361-399.
- Sarnthein, M., Winn, K., Jung, S.J.A., Duplessy J.-C., Labeyrie, L., Erlenkeuser, H., and Ganssen, G., 1994. Changes in east Atlantic deepwater circulation over the last 30,000 years. *Paleoceanography*, 9:209-267.
- Schneider, R., 1991. Spätquartäre Produktivitätsänderungen im östlichen Angola-Becken:

- Reaktion auf Variationen im Passat-Monsun-Windsystem und in der Advektion des Benguela-Küstenstroms. Berichte Fachbereich Geowissenschaften Nr. 21, 198 S.
- Schneider, R., Dahmke, A., Kölling, A., Müller, P.J., Schulz, H.D., and Wefer, G., 1992. Strong deglacial minimum in the $\delta^{13}\text{C}$ record from planktonic foraminifera in the Benguela Upwelling Region: palaeoceanographic signal or early diagenetic imprint. In Summerhayes, C.P., Prell, W.L., Emeis, K.C. (Eds.), *Upwelling Systems: Evolution Since the Early Miocene*. Geol. Society Spec. Publ., 64:285-297.
- Schneider, R.R., Müller, P.J., Ruhland, G., Meinecke, G., Schmidt, H., and Wefer, G., 1996. Late Quaternary Surface Temperatures and Productivity in the East-Equatorial South Atlantic: Response to Changes in Trade/Monsoon Wind Forcing and Surface Water Advection. In Wefer, G., Berger, W.H., Siedler, G., and Webb, D. (Eds.), *The South Atlantic: Present and Past Circulation*, Springer-Verlag, 527-551.
- Schulz, H.D., et al., 1992. Bericht und erste Ergebnisse über die Meteor-Fahrt M 20/2, Abidjan - Dakar, 27.12.1991 - 3.2.1992. Berichte, Fachbereich Geowissenschaften, Universität Bremen, Nr. 25, 173.
- Shackleton, N.J., et al., 1984. Oxygen isotope calibration of the onset of ice-rafting and history of glaciation in the North Atlantic region. *Nature*, 307:620-623.
- Shackleton, N.J., Hall, M.A., and Shuxi, C., 1983. Carbon isotope data in core V19-30 confirm reduced carbon-dioxide concentration in the ice age atmosphere. *Nature*, 306:319-322.
- Suess, E., von Huene, R., et al., 1990. *Proc. ODP, Sci. Results*, 112: College Station, TX (Ocean Drilling Program).
- Sundquist, E.T., and Broecker, W.S., 1985. The carbon cycle and atmospheric CO_2 : natural variations Archean to Present. *American Geophysical Union, Monograph*, 32.
- Sverdrup, H.U. Johnson, M.W., Fleming, R.H., 1942. *The Oceans*. Prentice-Hall, Englewood Cliffs, 1087 p.
- Vincent, E., and Berger, W.H., 1985. Carbon dioxide and polar cooling in the Miocene: the Monterey hypothesis. In Sundquist, E.T., and Broecker, W.S. (Eds.), *The Carbon Cycle and Atmospheric CO_2 : Natural Variations Archean to Present, Geophysical Monograph 32*, 455-468.
- Wefer, G. et al., 1988. Bericht über die Meteor-Fahrt M 6-6, Libreville - Las Palmas, 18.2.1988-

- 23.3.1988. Berichte, Fachbereich Geowissenschaften, Universität Bremen, Nr. 3.
- Wefer, G., Berger, W.H., Bickert, T., Donner, B., Fischer, G., Kemle-von Mücke, S., Meineck, G., Müller, P.J., Mulitza, S., Niebler, H.-S., Pätzold, J., Schmidt, H., Schneider, R.R., and Segl, M., 1996. Late Quaternary surface circulation in the South Atlantic: The stable isotope record and implications for heat transport and productivity. *In* Wefer, G., Berger, W.H., Siedler, G., and Webb, D.,(Eds.), *The South Atlantic: Present and Past Circulation*, Springer-Verlag, 461-502.
- Woods, J., 1981. The memory of the ocean. *In* Berger, A. (Ed.), *Climatic Variations and Variability: Facts and Theories*, D.-Reidel Publ. Company, Dordrecht, 63-83.

FIGURE CAPTIONS

Figure 1. Overview map showing general areas of planned sites (black rectangles).

Figure 2. Carbon dioxide concentrations in the Vostok ice core from Antarctica (Barnola et al., 1987; Jouzel et al., 1993; solid triangles), compared with a productivity-related carbon isotope signal from the eastern tropical Pacific (difference between the $\delta^{13}\text{C}$ values of planktonic and benthic foraminifera; Shackleton et al., 1983; heavy line), show that ocean productivity and atmospheric CO_2 tend to vary together. Time scale of Barnola et al. is adjusted to the one of Shackleton et al. by correlation of the deuterium signal in the ice with the oxygen isotope signal in the sediment (from Berger et al., 1996).

Figure 3. Comparison of ice-core CO_2 record of Barnola et al. (1987) with surface water pCO_2 estimates for Core GeoB 1016-3, using conversion for $\delta^{13}\text{C}$ of organic matter to CO_2 pressure as proposed by Popp et al. (1989) and Rau et al. (1991). Time scale of Vostok ice core adjusted for best fit. (From Müller et al., 1994).

Figure 4. Relationship between $\delta^{18}\text{O}$ record and $\delta^{13}\text{C}$ record of benthic foraminifera, DSDP Site 216, tropical Indian Ocean. It suggests that extraction of organic carbon in upwelling regions during Monterey time eventually resulted in cooling because of drawdown of atmospheric pCO_2 (after Vincent and Berger, 1985).

Figure 5. The cooling step observed between 2.5 and 3 Ma (as seen in the shift to more positive values of $\delta^{18}\text{O}$) marks a change toward greater instability of climate, as seen in increased fluctuations of $\delta^{18}\text{O}$ values of planktonic foraminifera (*Neogloboquadrina pachyderma*, *Globigerina bulloides*) and benthic foraminifera (*Cibicides* spp.) sampled near the boundary between the South Atlantic and Southern oceans. From Hodell and Venz (1992).

Figure 6. Angola/Namibia upwelling system off southern Africa, as seen in productivity distributions. Modified from Berger (1989). Numbers are the primary production in $\text{g}^\circ\text{C}/\text{m}^2\text{yr}$; black areas have primary production values greater than $180 \text{ g}^\circ\text{C}/\text{m}^2\text{yr}$.

Figure 7. Conceptual model showing areas where low-oxygen water is formed in the Southeast Atlantic and the inferred movement of this water (dashed arrows; from Chapman and Shannon, 1987).

Figure 8. Schematic representation of the large-scale, upper-level geostrophic currents and fronts in the South Atlantic Ocean. After Peterson and Stramma (1991) with minor additions from several other compilations.

Figure 9. Depositional cycles in biogenous sediments on Walvis Ridge, in the Benguela System. Note overall trend in diatom abundance, with maximum in early Quaternary (after Dean and Gardner [1985] and Hay and Brock [1992]).

Figure 10. Meridional heat transport in the Indian and Atlantic Oceans (from Woods, 1981, modified). Note the major transfer of heat from the Indian Ocean to the Atlantic, which can be modulated through time by changing the position of the subantarctic frontal system.

Figure 11. Estimates for annual heat transports for the present world ocean (heavy solid line with thin lines showing the approximate error bounds) and for the Atlantic (present conditions and last glacial maximum, as labeled). From Berger and Wefer (1996a), after Miller and Russell (1989), modified. Note the anomalous pattern for the present South Atlantic and the more symmetric pattern for glacial conditions.

Figure 12. Deep-water patterns and flow in the South Atlantic during the present (a, b) and the last glacial maximum (c). **(a)** Present salinity distributions; from Sverdrup et al. (1942), according to G. Wüst. **(b)** Distribution of $\delta^{13}\text{C}$ values in dissolved inorganic carbon; from Kroopnick (1980). **(c)** Distribution of $\delta^{13}\text{C}$ values in dissolved inorganic carbon, 20 k.y. ago, inferred from $\delta^{13}\text{C}$ values in benthic foraminifers of that age; source: Berger and Wefer (1996a), after Duplessy et al. (1988) and Sarnthein et al. (1994).

Figure 13. Plot of carbon isotope records, measured on the benthic foraminifer taxon *C. wuellerstorfi*, as a function of time. From Bickert and Wefer (1996). Numbers are core labels, in the GeoB collections.

Figure 14. Zonal section of nitrate concentrations at 11°20'S. From Siedler et al. (1996).

Figure 15. Seismic lines and proposed drill sites in the Lower Congo Basin (LCB).

Figure 16. Seismic lines and proposed drill sites in the Mid-Angola Basin (MAB).

Figure 17. Seismic lines and proposed drill Site SAB-2 in the Southern Angola Basin (SAB).

Figure 18. Seismic lines and proposed drill sites in the Walvis Basin (WB).

Figure 19. Seismic lines and proposed drill Site NCB-2B in the Northern Cape Basin (NCB).

Figure 20. Seismic lines and proposed drill Site MCB-A in the Mid-Cape Basin (MCB).

Figure 21. Seismic lines and proposed drill Site SCB-1 in the Southern Cape Basin (SCB).




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8

532+532B




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15




Figure 16




Figure 17




Figure 19




Figure 20




Figure 21

TABLE 1. PRIMARY SITE TIME ESTIMATES

Site	Latitude Longitude	Water Depth (m)	Penetration (mbsf)	Drilling Operations (days)	Logging (days)	Transit Time (days)
<i>Transit Las Palmas to LCB-1</i>						12.4
LCB-1	5°04.1'S 11°06.1'E	1408	200	2.8	0.0	
<i>Transit from LCB-1 to LCB-4</i>						0.3
LCB-4	4°47.1'S 10°04.5'E	3000	200	3.8	0.0	
<i>Transit from LCB-4 to MAB-1</i>						1.9
MAB-1	11°55.2'S 13°24.0'E	432	200	1.9	0.0	
<i>Transit from MAB-1 to MAB-2</i>						0.1
MAB-2	11°55.8'S 13°02.3'E	722	120	1.5	0.0	
<i>Transit from MAB-2 to SAB-2</i>						1.2
SAB-2	16°33.7'S 10°49.3'E	2843	600	7.0	0.7	
<i>Transit from SAB-2 to WB-B</i>						1.1
WB-B	21°05.6'S 11°49.2'E	1290	600	5.1	0.7	
<i>Transit from WB-B to NCB-2B</i>						1.1
NCB-2B	25°30.8'S 13°01.7'E	2004	600	6.1	0.7	
<i>Transit from NCB-2B to SCB-1</i>						1.1
SCB-1	31°25.0'S 15°17.0'E	1350	600	5.2	0.7	
<i>Transit from SCB-1 to Cape Town</i>						0.9
Total				33.4	2.8	20.1

Total Days at Sea = 55.9 (Available Time = 56.0)

TABLE 2. ALTERNATE SITE TIME ESTIMATES

Site	Latitude Longitude	Water Depth (m)	Penetration (mbsf)	Drilling Operations (days)	Logging (days)
LCB-2	5°6.6'S 10°51.1'E	1811	200	3.2	0.0
LCB-3A	5°10.8'S 10°26.2'E	2392	200	3.5	0.0
LCB-3B	5°11.4'S 10°22.3'E	2500	200	3.5	0.0
LCB-3C	4°45.0'S 10°29.3'E	2435	200	3.5	0.0
SAB-1	16°39.6'S 10°57.0'E	2588	600	6.6	0.7
WR-1A	19°37.0'S 11°20.4'E	763	600	8.1	0.7
WB-A	21°29.0'S 11°15.1'E	2707	600	7.0	0.7
WB-C	20°53.9'S 11°13.4'E	2201	600	6.1	0.7
MCB-A	29°22.5'S 13°59.4'E	1726	600	5.7	0.7
SCB-A	31°54.4'S 15°14.1'E	2234	600	6.2	0.7
SCB-B	31°47.1'S 15°30.0'E	1507	600	5.5	0.7
SCB-C	31°41.4'S 15°41.9'E	887	600	4.7	0.7
SCB-D	31°21.3'S 15°36.5'E	778	600	4.6	0.1
Total				68.2	5.7

Orientation & T measurement times have been incorporated into drilling operations

Site: LCB-1

Priority: 1

Position: 5°04.1'S, 11°06.1'E

Water Depth: 1408 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 200 mbsf

Seismic Coverage: High-resolution multichannel seismic (MCS) survey, seismic Line GeoB 93-001 and 93-009

Objectives: The objectives of LCB-1 are to determine the:

1. variability of riverine input (Congo), and
2. history of opal, carbonate, and organic-matter deposition off Zaire (Congo).

Drilling Program: APC Holes A, B, C to 200 m or to refusal

Logging and Downhole: None

Nature of Rock Anticipated: Hemipelagic calcareous silty mud.

Site: LCB-2

Priority: 2

Position: 5°6.6'S, 10°51.1'E

Water Depth: 1832 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 200 m

Seismic Coverage: High-resolution MCS, seismic Line GeoB 93-001 and -007

Objectives: The objectives of LCB-2 are to determine the:


1. variability of riverine input (Congo), and
2. history of opal, carbonate, and organic-matter deposition off Zaire (Congo).

Drilling Program: APC Holes A, B, C to 200 m or to refusal

Logging and Downhole*: None

Nature of Rock Anticipated: Hemipelagic calcareous silty mud.

***Note:** It is anticipated that this site may not be drilled, or only drilled for APC, depending on time available. In the event that there is less than 250 m penetration, no logging or other downhole measurements will be attempted.



Site LCB-2

Site: LCB-3A

Priority: 2

Position: 5°10.8'S, 10°26.2'E

Water Depth: 2392 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 200 m

Seismic Coverage: High-resolution MCS survey, seismic Lines GeoB 93-001 and -006


Objectives: The objectives of LCB-3A are to determine the:

1. variability of riverine input (Congo), and
2. history of opal, carbonate, and organic matter deposition off Zaire (Congo).

Drilling Program: APC core Holes A, B, C to 200 m or to refusal

Logging and Downhole: None

Nature of Rock Anticipated: Hemipelagic calcareous silty mud



Site: LCB-3B

Priority: 2

Position: 5°11.4'S, 10°22.0'E

Water Depth: 2506 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 200 m

Seismic Coverage: High-resolution MCS, seismic Lines GeoB 93-001 and -004

Objectives: The objectives of LCB-3B are to determine:


1. the variability of riverine input (Congo), and
2. the history of opal, carbonate, and organic-matter deposition off Zaire (Congo).

Drilling Program: APC Holes A, B, C to 200 m or to refusal

Logging and Downhole*: None

Nature of Rock Anticipated: Hemipelagic calcareous silty mud.

***Note:** It is anticipated that this site may not be drilled, or only drilled for APC, depending on time available. In the event that there is less than 250 m penetration, no logging or other downhole measurements will be attempted.



Site LCB-3B

Site: LCB-3C

Priority: 2

Position: 4°45.0'S, 10°29.3'E

Water Depth: 2430 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 200 m

Seismic Coverage: High-resolution MCS, seismic Lines GeoB 93-002 and -007

Objectives: The objectives of LCB-3C are to determine:


1. the variability of riverine input (Congo), and
2. history of opal, carbonate, and organic matter deposition off Zaire (Congo).

Drilling Program: APC Holes A, B, C to 200 m or to refusal

Logging and Downhole*: None

Nature of Rock Anticipated: Hemipelagic calcareous silty mud.

***Note:** It is anticipated that this site may not be drilled, or only drilled for APC, depending on time available. In the event that there is less than 250 m penetration, no logging or other downhole measurements will be attempted.



Site: LCB-4

Priority: 1

Position: 4°47.1'S, 10°04.5'E

Water Depth: 3000 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 200 m

Seismic Coverage: High-resolution MCS survey, seismic Line GeoB 93-002


Objectives: The objectives of LCB-4 are to:

1. determine the variability of riverine input (Congo), and
2. determine the history of opal, carbonate, and organic-matter deposition off Zaire (Congo).

Drilling Program: APC core Holes A, B, C to 200 m or to refusal

Logging and Downhole: None

Nature of Rock Anticipated: Hemipelagic calcareous silty mud.



Site: MAB-1

Priority: 1

Position: 11°55.2'S, 13°24.0'E

Water Depth: 432 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 200 mbsf

Seismic Coverage: High-resolution MCS survey, seismic Lines GeoB 93-015 and -018

Objectives: The objectives of MAB 1 are to determine the history of "nearly normal" sedimentation, being influenced neither by riverine input nor by sustained year-round upwelling.

Drilling Program: APC core Holes A, B, C to 200 m or to refusal

Logging and Downhole: None

Nature Of Rock Anticipated: Hemipelagic calcareous silty mud.

Site: MAB-2

Priority: 1

Position: 11°55.8'S, 13°02.3'E

Water Depth: 722 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 120 m

Seismic Coverage*: High-resolution MCS survey, seismic Lines GeoB 93-015 and -020

Objectives: The objectives of MAB-2 are to determine the history of "nearly normal" sedimentation, being influenced neither by riverine input nor by sustained year-round upwelling.

Drilling Program: APC core Holes A, B, C to 120 m or to refusal

Logging and Downhole: None

Nature of Rock Anticipated: Hemipelagic calcareous silty mud.

* See seismic line for Site MAB-1

Site: SAB-1

Priority: 2

Position: 16°39.6'S, 10°57.0'E

Water Depth: 2588 m

Sediment Thickness: > 1000 m

Approved Maximum Penetration: 600 m

Seismic Coverage: High-resolution MCS survey, seismic Lines GeoB 93-030 and -034

Objectives: The objective of SAB-1 is to determine the history of the northern end of Angola/Namibia coastal upwelling and that of the climate of southern Africa, as reflected in terrigenous sediments.

Drilling Program:


Hole A: APC/XCB to 600 m

Holes B and C: APC to refusal

Logging and Downhole*: Triple-combo, GHMT, FMS

Nature of Rock Anticipated: Hemipelagic calcareous silty mud.

***Note:** It is anticipated that this site may not be drilled, or only drilled for APC, depending on time available. In the event that there is less than 250 m penetration, no logging or other downhole measurements will be attempted.



Site SAB-1

Site: SAB-2

Priority: 1

Position: 16°33.7'S, 10°49.3'E

Water Depth: 2843 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 600 m

Seismic Coverage: High-resolution MCS survey, seismic Lines GeoB 93-030 and -033

Objectives: The objectives of SAB-2 are to determine the history of the northern end of Angola/Namibia coastal upwelling, migration of Benguela-Angola Front, and climatic history of southern Africa.


Drilling Program:

Hole A: APC/XCB to 600 m

Holes B and C: APC to refusal

Logging and Downhole: Triple-combo, GHMT, FMS

Nature of Rock Anticipated: Hemipelagic calcareous silty mud.



Site: WR-1A

Priority: 2

Position: 19°37.0'S, 11°20.4'E

Water Depth: 763 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 600 mbsf

Seismic Coverage: High-resolution MCS survey, seismic Lines GeoB 96-024 and -026

Objectives: The objective of WR-1A is to expand existing information from DSDP Sites 532 and 362 (Legs 75 and 40) providing for a cross-current transect. This site would help resolve the Walvis Ridge opal paradox, that is, the observation that the opal accumulation during glacials is much less than expected from productivity proxies.

Drilling Program:

Hole A: APC, XCB to refusal

Holes B and C: APC to refusal

Logging and Downhole*: Triple-combo, GHMT, FMS

Nature Of Rock Anticipated: Hemipelagic calcareous silty mud.

***Note:** It is anticipated that this site may not be drilled, or only drilled for APC, depending on time available. In the event that there is less than 250 m penetration, no logging or other downhole measurements will be attempted.

Site: WB-A

Priority: 2

Position: 21°29.0'S, 11°15.1'E

Water Depth: 2707 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 600 mbsf

Seismic Coverage: High-resolution MCS survey, seismic Lines GeoB/AWI 96-015 and -020

Objectives: The objective of WB-A is to determine:

1. the history of the Benguela Current, and strength of coastal upwelling, as seen in messages from filaments and in redeposited material from upslope; and
2. the climatic history of South Africa, as seen in terrigenous components.

Drilling Program:

Hole A: APC, XCB to refusal

Holes B and C: APC to refusal

Logging and Downhole*: Triple-combo, GHMT, FMS

Nature Of Rock Anticipated: Hemipelagic calcareous silty mud.

***Note:** It is anticipated that this site may not be drilled, or only drilled for APC, depending on time available. In the event that there is less than 250 m penetration, no logging or other downhole measurements will be attempted.

Site: WB-B

Priority: 1

Position: 21°05.6'S, 11°49.2'E

Water Depth: 1290 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 600 m

Seismic Coverage: High-resolution MCS survey, seismic Lines GeoB/AWI 96-015 and -020

Objectives: The objectives of WB-B are to determine:

1. the history of the Benguela Current, and strength of coastal upwelling, as seen in filaments and in redeposited material from upslope; and
2. the climatic history of South Africa, as seen in terrigenous components.

Drilling Program:

Hole A: APC, XCB to refusal

Holes B and C: APC to refusal

Logging and Downhole: Triple-combo, GHMT, FMS

Nature of Rock Anticipated: Hemipelagic calcareous silty mud.

Site: WB-C

Priority: 2

Position: 20°53.9'S, 11°13.4'E

Water Depth: 2201 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 600 mbsf

Seismic Coverage: High-resolution MCS survey, seismic Lines GeoB/AWI 96-017 and -021

Objectives: The objectives of WB-C are to determine:

1. the history of the Benguela Current, and strength of coastal upwelling, as seen in messages from filaments and in redeposited material from upslope.
2. the climatic history of South Africa, as seen in terrigenous components.

Drilling Program:

Hole A: APC, XCB to refusal

Holes B and C: APC to refusal

Logging and Downhole*: Triple-combo, GHMT, FMS

Nature Of Rock Anticipated: Hemipelagic calcareous silty mud.

***Note:** It is anticipated that this site may not be drilled, or only drilled for APC, depending on time available. In the event that there is less than 250 m penetration, no logging or other downhole measurements will be attempted.

Site: NCB-2B

Priority: 1

Position: 25°30.8'S, 13°1.7'E

Water Depth: 2004 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 600 m

Seismic Coverage: High-resolution MCS, seismic Line GeoB/AWI 96-014

Objectives: The objectives of NCB-2B are to:

1. determine the history of Benguela Current, including northward excursions and fluctuations in the intensity of productivity;
2. document shoreward/seaward migration of the coastal upwelling center; and
3. reconstruct the history of oxygen supply, in particular of periods showing oxygen deficiency.

Drilling Program:

Hole A: APC, XCB to refusal

Holes B and C: APC to refusal

Logging and Downhole: Triple-combo, GHMT, FMS

Nature of Rock Anticipated: Hemipelagic calcareous silty mud.

Site: MCB-A

Priority: 2

Position: 29°22.5'S, 13°59.4'E

Water Depth: 1726 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 600 mbsf

Seismic Coverage: High-resolution MCS survey, seismic Lines GeoB/AWI 96-009 and -011

Objectives: The objectives of MCB-A are to:

1. explore early history of the Benguela Current in the southern Cape Basin, and
2. detect possible Angola Current influences.

Drilling Program:

Hole A: APC, XCB to refusal

Holes B and C: APC to refusal

Logging and Downhole*: Triple-combo, GHMT, FMS

Nature of Rock Anticipated: Hemipelagic calcareous silty mud.

***Note:** It is anticipated that this site may not be drilled, or only drilled for APC, depending on time available. In the event that there is less than 250 m penetration, no logging or other downhole measurements will be attempted.

Site: SCB-1

Priority: 1

Position: 31°25.0'S, 15°17.0'E

Water Depth: 1350 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 600 mbsf

Seismic Coverage: High resolution MCS, Lines GeoB/AWI 96-003 and -008

Objectives: The objectives of SCB 1 are to:

1. explore early history of the Benguela Current in the southern Cape Basin, and
2. detect possible Angola Current influences.

Drilling Program:

Hole A: APC, XCB to refusal

Holes B and C: APC to refusal

Logging and Downhole: Triple-combo, GHMT, FMS

Nature Of Rock Anticipated: Hemipelagic calcareous silty mud.

Site: SCB-A

Priority: 2

Position: 31°54.4'S, 15°14.1'E

Water Depth: 2234 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 600 mbsf

Seismic Coverage: High-resolution MCS survey, seismic Lines GeoB 96-001 and -004

Objectives: The objectives of SCB-A are to:

1. explore early history of the Benguela Current in the southern Cape Basin, and
2. detect possible Angola Current influences.

Drilling Program:

Hole A: APC, XCB to refusal

Holes B and C: APC to refusal

Logging and Downhole*: Triple-combo, GHMT, FMS

Nature of Rock Anticipated: Hemipelagic calcareous silty mud.

***Note:** It is anticipated that this site may not be drilled, or only drilled for APC, depending on time available. In the event that there is less than 250 m penetration, no logging or other downhole measurements will be attempted.

Site: SCB-B

Priority: 2

Position: 31°47.1'S, 15°30.0'E

Water Depth: 1507 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 600 mbsf

Seismic Coverage: High-resolution MCS survey, seismic Lines GeoB 96-001 and -008

Objectives: The objectives of SCB-B are to:

1. explore early history of the Benguela Current in the southern Cape Basin, and
2. detect possible Angola Current influences.

Drilling Program:

Hole A: APC, XCB to refusal

Holes B and C: APC to refusal

Logging and Downhole*: Triple-combo, GHMT, FMS

Nature of Rock Anticipated: Hemipelagic calcareous silty mud.

***Note:** It is anticipated that this site may not be drilled, or only drilled for APC, depending on time available. In the event that there is less than 250 m penetration, no logging or other downhole measurements will be attempted.

Site: SCB-C

Priority: 2

Position: 31°41.4'S, 15°41.9'E

Water Depth: 887 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 600 mbsf

Seismic Coverage: High-resolution MCS survey, seismic Lines GeoB 96-001 and -006

Objectives: The objectives of SCB-C are to:

1. explore early history of the Benguela Current in the southern Cape Basin, and
2. detect possible Angola Current influences.

Drilling Program:


Hole A: APC, XCB to refusal

Holes B and C: APC to refusal

Logging and Downhole*: Triple-combo, GHMT, FMS

Nature of Rock Anticipated: Hemipelagic calcareous silty mud.

***Note:** It is anticipated that this site may not be drilled, or only drilled for APC, depending on time available. In the event that there is less than 250 m penetration, no logging or other downhole measurements will be attempted.



NW

GeoB 96-006

SE

Site SCB-C

Site: SCB-D

Priority: 2

Position: 31°21.3'S, 15°36.5'E

Water Depth: 778 m

Sediment Thickness: >1000 m

Approved Maximum Penetration: 600 mbsf

Seismic Coverage: High-resolution MCS survey, seismic Lines GeoB 96-003 and -006

Objectives: The objectives of SCB-D are to:

1. explore early history of the Benguela Current in the southern Cape Basin, and
2. detect possible Angola Current influences.

Drilling Program:

Hole A: APC, XCB to refusal

Holes Band C: APC to refusal

Logging and Downhole*: Triple-combo, GHMT, FMS

Nature of Rock Anticipated: Hemipelagic calcareous silty mud.

***Note:** It is anticipated that this site may not be drilled, or only drilled for APC, depending on time available. In the event that there is less than 250 m penetration, no logging or other downhole measurements will be attempted.

SCIENTIFIC PARTICIPANTS

Co-Chief

Wolfgang H. Berger
Scripps Institution of Oceanography
University of California, San Diego
Geosciences Research Division
La Jolla, CA 92093-0215
U.S.A.
Internet: wberger@ucsd.edu
Work: (619) 534-1829
Fax: (619) 534-0784

Co-Chief

Gerold Wefer
Fachbereich Geowissenschaften
Universität Bremen
Bibliothekstraße
Postfach 33 04 40
Bremen 28334
Federal Republic of Germany
Internet: gwefer@alf.zfn.uni-bremen.de
Work: (49) 421-218-3389
Fax: (49) 421-218-3116

Staff Scientist

Carl Richter
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
U.S.A.
Internet: richter@tamu.edu
Work: (409) 845-2522
Fax: (409) 845-0876

Inorganic Geochemist

Richard W. Murray
Department of Earth Sciences
Boston University
675 Commonwealth Avenue
Boston, MA 02215
U.S.A.
Internet: rickm@bu.edu
Work: (617) 353-6532
Fax: (617) 353-3290

Organic Geochemist

Ioanna Bouloubassi
Département de Géologie et Océanographie
Université de Bordeaux I
Avenue des Facultés
Talence Cedex 33405
France
Internet: bouloubassi@geocean.u-bordeaux.fr
Work: (33) 5-56-84-89-60
Fax: (33) 5-56-84-08-48

Organic Geochemist

Philip A. Meyers
Department of Geological Sciences
University of Michigan
2534 C. C. Little Building
425 East University Avenue
Ann Arbor, MI 48109-1063
U.S.A.
Internet: pameyers@umich.edu
Work: (313) 764-0597
Fax: (313) 763-4690

JOIDES Logging Scientist

Bryce W. Hoppie
Department of Chemistry and Geology
Mankato State University
Mankato, MN 56002-8400
U.S.A.
Internet: bryce_hoppie@ms1.mankato.msus.edu
Work: (507) 389-1963
Fax: n/a

Paleomagnetist

Toshitsugu Yamazaki
Marine Geology Department
Geological Survey of Japan
1-1-3 Higashi
Tsukuba, Ibaraki 305
Japan
Internet: yamazaki@gsj.go.jp
Work: (81) 298-54-3591
Fax: (81) 298-54-3589

Paleontologist (Diatom)

Carina Beatriz Lange
Scripps Institution of Oceanography
University of California, San Diego
Geological Research Division
La Jolla, CA 92093-0215
U.S.A.
Internet: clang@ucsd.edu
Work: (619) 534-4605
Fax: (619) 534-0784

Paleontologist (Foraminifer)

Beth A. Christensen
Department of Geological Sciences
University of South Carolina
Columbia, SC 29208
U.S.A.
Internet: bac@geol.sc.edu
Work: (803) 777-8845
Fax: (803) 777-4525

Paleontologist (Foraminifer)

Jan Otto R. Hermelin
Geologiska Institutionen
Stockholms Universitet
Deep Sea Geology Division
Stockholm 106 91
Sweden
Internet: hermelin@geol.su.se
Work: (46) 8-164-734
Fax: (46) 8-164-734

Paleontologist (Nannofossil)

Jacques Giraudeau
Département de Géologie et Océanographie
Université de Bordeaux I
Avenue des Facultés
Talence Cedex 33405
France
Internet: giraudeau@geocean.u-bordeaux.fr
Work: (33) 5-56-84-88-63
Fax: (33) 5-56-84-08-48

Paleontologist (Radiolaria)

Isao Motoyama
Department of Physics and Earth Sciences
University of Ryukyus
Senbaru 1
Nishihara-cho
Okinawa 903-01
Japan
Internet: motoyama@sci.u-ryukyu.ac.jp
Work: (81) 98-895-8551
Fax: (81) 98-895-2414

Physical Properties Specialist,
Stratigraphic Correlator

Dyke J. Andreasen
Earth Sciences Board
University of California, Santa Cruz
Santa Cruz, CA 95064
U.S.A.
Internet: andreasn@aphrodite.ucsc.edu
Work: (408) 459-5061
Fax:

Physical Properties Specialist

Bernd Laser
Fachbereich Geowissenschaften
Universität Bremen
FB 5
Postfach 330440
Bremen 28334
Federal Republic of Germany
Internet: bel@mtu.uni-bremen.de
Work: (49) 218-7286
Fax: (49) 218-7179

Physical Properties Specialist

Volkhard Spiess
Fachbereich Geowissenschaften
Universität Bremen
Postfach 330440
Bremen 28334
Federal Republic of Germany
Internet: al3g@mtu.uni-bremen.de
Work: (49) 421-218-3387
Fax: (49) 421-218-3116

Sedimentologist

Linda Davis Anderson
Institute of Marine Sciences
University of California, Santa Cruz
Santa Cruz, CA 95064
U.S.A.
Internet: linda@cats.ucsc.edu
Work: (408) 459-3123
Fax: (408) 459-4882

Sedimentologist

Volker Brüchert
Department of Geological Sciences
Indiana University, Bloomington
Biogeochemical Laboratories
Bloomington, IN 47405
U.S.A.
Internet: vbrucher@indiana.edu
Work: (812) 855-8034
Fax: (812) 855-7961

Sedimentologist

J.H. Fred Jansen
Nederlands Instituut voor Onderzoek der Zee
(NIOZ)
PO Box 59
AB Den Burg, Texel 1790
The Netherlands
Internet: jansen@nioz.nl
Work: (31) 222-369396
Fax: (31) 222-319674

Sedimentologist

Hui-Ling Lin
Institute of Marine Geology and Chemistry
National Sun Yat-Sen University
Kaohsiung 804
Taiwan
Internet: hllin@mail.nsysu.edu.tw
Work: (886) 752-52000
Fax: (886) 752-55149

Sedimentologist

Mark A. Maslin
Environmental Change Research Centre
University College London
26 Bedford Way
London WC1H 0AP
United Kingdom
Internet: mmaslin@geog.ucl.ac.uk
Work: (44) 171-380-7556
Fax: (44) 171-380-7565

Sedimentologist

Maria Elena Pérez
Scripps Institution of Oceanography
University of California, San Diego
9500 Gilman Drive
La Jolla, CA 92093-0215
U.S.A.
Internet: mperez@ucsd.edu
Work: n/a
Fax: (619) 534-0784

Sedimentologist

Peir Kenneth Pufahl
Department of Earth and Sciences
University of British Columbia
6339 Stores Rd.
Vancouver, BC V6T 1Z4
Canada
Internet: ppufahl@eos.ubc.ca
Work: (604) 822-2449
Fax: (604) 822-6088

Sedimentologist

Ralph R. Schneider
Fachbereich Geowissenschaften
Universität Bremen
Postfach 330 440
Bremen 28334
Federal Republic of Germany
Internet: rschneid@zfn.uni-bremen.de
Work: (49) 421-218-3579
Fax: (49) 421-218-3116

LDEO Logging Scientist

Hervé Cambray
Institut Méditerranéen de Technologie
Technopole de Château Gombert
Marseille Cedex 20 13451
France
Internet: cambray@imtmer1.imt-mrs.fr
Work: (33) 4-91-05-45-01
Fax: (33) 4-91-05-43-43

Schlumberger Engineer

Jonathan Kreb
Schlumberger Offshore Services
369 Tristar Drive
Webster, TX 77598
U.S.A.
Internet: jkreb@webster.wireline.slb.com
Work: (713) 480-2000
Fax: (713) 480-9550

Operations Manager

Ron Grout
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
U.S.A.
Internet: ron_grout@odp.tamu.edu
Work: (409) 845-2144
Fax: (409) 845-2308

Laboratory Officer

Bill Mills
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
U.S.A.
Internet: bill_mills@odp.tamu.edu
Work: (409) 845-2478
Fax: (409) 845-2380

Assistant Laboratory Officer,
Marine Lab Specialist: X-Ray

Kuro Kuroki
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
U.S.A.
Internet: kuro_kuroki@odp.tamu.edu
Work: (409) 845-8482
Fax: (409) 845-2380

Marine Lab Specialist: Yeoperson

Jo Ribbens
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
U.S.A.
Internet: jo_ribbens@odp.tamu.edu
Work: (409) 845-8482
Fax: (409) 845-2380

Marine Lab Specialist: Chemistry

Tim Bronk
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
U.S.A.
Internet: tim_bronk@odp.tamu.edu
Work: (409) 845-2480
Fax: (409) 845-2380

Marine Lab Specialist: Chemistry

Anne Pimmel
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
U.S.A.
Internet: anne_pimmel@odp.tamu.edu
Work: (409) 845-8482
Fax: (409) 845-2380

Marine Lab Specialist: Curator

Erinn McCarty
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
U.S.A.
Internet: john_miller@odp.tamu.edu
Work: (409) 845-5056
Fax:

Marine Lab Specialist: Thin Sections,
Marine Lab Specialist: Downhole Tools

Chris Nugent
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845
U.S.A.
Internet: chris_nugent@odp.tamu.edu
Work: (409) 845-2481
Fax: (409) 845-2380

Marine Lab Specialist: Fantail,
Marine Lab Specialist: Underway
Geophysics

Monty Lawyer
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
U.S.A.
Internet: monty_lawyer@odp.tamu.edu
Work: (409) 845-2480
Fax: (409) 845-2380

Marine Lab Specialist: Paleomagnetism

Margaret Hastedt
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
U.S.A.
Internet: margaret_hastedt@odp.tamu.edu
Work: (409) 845-2483
Fax: (409) 845-2380

Marine Lab Specialist: Photographer

Roy Davis
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
U.S.A.
Internet: roy_davis@odp.tamu.edu
Work: (409) 845-8482
Fax: (409) 845-1026

Marine Lab Specialist: Storekeeper

Sandy Dillard
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
U.S.A.
Internet: sandy_dillard@odp.tamu.edu
Work: (409) 845-2480
Fax: (409) 845-2380

Marine Lab Specialist: X-Ray

Jaquelyn Ledbetter
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
U.S.A.
Internet: jaque_ledbetter@odp.tamu.edu
Work: (409) 845-8482
Fax: (409) 845-2380

Marine Computer Specialist

John Eastlund
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845
U.S.A.
Internet: john_eastlund@odp.tamu.edu
Work: (409) 845-3044
Fax: (409) 845-4857

Marine Computer Specialist

Chris Stephens
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
U.S.A.
Internet: chris_stephens@odp.tamu.edu
Work: (409) 862-4846
Fax: (409) 845-4857

Marine Electronics Specialist

Larry St. John
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
U.S.A.
Internet: larry_st.john@odp.tamu.edu
Work: (409) 845-2473
Fax: (409) 845-2308

Marine Electronics Specialist

Bill Stevens
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
U.S.A.
Internet: bill_stevens@odp.tamu.edu
Work: (409) 845-2454
Fax: (409) 845-2380

Marine Electronics Specialist

Mark Watson
Ocean Drilling Program
Texas A&M University
1000 Discovery Drive
College Station, TX 77845-9547
U.S.A.
Internet: mark_watson@odp.tamu.edu
Work: (409) 845-2473
Fax: (409) 845-2380